



HAL
open science

Sexual reproduction in Populus. I. Some physiological and biochemical events of the progamic phase

Marc M. Villar, M. Gaget, Christian Dumas

► **To cite this version:**

Marc M. Villar, M. Gaget, Christian Dumas. Sexual reproduction in Populus. I. Some physiological and biochemical events of the progamic phase. International symposium Forest Tree Physiology, Sep 1988, Nancy, France. <hal-02856015>

HAL Id: hal-02856015

<https://hal.inrae.fr/hal-02856015v1>

Submitted on 8 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Sexual reproduction in *Populus* I. Some physiological and biochemical events of the progamic phase

M. Villar¹, M. Gaget² and C. Dumas²

¹INRA, Station d'Amélioration des Arbres Forestiers, Ardon, F-45160 Olivet, and

²Laboratoire de Reconnaissance Cellulaire et Amélioration des Plantes, INRA 23879, Université Lyon I, 43, bd 11-Novembre-1918, F-69622 Villeurbanne Cedex, France

Introduction

Hybridization in *Populus* breeding programs is limited by sexual incompatibility barriers, whose cellular and molecular mechanisms are not yet known. In an attempt to understand the nature of inter-specific incompatibility in *Populus*, we have explored the interactions between male and female partners (pollen–pistil) in compatible and incompatible crosses *P. nigra* (female) x *P. nigra* (male) and *P. nigra* (female) x *P. alba* (male) (Villar, 1987).

Materials and Methods

P. nigra and *P. alba* branches were obtained from the INRA Forestry station in Orléans (France). Pollen was collected and stored in closed vials at -18°C . Kinetics of pollen tube growth *in vivo* were studied in growth chambers (20°C). Pollen tubes in the stylar tissues were visualized using the aniline blue fluorescence method (ABF method, Dumas and Knox, 1983).

Isoelectric focusing (IEF) of pollinic and stigmatic proteins, gel preparation, focusing condi-

tions, silver nitrate staining and homogenates of pollinated stigma were performed according to the procedures of Villar *et al.* (1988). Each of the 2 crosses was represented by 3 series of stigmatic extracts, in accordance with the growth kinetics (0.6 and 20 h after pollination). Protein patterns of these extracts were compared on a single gel. Glycoprotein revelation (concanavalin A-binding proteins) on IEF polyacrylamide gel was adapted from Hawkes (1982). β -Galactosidase visualization was optimized from the protocol of Singh and Knox (1985).

Results and Discussion

Kinetics of pollen tube growth (visualization using the ABF method) have demonstrated distinct behaviors of *P. nigra* and *P. alba* pollen tubes inside *P. nigra* pistils (Fig. 1). *P. alba* pollen tubes exhibit a unique S-shaped growth curve and an arrested growth near the stylodium. On the other hand, *P. nigra* pollen tubes exhibit 2 growth phases, respectively, in the stigmatic tissues and in the ovarian cavity. *P. nigra* and *P. alba* curves diverge 5 h after pollination (20°C), corresponding precisely to the deposition of the first




Fig. 1. Kinetics of pollen tube growth *in vivo*. (From Villar *et al.*, manuscript in preparation)

callose plug inside pollen tubes. This divergence could likely be related to a change in the physiology of the compatible pollen tube, shifting from an autotrophic to a heterotrophic type of nutrition.

Biochemical studies focused on pollinic and stigmatic proteins, known to be involved in male-female interactions (Gaude and Dumas, 1987). After silver nitrate staining, qualitative and quantitative differences could be observed, related to the presence of *P. nigra* and *P. alba* pollen tubes inside stigmatic tissues. However, increasing protein bands were detectable, 0–20 h after pollination, only in compatible pollinated stigmas. The concanavalin A-peroxidase reaction allowed the visualization of 15 stigmatic and pollinic

glycoproteins. Differences were observed, according to the type of cross: one distinct glycoprotein increases only in the compatible cross. Moreover, β -Galactosidase activities were revealed with a similar electrophoretic technique in pollinated stigma. This pollinic enzyme could play a role in heterotrophic pollen tube nutrition (Singh and Knox, 1985). An increase of its activity (one isoenzyme of isoelectric point about pH 4.2) from 6–20 h after pollination was detected only in the compatible cross.

Conclusion

The compatible progamic phase in *P. nigra*, *i.e.*, pollen tube growth up to the

embryo sac, could be related to pollinic enzymes involved in pollen tube metabolism, such as β -galactosidase. Its activity could be the final result of a series of interactions started by initial pollen-stigma communications. This dialogue probably implicates protein compounds, detected in pollen and pollen tube diffusates *in vitro* (Villar, 1987; Gaget, 1988).

References

- Dumas C. & Knox R.B. (1983) Callose and determination of pistil viability and incompatibility. *Theor. Appl. Genet.* 67, 1-10
- Gaget M. (1988) Incompatibilité interspécifique chez *Populus*: effet Mentor. Ph.D. Thesis, Université Lyon I, France
- Gaude T. & Dumas C. (1987) Molecular and cellular events of self-incompatibility. *Int. Rev. Cytol.* 107, 333-366
- Hawkes R. (1982) Identification of concanavalin A-binding proteins after sodium dodecyl sulfate-gel electrophoresis and protein blotting. *Anal. Biochem.* 123, 143-146
- Singh K. and Knox R.B. (1985) β -Galactosidase of *Lilium* pollen. *Phytochemistry* 24, 1639-1643
- Villar M. (1987) Incompatibilité interspécifique chez *Populus*: approches physiologique et biochimique. Ph.D. Thesis, Université Lyon I, France
- Villar M., Gaget M. & Dumas C. (1988) Microisoelectric focusing of proteins from single stigma in *Populus*. *Can. J. For.* 18, 1261-1264