



HAL
open science

The spore coat is essential for *Bacillus subtilis* spore resistance to pulsed light, and pulsed light treatment eliminates some spore coat proteins

Gérémy Clair, Julia Esbelin, Sabine Mallea, Isabelle Bornard, Frederic Carlin

► To cite this version:

Gérémy Clair, Julia Esbelin, Sabine Mallea, Isabelle Bornard, Frederic Carlin. The spore coat is essential for *Bacillus subtilis* spore resistance to pulsed light, and pulsed light treatment eliminates some spore coat proteins. *International Journal of Food Microbiology*, 2020, 323, pp.108592. 10.1016/j.ijfoodmicro.2020.108592 . hal-02862233

HAL Id: hal-02862233

<https://hal.inrae.fr/hal-02862233>

Submitted on 22 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 The spore coat is essential for *Bacillus subtilis* spore resistance to pulsed light, and pulsed light
2 treatment eliminates some spore coat proteins

3 G r my Clair^{a,b*}, Julia Esbelin^{a*}, Sabine Mall a^a, Isabelle Bornard^c, Fr d ric Carlin^{a**}

4

5 ^a INRAE, Avignon Universit , UMR SQPOV, 84914 Avignon, France

6 ^b Integrative Omics, Pacific Northwest National Laboratory, 902 Battelle Boulevard, Richland,

7 Washington, 99352, USA

8 ^c UR407 PV "Pathologie V g tale", INRA, 84000 Avignon, France

9 *G r my Clair and Julia Esbelin contributed equally to this study and should be considered both
10 as shared first co-authors

11 **Corresponding author :

12 Fr d ric Carlin. Mailing address: INRAE, Centre de Recherche PACA, UMR408 s curit  et Qualit 
13 des Produits d'Origine V g tale, 228 Route de l'A rodrome, CS40509, Domaine Saint-Paul - Site
14 Agroparc, 84914 Avignon Cedex 9, France.

15 Email: frederic.carlin@inra.fr. Phone: +33 (0)432 72 25 19

16 Declarations of interest : none.

17 **Abstract**

18 Microbial surface contamination of equipment or of food contact material is a recurring problem in the
19 food industry. Spore-forming bacteria are far more resistant to a wide variety of treatments than their
20 vegetative forms. Understanding the mechanisms underlying decontamination processes is needed to
21 improve surface decontamination strategies against endospores potentially at the source of foodborne
22 diseases or food-spoilage. Pulsed light (PL) with xenon lamps delivers high-energy short-time
23 pulses of light with wavelengths in the range 200 nm-1100 nm and a high UV-C fraction. *Bacillus*
24 *subtilis* spores were exposed to either PL or to continuous UV-C. Gel electrophoresis and
25 western blotting revealed elimination of various proteins of the spore coat, an essential outer
26 structure that protects spores from a wide variety of environmental conditions and inactivation
27 treatments. Proteomic analysis confirmed the elimination of some spore coat proteins after PL
28 treatment. Transmission electron microscopy of PL treated spores revealed a gap between the
29 lamellar inner spore coat and the outer spore coat. Overall, spores of mutant strains with
30 defects in genes coding for spore coat proteins were more sensitive to PL than to continuous
31 UV-C. This study demonstrates that radiations delivered by PL contribute to specific damage to
32 the spore coat, and overall to spore inactivation.

33 **Keywords**

34 Decontamination, UV, proteins, proteomics, microscopy

35

36

37 **1. Introduction**

38 The pulsed light (PL) technology inactivates harmful or food spoiling microorganisms
39 contaminating surfaces and clear liquids. PL delivered by commonly used xenon lamps consists
40 of short (typically < 250 ms) and intense flashes of white light (200 nm–1100 nm wavelengths)
41 rich in ultraviolet (200 nm–400 nm wavelengths) (Garvey and Rowan, 2019; Gomez-Lopez and
42 Bolton, 2016; Gomez-Lopez et al., 2007). PL delivers short-time light pulses that are several
43 thousand times more intense than UV-C delivered by a continuous source. The ability of PL to
44 kill microorganisms is well documented (Kramer et al., 2017). Nevertheless, understanding
45 intrinsic PL mechanisms leading to inactivation of resistant forms of bacteria, such as
46 endospores, could improve PL technology and expand applications.

47 Spores of bacteria are markedly more resistant than vegetative cells to adverse conditions that
48 prevail in natural or industrial environments (Setlow, 2007). This resistance favours spore
49 dispersion and transfer to food processing facilities. Elimination of bacterial spores in industrial
50 environments contributes to prevention of foodborne poisonings or food spoilage. Technologies
51 have been developed over years either to inactivate bacterial spores or to decontaminate
52 surfaces in contact with foods, such as processing equipment or packaging material (Carlin,
53 2011; Setlow, 2014; Soni et al., 2016; Wells-Bennik et al., 2016). Chemical biocides, such as
54 chlorine derivatives, peracetic acid or hydrogen peroxide, are highly efficient against bacterial
55 spores (Gopal et al., 2015; Maillard, 2011). Nevertheless, residues left after biocide application
56 are suspected of detrimental effects on environment or health (Carrasco and Urrestarazu, 2010;
57 Russell, 2003). UV-technologies are an alternative to chemical biocides for surface treatment

58 and inactivation of spore-forming bacteria. Continuous UV-C technology (low-pressure mercury
59 lamps emitting at 254 nm) requires relatively long exposure (typically few minutes) for spore
60 inactivation. Pulsed light (PL) is capable of achieving similar spore inactivation as continuous UV-
61 C (Gomez-Lopez et al., 2007; Koutchma et al., 2009; Levy et al., 2012) with much shorter
62 exposure times (Garvey et al., 2014) and therefore of accelerating high-flow industrial chains. PL
63 achieves up to 6 log-reduction on spores of many bacterial species with only one or two light
64 pulses (Levy et al., 2012). Application of PL on plastic polymers does not seem to generate
65 chemical residues or migrants (Castillo et al., 2013).

66 Resistance of spores lies in their specific structural organization. Spores are made of the
67 concentric layers of the coat and cortex surrounding the spore core. Spore core is characterized
68 by low water content, high content in dipicolinic acid and chelated divalent metal ions, and
69 small acid soluble proteins (SASPs) bond to DNA. DNA repair during outgrowth along with
70 SASPs, low water content of spore core and pigments allow spores to resist to UV treatments
71 (Nicholson et al., 2000; Setlow, 2014). In *Bacillus subtilis* the spore coat mainly consists of an
72 assembly of at least 70 proteins organized in a multi-layered structure (McKenney et al., 2013).
73 From the inside out, spore coat layers are the inner spore coat, the outer spore coat and the
74 crust (Imamura et al., 2011; McKenney et al., 2013). Several morphogenetic proteins, including
75 SpoIVA, SpoVID, SafA, CotE and CotXYZ, participate in the assembly of coat layers (McKenney
76 and Eichenberger, 2012; Ozin et al., 2000; Roels et al., 1992; Zheng et al., 1988).

77 Spore coat has a major role in protection against oxidative damage caused by oxidizing agents
78 (Moeller et al., 2011; Riesenman and Nicholson, 2000), or also caused by UV-A (Pattison and
79 Davies, 2006), and possibly, as shown in our previous work, against damage caused by PL

80 (Esbelin et al., 2016). For instance, spores of a CotE defective mutant (lacking some spore coat
81 external layers) or chemically decoated spores were more sensitive to PL than wild-type (WT) or
82 than untreated spores, while difference was marginal after exposure to a continuous UV-C
83 source. More generally, spore coat proteins are not identified as major factors of resistance to
84 UV-C (Nicholson et al., 2000; Setlow, 2014). This difference in sensitivity to PL and to
85 monochromatic UV-C source (emitting mostly at 254 nm) could be due to a wider range of UV
86 wavelengths emitted by PL and to higher irradiance (more than 1000 W/cm² with PL compared
87 to 1 x 10⁻³ W/cm² with low-pressure mercury lamps). Our work is aimed at understanding the
88 origin of damage caused by different UV-technologies on bacterial spores. Specifically our
89 objectives were to determine (i) whether PL has a specific degradation effect on coat proteins of
90 *Bacillus subtilis* using both biochemical and proteomic approaches and (ii) whether there is a
91 specific role of spore coat proteins in resistance to PL using several *B. subtilis* strains with
92 mutations in genes coding for coat proteins.

93

94 **2. Materials and Methods**

95 **2.1. *Bacillus subtilis* strains used and preparation of spores**

96 Origin and properties of *B. subtilis* strains 1A1 and PY79, wild-type background initially *B. subtilis*
97 168, and derived mutants strains are detailed in Table 1. Strains were routinely cultivated on
98 Luria-Bertani (LB) (Biokar, Beauvais, France) agar supplemented with antibiotic concentrations
99 recommended by donator (See Table 1) to ensure strain purity. Then, one colony was picked,
100 grown overnight at 30°C in LB and 200 µl were plated on double-strength Schaeffer sporulation

101 (2xSG) agar without antibiotics (Nicholson and Setlow, 1990). Plates were incubated for 7 days
102 at 30°C before spore harvest. Spore suspensions were independently prepared (at least three
103 replicates of each strain) and resistance of each suspension to PL and UV-C was tested. Spores
104 were purified by repeated washing as previously described (Levy et al., 2011). Spore
105 preparations were checked to be free (< 5 %) of growing cells, germinated spores and cell debris
106 using phase contrast microscopy under 1000x magnification (Olympus BX50, Rungis, France).
107 Purified spores were suspended in demineralized sterile water, heat-treated at 70°C for 10 min
108 to inactivate vegetative cells and stored in the dark at 4°C until use. Spore counts were
109 determined by spreading 100 µl volumes of appropriate decimal serial dilutions on duplicate LB
110 agar plates incubated at 30°C for 48 h. All spore suspensions contained 10⁸ to 10⁹ spore cfu/ml
111 and were checked at regular time intervals for retaining phase-brightness.

112 Table 1. *Bacillus subtilis* strains used in the present work

<i>B. subtilis</i> strain (<i>designation</i>)	Genotype	Properties	Origin (donor)	Reference
1A1 (<i>wild-type</i> , WT)	168, <i>trpC2</i>	Wild-type strain	BSCG ^a	
1A184 (<i>cotA</i>)	<i>cotA1 trpC2</i>	Deletion in <i>cotA</i> coding for a spore coat protein. Brownish pigmentation	BSCG	(Hullo et al., 2001)
1L45 (<i>gerE</i>)	(ϕ 105J45) - (<i>gerE36</i>) ⁺ <i>trpC2</i>	Incomplete spore coat structure	BSCG	(Moir, 1981)
1S105 (<i>cotE</i>)	<i>cotEΔ::cat trpC2</i>	Spores lacking the outer spore coat layer	BSCG	(Zheng et al., 1988)
1S46 (<i>spoIVA</i>)	<i>pheA12 spoIVA178</i>	SpoIVA is a global spore coat morphogenic protein	BGSC	(Coote, 1972)
AH1910 (<i>spoVID</i>)	<i>trpC2 metC3 spoVID</i> : : <i>pJMXE</i>	SpoVID is the basement morphogenic protein of the spore coat structure	A.O. Henriques	(Costa et al., 2006)
AOB68 (<i>safA</i>)	<i>trpC2 metC3 safA</i> : : <i>pOZ83</i>	SafA is a morphogenic protein involved in assembly of inner spore coat layer	A.O. Henriques	(Ozin et al., 2000)
MTB902 (<i>cotB</i>)	<i>trpC2 cotB</i> : : <i>pMutin3</i>	CotB is an outer spore coat component	H. Takamatsu	
1S103 (<i>cotC</i>)	<i>cotC::cat trpC2</i>	CotC is an outer spore coat component	BGSC	(Donovan et al., 1987)
1S104 (<i>cotD</i>)	<i>trpC2 cotD::cat</i>	CotD is a spore spore coat component	BGSC	(Donovan et al., 1987)
1S107 (<i>cotF</i>)	<i>trpC2 cotF::cat</i>	CotF is an inner spore coat component	BGSC	(Cutting et al., 1991)

<i>B. subtilis</i> strain (<i>designation</i>)	Genotype	Properties	Origin (donor) Reference
1A1 (<i>wild-type</i> , WT)	168, <i>trpC2</i>	Wild-type strain	BGSC ^a
MTB907 (<i>cotG</i>)	<i>trpC2 cotG</i> : : <i>pMutin3</i>	CotG is a spore coat protein that participates to the assembly of CotH (and CotB) protein in the outer spore coat layer	H. Takamatsu (Kuwana et al., 2004)
MTB908 (<i>cotH</i>)	<i>trpC2 cotH</i> : : <i>pMutin3</i>	CotH is an outer spore coat component	H. Takamatsu (Kuwana et al., 2004)
1S88 (<i>spoVIA</i>)	<i>spoVIA513 trpC2</i>	SpoVIA is an outer spore coat component	BGSC (Jenkinson, 1981)
PE594 (PY79)	Strain PY79	Wild Type strain. Prototrophic derivative of <i>B. subtilis</i> 168	P. Eichenberger
PE618 PY79- Δ <i>cotE</i>	<i>trpC2 cotE</i> : : <i>cat</i>	CotE is a morphogenic protein involved in assembly of the outer spore coat layer	P. Eichenberger (Arrieta-Ortiz et al., 2015; Zheng et al., 1988)
PE620 Δ <i>cotXcotYZ</i>	<i>trpC2 Δ(cotXcotYZ)</i> : : <i>Neo</i>	CotX, Y and Z proteins compose the outermost spore coat layer, the crust	P. Eichenberger (McKenney and Eichenberger, 2012)

113 ^a*Bacillus* Genetic Stock Center (BGSC) (Columbus, OH)

114

115

116 **2.2. Evaluation of spore resistance**

117 Methods used for evaluation of spore resistance were similar to the ones used in previous work
118 (Esbelin et al., 2016; Levy et al., 2012). Briefly spores were diluted to a A_{600} of 1.0 (10^8
119 spores/ml), heat-treated for 10 min at 70°C, then cooled and kept on ice to prevent germination
120 during experiments. For UV and PL resistance, volumes of 100 μ l of 6-7 decimal serial dilutions
121 of spore suspensions were spread with a rake on LB agar plates of 9 cm diameter. Treatments
122 immediately followed inoculum spreading. The number of spores on plates was therefore
123 comprised between approx. 10^7 and 1 spore cfu/plate. Phase-contrast microscopy at 1000x
124 magnification did not reveal any spore that turned phase-dark and therefore having germinated
125 within 2.5 min, i.e. longer than time between spreading and exposure. The distribution of
126 spores on the agar surface was homogeneous and spore shielding was not detected. PL and
127 continuous UV-C exposure of inoculated agar plates were performed within 30 s - 2 min
128 following spreading. LB agar plates were placed under radiation sources. Inoculated plates were
129 incubated at 30°C for at least 48 h prior to colony counting to ensure that all survivors had
130 formed visible colonies. Untreated samples were used as controls.

131 **2.3. Pulsed light and continuous UV-C devices**

132 PL resistance of *B. subtilis* spores was tested with a previously described lab-scale equipment
133 (Claranor, Avignon, France) (Levy, 2010; Levy et al., 2012). This device delivers pulses (duration
134 250 μ s) of polychromatic light (wavelengths between 200 nm and 1100 nm) produced by a
135 xenon flashlamp containing 18.5 % of fluence corresponding to UV (wavelengths between 200

136 nm and 400 nm) under a charging voltage of 2500 V. The emission spectrum of the xenon lamps
137 is shown in supplementary Fig 1. The UV-C proportion in PL was determined with
138 spectrophotometric methods (Claranor, unpublished data). PL fluence varied from 0.3 J/cm² to
139 10.8 J/cm² (from 0.3 J/cm² to 1.8 J/cm² for survival curves) according to the number of delivered
140 flashes (one to 16 at 1 s interval) and the distance to the lamp (between 8 cm and 25 cm).
141 Unflashed samples were used as controls. UV-C resistance was measured by using a UV-C
142 chamber equipped with three low-pressure mercury lamps (2.5 cm-diameter and 43.5 cm-long)
143 (model TUV-T8, 15 W, Philips, The Netherlands) placed in parallel at approx. 45 cm from the
144 target and emitting short-wave radiations with a peak at 253.7 nm according to manufacturer's
145 specifications. The irradiation strength of 1.07 mW/cm² was measured with a radiometer
146 (model VLX-3W, Vilber Lourmat, Marne-la-Vallée, France). Tested UV-C fluence ranged from
147 0.027 to 0.88 J/cm² (from 0.027 J/cm² to 0.15 J/cm² for survival curves) and were obtained by
148 times of treatment ranging from 25 s to 13 min 48 s (to 2 min 30 s for survival curves). Viability
149 reduction was expressed as log₁₀ N₀/N as a function of the applied fluence, where N₀ is the
150 initial count of the spread suspensions in cfu/plate and N the number of survivors in cfu/plate.

151 **2.4. SDS PAGE of coat protein extracts**

152 Samples (100 µl) of purified spores (at $A_{600} = 3.0$) of wild-type strain 1A1 were mixed with 14 ml
153 of sterile demineralized distilled water and poured into Petri dishes of 9 cm diameter giving a 3
154 mm deep water layer. Spores were then exposed to either PL (fluence between 1.8 J/cm² to
155 10.8 J/cm²) or UV-C (fluence between 0.15 J/cm² and 0.88 J/cm²). Exposed spores were
156 transferred to a new test tube and harvested by centrifugation (7000 X *g* for 5 min at 4°C).
157 Spore pellets were suspended again in 40 µl of SDS PAGE loading buffer (Biorad, Munich,
158 Germany) and boiled for 5 min, mixed vigorously for 1 min and boiled again for 5 min. Samples
159 were briefly centrifuged and 15 µl (approximately 40 µg of total proteins) were fractionated
160 using a large format 1-D electrophoresis systems (protean II Biorad) on a 12 % acrylamide gel
161 containing SDS (SDS PAGE, sodium dodecyl sulfate polyacrylamide gel electrophoresis). Gels
162 were stained with Coomassie brilliant blue R-250, and then destained.

163 **2.5. Western blot**

164 Samples were resolved by SDS-PAGE and electro-transferred onto nitrocellulose membranes
165 using the iBlot Dry Blotting System according to standard procedures (Invitrogen, Illkirch,
166 France). Membranes were probed with the appropriate antibody. Blotted membranes were
167 developed with a 1:15000 dilution of peroxidase-conjugated goat anti-rabbit immunoglobulin G
168 (Sigma, Saint-Quentin Fallavier, France) and an enhanced chemiluminescence substrate
169 (SuperSignal West Femto Chemiluminescent Substrate).

170 **2.6. Proteomic analysis**

171 For proteomics, a 100 µl sample of each spore suspension was treated as follows. The spores
172 were rinsed by suspension in 5 mL of cold sterile demineralized water and agitation with an
173 orbital shaker for 5 min at 4°C to remove loosely attached proteins. The spore pellet was
174 obtained by centrifugation of the suspension at 3500 x *g* and the supernatant was discarded.
175 This procedure was twice repeated. The spores were then suspended in 1X NuPAGE LDS sample
176 buffer (Invitrogen) supplemented with DTT (at 100 mmol/l final concentration) (Sigma) for
177 protein solubilisation and loaded onto the different wells of one 4–12 % gradient NuPAGE gel
178 (Invitrogen). The gel was operated with MES buffer, run at 150 V (Invitrogen) for 5 min resulting
179 into a short gel migration (5-8 mm). For each sample, the whole protein content from each well
180 was excised. Their protein contents were then treated with DTT and iodoacetamide and then
181 proteolysed with trypsin and proteasMAX™ (Promega, Madison, WI, USA) and analysed by LC-
182 MS/MS as previously described (Clair et al., 2012). Briefly, digested peptides were loaded and
183 desalted on line on a reverse-phase precolumn (C18 PepMap 100 column, LC Packings).
184 Peptides were separated using a 90 min gradient from 5 % to 60 % solvent B (0.1 % HCOOH / 80
185 % CH₃CN) in solvent A (0.1 % HCOOH / 100 % H₂O), at a flow rate of 0.3 µl/min and analysed
186 with an LTQ-Orbitrap XL mass spectrometer (ThermoFisher, Illkirch, France) in data-dependent
187 mode using the TOP7 strategy.

188 **2.7. Proteomic data analysis**

189 The generated raw files were analysed in MaxQuant (Cox et al., 2014) as previously described
190 (Clair et al., 2016). Briefly, the false discovery rate was set at 0.01 at the spectral, peptide and
191 protein levels. Proteins were identified with at least two tryptic or semi-tryptic peptides of a
192 minimum length of six amino acids by searching against the UniProt *Bacillus subtilis* strain 168

193 database (UniProtKB, downloaded in 2018). Carbamidomethylation was set as fixed
194 modification and N-terminal acetylation and oxidation of methionine. The match between run
195 algorithm was used and the unmodified intensities (note: LFQ intensities were not used, as they
196 comprise some global normalization steps that would have impaired the cumulative intensity
197 values). Normalization and statistical test were performed with the 'stat' package of R. The
198 intensities were then \log_2 transformed and median centred normalized within a given sample
199 group (e.g. the samples treated with 1.8 J/cm^2 were median centred between them but not
200 against untreated control samples). Only proteins with measured intensities in at least 2-3
201 replicates of a given condition were considered for quantification and missing values were
202 imputed using the minimal normalized intensity of the table divided by two. Two-tailed
203 distribution heteroscedastic Student's t-tests were used to identify proteins with lower
204 abundance in treated samples.

205

206 **2.8. Transmission Electron Microscopy**

207 Freshly prepared spores were centrifuged at $3500 \times g$ during 5 min and fixed for 2 h at room
208 temperature with 2.5 % glutaraldehyde (v/v) in a 0.1 mol/l sodium cacodylate buffer (pH 7.1)
209 containing 0.5 mg/ml ruthenium red. Spores were washed by three centrifugations (5 min at
210 $3500 \times g$) in 0.2 mol/l sodium cacodylate and fixed for 1 h at room temperature with 2 %
211 osmium tetroxide. Next, samples were washed three times with water. After centrifugation,
212 pellets were embedded in 3 % (wt/v) agar and submitted to successive agitated dehydration
213 baths with increasing concentrations of ethanol (30 %, 50 %, 70 %, 90 %, and 100 %). Ethanol

214 was replaced with propylene oxide and sequentially exchanged with araldite resin. Samples
215 were polymerized for 48 h at 60°C. Thin sections (60 - 80 nm thick) were obtained with an
216 ultramicrotome (Leica) and stained with 32 % uranyl acetate and lead citrate and were observed
217 by transmission electron microscopy (TEM) (TE Microscope FEI-Philips CM10). Thin sections of
218 at least 100 spores of one batch of spores were carefully examined with TEM by two
219 experimenters to detect possible differences in structures between control unexposed and PL-
220 exposed spores of *B. subtilis* strain 1A1 (fluence 7.1 J/cm² delivered in 8 flashes). Representative
221 micrographs of the observed differences were selected.

222

223 **2.9. Statistical analysis**

224 Parameters describing inactivation of spores as a function of PL or UV-C fluence were calculated
225 for each survival curve. Fluence leading to 3-fold log reduction (F_3) (Albert and Mafart, 2005)
226 was estimated using non-linear mathematical functions and Microsoft® Excel® 2010 solver
227 function. R_{max} was the observed log₁₀-reduction at the highest tested fluence. TMean values and
228 standard error (SE) of the mean were calculated from data obtained from at least three
229 independent experiments (different dates, different cells cultures and spore preparations). The
230 significance of differences between strains 1A1 or PY79 and derivative mutants was determined
231 by a Student's t-test using the Microsoft® Excel data Analysis tool.

232

233

234 **3. Results**

235 **3.1. Elimination of *B. subtilis* spore coat proteins by PL**

236 Spores of *B. subtilis* strain 1A1 were exposed to increasing fluence emitted by PL or a continuous
237 UV-C source ($\lambda = 254$ nm). Spores were rinsed in cold water to remove loosely attached
238 proteins. Coat proteins were extracted, then separated by SDS-PAGE (see section 2.4). A fluence
239 of 1.8 J/cm^2 delivered by PL caused a 2-3 log reduction in spore counts; PF fluence of 7.1 J/cm^2
240 or 10.8 J/cm^2 caused a log-reduction greater than 6. A UV-C fluence of 0.15 J/cm^2 delivered by
241 the continuous UV-C source caused a 4-log reduction; UV-C fluence of 0.59 J/cm^2 or 0.88 J/cm^2
242 caused a log-reduction greater than 6. Controls were spore coat proteins extracts from
243 unexposed spores. Number and intensity of electrophoresis gel bands of proteins of molecular
244 weight (MW) between 10 kDa and 50 kDa remained unchanged whatever UV-C fluence. In
245 contrast, intensity of bands of almost all 10 kDa - 50 kDa MW proteins decreased as PL fluence
246 increased (Supplementary Fig 2).

247 Proteins of spores treated by PL at a fluence of 1.8 J/cm^2 or by UV-C at a fluence of 0.15 J/cm^2
248 were analysed by western blotting using antibodies targeted against spore coat proteins CotA,
249 CotSA, CotC, CotE, CotH, and CotJ of MW comprised between 25 kDa and 70 kDa (Fig 1 and
250 supplementary Fig 3). The intensity of bands of untreated and UV-C treated spores was highly
251 similar for all tested coat proteins. In contrast, binding to their specific antibodies of spore coat
252 proteins from PL-exposed spores showed diverse patterns. Strikingly, while the applied PL
253 fluence resulted in a lower reduction in spore counts than the applied UV-C fluence, there was a

254 total disappearance on western blots of immunoreactive bands of anti-CotA and anti-CotSA
255 antibodies against the PL-exposed spore protein extract. Binding of anti-CotC and anti-CotE
256 antibodies was slightly lower for the PL-exposed spore coat protein extract than for the ones of
257 the control. Binding with anti-CotJC and anti-CotH antibodies on PL-exposed and control spore
258 protein extract was highly similar. Nevertheless, cotH abundance was lower in PL-treated spores
259 (Fig. 2). Overall, these differences in patterns of immunoreactive bands of coat proteins from
260 spores exposed to PL or UV-C suggests that (i) PL has a stronger effect than UV-C on coat
261 proteins located within the spore, and (ii) that effects of PL may be markedly different for
262 diverse epitopes/proteins (Fig 1). The disappearance of bands on electrophoresis gels or of
263 some bands on western blots could be either due to (i) a detachment of coat proteins from
264 spore as direct consequence of treatment and elimination by washing steps or (ii) of
265 degradation of proteins, at least of site binding antibody.

266

267 **3.2. Proteomic analysis of proteins extracted from *B. subtilis* spores treated by PL**

268 To evaluate in depth how PL treatment affects coat proteins, proteome of coat of *B. subtilis*
269 spores exposed to PL at fluence of 1.8 J/cm² or 7.1 J/cm² (the latter resulting in a log reduction
270 greater than 6) was analysed and compared to coat proteome of non-exposed spores. Same
271 amount of either treated or untreated (control) spores was rinsed in cold water to remove
272 potentially detached proteins. Then coat proteins were extracted and loaded on a SDS page gel.
273 After a short migration of proteins in the gel, the whole protein content from each well was

274 sampled as a sole band and digested with trypsin. Resulting peptides were analysed by LC-
275 MS/MS (See section 2.6).

276 When only 2.9 % of genes are known to code for spore coat proteins (Abhyankar et al., 2015;
277 Abhyankar et al., 2016; Lai et al., 2003; McKenney et al., 2010; McKenney and Eichenberger,
278 2012), over 79 % of the 34,936 spectra attributed to *B. subtilis* proteins (i.e. 27,644 spectra)
279 were belonging to proteins previously identified as located in spore coat, therefore
280 demonstrating that the preparation protocol efficiently enriched for coat proteins
281 (Supplemental datasets 1-2). These spectra were attributed to 1,157 *B. subtilis* peptides
282 (Supplemental dataset 3). In order to determine whether the global amount of detected
283 peptides was affected by PL treatment, peptide intensities attributed to *B. subtilis* proteins were
284 summed within each sample and compared. As shown in Fig 2A, summed peptide intensities
285 were more than 2-fold lower in treated samples than in untreated control (Student's t-test p -
286 value < 0.05). Furthermore, the proportion of the signal attributed to peptides not resulting
287 from trypsin digestion (i.e. non-tryptic peptides) was significantly increased (Student's t-test p -
288 value < 0.01) in treated samples (Fig 2B). Altogether, these results suggest that PL reduces the
289 amount of spore coat proteins likely by cleaving amino acid bonds. In total 149 *Bacillus subtilis*
290 proteins were identified (i.e. with at least two peptides and a false discovery rate < 1 %).

291 Proteins intensities were \log_2 transformed and normalized within each treatment group (to
292 avoid an artificial increase for treated samples of protein amount). The proteomics profile data
293 were classified by applying a principal component analysis (PCA) (Fig 2C). First principal
294 component (PC1) explained 49 % of the overall variance and showed a clear discrimination of
295 PL-treated and untreated samples. Neither second component nor third component of PCA

296 explaining 18.0 % and 12.5 % of the variance respectively (data not shown) distinguished the
297 two treatments. The abundance of 59 proteins was significantly lower in treated samples than
298 in controls (Student's t-test, p -value < 0.05) (Fig 2D and Supplemental dataset 4). These 59
299 proteins include 38 proteins known to be part of spore coat. In most instances, the two applied
300 PL fluences reduced abundance of the same proteins. As expected from western blots, CotA,
301 CotSA, CotC and CotE were reduced in abundance in treated samples and abundance of CotJC
302 was not significantly affected by PL (Supplementary Fig 4). Conversely, while western blot
303 analysis suggested that abundance of CotH in spores was not affected by PL, proteomics
304 measurement indicates the opposite, as indicated in section 3.1. This discrepancy either could
305 result from a lack of specificity of the CotH antibody, or from a degradation, revealed by
306 proteomic analysis of CotH, that does not affect the epitope targeted by the anti-CotH antibody.
307 Interestingly, among spore coat proteins reduced in abundance were some proteins belonging
308 to diverse layers of coat including crust component CotY, outer coat proteins CotE, CotA and
309 CotS (McKenney et al., 2013), or inner coat proteins SafA (Ozin et al., 2000) and YaaH (Imamura
310 et al., 2010). Overall proteome analysis suggests a global degradation or detachment after PL
311 treatment of proteins composing spore coat.

312

313 **3.3. Inactivation of spores of *B. subtilis* strains 1A1 or PY79 and of isogenic mutants with** 314 **defects in genes coding for spore coats**

315 Fig 3 shows typical survival curves of WT strain and derivative mutants exposed to PL. Survival
316 curves were sigmoidal in most instances. The \log_{10} reduction at the highest tested UV-C fluence

317 was comprised, for spores of all strains, between 2.6 and 5.7. The \log_{10} reduction was 1.9 at the
318 highest tested PL fluence with spores of *B. subtilis* strain 1A1 and between 1.4 and 5.3 for
319 spores of its isogenic strains (data not shown). The \log_{10} reduction was 4.8 at the highest tested
320 PL fluence with spores of *B. subtilis* strain PY79 and between 3.2 and 6.0 for spores of its
321 isogenic strains (Fig 4). Curves fitted with the model of Albert and Mafart (Albert and Mafart,
322 2005) satisfactorily followed experimental data for both PL and continuous UV-C treatments and
323 spores of all tested strains, as illustrated for instance in Fig 3. Fluence leading to a three-fold log
324 reduction (F_3) and maximal \log_{10} reduction for the highest tested fluence (R_{\max}) were estimated
325 for each survival curve (Fig 4). Typical inactivation patterns were observed. Spores of some
326 strains, such as *cotG*, behaved similarly to WT spores, with no significant difference for both F_3
327 and R_{\max} ($p > 0.25$). In contrast, spores of strains *spoVID* and *cotE* showed a marked difference in
328 inactivation (Fig 3), illustrated by R_{\max} and F_3 values significantly different compared to the ones
329 of the WT ($p < 0.05$) (Fig 4). Both parameters R_{\max} and F_3 were descriptors of sensitivity of spores
330 of the two WT strains and of the 15 derivative-mutant strains with defects in spore coat-
331 proteins (Fig 4).

332 Overall, there were more derivative-mutant strains with spores showing a higher sensitivity to
333 PL than to continuous UV-C. Spores of 10 of these strains exposed to PL showed a significantly
334 higher sensitivity than WT strains, according to their R_{\max} values while spores of only three
335 strains exposed to continuous UV-C were more sensitive. In the same way, F_3 of spores of six
336 strains exposed to PL were significantly lower than F_3 of WT ($p < 0.05$), while F_3 of spores of only
337 one strain exposed to continuous UV-C was significantly lower than F_3 of WT. Among those
338 strains, F_3 values and R_{\max} values of spores of strains *spoIVA*, *spoVID*, *safA*, *cotE*, *cotF*, and

339 *spoVIA* were significantly different of the ones of WT. Spores of none of the strains exposed to
340 UV-C were significantly affected for both R_{max} and F_3 when compared to the WT strains. Spores
341 of some strains, *cotB*, *cotD*, *cotG* and *cotH*, were not affected in their sensitivity to PL.
342 Surprisingly R_{max} of spores of PY79 *cotXYZ* was lower than R_{max} of spores of WT PY79 strain when
343 exposed to both PL and continuous UV-C, suggesting a higher resistance to both treatments.
344 Spores of a few strains showed higher R_{max} and/or F_3 values than the ones of WT when exposed
345 to continuous UV-C.

346

347 **3.4. Changes in coat arrangement in PL-exposed spores**

348 Thin sectioning transmission electron micrographs (Fig 5) reveal differences in ultrastructure of
349 control spores and spores exposed to PL at a fluence of 7.1 J/cm² resulting in log-reduction
350 greater than 6. A gap between lamellar inner coat and outer coat and its electron dense
351 striations appears on at least 80 % of observed PL-exposed spores.

352

353

354 **4. Discussion**

355 Previous works have clearly established that UV-C wavelengths play a major role in the killing
356 effect of PL on micro-organisms. Filtering-out UV-C radiations dramatically decreases PL
357 efficiency (Bohrerova et al., 2008; Levy et al., 2012; Takeshita et al., 2003). Does this mean that
358 PL causes the same damage as continuous UV-C sources or that mechanisms implicated in
359 resistance to PL or to UV-C are identical? This question has been previously addressed and
360 protein leakage likely consecutive to membrane damage caused by PL and not observed on UV-
361 C treated has been reported for instance on bacteria or yeast cell (Garvey et al., 2016; Kramer et
362 al., 2017; Takeshita et al., 2003). PL applies a polychromatic source of very high energy with
363 wavelengths between 200 nm and 1100 nm. Irradiance of PL sources is several orders of
364 magnitude higher than irradiance of continuous UV-C sources. In the present work, we
365 compared damage caused by PL on *B. subtilis* spores to damage caused by UV-C. PL, but not
366 continuous UV-C treatment, caused in electrophoresis gels disappearance of spore coat proteins
367 bands. These observations were further confirmed by western-blot of specific spore coat
368 proteins, suggesting that PL can induce their elimination by direct degradation or detachment.
369 CotA and CotSA were not detected after exposure of spores to PL. Elimination of CotA and
370 CotSA and more generally of a large set of other proteins including CotB, CotY, YaaH, CotD, SafA,
371 and CotE was confirmed by proteomic analysis of spore coat (Fig 2). In this regard, TEM images
372 revealed unbound junction segments at many locations between inner coat and outer coat
373 because of PL treatment (Fig 5). Remarkably some proteins believed to be mainly localized in
374 the core of the spore such as SASPs SspN and SspO (Cabrera-Hernandez et al., 1999; Cabrera-
375 Hernandez and Setlow, 2000) as well as the 30S ribosomal protein S11 (RpsK) were also reduced

376 in abundance in PL-treated samples suggesting that PL might also impact internal spore core
377 proteins (Fig 2).

378 Subjected to damage, spore coat proteins are also deeply implicated in resistance to PL. Spores
379 of strains defective in spore coat proteins were more sensitive to inactivation by PL than to
380 inactivation by UV-C (Fig 4). This includes in particular spores of strains defective in proteins
381 SpoIVA, SpoVID, SafA, and CotE contributing to morphogenesis of the multi-layered
382 proteinaceous spore coat (McKenney et al., 2013). CotE importance was underlined in a
383 previous report (Eselin et al., 2016) and confirmed here. Spore coat integrity was also a major
384 determinant of the resistance of *B. subtilis* spores to the diversity of Mars UV radiations
385 (Moeller et al., 2012), blue light (\pm 400 nm wavelength) (Djouiai et al., 2018) and to low-pressure
386 plasma, which generates a range of UV wavelengths (a characteristic of the treatment shared
387 with PL) together with reactive chemical species (Raguse et al., 2016), but not to UV-C at 254
388 nm wavelength (Riesenman and Nicholson, 2000). Absorption of radiation by spore coat and/or
389 neutralization of reactive chemical species generated by process are the main hypotheses to
390 explain higher sensitivity of spore coat-defective mutant strains (Djouiai et al., 2018; Raguse et
391 al., 2016). Spore coat proteins are also major determinants of the resistance of spores to
392 oxidative chemical biocides including peroxyacetic acid alone, or in combination with
393 supercritical CO₂, peroxides or iodine (Leggett et al., 2015; Li et al., 2017; Setlow et al., 2016;
394 Setlow et al., 2014).

395 Which properties of PL could generate damage not observed with a continuous UV-C source?
396 Photochemical damage to DNA is generally reported as a major cause of cell death (Kramer et
397 al., 2017). Photochemical damage was not examined in the present work, but is expected

398 because of the high proportion of UV in PL. Previous work already suggested that damage may
399 depend on UV light source: systems allowing DNA repair of damage to DNA are unequally
400 important for spores treated either by PL or by continuous UV-C (Esbelin et al., 2016).
401 Inactivation of micro-organisms has been observed at diverse wavelengths in the UV-C
402 spectrum, certainly to different extents. Spores of *B. pumilus* were 8-9 times less sensitive to a
403 medium pressure UV-source (with peak emission at several wavelengths in the UV-C range) as
404 wavelengths lower than 244 nm were filtered (Beck et al., 2015). In general terms, the relative
405 action spectra of UV radiation against *B. subtilis* spores do not match DNA absorption spectra.
406 Wavelengths below 250 nm are emitted at a lower intensity by the PL source used in this work
407 (Supplementary Fig 1.) but Chen et al. (2009) showed that these wavelengths are more active
408 against *B. subtilis* spores. External layers may not be the primary cause of difference in
409 absorption spectra of spores and DNA, as spectra of decoated spores matched pretty well
410 spectra of intact *B. subtilis* spores. PL has also shown photochemical effects on a range of
411 biological molecules. PL degrades mycotoxins made of conjugated heterocycles such as
412 aflatoxin, zearalenone, ochratoxin, or deoxynivalenol (Moreau et al., 2013). Proteins, in
413 particular side chains of tryptophan, tyrosine, phenylalanine, histidine, cysteine and cysteine,
414 are sensitive to photo-oxidation by UV-A and UV-B, also delivered by PL (Pattison and Davies,
415 2006). Wavelengths of absorption of peptide bonds are in the range 190 – 230 nm and light of
416 these wavelengths also emitted by PL have therefore the potential to degrade proteins.
417 Consequences of protein photo-oxidation are diverse and include unfolding, aggregation, or
418 modification of protein network organization. Analyses of UV and fluorescence spectra of milk
419 proteins specifically exposed to PL indicated protein aggregation and modification of polarity at

420 the proximity of tryptophan residues (Elmnasser et al., 2008). Additionally, photo-oxidized
421 proteins can cause secondary damage through the formation of oxidative products (Manzocco,
422 2015; Pattison and Davies, 2006). Does high energy significantly contribute to inactivation by
423 PL? Second law of photochemistry states that photochemical effect depends only on number of
424 photons absorbed and, in other words, that same inactivation can be achieved with high-
425 fluence – short time and low fluence - long-time exposures (Gomez-Lopez and Bolton, 2016).
426 For instance *B. subtilis* spore inactivation with sources emitting at ± 250 nm was only dependent
427 on fluence and independent of irradiance despite a $1: 10^8$ difference between sources (Rice and
428 Ewell, 2001). Demonstrating specific effects of high irradiance remains an issue.

429 In terms of application, in addition to the acceleration of decontamination processes in food
430 industry lines, the diversity of wavelengths emitted by PL multiplies cellular targets and
431 subsequently cellular damage causing injury and death. Hence, antimicrobial efficiency of PL is
432 potentially higher than antimicrobial efficiency of other UV-radiation sources, and not solely on
433 spores of bacteria. For instance, conidiospores of *Aspergillus niger* were, at equivalent fluence,
434 strongly inactivated by PL and were almost non-affected by a continuous UV-C source (Levy et
435 al., 2012). The spectrum emitted by xenon lamps used for PL decontamination is determined by
436 lamps themselves and by operating conditions (charging voltage, pulse duration...) (Schaefer et
437 al., 2007). In other terms spectra emitted by PL could be adjusted to some extent towards
438 emission of the most efficient wavelengths. A better understanding of the relative contribution
439 of the diverse wavelengths to photochemical damage induced by PL on nucleic acids, proteins
440 and other cell components is necessary for wise evolution and design of future PL devices.

441

442 **Acknowledgments**

443 Antibodies were donated by Pr. Takamatsu and Pr Watabe (Setsunan University, Japan) and Dr.
444 Adriano O. Henriques (ITQB, Lisbon, Portugal). Strains were received from the *Bacillus* Genetic
445 Stock Center (Columbus, Ohio) and from laboratory of Pr. Patrick Eichenberger (New York
446 University, USA). Proteomic analysis was performed in Dr. Jean Armengaud's laboratory at CEA,
447 Institute of Environmental Biology and Biotechnology (Marcoule, France). Dr. Joshua Adkins
448 from Integrative omics group at the Pacific Northwest National Laboratory kindly gave his input
449 on the manuscript.

450

451 **References**

- 452 Abhyankar, W., Pandey, R., Ter Beek, A., Brul, S., de Koning, L.J., de Koster, C.G., 2015. Reinforcement of
453 *Bacillus subtilis* spores by cross-linking of outer coat proteins during maturation. Food Microbiol. 45, 54-
454 62.
- 455 Abhyankar, W.R., Kamphorst, K., Swarge, B.N., van Veen, H., van der Wel, N.N., Brul, S., de Koster, C.G.,
456 Koning, L.J., 2016. The Influence of sporulation conditions on the spore coat protein composition of
457 *Bacillus subtilis* spores. Front. Microbiol. 7, 1-10.
- 458 Albert, I., Mafart, P., 2005. A modified Weibull model for bacterial inactivation. Int. J. Food Microbiol.
459 100, 197-211.
- 460 Arrieta-Ortiz, M.L., Hafemeister, C., Bate, A.R., Chu, T., Greenfield, A., Shuster, B., Barry, S.N., Gallitto, M.,
461 Liu, B., Kacmarczyk, T., Santoriello, F., Chen, J., Rodrigues, C.D.A., Sato, T., Rudner, D.Z., Driks, A.,
462 Bonneau, R., Eichenberger, P., 2015. An experimentally supported model of the *Bacillus subtilis* global
463 transcriptional regulatory network. Mol. Syst. Biol. 11, 839.
- 464 Beck, S.E., Wright, H.B., Hargy, T.M., Larason, T.C., Linden, K.G., 2015. Action spectra for validation of
465 pathogen disinfection in medium-pressure ultraviolet (UV) systems. Water Res. 70, 27-37.
- 466 Bohrerova, Z., Shemer, H., Lantis, R., Impellitteri, C.A., Linden, K.G., 2008. Comparative disinfection
467 efficiency of pulsed and continuous-wave UV irradiation technologies. Water Res. 42, 2975-2982.
- 468 Cabrera-Hernandez, A., Sanchez-Salas, J.L., Paidhungat, M., Setlow, P., 1999. Regulation of four genes
469 encoding small, acid-soluble spore proteins in *Bacillus subtilis*. Gene 232, 1-10.
- 470 Cabrera-Hernandez, A., Setlow, P., 2000. Analysis of the regulation and function of five genes encoding
471 small, acid-soluble spore proteins of *Bacillus subtilis*. Gene 248, 169-181.
- 472 Carlin, F., 2011. Origin of bacterial spores contaminating foods. Food Microbiol 28, 177-182.
- 473 Carrasco, G., Urrestarazu, M., 2010. Green chemistry in protected horticulture: the use of peroxyacetic
474 acid as a sustainable strategy. Int J. Mol. Sci. 11, 1999-2009.

475 Castillo, R., Biedermann, M., Riquet, A.M., Grob, K., 2013. Comprehensive on-line HPLC-GC for screening
476 potential migrants from polypropylene into food: The effect of pulsed light decontamination as an
477 example. *Polym. Degrad. Stabil.* 98, 1679-1687.

478 Chen, R.Z., Craik, S.A., Bolton, J.R., 2009. Comparison of the action spectra and relative DNA absorbance
479 spectra of microorganisms: Information important for the determination of germicidal fluence (UV dose)
480 in an ultraviolet disinfection of water. *Water Res.* 43, 5087-5096.

481 Clair, G., Armengaud, J., Duport, C., 2012. Restricting fermentative potential by proteome remodeling.
482 An adaptive strategy evidenced in *Bacillus cereus*. *Mol Cell Proteomics* 11.

483 Clair, G., Piehowski, P.D., Nicola, T., Kitzmiller, J.A., Huang, E.L., Zink, E.M., Sontag, R.L., Orton, D.J.,
484 Moore, R.J., Carson, J.P., Smith, R.D., Whitsett, J.A., Corley, R.A., Ambalavanan, N., Ansong, C., 2016.
485 Spatially-resolved proteomics: rapid quantitative analysis of laser capture microdissected alveolar tissue
486 samples. *Sci. Rep.* 6.

487 Coote, J.G., 1972. Sporulation in *Bacillus subtilis*. Characterization of oligosporogenous mutants and
488 comparison of their phenotypes with those of asporogenous mutants. *J. Gen. Microbiol.* 71, 1-15.

489 Costa, T., Isidro, A.L., Moran, C.P., Henriques, A.O., 2006. Interaction between coat morphogenetic
490 proteins SafA and SpoVID. *J. Bacteriol.* 188, 7731-7741.

491 Cox, J., Hein, M.Y., Lubner, C.A., Paron, I., Nagaraj, N., Mann, M., 2014. Accurate proteome-wide label-free
492 quantification by delayed normalization and maximal peptide ratio extraction, termed maxLFQ. *Mol. Cell.*
493 *Proteomics* 13, 2513-2526.

494 Cutting, S., Zheng, L.B., Losick, R., 1991. Gene encoding 2 alkali-soluble components of the spore coat
495 from *Bacillus subtilis*. *J. Bacteriol.* 173, 2915-2919.

496 Djouia, B., Thwaite, J.E., Laws, T.R., Commichau, F.M., Setlow, B., Setlow, P., Moeller, R., 2018. Role of
497 DNA repair and protective components in *Bacillus subtilis* spore resistance to inactivation by 400-nm-
498 wavelength blue light. *Appl. Environ. Microbiol.* 84.

499 Donovan, W., Zheng, L., Sandman, K., Losick, R., 1987. Genes encoding spore coat polypeptides from
500 *Bacillus subtilis*. J. Mol. Biol. 196, 1-10.

501 Elmnasser, N., Dalgarrondo, M., Orange, N., Bakhrouf, A., Haertle, T., Federighi, M., Chobert, J.M.,
502 2008. Effect of pulsed-light treatment on milk proteins and lipids. J. Agric. Food Chem. 56, 1984-1991.

503 Esbelin, J., Malléa, S., Clair, G., Carlin, F., 2016. Inactivation by pulsed light of *Bacillus subtilis* spores with
504 impaired protection factors. Photochem. Photobiol. 92, 301-307.

505 Garvey, M., Rowan, N.J., 2019. Pulsed UV as a potential surface sanitizer in food production processes to
506 ensure consumer safety. Curr. Opin. Food Sci. 26, 65-70.

507 Garvey, M., Stocca, A., Rowan, N., 2016. Use of a Real Time PCR assay to assess the effect of pulsed light
508 inactivation on bacterial cell membranes and associated cell viability. Water Env. Res. 88, 168-174.

509 Garvey, M., Thokala, N., Rowan, N., 2014. A comparative study on the pulsed UV and the low-pressure
510 UV inactivation of a range of microbial species in water. Water Environ. Res. 86, 2317-2324.

511 Gomez-Lopez, V.M., Bolton, J.R., 2016. An approach to standardize methods for fluence determination in
512 bench-scale pulsed light experiments. Food Bioprocess Technol. 9, 1040-1048.

513 Gomez-Lopez, V.M., Ragaert, P., Debevere, J., Devlieghere, F., 2007. Pulsed light for food
514 decontamination: a review. Tr. Food Sci. Technol. 18, 464-473.

515 Gopal, N., Hill, C., Ross, P.R., Beresford, T.P., Fenelon, M.A., Cotter, P.D., 2015. The prevalence and
516 control of *Bacillus* and related spore-forming bacteria in the dairy industry. Front Microbiol. 6, 1418.

517 Hullo, M.F., Moszer, I., Danchin, A., Martin-Verstraete, I., 2001. CotA of *Bacillus subtilis* is a copper-
518 dependent laccase. J. Bacteriol. 183, 5426-5430.

519 Imamura, D., Kuwana, R., Takamatsu, H., Watabe, K., 2010. Localization of proteins to different layers
520 and regions of *Bacillus subtilis* spore coats. J. Bacteriol. 192, 518-524.

521 Imamura, D., Kuwana, R., Takamatsu, H., Watabe, K., 2011. Proteins involved in formation of the
522 outermost layer of *Bacillus subtilis* spores. J. Bacteriol. 193, 4075-4080.

523 Jenkinson, H.F., 1981. Germination and resistance defects in spores of a *Bacillus subtilis* mutant lacking a
524 coat polypeptide J. Gen. Microbiol. 127, 81-91.

525 Koutchma, T.N., Forney, L., Moraru, C.I., 2009. Ultraviolet Light in Food Technology. Principles and
526 Applications. CRC Press, Boca Raton.

527 Kramer, B., Wunderlich, J., Muranyi, P., 2017. Recent findings in pulsed light disinfection. J. Appl.
528 Microbiol. 122, 830-856.

529 Kuwana, R., Ikejiri, H., Yamamura, S., Takamatsu, H., Watabe, K., 2004. Functional relationship between
530 SpoVIF and GerE in gene regulation during sporulation of *Bacillus subtilis*. Microbiology-(UK) 150, 163-
531 170.

532 Lai, E.M., Phadke, N.D., Kachman, M.T., Giorno, R., Vazquez, S., Vazquez, J.A., Maddock, J.R., Driks, A.,
533 2003. Proteomic analysis of the spore coats of *Bacillus subtilis* and *Bacillus anthracis*. J. Bacteriol. 185,
534 1443-1454.

535 Leggett, M.J., Schwarz, J.S., Burke, P.A., McDonnell, G., Denyer, S.P., Maillard, J.Y., 2015. Resistance to
536 and killing by the sporicidal microbicide peracetic acid. J. Antimicrob. Chemother. 70, 773-779.

537 Levy, C., 2010. Principaux facteurs influençant l'efficacité de la lumière pulsée pour la décontamination
538 des microorganismes pathogènes et d'altération des denrées alimentaires. , Thèse d'Université.
539 Université d'Avignon et des Pays de Vaucluse. available at <http://tel.archives-ouvertes.fr/tel-00747302>.

540 Levy, C., Aubert, X., Lacour, B., Carlin, F., 2012. Relevant factors affecting microbial surface
541 decontamination by pulsed light. Int. J. Food Microbiol. 152, 168-174.

542 Levy, C., Bornard, I., Carlin, F., 2011. Deposition of *Bacillus subtilis* spores using an airbrush-spray or spots
543 to study surface decontamination by pulsed light. J. Microbiol. Meth. 84, 223-227.

544 Li, Q., Korza, G., Setlow, P., 2017. Killing the spores of *Bacillus* species by molecular iodine. J. Appl.
545 Microbiol. 122, 54-64.

546 Maillard, J.Y., 2011. Innate resistance to sporicides and potential failure to decontaminate. *J. Hosp. Inf.*
547 *77*, 204-209.

548 Manzocco, L., 2015. Photo-induced modification of food protein structure and functionality. *Food Eng.*
549 *Rev. 7*, 346-356.

550 McKenney, P.T., Driks, A., Eichenberger, P., 2013. The *Bacillus subtilis* endospore: assembly and functions
551 of the multilayered coat. *Nat. Rev. Microbiol.* *11*, 33-44.

552 McKenney, P.T., Driks, A., Eskandarian, H.A., Grabowski, P., Guberman, J., Wang, K.H., Gitai, Z.,
553 Eichenberger, P., 2010. A distance-weighted interaction map reveals a previously uncharacterized layer
554 of the *Bacillus subtilis* spore coat. *Curr. Biol.* *20*, 934-938.

555 McKenney, P.T., Eichenberger, P., 2012. Dynamics of spore coat morphogenesis in *Bacillus subtilis*. *Mol.*
556 *Microbiol.* *83*, 245-260.

557 Moeller, R., Schuerger, A.C., Reitz, G., Nicholson, W.L., 2012. Protective role of spore structural
558 components in determining *Bacillus subtilis* spore resistance to simulated Mars surface conditions. *Appl.*
559 *Environ Microbiol.* *78*, 8849-8853.

560 Moeller, R., Wassmann, M., Reitz, G., Setlow, P., 2011. Effect of radioprotective agents in sporulation
561 medium on *Bacillus subtilis* spore resistance to hydrogen peroxide, wet heat and germicidal and
562 environmentally relevant UV radiation. *J. Appl. Microbiol.* *110*, 1485-1494.

563 Moir, A., 1981. Germination properties of a spore coat-defective mutant of *Bacillus subtilis*. *J. Bacteriol.*
564 *146*, 1106-1116.

565 Moreau, M., Lescure, G., Agoulon, A., Svinareff, P., Orange, N., Feuilloley, M., 2013. Application of the
566 pulsed light technology to mycotoxin degradation and inactivation. *J. Appl. Toxicol.* *33*, 357-363.

567 Nicholson, W.L., Munakata, N., Horneck, G., Melosh, H.J., Setlow, P., 2000. Resistance of *Bacillus*
568 endospores to extreme terrestrial and extraterrestrial environments. *Microbiol. Mol. Biol. Rev.* *64*, 548-
569 572.

570 Nicholson, W.L., Setlow, P., 1990. Sporulation, germination and outgrowth, in: Harwood, C.R., Cutting,
571 S.M. (Eds.), *Molecular Biology Methods for Bacillus*. John Wiley and Sons Ltd, Chichester, England, pp.
572 391-450.

573 Ozin, A.J., Henriques, A.O., Yi, H., Moran, C.P., Jr., 2000. Morphogenetic proteins SpoVID and SafA form a
574 complex during assembly of the *Bacillus subtilis* spore coat. *J Bacteriol.* 182, 1828-1833.

575 Pattison, D.I., Davies, M.J., 2006. Actions of ultraviolet light on cellular structures, in: Bignold, L.P. (Ed.),
576 *Cancer: Cell Structures, Carcinogens and Genomic Instability*. Experientia Supplementum, vol 96.
577 Birkhäuser Basel, pp. 131-157.

578 Raguse, M., Fiebrandt, M., Denis, B., Stapelmann, K., Eichenberger, P., Driks, A., Eaton, P., Awakowicz, P.,
579 Moeller, R., 2016. Understanding of the importance of the spore coat structure and pigmentation in the
580 *Bacillus subtilis* spore resistance to low-pressure plasma sterilization. *J. Phys. D: Appl. Phys.* 49, 1-16.

581 Rice, J.K., Ewell, M., 2001. Examination of peak power dependence in the UV inactivation of bacterial
582 spores. *Appl. Environ. Microbiol.* 67, 5830-5832.

583 Riesenman, P.J., Nicholson, W.L., 2000. Role of the spore coat layers in *Bacillus subtilis* spore resistance
584 to hydrogen peroxide, artificial UV-C, UV-B, and solar UV radiation. *Appl. Environ. Microbiol.* 66, 620-626.

585 Roels, S., Driks, A., Losick, R., 1992. Characterization of *spoVIA*, a sporulation gene involved in coat
586 morphogenesis in *Bacillus subtilis*. *J. Bacteriol.* 174, 575-585.

587 Russell, A.D., 2003. Biocide use and antibiotic resistance: the relevance of laboratory findings to clinical
588 and environmental situations. *Lancet Infect. Dis.* 3, 794-803.

589 Schaefer, R., Grapperhaus, M., Schaefer, I., Linden, K., 2007. Pulsed UV lamp performance and
590 comparison with UV mercury lamps. *J. Environ. Eng Sci.* 6, 303-310.

591 Setlow, B., Korza, G., Blatt, K.M.S., Fey, J.P., Setlow, P., 2016. Mechanism of *Bacillus subtilis* spore
592 inactivation by and resistance to supercritical CO₂ plus peracetic acid. *J. Appl. Microbiol.* 120, 57-69.

593 Setlow, B., Parish, S., Zhang, P., Li, Y.Q., Neely, W.C., Setlow, P., 2014. Mechanism of killing of spores of
594 *Bacillus anthracis* in a high-temperature gas environment, and analysis of DNA damage generated by
595 various decontamination treatments of spores of *Bacillus anthracis*, *Bacillus subtilis* and *Bacillus*
596 *thuringiensis*. J. Appl. Microbiol. 116, 805-814.

597 Setlow, P., 2007. I will survive: DNA protection in bacterial spores. Tr. Microbiol. 15, 172-180.

598 Setlow, P., 2014. Spore resistance properties. Microbiol. Spectr. 2.

599 Soni, A., Oey, I., Silcock, P., Bremer, P., 2016. *Bacillus* spores in the food industry: a review on resistance
600 and response to rovel inactivation technologies. Compr. Rev. Food Sci. Food Saf. 15, 1139-1148.

601 Takeshita, K., Shibato, J., Sameshima, T., Fukunaga, S., Isobe, S., Arihara, K., Itoh, M., 2003. Damage of
602 yeast cells induced by pulsed light irradiation. Int. J. Food Microbiol. 85, 151-158.

603 Wells-Bennik, M.H.J., Eijlander, R.T., den Besten, H.M.W., Berendsen, E.M., Warda, A.K., Krawczyk, A.O.,
604 Groot, M.N.N., Xiao, Y.H., Zwietering, M.H., Kuipers, O.P., Abee, T., 2016. Bacterial Spores in Food:
605 Survival, Emergence, and Outgrowth, in: Doyle, M.P., Klaenhammer, T.R. (Eds.), Annual Review of Food
606 Science and Technology, Vol 7, pp. 457-482.

607 Zheng, L.B., Donovan, W.P., Fitz-James, P.C., Losick, R., 1988. Gene encoding a morphogenic protein
608 required in the assembly of the outer coat of the *Bacillus subtilis* endospore. Gen. Develop. 2, 1047-
609 1054.

610

611

612

613 **Figure captions**

614 **Fig 1.** Western blots with antibodies directed against spore coat proteins CotA, CotSA, CotC,
615 CotE, CotH and CotJC after exposure of *B. subtilis* spores to a fluence of 1.8 J/cm² delivered by
616 PL (allowing a 2-log reduction) or a UV-C fluence of 0.15 J/cm² (allowing a 4-log reduction).
617 Numbers on the left-hand are molecular weights (kDa); U: untreated controls; PL: pulsed-light
618 treated samples; UV: UV-C treated samples.


619 **Fig 2.** Proteomic characterization of *B. subtilis* strain 1A1 spores subjected to a pulsed light (PL)
620 treatment. (A) Cumulative intensity of peptides for PL-treated and control samples. (B)
621 Percentage of cumulative intensity attributed to semi-tryptic events. In (A) and (B) error bars
622 represent standard error and asterisks represent Student's t-test *p*-values (***, *p* <0.001; **,
623 *p* <0.01, *, *p* <0.05). (C) Principal Component Analysis projection of untreated (orange) and PL-
624 treated samples (blue). The first principal component separates out treated and untreated
625 sample. Replicates are identified by numbers (1 to 3). (D) Heat map of proteins significantly
626 changing in abundance in PL treated spore coat compared to untreated spore coat. The color
627 scale represent log₂ (reduction) of protein abundance in treated samples. Asterisks represent
628 the *p*-value cutoffs (***, *p* <0.001; **, *p* <0.01, *, *p* <0.05) of the comparison with untreated
629 samples. The green box shows results validating western blot results (Fig 1). The black box
630 around CotH indicates a possible decrease revealed by proteomic analysis, and conversely not
631 observed on the western-blot (Fig 1).

632 **Fig 3.** Inactivation by PL of *B. subtilis* strain 1A1 (◆) and of a selection of derivative mutants
633 with defect in genes *cotG* (●), *spoVID* (▲) or *cotE* (■) coding for morphogenetic and/or spore
634 coat proteins. Symbols show mean of at least three biological replications (different spore


635 preparations). Error bars show standard error. Color lines represent curves that have the best fit
636 to data points with the model of Albert and Mafart (2005). The model was otherwise used to
637 determine fluence to 3 log-reduction (F_3) and log-reduction at maximal tested fluence (R_{max})
638 presented in Fig 4.

639 **Fig 4.** Comparison of log-reduction at maximal tested fluence (R_{max}) (A, B) and of fluence to 3
640 log-reduction (F_3) (C, D) between *B. subtilis* strain WT 1A1 or PY79 (in bold characters) and their
641 respective derivative mutants with defects in genes coding for spore coat proteins. Strains were
642 treated either with pulsed light (A, C) or with a continuous UV-C source (B, D). Histogram bars
643 represent the arithmetic difference in R_{max} values of WT strains and respective mutants. Errors
644 bars represent standard error ($n \geq 3$). Shaded bands surrounding vertical axes show standard
645 error on R_{max} or F_3 values of WT strains. Negative values in R_{max} comparison graphs correspond
646 to inactivation greater for mutant strain than for WT. Values < 1.0 in F_3 comparison graphs (C, D)
647 correspond to inactivation greater for mutant strain than for WT. Blue histogram bars show a
648 significant difference in R_{max} or F_3 value at $p < 0.05$. The \log_{10} reduction was 1.8 at the highest
649 tested PL fluence with *B. subtilis* strain 1A1 and between 2 and 6 for its derivative strains. The
650 \log_{10} reduction was between 4 and 5 at the highest tested PL fluence with *B. subtilis* strain PY79
651 and between 2 and 6 for its derivative strains.


652 **Fig 5.** Transmission electron microscopy pictures and magnification of spore coat section of *B.*
653 *subtilis* strain 1A1 spores, control (A, C) and exposed to a PL fluence of 7.1 J/cm² (B, D).
654 Co= spore core; Cx = spore cortex; IC = inner spore coat, OC = outer spore coat. Red arrows
655 indicate position of gap between inner and outer spore coat observed in PL-exposed spores.




A



B





C






D

Protein significantly lower in abundance in the treated samples





A Pulsed Light R_{max} (WT) - R_{max} (tested strain)**B** UV-C R_{max} (WT) - R_{max} (tested strain)**C** Pulsed light F_3 (tested strain) / F_3 (WT)**D** UV-C F_3 (tested strain) / F_3 (WT)