



Cyanobacteria and cyanotoxins in estuarine water and sediment

M. Bormans, V. Savar, B. Legrand, E. Mineaud, E. Robert, E. Lance, Z. Amzil

► To cite this version:

M. Bormans, V. Savar, B. Legrand, E. Mineaud, E. Robert, et al.. Cyanobacteria and cyanotoxins in estuarine water and sediment. *Aquatic Ecology*, 2020, 54 (2), pp.625-640. 10.1007/s10452-020-09764-y . hal-02862833

HAL Id: hal-02862833

<https://univ-rennes.hal.science/hal-02862833>

Submitted on 25 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Cyanobacteria and cyanotoxins in estuarine water and sediment

Myriam Bormans^{*1}, Véronique Savar², Benjamin Legrand³, Emilien Mineaud¹,
Elise Robert², Emilie Lance⁴⁵, Zouher Amzil²

¹ Univ Rennes, CNRS, ECOBIO - UMR 6553, F-35000 Rennes, France

² IFREMER, Phycotoxins Laboratory, F-44311 Nantes, France

³ ATHOS Environnement, 112 avenue du Brézet, 63100 Clermont Ferrand, France

⁴ UMR-I 02 SEBIO, Bat 18, Campus du Moulin de la Housse, BP 1039, 51687 REIMS Cedex 2, France

⁵ UMR 7245 MNHN/CNRS MCAM, équipe CCE, Muséum National d'Histoire Naturelle, 12 rue Buffon, F-75231, Paris.

*Corresponding author: myriam.bormans@univ-rennes1.fr tel : +33 2 2323 5979

Key words: toxic cyanobacteria, microcystin, colonies, estuary, sediment

Abstract

While transfer of freshwater cyanobacteria to estuaries has been observed worldwide, the associated transfer of cyanotoxins is less often reported, in particular the sediment contribution. During fall 2018, we monitored the co-occurrence of cyanobacteria and microcystin (MC) in both the water column and in surface sediments at 5 stations along a river continuum, from a freshwater reservoir to the coastal area in Brittany, France. Cyanobacteria dominated the phytoplankton community in the water column with high densities at the freshwater sites. *Microcystis* cells and intracellular MC transfer to estuarine and marine sites were observed with decreasing concentrations in accordance with flow dilution. Extracellular MC showed the opposite trend and increased from upstream to downstream in accordance with the lysing of the cells at elevated salinities. Surface sediment samples contained high densities of colonial *Microcystis* in freshwater and with decreasing concentrations along the salinity gradient, similarly to cells concentrations in the water column. Intracellular MC was detected in sediment at all sites except at the marine outlet suggesting the survival of intact cells. Extracellular MC concentrations in sediment were up to 5 times higher than intracellular

concentrations suggesting incomplete MC degradation. *mcyB* genes were present at all sites while *mcyA* genes were absent at the marine outlet suggesting the presence of toxic strains along the estuary. The high densities of intact colonies of potentially toxic *Microcystis* in the estuarine sediment strongly suggest that sediments can act as an inoculum of cyanobacteria and cyanotoxins in estuaries.

1. Introduction

Cyanobacterial blooms have been reported worldwide (Merel et al. 2013) and their proliferations have been increasing in recent years as a result of anthropogenic activities including eutrophication and climate warming (O'Neil et al. 2012; Rigosi et al. 2014; Paerl 2018). Freshwater cyanobacteria produce a large variety of toxins (including hepatotoxins, neurotoxins, dermatotoxins) which have strong negative impact on animal and human health (Lance et al. 2010; Metcalf and Codd 2012; Wood 2016 for a review, Meriluoto et al. 2017). Among the diversity of cyanotoxins, microcystin (MC) is largely recognized as the most common and widespread in freshwater ecosystems (Harke et al. 2016). The general structure of those cyclic heptapeptides includes a specific beta amino acid-3-amino-9-methoxy-2,6,8-trimethyl-10-phenyl-4,6-decadienoic acid-Adda (Ortiz et al. 2017; Tillett et al. 2000) as well as two amino acids that can vary leading to the identification of more than 250 MC variants (Puddick et al. 2014). The regulation and synthesis of MC, as well as its ecological role, are complex and not yet fully understood (Omidi et al. 2017).

The transfer of freshwater cyanobacteria to estuaries has been observed worldwide (Preece et al. 2017 for a review), in Africa (Ndelela et al. 2016), USA (e.g. Lehman et al. 2005; Peacock et al. 2018), South America (e.g. Dörr et al. 2010), Australia (Robson and Hamilton 2003; Orr et al. 2004), Europe (e.g. Verspagen et al. 2006; Tonk et al. 2007; Paldavičiene et al. 2009; Bormans et al. 2019), and Turkey (Taş et al. 2006). The majority of these studies reported on the cyanobacterial transfer being dominated by *Microcystis aeruginosa* demonstrating a certain salt tolerance of that species. The associated transfer of intracellular or extracellular MC along the river continuum was less often reported (Preece et al. 2017). In particular MC transfer to the coastal environment resulting from freshwater discharge from an upstream

reservoir has only been reported in Italy (De Pace et al. 2014), Japan (Umehara et al. 2012; 2015) and recently in France (Bormans et al. 2019).

To our knowledge, MC occurrence in estuarine sediment has only been reported in Isahaya Bay, Japan (Umehara et al. 2012, 2015), and in sub-estuaries of the Baltic Sea (Paldaviciene et al. 2015) and Chesapeake Bay, USA (Bukaveckas et al. 2017) while cyanobacterial (e.g. *Microcystis*) occurrence in estuarine sediment has never been demonstrated. Hence, given the demonstrated contamination of estuarine and marine benthic and pelagic organisms (Preece et al. 2015; Gible et al. 2016; Bukaveckas et al. 2017, 2018), as well as the high potential for tidal derived sediment resuspension in estuaries (Labry et al. 2016) we wanted to investigate the sediment's contribution of a French estuary as a potential source/sink of both cyanobacteria and MC.

Here we report on the co-occurrence of both cyanobacteria and microcystin in the water column and in surface sediment along a river continuum from a freshwater reservoir discharge to the marine environment during fall 2018. This study is part of a larger project aiming at the evaluation of the potential risk of contamination of aquatic organisms (i.e. bivalves) by cyanotoxins during transfer from a freshwater reservoir to an estuary mouth in Brittany, France. We present results on phytoplankton/cyanobacterial species composition, and concentrations of different variants of MC both in intracellular and extracellular forms. The quantification of those two forms of toxins are necessary as we anticipate gradual cells lysing along the salinity gradient. We also report on the presence of microcystin producing genes *mcyA* and *mcyB* together with the *anaC* gene in surface sediment to assess the potential toxicity of estuarine sediment.

2. Materials and methods

2.1. Study site and sampling protocols

We chose to investigate a site in Brittany where most of the French agricultural lands are located and where reports of toxic blooms have been numerous in freshwater lakes and reservoirs (Vezie et al. 1998; Brient et al. 2009; Pitois et al. 2018). Moreover, in Brittany, most rivers have short residence times (Fraisie et al, 2013)

At each station, subsurface grab water sampling was carried out in the water column to: i) determine physico-chemical parameters (temperature, conductivity, dissolved oxygen, Phosphorus and Nitrogen) using a YSI 6920 multi-parameter probe (YSI Environmental, Anhydre) and physico-chemical analyses, ii) describe phytoplankton and cyanobacterial species (identification and enumeration), and (iii) quantify microcystins in the phytoplanktonic biomass and in the filtered water. At each station, surface sediment sampling (top 1 to 2 cm) was carried out to (i) determine sediment characteristics including percentage of organic matter (OM) and major elements, ii) identify and count phytoplankton species including cyanobacteria, (iii) quantify cyanotoxins and (iv) determine the presence of MC and anatoxin production genes. It is important to note that for the sampling of surface sediment, the site F1 was chosen in the river directly downstream of the reservoir, as no sediment was found in the reservoir itself at the site where the water column sampling was performed, the bottom there being dominated by larger rocks.

2.2. Samples Analyses

2.2.1. Water analyses

Water column samples were transported in dark and cold conditions and filtered upon arrival at the laboratory within few hours of sampling. Dissolved inorganic nutrient concentrations were measured from filtered (GF/F 0.7 μm) water using common colorimetric methods with a Bran and Luebbe Autoanalyser 3 (Axflow, Norderstedt, Germany). Nitrate was measured after reduction to nitrite on a cadmium-copper column (Henriksen and Selmer-Olsen 1970). Phosphate was measured following the method of Murphy and Riley (1962). Filtration was also done to determine total suspended solids (TSS). Whatman glass fiber filters (GF/F 0.7 μm) were dried at 105°C and weighed. Ashing to 550°C was performed to determine the organic contribution. Phytoplankton and cyanobacteria identification and counts were conducted on both fresh and lugolised samples under an optical microscope (100x magnification) using a Nageotte chamber within 48 hours of sampling as reported in Lance et al. (2010) and Pannard et al. (2018). This was preferred over a haemocytometer chamber for the larger volume sampled to limit counting errors as described in Brient et al. (2008).

2.2.1.1. *Microcystin extraction*

Water samples containing cyanobacteria were filtered upon arrival at the laboratory through a 0.1 µm GF filter to separate the cell pellet for the intracellular cyanotoxin analysis and the filtrate which may contain dissolved extracellular toxins. Both filters and filtrates were frozen at -20°C until chemical analysis. The filtrate was purified on a C₁₈ SPE cartridge (Solid Phase Extraction) according to the ISO 20179 standard method (Anon, 2005). The fraction containing the toxins was frozen until LC-MS/MS analysis. The GF filter was sonicated with 10 mL of MeOH for 15 minutes. The MeOH extract was evaporated under nitrogen and then the residue was diluted in 1.5 mL of methanol.

2.2.2. *Sediment samples*

Fresh surface sediment was kept in the dark and at 4°C until analyses. Sediment characteristics including major elements and % organic matter (OM) were analysed by the SARM (Service d'Analyse des Roches et des Minéraux) laboratory (Nancy, France) using an ICP-MS iCapQ instrument.

2.2.2.1. *Extraction of cyanobacteria*

Cyanobacteria were extracted from sediment using a density gradient with Ludox TM-50 (Sigma-Aldrich, Saint Louis, USA), as described in Legrand et al. (2016). 0.5g of fresh sediment was diluted with 9.5 ml of distilled water and 4 ml of Ludox. These solutions were sonicated for 30 seconds with a frequency of 50% and a power of 80 W (Sonoplus, Bandelin, Berlin, Germany). Finally, solutions were centrifuged at 10,000 x g for 30 minutes and after the first 6 ml was pipetted and kept in the dark at 4°C for further analysis.

2.2.2.2. *Estimation of cyanobacterial abundance*

Genus determination for vegetative cells and akinetes (Nostocale resting stage cells) was based on morphological criteria using reference books (Bourelly 1985; Komàrek and Anagnostidis 2005, 2013). 2 ml of Ludox solution was filtered through an 8-µm mesh (TETP filter; Merck Millipore, Tullagreen, Ireland). Counts were performed in triplicate for each sample in this study. Cyanobacteria were enumerated at a

magnification of x200 with an epifluorescence microscope (Axiovert 200 M; Zeiss, Oberkochen, Germany) and a minimum of 40 fields was counted for each filter. One sample (F2) was tested for cell viability with SYTOX green following Legrand et al. (2016). Briefly, 1 µL of SYTOX green (50 mM) was added for 1 mL of isolated cyanobacterial solution. Resulting solution was incubated in the dark at least for 30 min. 1 mL of this solution was filtered using an 8 µm filter (TETP, Merck Millipore, Tullagreen, Ireland). The sample was enumerated with the same epifluorescence microscope than previously. Two separate counts were performed: the first at 546 nm to determine intact cells still containing chlorophyll pigment; and the second at 448 nm to discriminate damaged cells with SYTOX green staining.

2.2.2.3. DNA extraction and target genes amplification

Genomic DNA from sediment samples was extracted with the FastDNA® Spin kit for soil (MP biomedical, Santa Ana, USA) following the manufacturer's instructions. The DNA concentration in each sample was estimated using a UV spectrophotometer (Nanodrop® ND 2000). The cyanobacterial 16S rRNA gene was amplified in sediment samples using the cya359F and cya781R set of primers from Nubel et al. (1997). Then, two genes involved in the biosynthesis of microcystin, *mcyA* and *mcyB* genes and one gene involved in the biosynthesis of anatoxin-a, *anaC*, were also investigated in this study, given the cyanobacterial species found in sediment. *anaC* and *mcyB* were amplified with nested PCRs whereas classic PCR was used to detect *mcyA*. For *mcyB* nested PCR, MCY F1/R1 then MCY F1/R2 were used and *mcyA*_Cd1F and *mcyA*-Cd1R for *mcyA*. *anaC* was detected with the sets of primers anxgen-F2/R and *anaC* 80F/706R for nested 1 and 2 PCR respectively. All PCR mixtures and PCR programs were the same as described in Legrand et al. (2017). All PCR products were revealed with 0.5% agarose gel with 0.3 mg L⁻¹ ethidium bromide and migrated in a TAE buffer 0.5X at 100V for 30 minutes.

2.2.2.4. Microcystin extraction

To analyse total MC (intracellular plus extracellular), the crude sediment (approximately 2 g) were extracted, according to the method of Umehara et al (2012), 4 times with 4 ml of MeOH/H₂O (30%/70%), each time using sonication for 45 min and then centrifugation at 3500 g at 4 °C for 10 min. To analyse intracellular

MC, the ludox supernatant containing cyanobacteria was filtered under pressure (around 53kPa) using 25 µm nylon filtration tissues in order to eliminate akinetes and extracellular MC. These tissues were cleaned with distilled water under pressure 3 times. Then, tissues containing *Microcystis* were cleaned with methanol 100%. These solutions were kept in dark conditions at 4°C until further analysis.

2.2.3 Chemical analysis of microcystins by LC-MS / MS

All extracts obtained from the water column and surface sediment were filtered through a 0.2 µm filter and analyzed by Ultra Fast Liquid Chromatography (Shimadzu, Marne la Vallée, France) coupled to API-5500 QTrap tandem mass spectrometry (ABSciex, Villebon sur Yvette, France). Toxins were separated on a Kinetex XB C18 column (100 x 2.1 mm, 2.6 µm, Phenomenex) at 25°C, with water (A) and acetonitrile (B), both containing 0.1% formic acid at 0.3 mL min⁻¹ flow rate. The gradient was raised from 30 to 80% B in 5 min and was held during 1 min before dropping down during 0.5 min to the initial conditions.

Mass spectrometry detection was carried out in multiple reactions monitoring (MRM) mode. The electrospray ionization interface (ESI) was operated in positive mode using source setting: curtain gas set at 30 psi, ion spray at 5000 V, a turbogas temperature of 300°C, gas 1 and 2 set at 30 and 40 psi respectively and an entrance potential of 10 V. Each toxin was identified and quantified with two transitions as follow (Table S1): The toxin concentrations were determined using certified standards provided by CNRC (Halifax, NS, Canada). The method was developed and validated internally in the IFREMER Phycotoxins laboratory.

Table S1: LC-MS/MS transitions for the 9 MC variants tested with standards

Toxin	Precursor ion (<i>m/z</i>)	Transition (<i>m/z</i>) - Quantification	Transition (<i>m/z</i>) - identification
MC-LR	995.6	213.2	374.5
MC-LW	1025.6	375.2	135.2
MC-LF	986.6	375.2	135.2

MC-LY	1002.6	375.2	135.2
dmMC-LR	981.4	103.0	135.2
MC-RR	520.1	135.2	213.2
dmMC-RR	512.8	135.0	103.0
MC-LA	910.7	375.2	135.2
MC-YR	1045.6	213.2	375.2

3. Results

3.1. In the water column

3.1.1. Physico-chemical parameters

The physico-chemical characteristics of the water sampled at each site along the continuum are presented in Table 1. While temperature was constant along the continuum, conductivity (as a proxy of salinity) increased from upstream to downstream. Dissolved oxygen exhibited a decreasing concentration associated with a simultaneous decrease in Chl a concentration. Nitrate presented an upstream to downstream decrease while phosphate values showed the opposing trend with higher values downstream. The total suspended solids in the water column exhibited higher concentrations downstream in the estuary and revealed a decreasing gradient in percentage of organic matter (OM) varying from 100% in freshwater to 24% at the marine site.

Table 1 : Physico-chemical parameters measured in the water at the 5 sampling stations

Sites	Temp (°C)	Cond (mS/cm)	O ₂ (mg/L)	Chl a (µg/L)	pH	P-PO ₄ (mg/L)	N-NO ₃ (mg/L)	TSS (mg/L)	OM (%)
F1	14.80	0.24	13.67	22.70	8.02	0.070	1.51	34	100.0
F2	14.52	0.26	10.91	13.54	7.59	0.017	1.17	25	85.3
E1	14.93	12.30	9.61	6.29	7.54	0.044	0.98	121	35.9
E2	14.98	45.35	8.63	0.07	7.78	0.042	0.47	148	30.5
M	14.96	40.21	7.84	1.06	7.77	0.055	0.48	135	24.7

3.1.2. Phytoplankton/Cyanobacteria community

In the water column, the phytoplankton community was largely dominated by cyanobacteria in cells density (Fig. 2). In particular several species of the *Microcystis* genus (*M. aeruginosa*, *M. botrys*, *M. viridis* and *M. flos-aquae*) accounted for most of the cells density at the freshwater sites. *M. aeruginosa* was the only cyanobacterial species observed in the estuary with a concentration of 3.7×10^4 cells mL⁻¹ at E1. Several Chlorophyceae and pennate diatoms were also recorded in the estuary in low numbers ($< 10^3$ cells mL⁻¹).




Fig 2: Species composition and phytoplankton density in the water column at the 5 stations along the freshwater-marine continuum on October 10, 2018

The biovolumes of the phytoplankton groups present together with the percentage of cyanobacteria in the total biovolume of phytoplankton at each site are presented in Table 2, demonstrating the strong dominance of cyanobacteria at the three most upstream sites.

Table 2 : Biovolume of phytoplankton groups at each sampling site

Biovolume (mm ³ L ⁻¹)	F1	F2	E1	E2	M
Cyanobacteria	21.655	4.721	2.434	0.000	0.000
Chlorophyceae	0.079	0.035	0.037	0.000	0.000
Bacillariophyceae	0.061	1.147	0.258	0.008	0.005

Euglenophyceae	0.201	0.060	0.000	0.000	0.000
% Cyanobacteria	0.984	0.792	0.892	0.000	0.000

3.1.3. Intracellular and extracellular microcystins

We observed a gradual decrease of intracellular MC along the continuum from 28 $\mu\text{g/L}$ in the freshwater reservoir, 3 $\mu\text{g L}^{-1}$ in the estuary and down to 0.21 $\mu\text{g L}^{-1}$ at the marine station (Fig. 3A). This gradual decrease was mostly associated with the decrease in cyanobacterial biovolume, in particular that of *M. aeruginosa*. Indeed, based on *M. aeruginosa* densities, the cellular quotas corresponded to 88 fg cell⁻¹ at F1, 69 fg cell⁻¹ at F2 and 80 fg cell⁻¹ at E1. Nine MC variants were recorded with the dominance of MC-LR followed by MC-RR and MC-YR among all sites. No particular selection of variants was observed with increasing salinity.




Fig 3A) Intracellular MC present in the cyanobacterial biomass in the water at the investigated sites

The extracellular MC measured in the filtrate was much smaller (100 to 400 times) than the intracellular fraction, varying between 0.1 and 0.3 $\mu\text{g L}^{-1}$, but increased from upstream to downstream (Fig. 3B). In particular, at the two most downstream sites, the proportion of extracellular/total MC reached 90% and 55% respectively.




Fig. 3B) Extracellular MC present in dissolved form in the water

Similarly to the intracellular fraction, 9 MC variants were recorded in the extracellular fraction and the dominant MC variant was MC-LR. It was followed by MC-YR and MC-RR at low salinities while MC-RR became more abundant at higher salinities.

3.2. In surface sediments

3.2.1. Chemical characteristics

Overall, the surface sediment characteristics were not drastically different along the estuary (Table 3). Surface sediment showed a dominance of inorganic material at all 5 sites with a maximum of 17% of organic matter (OM) at the marine site. They contained a high proportion of silicate (SiO_2) (50 to 73%), aluminum (Al_2O_3) (up to 15%), calcium (CaO) (up to 9.8%), iron (Fe_2O_3) (up to 6.7%), potassium (K_2O) and sodium (Na_2O) (up to 4%), magnesium (MgO) (up to 2.3%) and phosphorus (P_2O_5) (up to 0.3%). The OM was maximal at F2 and M where sediment was fine and muddy while sediment at F1 and E2 was coarser and sandy. The water content of sediment varied between 30 and 71% being higher in muddy sediment. Coarser sandy sediment was also lower in Fe, Mn, Mg and OM.

Table 3 : Surface sediment characteristics along the continuum

Sites	SiO ₂	Al ₂ O ₃	Fe ₂ O ₃	MnO	MgO	CaO	Na ₂ O	K ₂ O	TiO ₂	P ₂ O ₅	H ₂ O	OM
	%	%	%	%	%	%	%	%	%	%	%	%
F1	73.04	12.03	2.43	0.032	0.53	0.59	1.61	4.37	0.40	0.20	48	3.98
F2	53.54	15.17	6.72	0.22	1.76	1.97	0.75	2.63	0.75	0.34	71	15.45
E1	60.23	12.17	4.09	0.056	1.52	4.22	2.79	3.16	0.47	0.15	65	11.09
E2	73.71	2.76	0.92	0.018	0.44	9.82	0.74	0.78	0.12	< L.D.	30	9.92
M	51.32	12.25	4.95	0.11	2.30	5.67	3.74	2.40	0.61	0.20	68	16.49

3.2.2. Cyanobacteria biomass

There was a dominance of *Microcystis* sp. (> 90%) with some akinetes of *Dolichospermum* sp, (Komàrek 2016) and few filaments of *Planktothrix* sp. at F2 in surface sediment. The presence of *Microcystis* cells in surface sediment was recorded at all 5 sites with concentrations varying between 3×10^3 and 3×10^6 cells g⁻¹ dry weight (DW) of sediment (Fig 4).




Fig 4: *Microcystis* biomass in surface sediment at the 5 sampling sites along the continuum.

Photographs from epifluorescence microscopy together with counting suggested that the cells were not lysed (Fig. 5).




Fig 5: Photomicrographs of extracted *Microcystis* from studied sediment (Epifluorescence light): colonies with filaments and diatoms (x100), focus on colony (x200), focus on *Microcystis* intact cells (x400)

3.2.3. Intracellular and extracellular microcystins

Intracellular MC varied between $193 \mu\text{g kg}^{-1}$ wet weight (WW) at F2, $14.4 \mu\text{g kg}^{-1}$ WW of sediment at E1, $0.2 \mu\text{g kg}^{-1}$ WW at E2 and non detectable at site M (Fig. 6A). These intracellular concentrations corresponded respectively to $482 \mu\text{g kg}^{-1}$ DW at F2, $36 \mu\text{g kg}^{-1}$ DW at E1 and $0.5 \mu\text{g kg}^{-1}$ DW at E2. The quantification of intracellular MC in the estuarine sediment indicates that cells of *M. aeruginosa* were not lysed and that sediment preserved at least some cells intact. Different MC variants were recorded in the sediments with the dominance of MC-LR, RR and YR similarly to the proportions observed in the water column.




Fig 6A: Intracellular MC in surface sediment along the continuum

Extracellular MC concentrations were inferred from subtracting the intracellular concentrations from the total MC. Extracellular MC in surface sediment varied also along the estuary between 733 $\mu\text{g kg}^{-1}$ WW at F2 (1832 $\mu\text{g kg}^{-1}$ DW), 70 $\mu\text{g kg}^{-1}$ WW at E1 (175 $\mu\text{g kg}^{-1}$ DW) and non detectable further downstream (Fig. 6B).




Fig 6B: Extracellular MC in surface sediment along the continuum

3.2.4. Genes presence

Table 4: Detection of the targeted genes in studied surface sediments

	Cyanobacterial 16S rRNA gene	<i>mcyA</i>	<i>mcyB</i>	<i>anaC</i>
PCR	Classic	Classic	Nested	Nested
F1	+	+	+	-
F2	+	+	+	+
E1	+	+	+	-
E2	+	+	+	-
M	+	-	+	-

The PCR analyses using both classic and nested approaches confirmed the presence of cyanobacterial species in sediment at all sites. Microcystin genotypes were omnipresent. Indeed, microcystin biosynthesis genes were detected in the sediment of all studied sites except *mcyA* at the most downstream station (Table 4). In the freshwater section at F2, the *anaC* gene was also detected suggesting the presence of species potentially producing anatoxin. This site exhibited the highest cyanobacterial biomass including akinetes of *Dolichospermum sp.* and long filaments of *Planktothrix sp.* (data not shown).

4. Discussion

This is the first study reporting on co-occurrence of both cyanobacteria and MC in estuarine/marine waters and sediments, resulting from a freshwater discharge of an upstream reservoir. Moreover we report separately the quantification of intracellular and extracellular forms of MC.

4.1. In the water column

Cyanobacteria dominated the phytoplankton community in the water column with high densities and biovolumes at the freshwater sites and decreasing concentrations along the salinity gradient. Temperatures during sampling around 15°C at all 5 sites were much lower than the growth optimal temperature of cyanobacteria in general

and of *Microcystis* in particular (Carey et al. 2012) but still within growth potential observed in freshwater lakes and reservoirs (Kruk et al. 2017). Spatial variation of cyanobacterial biomass was most likely due to salinity as nutrients were unlikely to limit cyanobacterial growth (Reynolds 2006). Nitrate showed an upstream to downstream decrease suggesting a freshwater dominant source consistent with high nitrate from agricultural lands in the upstream catchment (Le Moal et al. 2019). Phosphate values showed the opposing trend with higher values downstream in the estuary where we expect sediment resuspension to be the largest, based on the observed high concentrations of total suspended solids and high tidal energy flows. Although the freshwater discharge was not monitored and therefore not quantified, the progressive increase in salinity near the surface was a good indicator of dilution. The short residence time (of the order of one day), inferred from observations of the surface velocities during sampling, would not permit *in situ* growth. Hence we suggest that the lower biomass in the estuary results from freshwater discharge dilution with estuarine water even though we minimized that influence by sampling within one hour of low tides.

The dominant blooming genus in the upstream freshwater reservoir was overwhelmingly *Microcystis*, consistent with the most widespread cyanobacterial occurrence of *Microcystis* in freshwaters worldwide (Harke et al. 2016) and indeed in Brittany (Pitois et al. 2014). While several species of *Microcystis* co-occurred at the two freshwater sites, *M. aeruginosa* was the only species found in the estuary. This observation is in accordance with several other studies worldwide which report on *M. aeruginosa* as the dominant species in estuaries (Preece et al. 2017) in relation with its highest salinity tolerance among cyanobacteria (Verspagen et al. 2006). Conductivities in the estuary and at the marine outlet ranged during sampling from 12 to 45 mS cm⁻¹ corresponding to salinities from 7 to 27, the maximum value beyond reported values for *M. aeruginosa* salinity tolerance (Verspagen et al. 2006) and justifying that no *M. aeruginosa* cells were found in the water column at the two most downstream stations. The mechanisms of salinity tolerance in *M. aeruginosa* are still unclear and current hypotheses include the presence of sucrose genes acting as osmolytes (Tanabe et al. 2018) or the role of mucilage protection against osmotic shock (Kruk et al. 2017).

M. aeruginosa density was high in the estuarine section (at E1) at intermediate salinity reaching 3.7×10^4 cells mL⁻¹ and associated with intracellular MC of 3 µg L⁻¹, resulting in a MC quota of 80 fg cell⁻¹, if we assume that MC was only due to M. aeruginosa cells and that all M. aeruginosa cells were toxic. This quota is similar to that measured in freshwater at F1 and F2, suggesting no particular change with salinity. Very few studies report on MC quotas in natural environments especially in estuaries. Moreover according to Wood et al 2011, MC quotas *in situ* can vary by 20 fold in 5 hours, and potentially are the response to a potential stress.

The toxins profiles in the cyanobacterial biomass (intracellular form) and dissolved in the water (extracellular form) were dominated by MC-LR, RR and YR similarly to those of a previous study when *Microcystis* was the dominant genus at the same sites (Bormans et al. 2019). Both intracellular and extracellular MC exhibited here higher concentrations than previously reported in the literature. The downstream gradient of intracellular MC was consistent with the downstream gradient in *Microcystis* biovolume in the water column while extracellular MC showed a reverse trend and increased at higher salinities suggesting some cells lysing. In accordance with literature results the dominant form of MC in the freshwater part was intracellular (Orr and Jones 1998; Briand et al. 2012), while the dominant form became extracellular as cells lysis increased at elevated salinity (Tonk et al. 2007; Bormans et al. 2019). Both intracellular and extracellular toxins were measured in the water (at E2 and M) while cyanobacterial cells were not counted at those sites. This result confirms previous reports that toxins can persist as extracellular form in the water after the presence of cyanobacteria (Briand et al. 2009). Concerning the presence of intracellular MCs in the water column, cyanobacterial cells may have been too scarce at the marine site to be visualised. Bukaveckas et al. 2018 also reported MC concentrations in the water column at the river estuarine interface of the James River while no cyanobacterial cells were recorded. However their values of total MC were very low (< 0.3 µg L⁻¹) and only measureable in the freshwater section (salinity < 0.5). While Umehara et al. (2015) reported in Isahaya Bay total MC values of 0.06 µg L⁻¹ at salinities approaching 30, we found a much higher concentration with total MC = 0.4 µg L⁻¹, most likely due to the high toxin concentration in the upstream reservoir.

4.2. In surface sediments

Most bottom sediments that accumulate in the estuarine zone consist of terrestrial detritus and biogenic debris. The surface sediments were dominated by inorganic material with a high proportion of silicate, which can have both a riverine and marine origin, if associated with biogenic production by diatoms (Conley 1997; Maavara et al. 2014). The organic matter percentage was maximum at F2 and M where the sediments were fine and muddy while sediments at F1 and E2 were coarse and sandy. Coarser sandy sediments were also lower in Fe, Mn, Mg and OM as is often found in estuaries (Folger 1972).

Quantifying cyanobacterial cells in sediments is difficult as reported by Pham and Utsumi (2018). The presence of *Microcystis* was observed in surface sediments at all 5 sites along the freshwater marine continuum in relatively high densities even at the marine outlet. *Microcystis* abundance was the highest at F2 where *Microcystis* concentration reached 3×10^6 cells g^{-1} DW of sediment corresponding to bloom concentrations. The higher number of cells at F2 compared to F1 is not surprising given that the sediment sampling at F1 was done in the flowing river downstream of the reservoir, while F2 corresponded to a much more stagnant water, associated with muddy sediments rich in OM. Hence F2 acted as a sediment trap accumulating *Microcystis* cells. The presence of *Microcystis* in sediments is consistent with *Microcystis* being the dominant species being transferred in the water column from the upstream reservoir freshwater discharge. Moreover, *Microcystis* concentration gradient in surface sediments along the salinity gradient reflected *Microcystis* concentration gradient in the water column over several months to years (Bormans et al. 2019) rather than instantaneously (this study).

Microcystis cells in the sediments were found intact and in high numbers suggesting a preservation role of sediments even at high salinities. The exact quantification of abundance of colonial cells is always a challenge. The presence of colonies in the sediments indicated that the mucilage was probably preserving cells from lysing (Misson et al 2012; Xiao et al. 2018) and that these cells had been deposited in the sediments during freshwater discharge (Kruk et al. 2017). At the two most downstream sites where no *Microcystis* cells were found in the water column due to high salinities, some *Microcystis* colonies were still intact in the sediments. While the

filtration at 8 μm did not allow the observation of individual *Microcystis* cells, it is very unlikely that cells without the protection of the mucilage would survive degradation and salinity osmotic shock as demonstrated for unicellular *M. aeruginosa* (Georges des Aulnois et al. 2019). Cyanobacterial cells can survive in sediments depending on numerous factors such as river flow, depth, bathymetry, sediment composition or redox conditions (Kravchuk et al. 2011; Legrand et al. 2017; Grutzmacher et al. 2010). Hence, some *Microcystis* cells can be punctually released in the estuary explaining the decreasing gradient from the beginning of the estuary to the marine section. These phenomena of primary and secondary sedimentation have already highlighted in lacustrine ecosystems (Evans 1994; Verspagen et al. 2005; Cirés et al. 2013; Legrand et al. 2017).

Concentrations of both extracellular and intracellular toxins measured in the sediments were relatively high compared to literature values. We measured total MC concentrations of 926 $\mu\text{g/kg}$ WW (or 2315 $\mu\text{g kg}^{-1}$ DW) at the freshwater site F2 and 70 $\mu\text{g kg}^{-1}$ WW (or 175 $\mu\text{g kg}^{-1}$ DW) in the estuary at E1. These values were higher than the value of 3 $\mu\text{g kg}^{-1}$ WW reported by Umehara et al. (2012) in Isahaya Bay, Japan associated with MC concentration of 100 $\mu\text{g kg}^{-1}$ WW in the freshwater reservoir upstream. High values of MC in surface silty sediments, reaching 40 $\mu\text{g kg}^{-1}$ DW, were also reported in Curonian lagoon after a large bloom of *Microcystis* (Paldavieciene et al 2015; Bukaveckas et al 2017). Intracellular MC measured in the sediments were most likely due to *Microcystis* species (*M. aeruginosa* in particular), given its dominance in the sediments. Assuming that *Microcystis* was the only MC producer, these intracellular concentrations corresponded at both F2 and E1 to cellular quotas of 160 fg cell⁻¹. Higher MC quotas in sediments compared with the overlying water column has already been observed in lakes where the authors have suggested that MC may intervene in cells preservation reducing grazing by benthic invertebrate or reducing microbial activities (Misson et al 2012). In surface sediments, extracellular MC concentrations were almost 5 times higher than intracellular concentrations, in contrast with the water column. This result suggests either some cells lysis from individual cells or small colonies and subsequent release in the porous sediments, or active transport of MC from intracellular to extracellular form possibly associated with the signaling of stress conditions. Relatively high

values of extracellular MC also indicated that MC degradation by bacteria or other abiotic factors was not completely occurring. Hence our results indicate that estuarine sediments are preserving both colonial cyanobacteria from cells lysing and probably also MC from degradation. Nevertheless, these results are still preliminary and there is still a need to investigate the duration of the toxins presence in sediments after the bloom period as well as their presence deeper in the sediments. Our results are in line with reports from Curonian lagoon where relatively high sediment MC concentrations have been measured many years after a large *Microcystis* bloom (Paldavieciene et al 2015). These results have important consequences for contamination of aquatic food webs as reported for zebra mussels by Paldavieciene et al (2015) and planktivorous fish by Bukaveckas et al (2017). In lacustrine systems, MC can persist in sediments several years (Misson et al. 2011) probably due to of the preservation conditions found in these environments (no light, low temperature, anoxic condition). Biodegradation in sediments of MC-LR was indeed found more rapid in aerobic than in anaerobic conditions (Grutzmacher et al 2010). Interestingly, Misson et al. (2012) have highlighted that benthic *Microcystis* from 6-years sediments were able to initiate MC production implying that MC could be involved in the benthic survival mechanisms of *Microcystis*. Here, it is evident that salinity may affect significantly intra or extracellular MC in terms of degradation or inducing *Microcystis* metabolism stimuli but no studies has examined these factors.

Generally, we observed a decrease of MC concentrations along the salinity gradient. A gradient of MC concentration in surface sediments was also observed in Isahaya Bay, Japan (Umehara et al. 2012; 2017) related to concentrations in the water column with maximum values during bloom periods. Temporal variability in MC concentrations in sediment is expected as demonstrated by Umehara et al. (2012; 2015) where the dynamics is governed by variations in the discharge from the reservoir, as well as sedimentation and resuspension processes associated with tidal flows. However, while we did not investigate the temporal variability, we demonstrated unambiguously the presence of *Microcystis* colonies along the entire freshwater-marine continuum, at the time of a toxic bloom in the upstream reservoir. Moreover, at F2 all toxic genotypes and a high MC concentration were detected in the sediment. The same applies at E1 and E2 except with a smaller MC concentration. Only at the most downstream site (marine outlet) did we not detect

any MC in the sediment in accordance with the genes analysis that suggested the presence of a very small number of toxic genotypes at the time of sampling. This may confirm the previous hypothesis, about the importance of F2 station to transfer *Microcystis* in the estuary. Our results are in line with the findings of Via-Ordorika et al. (2004) who reported that most *M. aeruginosa* colonies from several freshwater lakes contained the *mcyA* and *mcyB* genes and found a positive relationship between genes frequency and colony size. Our results also support the measure of MC production genes as a proxy of toxic genotypes as tested by Martinez et al. (2017) and Otten et al. (2015) to evaluate the toxic potential of cyanobacteria along a freshwater marine continuum.

Interestingly, *anaC* was also detected at F2 revealing the presence of potentially anatoxin-a producer such as *Planktothrix* sp. (Viaggiu et al. 2004) or *Dolichospermum* sp. (Delvin et al. 1977). These results suggest a potential co-occurrence between MC and anatoxin-a which has already observed in several lakes (Graham et al. (2006); Sabart et al. (2015) and the co-occurrence of several cyanotoxins in coastal areas (Tatters et al. 2017). This potential co-occurrence may link to a high number of toxigenic species present here.

5. Conclusions

The results of this study confirm the transfer of cyanobacteria and MC from a freshwater reservoir discharge, through an estuary and up to the marine outlet together with the co-occurrence of both cyanobacterial cells and MC in the estuary's sediments. Similarly to studies in freshwater lakes both *Microcystis* cells and MC were found in sediments even when no cells or MC was recorded in the water column at the same site, indicating that sediments integrate over time as well as preserve cells from dilution and degradation. Hence the sediments act as a reservoir of cyanobacteria and MC. The majority of cyanobacterial cells in sediments were found in colonial form suggesting that the mucilage plays a crucial role in maintaining colonies in sediments. This study highlights the survival of *Microcystis* cells in the water column at salinities in the range 5 to 10, and in surface sediments up to a salinity of 27. MC concentrations measured here in estuarine sediments were much higher than reported elsewhere in the literature. This is the first study reporting on the

presence of *mcy* genes in estuarine sediments, indicating the toxicity potential of estuarine sediments for benthic organisms and also as potential source to be resuspended in the water column.

Acknowledgments

The authors thank Maxime Georges des Aulnois for help with field work, Marion Chorin for nutrient analyses and Gorenka Bojadzija Savic for finalisation of the figures. Microscopy and biochemical analyses were performed at the Experimental Ecology (ECOLEX) and the Analytical (PLAY) platforms of the UMR Ecobio. The physicochemical characteristics of the sediments were analysed by the SARM laboratory of the CRPG-CNRS in Nancy, France. This project was financially supported by the ANSES (Agence Nationale de Sécurité Sanitaire de l'Alimentation, de l'Environnement et du Travail) under the Bieautox project EST-2015-191. The helpful suggestions from the reviewers are also acknowledged.

References

- Anon (2005) Anonyme ISO 20179:2005(E). Water quality – Determination of microcystins – Method using solid phase extraction (SPE) and high performance liquid chromatography (HPLC) with ultraviolet (UV) detection. International Standard ISO 20179, first edition 2005-10-01, 17 p
- Bormans M, Amzil Z, Mineaud E, Briant L, Savar V, Robert E, Lance E (2019) Demonstrated transfer of cyanobacteria and cyanotoxins along a freshwater-marine continuum in France. *Harmful Algae* 87: 101639
- Bourrelly P (1985) Les algues d'eau douce: les algues bleues et rouges (Vol. 3). Lubrecht & Cramer Ltd.
- Briand E, Escoffier N, Straub C, Sabart M, Quiblier C (2009) Spatiotemporal changes in the genetic diversity of a bloom-forming *Microcystis aeruginosa* (cyanobacteria) population. *ISME J* 3: 419–429
- Briand E, Bormans M, Quiblier C, Salençon MJ, Humbert JF (2012) Evidence of the cost of the production of microcystins by *Microcystis aeruginosa* under differing light and nitrate environmental conditions. *PLoS One* 7 (1): e29981
- Briant L, Lengronne M, Bertrand E, Rolland D, Sipel A, Steinmann D, Baudin I, Legeas M, Le Rouzic N, Bormans M (2008) A phycocyanin probe as a tool for monitoring cyanobacteria in freshwater bodies. *J Environ Monit* 10: 248–55

- Brient L, Lengronne M, Bormans M, Fastner J (2009) First occurrence of cylindrospermopsin in freshwater in France. *Environmental Toxicology* 24(4): 415-420
- Bukaveckas PA, Lesutiene J, Gaisunaite ZR, Lozys L, Olenina I, Pilkaityte R, Putys Z, Tassone S, Wood J (2017). Microcystin in aquatic food webs of the Baltic and Chesapeake Bay regions. *Estuarine, Coastal and Shelf Science* 191 :50-59
- Bukaveckas PA, Franklin R, Tassone S, Trache B, Egerton T (2018). Cyanobacteria and cyanotoxins at the river-estuarine transition. *Harmful Algae* 76: 11-21
- Carey CC, Ibelings BW, Hoffmann EP, Hamilton DP, Brookes JD (2012) Ecophysiological adaptations that favour freshwater cyanobacteria in a changing climate. *Water Res* 46:1394–1407
- Cires S, Wörmer L, Agha R, Quesada A (2013) Overwintering populations of *Anabaena*, *Aphanizomenon* and *Microcystis* as potential inocula for summer blooms. *J Plankton Res* 35(6): 1254–1266
- Conley DJ (1997) Contribution of biogenic silica to the oceanic silica budget. *Limnol Oceanogr* 42(4): 774–777
- De Pace R, Vita V, Bucci MS, Gallo P, Bruno M (2014) Microcystin contamination in sea mussel farms from the Italian southern Adriatic coast following cyanobacterial blooms in the artificial reservoir. *J. of Ecosystems* ID 374027
<http://dx.doi.org/10.1155/2014/374027>
- Devlin JP, Edwards OE, Gorham PR, Hunter NR, Pike RK, Stavric B (1977) Anatoxin-a, a toxic alkaloid from *Anabaena flos-aquae* NRC-44h. *Canadian Journal of Chemistry* 55(8): 1367-1371
- Dörr FA, Pinto E, Soares RM, Azevedo SMFO (2010) Microcystins in South American aquatic ecosystems: occurrence, toxicity and toxicological assays. *Toxicon* 56 : 1247–1256
- Evans RD (1994) Empirical evidence of the importance of sediment resuspension in lakes. *Hydrobiologia* 284:5–12
- Folger D (1972) Characteristics of estuarine sediments of the United States USGS report 2482
- Fraisse S, Bormans M, Lagadeuc Y (2013) Morphofunctional traits reflect differences in phytoplankton community between rivers of contrasted flow regime. *Aquat Ecol* 47: 315–327 doi:10.1007/s10452-013-9446-z
- Georges des Aulnois M, Roux P, Caruana A, Réveillon D, Briand E, Hervé F, Savar V, Bormans M, Amzil Z (2019) Physiological and metabolic responses of freshwater and brackish waters strains of *Microcystis aeruginosa* acclimated to a salinity gradient: insight into salt tolerance. *Applied and Env Microbiology*
DOI: 10.1128/AEM.01614-19

- Gibble CM, Peacock MB, Kudela RM (2016) Evidence of freshwater algal toxins in marine shellfish: implications for human and aquatic health. *Harmful Algae* 59: 59–66
- Graham JL, Loftin KA, Meyer MT, Ziegler AC (2006) Co-occurrence of Toxins and Taste-and-Odor Compounds in Cyanobacterial Blooms from the Midwestern United States. *Environ Sci Technol* 44: 7361–7368
- Grutzmacher G, Wessel G, Klitzke S, Chorus I (2010) Microcystin elimination during sediment contact. *Environ Sci Technol* 44: 657–662
- Harke MJ, Steffen MM, Gobler CJ, Otten TG, Wilhelm SW, Wood SA, Paerl HW (2016) A review of the global ecology, genomics, and biogeography of the toxic cyanobacterium *Microcystis* spp. *Harmful Algae* 54: 4–20
- Henriksen A, Selmer-Olsen AR (1970) Automatic methods for determining nitrate and nitrite in water and soil extracts. *Analyst* 95: 514–518
- Komárek J, Anagnostidis K (2005) Süßwasserflora von Mitteleuropa, bd. 19/2: Cyanoprokaryota: Oscillatoriales (Vol. 19). Spektrum Akademischer Verlag.
- Komárek J, Anagnostidis K (2013) Süßwasserflora von Mitteleuropa, bd. 19/2: Cyanoprokaryota: Heterocytous Genera (Vol. 19/3). Spektrum Akademischer Verlag.
- Komárek J (2016) Review of the cyanobacterial genera implying planktic species after recent taxonomic revisions according to polyphasic methods: state as of 2014. *Hydrobiologia* 764(1) : 259–270
- Kravchuk ES, Ivanova EA, Gladyshev MI (2011) Spatial distribution of resting stages (akinetes) of the cyanobacteria *Anabaena flos-aquae* in sediments and its influence on pelagic populations. *Mar Fresh Res* 62: 450–461
- Kruk C, Segura AM, Nogueira L, Alcántara I, Calliaria D, Martínez de la Escalera G, Carballoa C, Cabreraa C, Sarthoua F, Scavone P, Piccini C (2017) A multilevel trait-based approach to the ecological performance of *Microcystis aeruginosa* complex from headwaters to the ocean. *Harmful Algae* 70: 23–36
- Labry C, Delmas D, Youenou A, Quere J, Leynaert A, Fraisse S, Raimonet M, Ragueneau O (2016) High alkaline phosphatase activity in phosphate replete waters: The case of two macrotidal estuaries. *Limnol Oceanogr* 61:1513–1529
- Lance E, Brient L, Carpentier A, Acou A, Marion L, Bormans M, Gérard C (2010) Impact of toxic cyanobacteria on gastropods and microcystin accumulation in a eutrophic lake (Grand-Lieu, France) with special reference to *Physa* (= *Physella*) *acuta*. *Science of the Total Environment* 408: 3560–3568
- Legrand B, Lamarque A, Sabart M, Deltour D (2016) Characterization of akinetes from cyanobacterial strains and lake sediment: A study of their resistance and toxic potential. *Harmful Algae* 59: 42–50

- Legrand B, Le Jeune AH, Colombet J, Thouvenot A, Latour D (2017) Akinetes May Be Representative of Past Nostocalean Blooms: a Case Study of Their Benthic Spatiotemporal Distribution and Potential for Germination in a Eutrophic Lake. *Appl Environ Microbiol* 83:e01571-17
- Legrand B, Miras Y, Beauger A, Dussauze M, Latour D (2019) Akinetes and ancient DNA reveal toxic cyanobacterial recurrences and their potential for resurrection in a 6700-year-old core from a eutrophic lake. *Science of the Total Environment* 687: 1369-1380
- Lehman PW, Boyer G, Hall C, Waller S, Gehrts K (2005) Distribution and toxicity of a new colonial *Microcystis aeruginosa* bloom in the San Francisco Bay Estuary, California. *Hydrobiologia* 541: 87–99
- Le Moal M, Gascuel-Oudou C, Menesquen A, Souchon Y, Etrillard C, Levain A, Moatar F, Pannard A, Souchu P, Lefebvre A, Pinay G (2019) Eutrophication: A new wine in an old bottle? *Science of the Total Environment* DOI:[10.1016/j.scitotenv.2018.09.139](https://doi.org/10.1016/j.scitotenv.2018.09.139)
- Maavara T, Dürr HH, Van Cappellen P (2014) Worldwide retention of nutrient silicon by river damming: From sparse data set to global estimate. *Global Biogeochem Cycles* 28 : 842–855 doi:10.1002/2014GB004875
- Martínez de la Escalera G, Kruk C, Segurad AM, Nogueira L, Alcántara I, Piccini C (2017) Dynamics of toxic genotypes of *Microcystis aeruginosa* complex (MAC) through a wide freshwater to marine environmental gradient. *Harmful Algae* 62 :73–83
- Merel S, Walker D, Chicana R, Snyder S, Baurés E, Thomas O (2013) State of knowledge and concerns on cyanobacterial blooms and cyanotoxins. *Environ Int* 59: 303–327
- Meriluoto J, Blaha L, Bojadzija G, Bormans M, Brient L, Codd GA, Drobac D, Faassen EJ, Fastner J, Hiskia A, Ibelings BW, Kaloudis T, Kokocinski M, Kurmayer R, Pantelić D, Quesada A, Salmaso N, Tokodi N, Triantis TM, Visser PM, Svirčev Z (2017) Toxic cyanobacteria and cyanotoxins in European waters - recent progress achieved through the CYANOCOST Action and challenges for further research. *Advances in Oceanography and Limnology* 8(1) DOI: 10.4081/aiol.2017.6429
- Misson B, Sabart M, Amblard C, Latour D (2011) Involvement of microcystins and colony size in the benthic recruitment of the cyanobacterium *Microcystis* (Cyanophyceae). *Journal of Phycology* 47:42-51
- Misson B, Sabart M, Amblard C, Latour D (2012) Benthic survival of *Microcystis*: long-term viability and ability to transcribe microcystin genes. *Harmful Algae* 13: 20-25.
- Ndlela LL, Oberholster PJ, Van Wyk JH, Cheng PH (2016) An overview of cyanobacterial bloom occurrences and research in Africa over the last decade. *Harmful Algae* 60 : 11–26

- Nubel U, Garcia-Pichel F, Muyzer G (1997) PCR primers to amplify 16SrRNA genes from cyanobacteria. *Appl Environ Microbiol* 63: 3327–3332
- Omidi A, Esterhuizen-Londt M, Pflugmacher S (2017) Still challenging: the ecological function of the cyanobacterial toxin microcystin – What we know so far. *Toxin Reviews* 37: Issue 2.
- O’Neil JM, Davis TW, Burford MA, Gobler CJ (2012) The rise of harmful cyanobacteria blooms: The potential roles of eutrophication and climate change *Harmful Algae* 14 : 313–334
- Orr PT, Jones GJ (1998) Relationship between microcystin production and cell division rates in nitrogen-limited *Microcystis aeruginosa* cultures. *Limnol Oceanogr* 43: 1604–1614
- Orr PT, Jones GJ, Douglas GB (2004) Response of cultured *Microcystis aeruginosa* from the Swan River, Australia, to elevated salt concentration and consequences for bloom and toxin management in estuaries. *Mar Freshw Res* 55 (3) : 277– 283
- Ortiz et al., (2017) A high throughput targeted and non-targeted method for the analysis of microcystins and anatoxin-A using on-line solid phase extraction coupled to liquid chromatography–quadrupole time-of-flight high resolution mass spectrometry. *Analytical and Bioanalytical Chemistry* 409 (21): 4959
- Otten TG, Crosswell JR, Mackey S, Dreher TW (2015) Application of molecular tools for microbial source tracking and public health risk assessment of a *Microcystis* bloom traversing 300 km of the Klamath River. *Harmful Algae* 46 : 71-81
- Paerl HW (2018) Mitigating Toxic Planktonic Cyanobacterial Blooms in Aquatic Ecosystems Facing Increasing Anthropogenic and Climatic Pressures. *Toxins* 10: 76 doi:10.3390/toxins10020076
- Paerl HW, Otten TG, Kudela R (2018) Mitigating the Expansion of Harmful Algal Blooms Across the Freshwater-to-Marine Continuum. *Environ Sci Technol* 52 : 5519–5529
- Paldavičiene A, Mazur-Marzec H, Razinokovas A (2009) Toxic cyanobacteria blooms in the Lithuanian part of the Curonian Lagoon. *Oceanologia* 51 (2) : 203– 216
- Paldavičiene A, Zaiko A, Mazur-Marzec H, Razinokovas-Baziukas A (2015) Bioaccumulation of microcystins in invasive bivalves: a case study from the boreal lagoon ecosystem. *Oceanologia* 57: 93– 101
- Pannard A, Guislain A, Chorin M, Mahé S, Bouger G, Crave A, Le Rouzic B, Bormans M (2018) Phosphorus more than temperature controls the phytoplankton community in a deep quarry lake: a combined field and laboratory approach. *Inland Waters*, DOI:10.1080/20442041.2018.1424097

- Peacock MB, Gobble CM, Senn DB, Cloern JE, Kudela RM (2018) Blurred lines: Multiple freshwater and marine algal toxins at the land-sea interface of San Francisco Bay, California. *Harmful Algae* 73 : 138-147
- Pham TL, Utsumi M (2018) An overview of the accumulation of microcystins in aquatic ecosystems. *Journal of Environmental Management* 213: 520-529
- Pitois F, Thoraval I, Baurès E, Thomas O (2014) Geographical Patterns in Cyanobacteria Distribution: Climate Influence at Regional Scale. *Toxins* 6: 509-522
- Pitois F, Fastner J, Pagotto C, Dechesne M (2018) Multi-Toxin Occurrences in Ten French Water Resource Reservoirs. *Toxins* 10: 283
- Preece EP, Moore BC, Hardy FJ (2015) Ecotoxicology and Environmental Safety Transfer of microcystin from freshwater lakes to Puget Sound, WA and toxin accumulation in marine mussels (*Mytilus trossulus*). *Ecotoxicology and Environmental Safety* 122: 98–105
- Preece EP, Hardy FJ, Moore BC, Bryan M (2017) A review of microcystin detections in Estuarine and Marine waters: Environmental implications and human health risk. *Harmful Algae* 61 : 31–45
- Puddick J, Prinsep MR, Wood SA, Kaufononga SAF, Cary SC, Hamilton DP (2014) High Levels of Structural Diversity Observed in Microcystins from *Microcystis* CAWBG11 and Characterization of Six New Microcystin Congeners. *Mar Drugs* 12 : 5372-5395; doi:10.3390/md12115372
- Reynolds CS (2006) Ecology of Phytoplankton. Cambridge University Press, Cambridge, UK
- Rigosi A, Carey CC, Ibelings BW, Brookes JD (2014) The interaction between climate warming and eutrophication to promote cyanobacteria is dependent on trophic state and varies among taxa. *Limnol Oceanogr* 59(1): 99–114
- Robson BJ, Hamilton DP (2003) Summer flow event induces a cyanobacterial bloom in a seasonal western Australia estuary. *Mar Freshw Res* 54: 139–151
- Sabart M, Crenn K, Perrière F, Abila A, Lereboure M, Colombet J, ... Latour D (2015) Co-occurrence of microcystin and anatoxin-a in the freshwater lake Aydat (France): Analytical and molecular approaches during a three-year survey. *Harmful Algae* 48: 12-20
- Taş S, Okus E, Aslan-Yilmaz A (2006) The blooms of a cyanobacterium, *Microcystis cf. aeruginosa* in a severely polluted estuary, the Golden Horn, Turkey. *Estuar Coast. Shelf Sci* 68, : 593–599
- Tatters AO, Howard MDA, Nagoda C, Busse L, Gellene AG, Caron DA (2017) Multiple stressors at the land-sea interface: cyanotoxins at the land-sea interface in the Southern California Bight. *Toxins* doi:http://dx.doi.org/10.3390/toxins9030095.

- Tillett D, Dittmann E, Erhard M, von Döhren H, Börner T, Neilan BA (2000) Structural organization of microcystin biosynthesis in *Microcystis aeruginosa* PCC7806: an integrated peptide polyketide synthetase system. *Chemistry & Biology* 7:753-764
- Tonk L, Bosch K, Visser PM, Huisman J (2007) Salt tolerance of the harmful cyanobacterium *Microcystis aeruginosa*. *Aquat Microb Ecol* 46: 117–123
- Umehara A, Tsutsumi H, Takahashi T (2012) Blooming of *Microcystis aeruginosa* in the reservoir of the reclaimed land and discharge of microcystins to Isahaya Bay (Japan). *Environ Sci Pollut Res* 19: 3257–3267
- Umehara A, Komorita T, Tai A, Takahashi T, Orita R, Tsutsumi H (2015) Short-term dynamics of cyanobacterial toxins (microcystins) following a discharge from a coastal reservoir in Isahaya Bay, Japan. *Marine Pollution Bulletin* 92: 73–79.
- Umehara A, Takahashi T, Komorita T, Orita R, Choi JW, Takenaka R, Mabuchi R, Park HD, Tsutsumi H (2017) Widespread dispersal and bio-accumulation of toxic microcystins in benthic marine ecosystems. *Chemosphere* 167 : 492-500
- Verspagen JMH, Snelder EOFM, Visser PM, Jöhnk KD, Ibelings BW, Mur LR, Huisman J (2005) Benthic-pelagic coupling in the population dynamics of the harmful cyanobacterium *Microcystis*. *Freshw Biol* 50: 854–867
<https://doi.org/10.1111/j.1365-2427.2005.01368.x>.
- Verspagen JMH, Passarge J, Jöhnk KD, Visser PM, Peperzak L, Boers P, Laanbroek HJ, Huisman J (2006) Water management strategies against toxic *Microcystis* blooms in the Dutch delta. *Ecol Appl* 16 : 313–327
- Vezie C, Briant L, Sivonen K, Bertru G, Lefeuvre JC, Salkinoja-Salonen M (1998) Variation of microcystin content of cyanobacterial blooms and isolated strains in Lake Grand-Lieu (France). *Microb Ecol* 35:126–35
- Viaggiu E, Melchiorre S, Volpi F, Di Corcia A, Mancini R, Garibaldi L, Bruno M (2004) Anatoxin-a toxin in the cyanobacterium *Planktothrix rubescens* from a fishing pond in northern Italy. *Environmental Toxicology: An International Journal* 19(3): 191-197
- Via-Ordorika L, Fastner J, Kurmayer R, Hisbergues M, Dittmann E, Komarek J, Erhard M, Chorus I (2004) Distribution of Microcystin-Producing and Non-Microcystin-Producing *Microcystis* sp. in European Freshwater Bodies: Detection of Microcystins and Microcystin Genes in Individual Colonies. *Systematic and Applied Microbiology* 27(5): 592-602
- Wood R (2016) Acute animal and human poisonings from cyanotoxin exposure — A review of the literature. *Environment International* 91: 276-282
- Wood SA, Rueckert A, Hamilton DP, Cary SC, Dietrich DR (2011). Switching toxin production on and off : intermittent microcystin synthesis in a *Microcystis* bloom. *Env. Microbiol. Report* 3(1) : 118-124

979 Xiao M, Li M, Reynolds CS (2018) Colony formation in the cyanobacterium
980 Microcystis. Biol Rev 93: 1399-1420
981

Accepted manuscript