



**HAL**  
open science

## ICTV Virus Taxonomy Profile: Alphaflexiviridae

Jan Kreuze, Anna Maria Vaira, Wulf Menzel, Thierry T. Candresse, Sergey K  
Zavriev, John Hammond, Ki Hyun Ryu, Ictv Report Consortium

► **To cite this version:**

Jan Kreuze, Anna Maria Vaira, Wulf Menzel, Thierry T. Candresse, Sergey K Zavriev, et al.. ICTV  
Virus Taxonomy Profile: Alphaflexiviridae. *Journal of General Virology*, 2020, 101 (7), pp.699-700.  
10.1099/jgv.0.001436 . hal-02881958

**HAL Id: hal-02881958**

**<https://hal.inrae.fr/hal-02881958>**

Submitted on 26 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



## ICTV Virus Taxonomy Profile: *Alphaflexiviridae*

Jan F. Kreuze<sup>1,\*</sup>, Anna Maria Vaira<sup>2</sup>, Wulf Menzel<sup>3</sup>, Thierry Candresse<sup>4</sup>, Sergey K. Zavriev<sup>5</sup>, John Hammond<sup>6</sup>, Ki Hyun Ryu<sup>7</sup> and ICTV Report Consortium

### Abstract

The family *Alphaflexiviridae* includes viruses with flexuous filamentous virions that are 470–800 nm in length and 12–13 nm in diameter. Alphaflexiviruses have a single-stranded, positive-sense RNA genome of 5.5–9 kb. They infect plants and plant-infecting fungi. They share a distinct lineage of alphavirus-like replication proteins that is unusual in lacking any recognized protease domain. With a single exception, cell-to-cell and long-distance movement is facilitated by triple gene block proteins in plant-infecting genera. This is a summary of the International Committee on Taxonomy of Viruses (ICTV) Report on the family *Alphaflexiviridae*, which is available at [www.ictv.global/report/alphaflexiviridae](http://www.ictv.global/report/alphaflexiviridae).

**Table 1.** Characteristics of members of the family *Alphaflexiviridae*

<b>Typical member:</b>	<b>shallot virus X, Kanyuka (M97264), species <i>Shallot virus X</i>, genus <i>Allexivirus</i></b>
Virion	Flexuous filaments, usually 12–13 nm (range 10–15 nm) in diameter and from 470 to 800 nm in length
Genome	Single molecule of linear, single-stranded, positive-sense RNA of 5.5–9.0 kb
Replication	Cytoplasmic, virus-encoded RNA-directed RNA polymerase
Translation	From capped and polyadenylated genome length and 3'-terminal subgenomic mRNAs
Host range	Plants and fungi
Taxonomy	Realm <i>Riboviria</i> , kingdom <i>Orthornavirae</i> , phylum <i>Kitrinoviricota</i> , class <i>Alsuviricetes</i> , order <i>Tymovirales</i> ; several genera with over 50 species

### VIRION

Virions are flexuous filaments, usually 12–13 nm in diameter (range 10–15 nm) and from 470 to about 800 nm in length, depending on the genus. The viral capsid is composed of a single polypeptide ranging in size from 18 to 43 kDa except for members of the genus *Lolavirus*, which have two carboxy-terminal capsid protein variants, and members of the genus *Sclerodarnavirus* in which no capsid protein has been identified [1, 2] (Fig. 1, Table 1).

### GENOME

Virions contain a single molecule of linear, positive-sense RNA of 5.5–9.0 kb, which is 5–6% by weight of the virion. The RNA is typically capped at the 5'-terminus with m<sup>7</sup>G and has a polyadenylated tract at the 3'-terminus. Smaller 3'-co-terminal sgRNAs are encapsidated in some, but not all, members of the genus *Potexvirus*. There are five to seven genes depending upon the genus, except for members of the genus *Sclerodarnavirus*, which have a single gene. The typical member shallot virus X has six genes (Fig. 2).

Received 15 April 2020; Accepted 29 April 2020; Published 11 June 2020


**Author affiliations:** <sup>1</sup>International Potato Center (CIP), Apartado 1558, Lima 12, Peru; <sup>2</sup>Institute for Sustainable Plant Protection – CNR – 73, Strada delle Cacce, 10135 Torino, Italy; <sup>3</sup>Leibniz-Institute DSMZ, Inhoffenstraße 7 B, 38124 Braunschweig, Germany; <sup>4</sup>INRA, Univ. Bordeaux, UMR BFP, CS20032, 33882 Villenave d'Ornon cedex, France; <sup>5</sup>Institute of Bioorganic Chemistry of the Russian Academy of Sciences, 117997 Moscow, Russia; <sup>6</sup>USDA-ARS, USDA, 10300 Baltimore Avenue, Beltsville MD 20705, USA; <sup>7</sup>Seoul Women's University, Seoul, Republic of Korea.

\*Correspondence: Jan F. Kreuze, [j.kreuze@cgiar.org](mailto:j.kreuze@cgiar.org)

**Keywords:** *Alphaflexiviridae*; ICTV; taxonomy.

**Abbreviations:** ICTV, International Committee on Taxonomy of Viruses.  
001436 © 2020





**Fig. 1.** Electron micrograph of negatively-stained virions of an isolate of shallot virus X. Bar=200 nm.

## REPLICATION

The protein encoded by ORF1 (or ORF2 in members of the genus *Platypuvirus*) has homologies with replication-associated proteins of the ‘alphavirus-like’ supergroup of RNA viruses [1]. This protein (150–195 kDa) contains conserved methyltransferase, helicase and RNA-directed RNA polymerase motifs [3]; some also include an AlkB domain (alkylated RNA repair protein) [4]. In all plant-infecting members of the family, with the exception of members of the genus *Platypuvirus*, ORFs 2–4 encode the ‘triple gene block’ proteins involved in cell-to-cell movement [5], and ORF5 encodes the viral capsid protein. Some genera have 1–3 additional ORFs downstream of the capsid protein ORF. ORFs downstream of that encoding the replication-associated proteins are translated from 3′-terminal sGRNAs. Replication is cytoplasmic.

## TAXONOMY

The family *Alphaflexiviridae* is assigned to the order *Tymovirales*. The genera *Allxivirus*, *Botrexvirus*, *Lolavirus*, *Mandari-virus*, *Platypuvirus*, *Potexvirus* and *Sclerodarnavirus* include more than 50 species. Alphaflexiviruses infect a wide range of mono- and dicotyledonous plant species. Two members of the family have been discovered that infect the plant pathogenic fungi *Botrytis cinerea* and *Sclerotinia sclerotiorum*, respectively.

## RESOURCES


Current ICTV Report on the family *Alphaflexiviridae*: [www.ictv.global/report/alphaflexiviridae](http://www.ictv.global/report/alphaflexiviridae)

### Funding information

Production of this summary, the online chapter, and associated resources was funded by a grant from the Wellcome Trust (WT108418AIA).

### Acknowledgements

Members of the ICTV Report Consortium are Stuart G. Siddell, Andrew J. Davison, Elliot J. Lefkowitz, Peter Simmonds, Sead Sabanadzovic, Donald B. Smith, Richard J. Orton and F. Murilo Zerbini.



**Fig. 2.** Diagram showing genome organization of members of genera in the family *Alphaflexiviridae*. Blocks represent predicted ORFs. The replicase proteins are shown in blue, triple gene block proteins in yellow, 3A-like movement protein in orange, capsid proteins in red and RNA-binding proteins in purple. Other ORFs are white. The methyltransferase (M), AlkB (A), helicase (H) and RNA-directed RNA polymerase (R) domains of the replicase are also shown. Brackets indicate ORFs may be missing from some members of the genus and a dashed ORF outline indicates a missing start codon.

### Conflicts of interest

The authors declare that there are no conflicts of interest.

### References

- Martelli GP, Adams MJ, Kreuze JF, Dolja VV. Family *Flexiviridae*: a case study in virion and genome plasticity. *Annu Rev Phytopathol* 2007;45:73–100.
- Vaira AM, Kreuze J, Hammond J, Pearson MN, Martelli GP et al. *Alphaflexiviridae*. In: King AMQ, Lefkowitz EJ, Adams MJ, Carstens EB (editors). *Virus Taxonomy, Ninth Report of the International Committee on Taxonomy of Viruses*. Oxford: Elsevier; 2011. pp. 904–919.
- Batten JS, Yoshinari S, Hemenway C. Potato virus X: a model system for virus replication, movement and gene expression. *Mol Plant Pathol* 2003;4:125–131.
- van den Born E, Omelchenko MV, Bekkelund A, Leihne V, Koonin EV et al. Viral AlkB proteins repair RNA damage by oxidative demethylation. *Nucleic Acids Res* 2008;36:5451–5461.
- Verchot-Lubicz J, Torrance L, Solovyev AG, Morozov SY, Jackson AO et al. Varied movement strategies employed by triple gene block-encoding viruses. *Mol Plant Microbe Interact* 2010;23:1231–1247.