

Geographical Distribution of Ljungan Virus in Small Mammals in Europe

Cristina Fevola, Chiara Rossi, Fausta Rosso, Matteo Girardi, Roberto Rosà, Mattia Manica, Luca Delucchi, Duccio Rocchini, Carol Garzon-Lopez, Daniele Arnoldi, et al.

▶ **To cite this version:**

Cristina Fevola, Chiara Rossi, Fausta Rosso, Matteo Girardi, Roberto Rosà, et al.. Geographical Distribution of Ljungan Virus in Small Mammals in Europe. Vector-Borne and Zoonotic Diseases, 2020, 20 (9), pp.692-702. 10.1089/vbz.2019.2542 . hal-02884889

HAL Id: hal-02884889
https://hal.inrae.fr/hal-02884889

Submitted on 13 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

This document is the accepted manuscript version of the following article:
Branco, S., Faccoli, M., Brockerhoff, E. G., Roux, G., Jactel, H., Desneux, N., … Branco, M. (2021). Preventing invasions of Asian longhorn beetle and citrus longhorn beetle: are we on the right track? Journal of Pest Science. https://doi.org/10.1007/s10340-021-01431-x

Title

Preventing invasions of Asian longhorn beetle and citrus longhorn beetle: Are we on the right track?

Authors

Sofia Branco, Massimo Faccoli, Eckehard G. Brockerhoff, Géraldine Roux, Hervé Jactel, Nicolas Desneux, Emmanuel Gachet, Raphaelle Mouttet, Jean-Claude Streito, Manuela Branco

Author information

Sofia Branco: Centro de Estudos Florestais, Instituto Superior de Agronomia, Universidade de Lisboa, Tapada da Ajuda, 1349-017, Lisbon, Portugal, ORCID 0000-0002-2867-4950, sofiabranco@isa.ulisboa.pt (corresponding author)

Massimo Faccoli: Department of Agronomy, Food, Natural resources, Animals and Environment (DAFNAE), University of Padova, Viale dell'Università 16, 35020 Legnaro (PD), Italy, ORCID 0000-0002-9355-0516

Eckehard G. Brockerhoff: Swiss Federal Research Institute WSL, Zürcherstrasse 111, 8903 Birmensdorf, Switzerland, ORCID 0000-0002-5962-3208

Géraldine Roux: INRAE, INRAE, UR633 Zoologie Forestière, F-45075 Orléans, France, COST, Université d'Orléans, France

Hervé Jactel: INRAE, University of Bordeaux, BIOGECO, 33610, Cestas, France, ORCID 0000-0002-8106-5310

Nicolas Desneux: INRAE, Université Côte d'Azur, CNRS, UMR 1355-7254, Institute Sophia Agrobiotech, 06903 Sophia-Antipolis, France, ORCID 0000-0002-8171-3154

Emmanuel Gachet: ANSES, Laboratoire de la santé des végétaux, 7 rue Jean Dixméras, 49044 Angers, France

Raphaelle Mouttet: ANSES - Laboratoire de la santé des végétaux, CBGP, 755 avenue du Campus Agropolis, 34988 Montferrier sur Lez, France

Jean-Claude Streito: INRAE, CIRAD, IRD, Montpellier SupAgro, University of Montpellier, CBGP, 755 avenue du Campus Agropolis, 34988 Montferrier sur Lez, France

Manuela Branco: Centro de Estudos Florestais, Instituto Superior de Agronomia, Universidade de Lisboa, Tapada da Ajuda, 1349-017, Lisbon, Portugal, ORCID 0000-0002-8140-1257

DRAFT

Abstract

Two Asian longhorn beetles, *Anoplophora glabripennis* and *Anoplophora chinensis* are among the most serious alien invasive species attacking forest and urban trees, both in North America and Europe. Major efforts have been put into preventing further entry and establishment of the two species as well as promoting their successful eradication. Here we review these efforts, their progress and outcome, and scientific advancements in monitoring and control methods. The combined international activities and harmonizing legislative changes in detection and eradication methods have proven worthwhile, with more than 45% of eradication programmes successful in the last 12 years. Some countries were able to completely eradicate all populations and others managed to reduce the area affected. Although the costs of the eradication programmes can be very high, the benefits outweigh inaction. Attempts to eradicate *A. chinensis* have been more challenging in comparison with those targeting *A. glabripennis*. For both species, efforts are hampered by the ongoing arrival of new beetles, both from their native regions in Asia and from other invaded regions via bridgehead effects. The methods used for eradication have not changed much during the last decade, and host removal is still the method most commonly used. On the other hand, detection methods have diversified during the last decade with advances in semiochemical research and use of detection dogs. The next decade will determine if eradications continue to be successful, particularly in the case of *A. chinesis*, which has been targeted in some countries for containment instead of eradication.

Keywords: Biological invasions; *Anoplophora* spp.; eradication; management strategies; pest detection; surveillance

DRAFT

Declarations

Funding: This study was supported by the HOMED project (http://homed-project.eu/), which received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No. 771271. Both S. Branco and M. Branco are supported by CEF, a research unit funded by Foundation for Science and Technology (FCT), Portugal (UID/AGR/00239/2019 and (UIDB/00239/2020). M. Faccoli was partially funded by the DOR program of the University of Padua. This study was partly supported by the National Socio-Environmental Synthesis Center (SESYNC) under funding received from the National Science Foundation DBI-1639145.

Conflicts of interest/Competing interests: The authors declare that there is no conflict of interest/competing interests.

Availability of data and material: The datasets generated and analysed during the current study are available within the article and its supplementary materials

Ethics approval: This study does not contain any experiments using any animal species that require ethical approval.

Consent to participate: Not applicable.

Consent for publication: All authors consent to the publication of this manuscript in Journal of Pest Science.

Key Message

- *Anoplophora glabripennis* and *Anoplophora chinensis* are invasive wood borers native to Asia
- Both species are serious pests in their invaded range, attacking healthy forest and urban trees
- We analyze data from 2008-2020, regarding interceptions, establishments and eradications
- In Europe and North America more than 45% of eradication programmes were successful
- Innovations on management strategies and recent scientific achievements are reviewed

Author Contribution Statement

MB and HJ had the idea for the review. MB, HJ, EB and ND set the main structure of the work. SB, MF and GR conducted the literature search. SB conducted the data analysis and drafted the manuscript. All authors contributed to writing the final version of the manuscript and critically revised the work.

DRAFT

1. Introduction

In the last decades, increasing international trade resulting from globalisation has facilitated the introduction of non-native species to new environments and thus boosted the problems with biological invasions worldwide (Brockerhoff and Liebhold 2017; Liebhold and Kean 2019; Lesieur et al. 2019; Zhao et al. 2020). Invasive species have considerable ecological and economic impacts on agricultural, urban and forest systems, compromising their sustainability and the ecosystem services they provide (e.g., Boyd et al. 2013; de la Vega et al. 2020; Gugliuzzo et al. 2021). The Asian longhorn beetle (ALB) *Anoplophora glabripennis* (Motschulsky) and the citrus longhorn beetle (CLB) *Anoplophora chinensis* (Förster) (synonym *Anoplophora malasiaca* (Thomson)) (Lingafelter and Hoebeke 2002) are two emblematic examples of such alien invasive species.

Both ALB and CLB are highly polyphagous wood borers developing in dozens of deciduous tree species, with CLB having a wider host range than ALB (Lingafelter and Hoebeke 2002; Haack et al. 2010; Van der Gaag et al. 2010; Van der Gaag and Loomans 2014; Sjöman et al. 2014; EFSA et al. 2019a,b). In Europe, *Acer* is the most commonly attacked genus by both species (e.g. EFSA et al. 2019a,b). However, the two species differ regarding plant part on which oviposition and larval development take place. In ALB, oviposition and larval development occur on the upper trunk and main branches, whereas CLB mainly oviposits on the lower trunk, root collar region and on exposed roots, and larvae develop in the lower trunk and roots. This crucial difference translates into different pathways of introduction. ALB introductions are largely associated with the use of solid wood packing material (WPM) in international trade of goods, whereas CLB is rarely introduced with cut wood. CLB introductions are mainly associated with imports of live plants such as small maple trees and bonsais (e.g. Hérard and Maspero 2019).

ALB is native to China and the Korean Peninsula (Lingafelter and Hoebeke 2002; Williams et al. 2004a). Non-native breeding populations of ALB have been reported in many locations in the USA, Canada, Europe and Japan (Makihara 2002; Takahashi and Ito 2005; Hu et al. 2009; Haack et al. 2010), making ALB one of the most successful and most feared invasive insect species worldwide. CLB is native to eastern Asia, where it is widely distributed in China, Korea, and Japan. CLB has also been reported from Indonesia, Malaysia, Philippines, Taiwan, and Vietnam (Lingafelter and Hoebeke 2002; EFSA et al. 2019a). Contrary to ALB, established populations of CLB outside its native range have only been reported in a few countries in Europe. Both species have accidentally arrived in North America and Europe several times independently as documented by molecular genetic studies, numerous interceptions, and infestation hotspots (e.g., Haack et al. 2010; Hérard and Maspero 2019). These successive arrivals may hamper eradication attempts in a given region. Due to their potential impacts on ecosystems and many economically important tree species, these two species have been regulated as

DRAFT

priority quarantine pests in Europe, the United States and other countries (EU 2019; USDA-APHIS 2020a).

Haack et al. (2010) reported an extensive analysis of interceptions, establishments, eradications and management strategies used to deal with ALB and CLB in the invaded range, covering the period up to 2008. The authors also challenged the scientific community to respond to the needs identified by the difficulties associated with mitigating the threat posed by these beetles and with eradicating local established populations. Since then, 12 years have passed, but the two beetles still remain a menace for an increasing number of countries, and a large number of eradication programmes are still in progress. The aims of the present work are (i) to update the interception records which indicate ongoing transport with international trade, (ii) to review the eradication programmes carried out during the last 12 years, and (iii) to analyse the current status at the country level in order to understand the successes and failures of measures to mitigate invasions by the two beetles. Further objectives are to analyse the scientific achievements that occurred in the last 12 years, especially with regard to efforts in developing novel tools and methods for detection, monitoring and control, and to understand how the scientific community and managers have dealt with the challenges posed by these two species

1.1. Terminology and data sources

Interception. We follow the definition of interception provided in Haack et al. (2010), which further differentiates *entry interceptions* from *post-entry interceptions*. For the period prior to 2008, data from Haack et al. (2010) were used. After this period, interception data were retrieved from EPPO via Europhyt for Europe, thus representing the EU member states and Switzerland (data kindly provided by Françoise Petter, assistant director of EPPO) and for North America via USDA-APHIS (see Turner et al. 2020, 2021).

Establishment. The International Standard for Phytosanitary Measures (ISPM) No. 5 definition of establishment was adopted (FAO 2019). We consider a new establishment when located at least 5 km distant from infested trees detected in previous delimiting surveys or when findings occurred in a previously infested area, but where the population was officially declared eradicated by the relevant authorities (e.g., Toronto in 2013).

Demarcated area. The *demarcated area* corresponds to the area legally established by each national plant protection organization (NPPO) as subject to eradication and containment measures, and usually comprises an infested zone, where the pest is present, and a buffer zone around the infested zone (FAO 2019).

DRAFT

In order to obtain the temporal and geographical data of ALB and CLB establishments, demarcated areas and buffer zones, the main sources consulted were the EPPO Global Database (https://gd.eppo.int), GERDA - Global Eradication and Response DAtabase (Kean et al. 2015) and the USDA-APHIS website (https://www.aphis.usda.gov/). This information was complemented by a search of the scientific and grey literature, including works published in scientific journals, conference proceedings, presentations, and eradication reports and other technical reports of national and regional plant protection organisations. When only distribution maps were available, affected areas were extrapolated using ArcGis online tools. For data regarding the period up to 2008, this information was retrieved from Haack et al. (2010).

For most analyses we used two similar twelve-year periods, comparing data from 1997 to 2008 and from 2009 to 2020. For interceptions we used data from 1998 until 2019 (i.e., two eleven-year periods).

2. Interceptions and preventive measures

2.1 Regulation and legislation

In international trade, the International Standards for Phytosanitary Measures No. 15 (ISPM-15), which was adopted in 2002 and revised in 2009, provides treatment standards for WPM to be used in international trade and was intended "*to reduce significantly the risk of introduction and spread of most quarantine pests that may be associated with that material"* (IPPC 2009). Nevertheless, several factors can theoretically impact the effectiveness of ISPM 15: i) possibility of colonization after treatment, ii) insect tolerance to treatment, iii) fraudulent use of the ISPM 15 mark; and iv) unintentional noncompliance, which may occur when operators attempt to treat WPM according to ISPM 15, yet the minimum required doses of fumigant or heat are not achieved (Haack et al. 2010; Haack et al. 2014). Still, ALB and CLB are highly unlikely to colonise sawn timber as in WPM, and survival of appropriately applied ISPM 15-compliant treatment is also very unlikely (e.g., Myers and Bailey 2011). So, in most cases, ISPM 15 failure can probably be attributed to fraudulent use of the ISPM 15 mark and unintentional noncompliance (factors iii and iv).

Regarding introductions in association with live plants, a new EU regulation was adopted in October 2016 and implemented since December 2019 (regulation (EU) 2016/2031), on protective measures against pests of plants (repealing Council Directive 2000/29/EC), which completely bans the import of high-risk plants and selected plant products from countries outside of the EU (EU 2016). This regulation is expected to reduce the number of introductions/interceptions of *Anoplophora* spp., particularly of CLB.

DRAFT

Emergency measures to prevent the introduction into and the spread within the EU of ALB and CLB are defined in Commission Implementing Decisions 2015/893/EU and 012/138/EU, respectively (EU 2012, 2015). These include mandatory annual surveys to be conducted by each member state.

In the last decade, changes to protocols for inspection at ports of entry have also been adopted: the standard "Methodologies for Sampling of Consignments" (ISPM 31) was adopted in 2008. This standard outlines different types of sampling methods that NPPOs may use to verify compliance of consignments with phytosanitary requirements and the sample sizes required for general phytosanitary inspection (IPPC 2008). It complements ISPM 23 "Guidelines for Inspection", adopted in 2005, where the general procedures for inspection of consignments are described (IPPC 2005).

2.1. Interceptions

In Europe, ALB and CLB were intercepted 140 and 95 times, respectively, from 1980 until 2019. Considering the periods from 1998 to 2008 and from 2009 to 2019, the number of CLB interceptions decreased, with 48 *vs* 30 cases, whereas the number of ALB interceptions almost doubled (48 *vs* 90 cases) (Fig. 1).

A sharp difference was observed between time periods for both species regarding the site of interception, i.e., whether the interceptions occurred at "entry" or "post-entry" such as nurseries, warehouses, private residences, etc. For ALB, during 1998-2008, 97% of interceptions occurred "post-entry", whereas during 2009-2019 these proportions reversed, with 94% of interceptions reported during "entry" inspections. This increase in interceptions during border inspections is possibly a result of changes in legislation, namely the implementation of ISPM 31 in 2008. For CLB, the percentage of interceptions at "entry" also increased during 2009-2019, albeit more moderately (19% *vs* 57%).

For the period from 2009-2019, information on the origin of the infested material arriving at EPPO region was available in 88% of the cases (98% for ALB and 60% for CLB), mostly obtained during border inspections. For ALB, all infested consignments arrived from China while for CLB, in addition to China (83%), infested material originating from Japan was intercepted twice (11%) and an infested bonsai of unknown origin was shipped from the Netherlands. ALB interceptions were associated with wood packaging material (WPM) in 96% of cases (mostly linked to stone and tile products) and once to an object with wooden parts (1%). On the other hand, CLB was found in WPM only once. In 20% of cases, CLB was found in bonsais and in 70% it was found in other trees for planting. In two cases, each of ALB and CLB, only adults were found and the associated material could not be identified.

Excluding border inspections, 87% of ALB detections outside of their native range occurred after establishment (54/62). In the remaining 13%, which corresponded to "post-entry" interceptions, only adults and/or infested WPM were found. Contrasting this with CLB, the corresponding value is much

DRAFT

lower, with only 44% (20/45) of detections relating to established populations. These values reflect the different introduction pathways of each species: CLB is usually introduced in imported live plants whereas ALB is introduced in association with wood packaging materials (e.g. Eyre and Haack 2017). Live plants are subject to more intense inspection or incidental observation, either at nurseries or by the final consumer. Of the 25 post-entry interceptions of CLB approximately half (48%) occurred in nurseries, 40% at private residences, and three cases were detected in public parks and street trees.

Despite the adoption of ISPM 15 in 2002 which set strict standards for heat treatment and fumigation of WPM to be used in international trade (IPPC 2009), the number of reported interceptions with wood packaging in Europe has increased. Although this may be related to changes in inspection practices resulting from the implementation of ISPM 31, it still emphasises that ISPM 15 does not provide a guarantee that WPM is entirely pest-free, and that further improvements may be needed, especially to ensure prescribed treatments are indeed carried out (Haack et al. 2014).

In North America, a sharp decrease was observed in the number of interceptions of both ALB and CLB from 2009 to 2019 (18 ALB, zero CLB, Table 1), when compared to the period from 1998 to 2008 (72 ALB, 5 CLB, Haack et al. 2010).

In Europe, since 1998, three countries alone account for 70% *Anoplophora* spp. interceptions: The United Kingdom, the Netherlands and Germany. These three countries also ranked highest in international trade with East Asian countries during this period (WITS 2021), which may partially explain these results. In the United Kingdom, the number of interceptions of both species decreased in the last decade, by 88% for ALB and by 50% for CLB. By contrast, in Germany both increased (ALB by 108% and CLB by 40%). In the Netherlands, the number of ALB interceptions increased by 267% whereas CLB interceptions decreased by 55%. In Austria and Switzerland, while there were no ALB interceptions in the period 1998-2008, in the last decade, 18 and 17 cases were reported, respectively (Table 1). The different interception frequencies reported for each EU importing country are likely to reflect differences in inspection practices and differences in the reliance on ISPM 15 having solved the problem. Eyre and colleagues (2018) observed that the highest detection rates were achieved in Austria and France, whereas in Spain and Poland, despite the inspection of more than 500 consignments, no harmful organisms were detected. The authors suggested that harmonizing the inspection procedures to the most effective methodology may lead to an approximate sevenfold increase in the number of interceptions of invasive pests across all member states (Eyre et al. 2018). A study on relationships between interceptions and establishments of Cerambycidae (including ALB and CLB) found that there is a significant positive relationship overall between these parameters (Brockerhoff et al. 2014), which highlights the potential usefulness of recording interception data from inspections of relevant imports.

DRAFT




Figure 1 Temporal trend of the number of interceptions of *Anoplophora* spp. in Europe from 1998 to 2019.

Table 1 Interception data for *Anoplophora* spp. from 2009 to 2019

Year	2009		2010		2011		2012		2013		2014		2015		2016		2017		2018		2019		Total	
	A	C	A	C	A	C	A	C	A	C	A	C	A	C	A	C	A	C	A	C	A	C	A	C
Region/country																								
North America			5		1		6		3		2										1		18	
USA			5		1		6		3		2										NA	NA	17	
Canada																					1		1	
Europe	5	9	6	9	5		19	2	18	1	6	4	8	1	14	1	7	2	2				90	30
Austria									4		1		3		5		4		1				18	
Belgium					1			1															1	1
Czech Republic									2														2	
Cyprus		1																						1
Denmark						1*																		1
Estonia													1		1				1				3	
Finland													2				1						3	
France		2									1				1								2	2
Germany		1/2 *	4	1	3		10	1	7		1	1*	1		1			1					27	7
Netherlands	3	3	1	4			3		1		1			1	2	1							11	9
Slovakia									1*														1	
Sweden	2																						2	
Switzerland					1*		6		3	1*	2	1*	1		3		1						17	2
Turkey												1*						1*						2
United Kingdom			1*	4*		1*						1*			1		1						3	5
Total	5	9	11	9	6	2	25	1	21	1	8	4	8	1	14	1	7	2	2	0	1	0	108	30

A=ALB, C=CLB. Numbers indicate interceptions at ports of entry or transitional facilities ("entry" interceptions). "Post-entry" interceptions are indicated by "*". Specimens identified as *Anoplophora* spp. from wood packaging material were designated ALB and those from live plants were designated CLB.

3. Establishments

3.1. Spatial and temporal patterns of establishments

At the continental scale

Since the first detection of an established population in New York in 1996 until the end of 2020, 56 ALB and 20 CLB established populations were reported worldwide. From 2009 to 2020, 37 ALB and 10 CLB establishments were detected in North America and Europe (Fig. 2). More recently, an established population has also been reported from Hyogo Prefecture in Japan (Akita et al. 2021). Until now, CLB breeding populations outside their native range were detected only in Europe. For CLB, the number of new detected establishments was identical to the previous period (1997- 2008). However, a sharp difference was observed in the number of ALB establishments, which have more than doubled from 2009 to 2020. Furthermore, out of the 37 ALB establishments detected in the last 12 years, 62% were detected between 2012 and 2016.

A summary table with all the identified establishments of ALB and CLB by detection date and geographical location is shown in Supplement S1. For ALB, a brief description of the last decade of establishments by region and country is presented in Supplement S2. The detailed invasion history of CLB in Europe has recently been reviewed by Hérard and Maspero (2019) and is thus not covered in detail in the present work.

For ALB, the number of new establishments detected in Europe has increased more than fourfold in the period 2009-2020 relative to the period from 1997 to 2008. Out of the 37 ALB establishments detected in the last period, 84% were in Europe (Fig. 3, 4 and 5). The increase in the geographical distribution of ALB establishments in Europe mainly reflects the high number of establishments detected in Germany (9) and Italy (8). In contrast, in North America, until 2020 only six new establishments were detected since 2009 (three in Ohio, one in Boston, one in South Carolina and one in Ontario), which is approximately half of the number reported from 1997 to 2008. New CLB establishments were detected in Italy, Turkey, Croatia, France and the Netherlands (Fig. 4). Despite the high number of establishments detected in Italy, no interceptions have ever been reported there (see above).

By the end of 2020, the total demarcated area in Europe affected by ALB was about 630 km^2. This area corresponds to a 10-fold increase compared with the area affected by 2008 (62 km^2). This expansion reflects the large increase in the number of active establishments. By comparison, the total affected area changed little in North America, with an increase from 580 km^2 in 2008 to 770 km^2 by 2020. Still, despite the number of total ALB establishments detected in Europe being higher than those in North America, the current demarcated areas are similar in the two regions. The demarcated areas in Europe are mainly concentrated in three countries, Italy, Germany and France (Supplement S1).

For CLB, although the number of detected establishments was identical from 1997-2008 to 2009-2019 (10 establishments, Fig. 4), the total demarcated areas of all active establishments (including establishments detected before 2009), almost quadrupled in Europe (150 km^2 in 2008 *vs* 590 km^2 in 2020). The demarcated areas by country expanded mostly in Italy (from 140 to 510 km^2), Croatia (from 0 to 55 km^2) and France (from 3.1 to 8.9 km^2) (Supplement S1).

DRAFT




Figure 2 Number of *Anoplophora* spp. establishments detected by year, from 1996 to 2020.




Figure 3 Number *Anoplophora* spp. establishments detected by time period and country.

DRAFT




Figure 4 Geographical distribution of established populations of ALB in Europe by year of detection. a) Status of establishments up to 2008, b) status of establishments from 2009 to 2020. Red dots represent active establishments, green dots eradicated establishments (as of April 2021): 2001: Braunau, Austria (1); 2002: Gien, France (2); 2003: Sainte-Anne-sur-Brivet, France (3); 2004: Neukirchen, Germany (4); 2005: Bornheim, Germany (5); 2007: Corbetta, Italy (6); 2008: Strasbourg, France (7); 2009: Cornuda, Italy (8); 2010: Maser, Italy (9), Almere, Netherlands (10); 2011: Brünisried, Switzerland (11); 2012: Geinberg, Austria (12), Feldkirchen, Germany (13),Winterswijk, Netherlands (14), Winterthur, Switzerland (15), Paddock Wood, UK (16); 2013: Gallspach, Austria (17), Furiani, France (18), Grottazzolina, Italy (19); 2014: Magdeburg, Germany (20), Neubiberg, Germany (21), Ziemetshausen, Germany (22), Marly, Switzerland (23); 2015: Vantaa, Finland (24), Grenzach-Whylen, Germany (25), Budva, Montenegro (26), Porto San Giorgio,Italy (27), Berikon, Switzerland (28) ; 2016: Divonne-les-Bains, France (29), Kelheim, Germany (30), Murnau, Germany (31), Hildrizhausen, Germany (32), Ostra and Senigalia, Italy (33); 2017: Trescore Balneario, Italy (34); 2018: Vaie, Italy (35), Cuneo, Italy (36); 2019: Civitanova, Italy (37), Miesbach, Germany (38).



Figure 5 Geographical distribution of established populations of ALB in North America, by year of detection. a) Status of establishments up to 2008, b) status of establishments from 2009 to 2020. Red dots represent active establishments, green dots eradicated establishments (as of April 2021) 1996: Brooklyn, New York, USA (1), Long Island, New York, USA (2); 1998: Chicago, Illinois, USA (3), Addison, Illinois, USA (4), Summit, Illinois, USA (5); 1999: Park Ridge, Illinois, USA (6); 2000: Islip, New York, USA (7), Chicago O´Hare, Illinois, USA (8); 2003: Vaughan, Ontario, Canada (9); 2004: Carteret and Linden (2006), New Jersey and Prall and Staten Island (2007), New York, USA (10); 2008: Worcester, Massachusetts, USA (11); 2010: Boston, Massachusetts, USA (12); 2011: Tate Township, Ohio, USA (13), Monroe Township, Ohio, USA (14), Batavia/Stonelick Townships, Ohio, USA (15); 2013: Mississauga, Ontario, Canada (16); 2020: Hollywood, South Carolina, USA (17).

At the local scale

All established populations of *Anoplophora* spp. were initially detected in urban/peri-urban environments (Fig. 6). For both species, infested trees and live beetles were initially detected in private

gardens in approximately half of the establishments (52% for ALB and 50% for CLB). Detections in public parks and street trees were also common, whereas detection in peri-urban forests was rare and occurred only once in one ALB and one CLB establishment, during official surveys.

In its native range in South Korea, ALB has been reported to be a riparian species adapted to the long edges of these habitats (Williams et al. 2004a). If this is the case, it might explain its adaptability to hedgerows (along roads, gardens, and parks) typical of urban habitats (Williams et al. 2004a; Faccoli et al. 2016). This is in accordance with the infestation pattern in Cornuda (Italy), where, although part of the quarantine area fell within a natural hardwood forest, infested trees have only been found along its edges (Faccoli et al. 2016). Similarly, in Chicago (USA) hundreds of Acer spp. were found infested along the edge of a 50-ha woodlot but not in the interior, suggesting a strong edge effect during the invasion (Sawyer et al. 2011). The infestation of hardwood stands in a large outbreak observed in Massachusetts (USA) has been pointed out as an exception (Dodds and Orwig 2011). However, the small size of the infested stands in Massachusetts, surrounded by city outskirts and streets, makes them comparable to urban parks and small rural stands (Faccoli et al. 2016).




Figure 6 Sites of initial detection(s) of Anoplophora spp. establishments. This information was available for 49/55 and 18/20 of ALB and CLB establishments, respectively.

Differences between the two species can be seen which are related to their pathways. For ALB, industrial/commercial sites (areas that are likely to receive imports in WPM or live plants from potential source regions) are commonly affected and detections at such sites occurred in 30% of establishments. For CLB, 50% of detections involved sightings of insects or infested trees at plant nurseries (28%) or near plant nurseries (22%).

Considering how detections occurred initially, 73% of ALB establishments (33/45) were detected by passive surveillance and 24% were detected during official surveys. For CLB, 76% (13/17) of cases were reported during official surveys and the remaining 24% were the result of either passive surveillance (one case) or detected during scientific research activities (3 cases). Passive surveillance

corresponded mostly to citizens who reported symptoms or sightings of adult insects to phytosanitary authorities, operators of nurseries and city parks and landowners. For most establishments, the first trees infested were maples (*Acer* spp.), corresponding to 90% and 95% of ALB and CLB cases, respectively (Fig. 7). However, while ALB was found infesting mostly local trees of *A. platanoides* and *A. pseudoplatanus*, CLB was mostly found infesting *A. palmatum* and *A. negundo*. For ALB, *Salix* sp., *Ulmus* sp. and *Aesculus hippocastanum* were also commonly found infested. For CLB, in addition to maples, the most common infested tree genera were *Carpinus*, *Corylus*, *Betula* and *Platanus* (Fig. 7). The host trees affected at each site are expected to be influenced by the host species available. Nevertheless, affected hosts may also reflect the origin of the local populations: in South Korea, for example, ALB riparian forest populations appear to display a different host usage when compared to urban populations, and the latter have been shown to result from recent invasions from China (Lee et al. 2020).




Figure 7 Tree species or genera in which establishments of ALB and CLB were detected either during passive surveillance or official or scientific surveys. This information was available for 49/55 and 20/20 of ALB and CLB establishments, respectively.

3.2. Pathways of introduction: reconstruction of invasion routes

ALB

The first studies on the intraspecific genetic diversity have focused mainly on the native Asian regions (An et al. 2004; Carter et al. 2009a). They reported that, although the Asian populations clustered roughly into two major groups, the population structure has been influenced by movement of beetles and consequent genetic admixture (Carter et al. 2009a). Javal et al. (2019a,b) highlighted signs of an ancestral structure in NE Asia, and a strong differentiation among most of the populations following a north-south gradient. These studies also considered human-mediated population translocations at large scale, especially those linked to afforestation projects initiated by the Chinese government since the 1960s in northern and eastern China (Li 2004; Haack et al. 2010).

Studies of North American establishments revealed a reduced genetic diversity within populations in either the USA or Canada due to genetic bottlenecks (Carter et al. 2009b, 2010). Separate introduction events were responsible for most North American populations, the founders of which likely originated from populations invasive within China (Carter et al. 2010; Javal et al. (2019a,b). In addition, some subsequent human-mediated regional spread occurred in the USA (e.g. New York City, Carter et al. 2010) and in Canada (Turgeon et al. 2015).

In Europe, mitochondrial DNA and microsatellite marker studies revealed a complex worldwide invasion scenario involving recurrent introductions coupled with a bridgehead event. The genetic structure observed suggests that European establishments originated mostly from multiple independent introductions from the native area in Asia (Fig. 8). The resulting genetic differentiation among European establishments may indicate limited gene flow between populations once established, mostly due to the poor dispersal behaviour of this species. A fine-scale study in Switzerland (Tsykun et al. 2019) showed that only one or a maximum of two genetic clusters were found within a given tree, suggesting that most ALB individuals remain in proximity to the tree from which they emerged when suitable host trees are available (Smith et al. 2001 and 2004; Javal et al. 2019a,b). Low levels of genetic diversity, high levels of inbreeding, small numbers of founders and large differences in the severity of bottlenecks encountered by introduced populations have shaped the genetic structure of invasive populations (Javal et al. 2019a,b). Natural dispersal and/or human-mediated transportation (e.g. hitch-hiking) at a small spatial scale were observed in some regions as in Corsica and in Switzerland (Javal et al. 2019a,b; Tsykun et al. 2019). Bridgehead events may have contributed to the worldwide spread of ALB (Javal et al. 2019). This appears to be the case for the French population in Gien that may have resulted from a bridgehead population from North America (Javal et al. 2019a,b) rather than Asia as suggested in previous publications (Cocquempot et al. 2003).

DRAFT




Figure 8 Distribution of some European ALB populations and their defined genetic clusters inferred by structure analysis. The numbers correspond to the sampled locations used in Javal et al. (2019a): 1 Gien, France; 2 Sainte-Anne-sur-Brivet, France; 3 Strasbourg, France; 4 Cornuda, Italy; 5 Brünisried, Switzerland; 6 Winterthur, Switzerland; 7 Feldkirchen, Germany; 8 Rapagnano, Italy; 9 Gallspach, Austria; 10 Arenau, Corsica; 11 Colast, Corsica; 12 Conouv, Corsica; 13 MCarlo, Corsica; 14 Costad, Corsica; 15 Neubiberg, Germany; 16 Ebersberg, Germany; 17 Marly, Switzerland; 18 Berikon, Switzerland; 19 Divonne les Bains, France. Each colour corresponds to a haplotype cluster.

CLB

The genetic structure and invasion pathways of CLB have not been studied as intensively as for ALB. Strangi et al. (2017) conducted a mitochondrial DNA analysis on native populations from East Asia and three Italian establishments. In Italy, a total of five haplotypes were identified in Lazio, Lombardy and Tuscany. Three of these haplotypes were only found in Tuscany, and these were closely related to haplotypes found in Chinese populations. The remaining two haplotypes, found in Lazio and Lombardy, corresponded to populations from North and Central Japan (Strangi et al. 2017). These results suggest that the Italian establishments originated from at least two separate events. CLB is known to show phenotypic polymorphism that allow for the distinction of two forms: *A. chinensis chinensis* and *A. chinensis malasiaca* (Ohbayashi et al. 2009). In the Lazio and Lombardy establishments, where the origin was traced back to Japan, specimens were identified as *A. chinensis malasiaca*, whereas in Tuscany, where the population origin was traced back to China, specimens were shown to be *A. chinensis chinensis*. In the recent establishment detected in Royan, the *A. chinensis chinensis* subspecies has been detected, and further studies are currently underway to uncover the origin of the specimens recovered at this location (ANSES 2019).

DRAFT

These recent studies have started to unravel the complexity of ALB´s and CLB´s invasion histories: multiple introductions have occurred, originating from several regions of Asia (China, Korea, Japan) and in some cases these appear to have included bridgeheads effects. Furthermore, studies have also shown that even genetic populations of extremely low genetic diversity can multiply to outbreak proportions in urban areas (Carter et al. 2009b).

4. Eradications

4.1. Methods of eradication

In the European Union, CLB and ALB infestations that affect a Member State are subject to specific management procedures defined by the European Commission (EC 2012 and 2015 respectively) and transcribed in national and regional decrees with the aim to eradicate all active infestations. Each country where an infestation is detected usually develops an eradication programme that incorporates activities focused on detecting infested trees, removing pest populations and limiting pest movement and spread, i.e. every eradication programme includes monitoring, control and containment components (see section 6 for their description). The first step is to delineate a demarcated area and forbid movement outside the demarcated area of infested or potentially infested wood material and host trees (EPPO 2013a,b). Whenever a new establishment is detected, an initial, intensive delimiting survey must be conducted to determine the extent of the infested area. Demarcated areas are then established including the infested area and a surrounding buffer zone of typically 2 km radius. Depending on the extent of the infestation and the site-specific characteristics, such as the local distribution of host plants, this buffer zone may be reduced to a radius of 1 km. The prescribed survey methodology is similar across all EU member states. Once trees are found to be infested by ALB/CLB, they are recorded and submitted to specific protocols aimed to eliminate insect populations, based on different types of measures including tree destruction, chemical and physical treatments (EPPO 2013a,b).

In North America, annual surveys to actively search for new infestations are, to our knowledge, not mandatory. However, once an infestation is detected, similar protocols apply: establishment of regulated areas consisting of a core area (0.8 km radius) and a buffer zone (1.6 km beyond the boundary of the core area). USDA APHIS (2020b) further outlines that the buffer zone should expand to a minimum of 4 km from areas of high ALB density (defined as presence of a cluster of trees with many exit holes or one or more trees with >100 exit holes). In North America, where only ALB establishments have been detected to date, the eradication procedures have been similar to those applied in Europe (Haack et al. 2010; USDA-APHIS 2014; Fournier and Turgeon, 2017). In the USA specific ALB response guidelines were published providing the technical and general information needed to implement any phase of an ALB eradication programme and the Federal Quarantine authority for ALB according to the US Federal Regulations 7 CFR 301.51 for eradication programs (USDA-APHIS 2014, 2020b).

DRAFT

4.2. Spatial and temporal pattern of successful eradications

ALB

Globally, as of December 2020, approximately half of all detected ALB establishments have been eradicated successfully (53%, since the first successful eradication of the establishment of Addison, Illinois, USA, in 2004) (Fig. 9). However, eradication success rates varied considerably among countries and continents. In Europe, all ALB establishments in Austria, the Netherlands, Switzerland and the UK have been declared eradicated. Austria, which had the highest affected area in 2008, succeeded to eradicate its largest establishment, with only a small area of active cases remaining in 2020, which was finally declared eradicated in January 2021 (Supplement S1). By contrast, in Italy, which ranks first in the number of active establishments, 80% of detected ALB establishments remain active.

A high rate of successful eradications was also achieved in North America. In the United States, where the highest number of ALB establishments (15) has been reported to date, 67% of eradication attempts have been successful so far. A few successful eradications have been achieved before 2008, in Jersey City and Illinois. The total area for which successful eradication of ALB was achieved in North America during the period from 2008 to 2020 was about double that in Europe (Fig. 10). A decrease in the area affected was achieved in both the USA and Canada.

CLB

Eradication of CLB establishments appear to be more challenging, as only 30% of detected establishments have been declared eradicated (until December 2020) (Fig. 9). Only six out of 20 established populations outside its native range have been eradicated successfully since 2008.

Furthermore, eradication of most CLB establishment sites in Lombardy has recently been declared as unachievable. Currently, of the eight active establishments in Italy, only two are still under eradication. With the exception of the small establishment in Sirmione, which is still under eradication, a policy of containment is thus now in place at all other sites in Lombardy (SFRL 2020).




Figure 9 *Anoplophora* spp. eradication attempts by country and their respective status, as of December 2020 (active or eradicated).

DRAFT




Figure 10 *Anoplophora* spp. demarcated areas (DA) by world region, Europe and North America: active in 2020 and eradicated in the period 2008-2020.

5. Spread

5.1. Methods to monitor and predict the spread

Information on how the invasive population will likely spread across the landscape is fundamental to delineate cost-effective monitoring and control strategies. Currently, certain distances from a discovered infestation of *Anoplophora* spp. are used in eradication programmes to define the boundaries of delimiting survey areas mandatory by law (EU 2012, 2015; USDA 2019). However, those boundaries need to be adjusted according to the available scientific knowledge on the beetles' dispersal ability. A number of dispersal studies have been published for ALB (e.g. Bancroft and Smith 2005; Li et al. 2010; Sawyer et al. 2011; Turgeon et al. 2015) whereas for CLB the information is scarce (Adachi 1990; Cavagna et al. 2013). Due to their morphological resemblances one may assume the dispersal ability of the two species to be similar.

When analysing different studies, we distinguish those based on insect dispersal ability and observations of population spread. Potential dispersal ability does not always match the observed spread due to landscape features and aspects of insect behaviour. Insect dispersal ability was studied by mark-release studies, flight mills and modelling whereas population spread is accessed by analysis of historical infestation cases, genetic analysis, and different kinds of models (see below).

5.2. Patterns of spread at local scale

For ALB, mark-release-recapture studies conducted in China reported mean dispersal distances during one season of 100 m to 270 m, with a 98% probability of beetle recapture within 560 m to 920 m and a maximum dispersal potential of 2,600 m (Wen et al. 1998; Smith et al. 2001, 2004; Williams et al. 2004b; Bancroft and Smith 2005; Li et al. 2010). Studies conducted with computerized flight mills have shown that some beetles can fly considerably longer distances, up to 14 km (Lopez et al. 2017; Javal et al 2018), although only 5% of individuals travelled more than 8 km within a 24-h period (Lopez et al.

DRAFT

2017). These extreme specimens may lead to infestations outside of quarantine zones (Javal et al. 2018b). However, it is important to note that the beetles' ability to fly long distances in flight mills does not necessarily translate into long distance flights in the field. There is some evidence that suggests ALB is reluctant to fly far even though they are physically able to do so in a flight mill situation.

By examining historical infestation cases, spread rates were seen to be highly variable both between infested sites and from one period to another within a given infested area. For example, Sawyer et al. (2011) observed in urban areas at Carteret (NJ; USA) and Chicago (IL, USA) that ALB spread slowly, concentrated within a few hundred metres during the first 5-6 years. Yet, in another location, in Linden (NJ, USA), the infestation spread much faster, about 3.2 km within five years. In a study conducted in southern England, it was estimated that ALB remained restricted to a small area for approximately 10 years near a heavily infested sycamore tree (Straw et al. 2016). Similar patterns of infestation, with the beetles remaining at or close to the natal tree have been observed in the early phases of infestation at other sites (Haack et al. 2010; Sawyer et al. 2011; Turgeon et al. 2015). The discrepancies between sites may in part be attributable to differences in the time until an established population was discovered, while landscape heterogeneity may also play a role. Some land cover types may offer lower resistance to beetle movement and low availability of suitable host trees, favouring longer dispersal flights (Keena 2018). ALB adults are assumed to move by walking in the vicinity of the natal tree and disperse by flying only when conditions become less favourable. Still, in North America long-range dispersers of up to ~1,400 m, were reported, even before the originally infested host trees were fully exploited (Hull-Sanders et al. 2017).

Climatic conditions may also play a role in the dispersal of ALB. The apparently lower rate of population increase and spread of ALB in southern England (Straw et al. 2016), when compared to Cornuda in Italy (Favaro et al. 2015) or Jersey City and Linden in the US (Sawyer et al. 2011), has been attributed to lower summer temperatures resulting in longer insect developmental times (Straw et al. 2016; Trotter and Keena 2016). In northern Italy, ALB dispersal was shown to be influenced by the distance of suitable hosts from the nearest infested trees ($p < 0.01$ for distances above 510 to 1,040 m, which varied among years) and the number of infested trees around uninfested ones (Favaro et al. 2015). In that study, although the probability of dispersing farther than 1,900 m from a previously attacked tree was very low ($p < 0.001$), one dispersal occurrence was registered at 2,224 m. The dispersal pattern was shown to be density-dependent, in accordance with previous mark-release studies.

Several modelling approaches have been developed to describe ALB dispersion patterns. Trotter and Hull-Sanders (2015) and Trotter et al. (2019) used graph theory to determine the topological connections between infested trees, which was then used to calculate dispersal patterns across the landscape in Massachusetts. Two scenarios were used in this study: one in which beetles only left the natal tree when

it was overcrowded (strict scenario) and one under which all infested trees could act as sources of dispersing beetles (relaxed scenario). The longest dispersal distance, within a 99th percentile, was over 2.3 km for the strict scenario, and 1.3 km under the relaxed scenario. Fragnière et al. (2018) used data from establishments in Switzerland to develop a density-dependent model that relies on field observations of beetles and infested trees to provide a risk index (RI) of the presence of ALB in a given location. The output for Marly, for example, resulted in RI > 0.001 up to about 600 m of the centre of the highly infested area and RI > 0.0001 up to about 820 m. Elmes et al. (2019) modelled dispersal pathways using circuit theory. Their results showed that ALB tends to use non-habitat land-cover types to connect suitable habitat patches and that for this species, circuit theory was a better predictor of dispersal spatial patterns than least-cost dispersal models. The non-habitat land-cover type that displayed the lowest resistance was sealed surfaces (such as roads) followed by bare soil, grassland, trees, buildings, and water, in increasing resistance order. Recently, Huang et al. (2020) used a geographically weighted regression model to analyse the spatial differentiation of environmental drivers on the occurrence of ALB in China. Temperature, wind speed, precipitation and population density were shown to affect ALB occurrence in China, yet a high spatial heterogeneity was reported on the influence of these factors.

Studies on CLB dispersal are scarce compared with the information available for ALB. Its spread capacity is reported to be low (EFSA et al. 2019a). Similar to ALB, most adults are assumed to disperse by walking and remain in the vicinity of their natal tree unless conditions are unfavourable, although some adults were shown to be able to travel distances of 2 km (Adachi 1990). In Lombardy, Italy, the maximum distances between infestations in urban and agricultural areas were calculated to be about 500 m and 663 m, respectively (Cavagna et al. 2013). However, 97.0% and 99.2% of new cases were found within 200 m and 400 m, respectively. EFSA et al. (2019a) estimated the maximum distance of natural spread in one year to be approximately 194 m (with a 95% uncertainty range of 42–904 m), for a population with a 2-year life cycle (EFSA et al. 2019a).

As mentioned above, human-mediated dispersal related to commerce and transport of infested plants, wood and other materials is the major route for spread of both species at the continental scale. However, even at shorter distances, human-mediated dispersal is an important component that needs to be considered as a cause of satellite infestations, as has been shown, for example, in Switzerland, the USA and Canada (Turgeon et al 2015; Tsykun et al. 2019).

6. Control and Containment - current and future perspectives

6.1. Monitoring methods

DRAFT

In Europe, a survey is carried out in each demarcated area at least once per year to detect and monitor infested trees (EC 2012, 2015). The methods used have been quite similar among countries and mainly based on visual surveys. Advancements in alternative monitoring methods are described below.

Visual surveys

Despite the advances in new detection methods in the last decade, visual surveys remain the standard procedure for *Anoplophora* spp. monitoring (EFSA et al. 2019a,b). These surveys are generally based on examination of potential host trees looking for signs of infestation (i.e., exit holes, larval frass on the ground, oviposition pits and adult feeding, plant and branch dieback). CLB infestation signs are searched on the lower part of the trunk (usually the basal 50 cm, but infestations up to two meters high have been documented, Doris Hölling, Pers. Commun.), the root collar zone, and roots exposed above ground, while searches for ALB symptoms are focussed on the upper part of the trunk and the main branches (EFSA et al. 2019a,b). ALB surveys are usually conducted by observers on the ground equipped with binoculars to detect known signs and symptoms of attack. Turgeon et al. (2010) demonstrated that the efficacy of ground inspections is higher when the density of oviposition is higher, when signs are located lower on the tree, and when they are positioned on the main trunk. Furthermore, the authors observed that most infested trees were detected within the first 2 min of survey, and that using a team of inspectors to survey each tree would be more time effective than the use of a single inspector per tree (Turgeon et al. 2010). The type of environment on which the trees are located also affects detectability: infested street trees are more easily detected than those located in parks or woodland, therefore affecting the time required for tree inspection at different sites (Yemshanov et al. 2019). In addition to surveys carried out inside the demarcated area, specific surveys are usually conducted also randomly outside the demarcated area at high-risk sites such as commercial and industrial areas that receive imports from potential source regions, particularly those receiving wood packaging material or live plants (EFSA et al. 2019a,b).

Semiochemicals

For ALB, pheromone-based trapping systems have been developed (Nehme et al. 2014). Males of ALB are known to emit a sex pheromone composed of equal parts of 4-(n-heptyloxy)butan-1-ol and 4-(n-heptyloxy)butanal (Zhang et al. 2002; Nehme et al. 2009). Intercept panel traps baited with a combination of the pheromone and a mixture of selected host plant volatiles, namely linalool, linalool oxide, cis-3-hexen-1-ol and trans-caryophyllene, proved attractive to females (primarily virgin females) in field trials (Nehme et al. 2010, 2014). CLB males were shown to emit the same two functionalized dialkylethers as ALB males. In field bioassays both sexes were attracted to 4-(n-heptyloxy)butan-1-ol, suggesting that this compound is an important component of the CLB sex pheromone (Hansen et al. 2015). However, the effectiveness of these male pheromone-based trapping systems for monitoring *Anoplophora* spp. is thought to be limited (EFSA et al. 2019a,b), not only because the lures used primarily attract only virgin females but it is also likely that at close range mate finding includes

DRAFT

additional visual and chemical cues, including those coded in specific host phytochemicals (particularly sesquiterpenes) which require further research (Nehme et al. 2014; Hoover et al. 2014; Xu and Teale 2021).

New possibilities may arise from the identification of female-produced pheromones. For ALB, female-produced aggregation (Wickman et al. 2012; Xu et al. 2020a,b), contact (Zhang et al. 2003) and trail pheromones (Hoover et al. 2014) have been reported. Wickham et al. (2012) identified an ALB female-produced aggregation pheromone composed of a blend of heptanal, nonanal and hexadecanal, which proved attractive when combined with host volatiles. Xu et al. (2020a) showed that α-longipinene is a major component in extracts of virgin ALB female genitalia and that in olfactometer bioassays, both sexes were attracted to this sesquiterpene. Although α-longipinene is also released by males and host twigs, the authors suggest that the ratios released by these different sources may encode information pertaining to multiple purposes such as aggregation, mate and host location, and that identification of the naturally produced enantiomer in ALB and its hosts is also needed (Xu et al. 2020b).

For CLB, the sesquiterpenes b-elemene, b-caryophyllene, a-humulene, and a-farnesene, released both by the beetles and by the host plant, *Citrus unshiu*, after beetle feeding or after mechanical wounding, proved attractive to males and are thought to act both as kairomones and sex pheromones (Yasui et al. 2007, 2008; Yasui 2009). A female-produced contact sex pheromone of CLB has also been described (Fukaya et al. 2000; Akino 2001; Yasui et al. 2003, 2007).

Sniffer dogs

Recently, “sniffer dogs” have been trained and used in several European countries to identify infested trees through the specific odours released by ALB/CLB larvae and their frass. The use of sniffer dogs specifically trained for the detection of *Anoplophora* spp. was pioneered in 2009 by the Austrian Federal Forest Office (Bundesforschungszentrum für Wald (BFW)) in Vienna (Hoyer–Tomiczek and Sauseng 2013). These detection dogs proved effective at detecting all developmental stages of ALB/CLB in wood packaging materials, imported plants and standing trees in areas where establishment had occurred (Hoyer–Tomiczek and Sauseng 2013). In field experiments, trained dogs displayed high levels of sensitivity in the order of 75–88% (correct positives out of all positives) and specificity of 85–96% (correct negatives out of all negatives) (Hoyer–Tomiczek et al. 2016). This method is already being used in addition to visual surveillance in several areas in Europe and good results have been obtained in Austria, France, Italy, Switzerland and Germany (Hoyer–Tomiczek et al. 2016; EFSA et al. 2019a,b). New dog training teams have now been established in Austria and Switzerland (EFSA et al. 2019a,b). In the US, canine detector units were also evaluated with success in Worcester, Massachusetts (Errico 2012). The downside of this method is that in order to maintain a high-performance level, these dogs

DRAFT

must continuously be stimulated with *Anoplophora* material such as frass and live or dead larvae that are still relatively fresh, and they can only be used for limited periods per day so that a large number of trained dogs is necessary to inspect all relevant imports and potentially infested sites (Hoyer–Tomiczek et al. 2016; EFSA et al. 2019a,b).

Other detection methods

Bioacoustic detection methods use portable detectors attached to trees to record the sounds and vibrations produced by larvae (Mankin et al. 2008; Sutin et al. 2019). The potential use of acoustic methods for *Anoplophora* spp. detection has been acknowledged by the international EPPO standards. However, so far, the use of acoustic sensors in the field is difficult and the sensitivity and measuring accuracy of these devices are strongly influenced by the nature of the sensor-substrate interface. These factors limit these methods practical applications (Zorović and Čokl 2015; Hérard and Maspero 2019). More recently, laser vibrometry has been developed for this purpose. With this method, a laser beam is used to detect the vibrations produced by larvae. Recording is carried out directly from the vibrating surface avoiding the need to mount detectors on the tested materials (Zorović and Čokl 2015). Although only laboratory tests have been conducted to date, the methods displayed high sensitivity and a high signal to noise ratio (Zorović and Čokl 2015; Hérard and Maspero 2019). However, a major drawback is that eggs, pupae, and diapausing insects cannot be detected by these methods.

Citizens' involvement in monitoring and surveillance have been proposed and carried on in a few countries, namely in Austria (EC 2010), France (EPPO RS 2017/005), Italy (Jucker et al. 2007), Germany (StMELF 2020) and Switzerland (EFSF 2020).

6.2. Control and Containment

Tree destruction and physical treatments

Eradication programmes include the removal (felling) and destruction (chipping or burning) of infested trees and possibly their replacement with non-host tree species. Whereas many countries fell and destroy only infested trees, other countries apply preventive tree destruction of all host plants, even if healthy, within in a certain radius around infested trees. This radius usually ranges between 20 m and 100 m (EPPO 2013a,b) (Supplement S3). Under current EU legislation preventive tree destruction of high-risk hosts trees is now mandatory (EC 2012, 2015). Other differences among countries in the management of the CLB infestations concern the treatment of stumps, which could be uprooted and destroyed, covered with metal nets to avoid adult emergence, or treated with herbicide to prevent regrowth (Supplement S3) (EPPO 2013a). These measures are effective in reducing ALB/CLB populations and can contribute to eradication, although they are very laborious, expensive, and time-consuming.

DRAFT

743 *Chemical methods*

744 In the past, trunk or soil injections with imidacloprid, a neonicotinoid systemic insecticide, were applied
745 in the USA and Japan to each potential host tree growing within an 800 m radius from infested trees to
746 reduce ALB population density and prevent infestation spread (Hu et al. 2009; Haack et al. 2010).
747 Chemical treatments of healthy trees were combined with removal of infested trees, which proved to be
748 effective. In China, ALB populations were controlled by spraying pyrethroids (cypermethrin) in the tree
749 canopies or coating trunks of host trees to kill adults. Another strategy was inserting wooden sticks
750 containing aluminium phosphide (generating phosphine) into larval galleries to kill ALB larvae, or
751 injecting trunks with organophosphate insecticides such as methamidophos (Wang et al. 2005; Hu et al.
752 2009). Most systemic insecticides were found to persist at lethal levels for several months after injection,
753 but they require new treatments year after year, and their uniform distribution within trees is still
754 uncertain. A potential alternative might be the use of emamectin benzoate trunk injections. In a study
755 recently conducted in an infested willow forest in Beijing, China, this compound proved effective at
756 reducing ALB larval populations by 89% in the first spring after application and by >99% during the
757 second year. Only in the third year after application did re-infestation occur (Wang et al. 2020).
758 Nevertheless, insecticides are costly and their use is labour intensive, making chemical control
759 economically and environmentally expensive (Hu et al. 2009).

760

761 In Europe, the use of chemical treatments has been rare (Supplement S3): it has long been acknowledged
762 that insecticides may cause significant negative externalities including biodiversity loss, ground and
763 surface water contamination (including off-field habitat contamination), impacts on non-target
764 organisms including biocontrol agents, pollinators and earthworms, bio-amplification of toxic
765 substances within the food web with potential effects on human and animal health and development of
766 resistance (Pimentel 2005; Pelosi et al. 2021). The severity of these impacts will depend on the
767 specificity and toxicity levels of the substances used. Therefore, in case eradication fails, additional
768 management options such as biological control are required.

769

770 *Genetic and cultural methods*

771 Research on the identification of tree species or clones resistant to ALB and CLB has not been successful
772 in the last decade. However, the increased use of non-host trees would be suitable for reducing new ALB
773 and CLB infestations. Under current EU legislation the planting of high-risk species in the infested areas
774 is prohibited (EC 2012, 2015).

775

776 *Biological control*

777 Many studies have been carried out on natural enemies that could be used as potential biocontrol agents
778 of ALB, including pathogens (bacteria, fungi, and nematodes), parasitoids and predators (reviewed by
779 Brabbs et al. 2015). Virulent strains of *Beauveria brongniartii* (Sacc.) (Hypocreales: Cordycipitaceae),

Beauveria. asiatica Rehner and Humber, and *Metarhizium brunneum* Petch (formerly *M. anisopliae* (Metschnikoff) (Hypocreales: Clavicipitaceae) are under development for control of ALB (Goble et al. 2014, 2016; Meng et al. 2015; Clifton et al. 2020a). *Beauveria brongniartii* has already been developed into a commercial product in Japan, and *M. brunneum* is available for commercial use in the US, both inducing high mortality rates (Brabbs et al. 2015, Clifton et al., 2020a,b). *Beauveria brongniartii* (Hypocreales: Cordycipitaceae) and *M. brunneum* have also been shown to infect CLB (Brabbs et al. 2015). Exposure to *M. brunneum* fungal infection synergize with neonicotinoid insecticides (Imidacloprid) used for tree protection resulted in accelerated host death (Fisher et al. 2017). However, the fungal virulence of *M. brunneum* is limited by unsuitable environmental conditions and its effectiveness is affected by adult age (Fisher and Hajek 2014, 2016).

Entomopathogenic nematodes belonging to the genera *Steinernema* and *Heterorhabditis* were also tested against ALB (Fallon et al. 2004; Pan 2005). Strains of *Steinernema carpocapsae* and *S. feltiae* have proven to be capable of infecting both *Anoplophora* species and they have potential for use as biopesticides as an alternative to chemical treatments. Of the different application methods tested, the most effective included using sponges or gauze to block or cover larval tunnels for CLB (90%-91% mortality rate) and directly spraying into tunnels for ALB (86%). Simple trunk applications were also effective when tested against CLB, albeit more moderately (60 to 77%) (Brabbs et al. 2015).

Two woodpecker species native to Eurasia, *Dendrocopos major* Beicki and *Picus canus* Gmelin, are the major predators of ALB in China (Brabbs et al. 2015) and they have been shown to be effective at controlling ALB in Chinese forests where nesting has been encouraged (Pan 2005, Golec et al. 2018). Nevertheless, the low levels of mortality attained (less than 16%) are unlikely to provide population control on their own. No detailed information on insect predators of ALB is available.

The main ALB parasitoids in Asia are larval ectoparasitoids in the genera *Dastarcus* (Coleoptera: Bothrideridae) and *Scleroderma* (Hymenoptera: Bethylidae) (Golec et al. 2018; Wang et al. 2021a). Nevertheless, more than 20 parasitoid species associated with ALB have been reported in China and Korea (Wang et al. 2021a). *Dastarcus helophoroides* (= *D. longulus*) is an important natural enemy of ALB, CLB and other long-horned beetles in China, Japan, and Korea (Golec et al. 2018). However, *Dastarcus* and *Scleroderma* species native to Asia that attack ALB and CLB have broad host ranges, and their release as biological control agents is unlikely to be approved in Europe or North America (Meng et al. 2015; Gould et al. 2018). In a recent survey using sentinel logs with ALB larvae, *Oxysychus* sp. (Hymenoptera: Pteromalidae) and *Bracon planitibiae* Yang, Cao et Gould (Hymenoptera: Braconidae) were the most abundant parasitoids species recovered (Li et al 2020). Further studies are underway to assess their potential as biological control agents against ALB.

DRAFT

Regarding parasitoids of non-Asian origin, Lupi et al. (2017) tested the reproductive performance of *Sclerodermus brevicornis* (Kieffer), a bethylid wasp native to Europe, reared on ALB and CLB larvae. Based on their results, the authors suggest that *S. brevicornis* has the potential to be efficiently mass-reared and actively deployed in the biological control of these two longhorn beetles (Lupi et al. 2017). Also in Europe, eight species of idiobiont ectoparasitoids were discovered attacking both CLB and ALB, all of which were already known from other cerambycid hosts (Hérard et al. 2013; Maspero, 2015). The two species most frequently found were *Spathius erythrocephalus* Wesmael (Hymenoptera: Braconidae) and *Trigonoderus princeps* Westwood (Hymenoptera: Pteromalidae) (Hérard et al. 2013; Brabbs et al., 2015). Their mass release was so far not considered due to their wide host range (Hérard et al. 2013). In North America, several groups of native braconid parasitoids were found to be capable of attacking ALB larvae in laboratory trials (Duan et al. 2016). *Ontsira mellipes* Ashmead was shown to be the most promising species: it can be reared continuously with short generation times and produces a high female-biased progeny with rapidly maturing eggs (Duan et al. 2016; Golec et al. 2016; Wang and Aparicio 2020; Wang et al. 2020). In a study conducted to assess the potential host range and preferences of *O. mellipes*, this braconid successfully attacked ALB and CLB as well as three of six tested longhorned beetles native to North America (Wang et al. 2019). Field trials to assess the potential of *O. mellipes* to effectively reduce ALB populations are being carried out in Worcester, Massachusetts (USDA-APHIS 2021).

An egg parasitoid native to Asia that attacks CLB, *Aprostocetus fukutai* (Hymenoptera: Eulophidae), was detected in Northern Italy in 2002 and initially described as a new species, *Aprostocetus anoplophorae* (Delvare et al. 2004; Hérard et al., 2017). The parasitoid is thought to have accidently been introduced in Italy from Japan with bonsais containing parasitized CLB eggs (Brabbs et al. 2015, Hérard et al., 2017). So far, *Aprostocetus fukutai* is regarded as the most promising biological control against CLB because i) it attains high rates of parasitism in the field of up to 72% of CLB eggs (Hérard et al. 2005a, 2013), ii) it is CLB specific and not able to parasitize ALB or the Italian native cerambycid *Saperda carcharias* L. (Coleoptera: Cerambycidae) (Herard et al. 2005 a,b), iii) it does not show specificity in terms of the host plant (Hérard et al. 2005a), iv) it is socially gregarious which facilitates the rearing procedures (Maspero 2015), and v) the host and its parasitoid have a high degree of developmental synchronicity (Hérard et al. 2013; Maspero 2015). Furthermore, the parasitoid persists even at the very low host densities that resulted from the extensive eradication efforts conducted in Northern Italy (Hérard et al., 2017; Wang et al. 2021b). For ALB, no egg parasitoids have been identified (Golec et al. 2018; Wang et al. 2021a). Under such circumstances, it has been suggested that biological control programmes should resort to the use of natural enemies native to regions where ALB has been introduced via novel associations and augmentative releases (e.g. Wang et al. 2019; Wang et al. 2021a)

DRAFT

7. Conclusion and future outlooks

Major efforts have been put into achieving successful eradication of establishments of ALB and CLB. International collaborative activities translated into legislative changes to harmonize detection and eradication as well as prevention methods towards a common goal. We conclude that these efforts have resulted in considerable success as more than 45% of eradication programmes were successful (and some are still ongoing). Several countries were able to completely eradicate all ALB and/or CLB populations, and other managed to reduce the area affected. Still, these efforts are hampered by the ongoing arrival of new beetles, both from their native regions in Asia and in some cases apparently also from other invaded regions via the bridgehead effect.

Several biological traits of ALB and CLB may have favoured eradication success, such as long-life cycles, relatively low fecundity, low spread rate and their tendency to remain in the vicinity of the natal trees unless conditions are unfavourable (Haack et al. 2010). Detectability has been identified by Tobin et al. (2014) as another factor relevant for the success of eradication programmes. Thus, the fact that ALB develops mostly in the upper part of trees and CLB in the lower trunk and roots, may translate into a higher relative detectability of ALB, which in some cases might facilitate early detection and consequently its eradication success.

Eradication campaigns have hitherto been expensive. For example, just for Lombardy in Italy, the costs of CLB eradication campaigns between 2008 and 2013 totalled almost 20 million Euros (Cavagna 2014, in Hérard and Maspero 2019). Nevertheless, although the costs of these eradication programmes can be extremely high, the benefits still outweigh inaction in most cases. For ALB, the costs of eradication campaigns undertaken between 1996 and 2013 in the USA were estimated to have exceeded US$537 million (Eyre and Haack 2017). However, estimations of potential economic loss in compensatory value, resulting from a widespread ALB outbreak could exceed US$670 billion (over one trillion US dollars, if adjusted to 2021 values) and a potential loss of approximately 35% of urban tree cover across the United States (Nowak et al. 2001). For the small ALB outbreak in Cornuda, Italy, Faccoli and Gatto (2016) estimated that during the first year of the eradication program, the ornamental value of the saved trees was six times higher than the eradication costs. Pedlar et al. (2020) estimated that the annual costs of inaction in an ALB outbreak in Eastern Canada could exceed CDN$12 billion (considering street tree-related costs, standing timber value and maple food products), which contrasts with an annual control expenditure of approximately 5% of this value (CDN$0.5 billion).

The methods used for eradication have not seen many changes during the last decade, and host removal is still the method most commonly used, with or without preventive felling. In North America, the use of preventive chemical treatment may have yielded good results in containing the spread of established populations and facilitating their eradication, yet the externalities arising from large-scale use of most

891 insecticides may outweigh the benefits of their use. On the other hand, detection methods have evolved
892 significantly during the last decade, even if visual surveys remain the "gold standard". In Fig. 11, a
893 summary of the known steps of invasions by the two longhorned beetles and the available management
894 strategies is presented.
895
896 Despite the advances of the last decade, prevention and management of ALB and CLB is still
897 challenging but not impossible. Research avenues that could be pursued further to improve prevention,
898 eradication and management include technical solutions such as sensors in containers to detect
899 infestations based on acoustic signals or VOCs signals, improving trapping methods based on the use of
900 semiochemicals, new models to predict spread particularly in urban areas, diversification of tree species
901 in urban and peri-urban areas, and citizen science programmes to improve detection and responses.

DRAFT




Figure 11 Summary of the steps of invasion and management strategies of Anoplophora spp. * in invaded range.

Acknowledgements

We also thank Françoise Petter for providing Anoplophora spp. interception data for Europe, and USDA-APHIS and CFIA for interception data for North America. This study was supported by the HOMED project (http://homed-project.eu/), which received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No. 771271. This work was

partially supported by the National Socio-Environmental Synthesis Center (SESYNC) under funding received from the National Science Foundation DBI-1639145. Both S. Branco and M. Branco are partially supported by CEF, a research unit funded by Foundation for Science and Technology (FCT), Portugal (UID/AGR/00239/2019 and (UIDB/00239/2020). M. Faccoli was partially funded by the DOR program of the University of Padua.

References

Adachi, I (1990) Population studies of *Anoplophora malasiaca* adults (Coleoptera: Cerambycidae) in a citrus grove. Res Popul Ecol *32*(1): 15-32. https://doi.org/10.1007/BF02512587

Akino T, Fukaya M, Yasui H et al (2001) Sexual dimorphism in cuticular hydrocarbons of the white-spotted longicorn beetle, *Anoplophora malasiaca* (Coleoptera: Cerambycidae). Entomol Sci 4: 271-277

Akita K, Katô T, Yanagi T, Kubota K (2021) Reports of the alien species Anoplophora glabripennis (Motschulsky, 1853) (Coleoptera, Cerambycidae) found in Hyogo pref., Japan. 月刊むし: a monthly journal of entomology (Japan) 601:41-5.

An Y, Baode W, Xiaojun Y et al (2004). Characterizing populations of *Anoplophora glabripennis* and related taxa with RAPD. Acta Entomol Sinica 47(2):229-235

Bancroft JS, Smith MT (2005) Dispersal and influences on movement for *Anoplophora glabripennis* calculated from individual mark-recapture. Entomol Exp Appl 116(2):83-92. https://doi.org/10.1111/j.1570-7458.2005.00320.x

Boyd IL, Freer-Smith PH, Gilligan CA et al (2013) The consequence of tree pests and diseases for ecosystem services. Science 15:342(6160). https://doi.org/10.1126/science.1235773

Brabbs T, Collins D, Hérard F et al (2015) Prospects for the use of biological control agents against *Anoplophora* in Europe. Pest Manag Sci 71(1):7-14. https://doi.org/10.1002/ps.3907

Brockerhoff EG, Liebhold AM (2017) Ecology of forest insect invasions. Biol Invasions 19(11):3141-3159. https://doi.org/10.1007/s10530-017-1514-1

Brockerhoff EG, Kimberley M, Liebhold AM et al (2014) Predicting how altering propagule pressure changes establishment rates of biological invaders across species pools. Ecology 95(3):594-601. https://doi.org/10.1890/13-0465.1

CABI (2020). Invasive Species Compendium - *Anoplophora chinensis*. Wallingford, UK: CAB International. www.cabi.org/isc. Accessed 2 February 2020

Carter ME, Smith MT, Harrison RG (2009a) Patterns of genetic variation among populations of the Asian longhorned beetle (Coleoptera: Cerambycidae) in China and Korea. Ann Entomol Soc Am 102(5):895-905. https://doi.org/10.1603/008.102.0516

Carter ME, Smith, MT., Turgeon JJ et al (2009b) Analysis of genetic diversity in an invasive population of Asian long-horned beetles in Ontario, Canada. Can Entomol 141(6): 582-594. https://doi.org/10.4039/n09-026

948 Carter M, Smith M, Harrison R (2010) Genetic analyses of the Asian longhorned beetle (Coleoptera,
949 Cerambycidae, *Anoplophora glabripennis*), in North America, Europe and Asia. Biol
950 Invasions 12(5):1165-1182. https://doi.org/10.1007/s10530-009-9538-9
951 Cavagna B, Ciampitti M, Bianchi A et al (2013) Lombardy Region experience to support the
952 prediction and detection strategies. J Entomol Acarol Res 45(1s):1-6
953 Clifton EH, Jaronski ST, Hajek AE (2020a) Virulence of commercialized fungal entomopathogens
954 against Asian longhorned beetle (Coleoptera: Cerambycidae). J Insect Sci. 20(2):1-6.
955 https://doi.org/10.1093/jisesa/ieaa006
956 Clifton EH, Gardescu S, Behle RW et al (2020b) Optimizing application rates of *Metarhizium*
957 *brunneum* (Hypocreales: Clavicipitaceae) microsclerotia for infecting the invasive Asian
958 Longhorned Beetle (Coleoptera: Cerambycidae). J Econ Entomol 113(6):2650-2656.
959 https://doi.org/10.1093/jee/toaa222
960 Cocquempot C, Carmignac D, Prost M (2003) Interceptions et introductions en France de Longicornes
961 asiatiques: cas des *Anoplophora glabripennis* (Motschulsky) et *chinensis* (Forster)(Coleoptera
962 Cerambycidae). Publications de la Société Linnéenne de Lyon 72(8):273-278.
963 https://doi.org/10.3406/linly.2003.13483
964 de la Vega GJ, Corley JC, Soliani, C (2020) Genetic assessment of the invasion history of *Drosophila*
965 *suzukii* in Argentina. J Pest Sci 93: 63–75. https://doi.org/10.1007/s10340-019-01149-x
966 Delvare G, Bon MC, Hérard F et al (2004) Description of *Aprostocetus anoplophorae* n. sp.
967 (Hymenoptera: Eulophidae), a new egg parasitoid of the invasive pest *Anoplophora chinensis*
968 (Förster)(Coleoptera: Cerambycidae). Ann Soc Entomol Fr 40 (3-4):227-233.
969 https://doi.org/10.1080/00379271.2004.10697421
970 Dodds KJ, Orwig DA (2011) An invasive urban forest pest invades natural environments—Asian
971 longhorned beetle in northeastern US hardwood forests. Can J For Res 41(9):1729-1742.
972 https://doi.org/10.1139/x11-097
973 Duan JJ, Aparicio E, Tatman D et al (2016) Potential new associations of North American parasitoids
974 with the invasive Asian longhorned beetle (Coleoptera: Cerambycidae) for biological control.
975 J Econ Entomol 109(2):699-704. https://doi.org/10.1093/jee/tov328
976 EFSA (European Food Safety Authority), Hoppe B, Schrader G et al (2019a) Pest survey card on
977 *Anoplophora chinensis*. EFSA Supporting Publications, EN-1749, 22 pp.
978 https://doi.org/10.2903/sp.efsa.2019.EN-1749
979 EFSA (European Food Safety Authority), Hoppe, B., Schrader G et al (2019b) Pest survey card on
980 *Anoplophora glabripennis*. EFSA Supporting Publications, 16(12): EN-1750, 30 pp.
981 https://doi.org/10.2903/sp.efsa.2019.EN-1750
982 EFSF (Etat de Fribourg, Staat Freiburg) (2020) https://www.fr.ch/sfn/energie-agriculture-et-
983 environnement/forets/le-capricorne-asiatique. Accessed 24 May 2020

Elmes A, Rogan J, Williams C et al (2019) Modeling the potential dispersal of Asian longhorned beetle using circuit theory. Prof Geogr 71(4):580-594. https://doi.org/10.1080/00330124.2019.1611458

EPPO - Global Database. https://gd.eppo.int/

EPPO (European and Mediterranean Plant Protection Organization) (2013a) PM 9/16 (1) *Anoplophora chinensis*: procedures for official control. EPPO Bull 43:518-526

EPPO (European and Mediterranean Plant Protection Organization) (2013b) PM 9/15 (1) *Anoplophora glabripennis*: Procedures for official control. EPPO Bull 43:510-517

EPPO RS (2001–2020) EPPO Reporting Service online archives http://archives.eppo.int/EPPOReporting/Reporting_Archives.html. Accessed January 2020- May 2021

Errico M (2012) Asian longhorned beetle detector dog pilot project. In: Proceedings, 23rd USDA Interagency Research Forum on Invasive Species, 10-13 January 2012, Annapolis, MD, pp 18

EU (European Commission) (2010) Final report of a mission carried out in Austria from 07 to 11 June 2010 in order to evaluate the situation and control for *Anoplophora glabripennis*. EC Health and Consumers Directorate-General, Directorate F-Food and Veterinary Office, 12 pp.

EU (European Commission) (2012) Commission Implementing Decision 2012/138/EU of 1 March 2012 as regards emergency measures to prevent the introduction into and the spread within the Union of *Anoplophora chinensis* (Forster). Off J EU 64:38–47

EU (European Commission) (2015) Commission implementing decision 2015/893 of 9 June 2015 as regards measures to prevent the introduction into and the spread within the Union of *Anoplophora glabripennis* (Motschulsky). Off J EU 146:16–28

EU (European Union) (2016) "Regulation (EU) 2016/2031 of the European Parliament of the Council of 26 October 2016 on protective measures against pests of plants, amending Regulations (EU) No 228/2013, (EU) No 652/2014 and (EU) No 1143/2014 of the European Parliament and of the Council and repealing Council Directives 69/464/EEC, 74/647/EEC, 93/85/EEC, 98/57/EC, 2000/29/EC, 2006/91/EC and 2007/33/EC." Off J EU 317: 4-104

EU (European Union) (2019) "Commission Delegated Regulation (EU) 2019/1702 of 1 August 2019 supplementing regulation (EU) 2016/2031 of the European Parliament and of the council by establishing the list of priority pests. Off J EU 260: 8–10

Eyre D, Haack RA (2017) Invasive cerambycid pests and biosecurity measures. Chapter 13. In: Wang Q (ed) Cerambycidae of the world: biology and pest management. CRC Press, Boca Raton, pp 563-607

Eyre D, Macarthur R, Haack RA et al (2018) Variation in inspection efficacy by member states of wood packaging material entering the European Union. J Econ Entomol 111(2):707-715. https://doi.org/10.1093/jee/tox357

DRAFT

1019 FAO (2019) Glossary of phytosanitary terms. International Standard for Phytosanitary Measures No.
1020 5. Published by FAO on behalf of the Secretariat of the International Plant Protection
1021 Convention (IPPC), Rome
1022 Faccoli M, Gatto P (2016) Analysis of costs and benefits of Asian longhorned beetle eradication in
1023 Italy. Forestry 89(3):301-309. https://doi.org/10.1093/forestry/cpv041
1024 Faccoli M, Favaro R, Concheri G et al (2016) Tree colonization by the Asian longhorn beetle,
1025 *Anoplophora glabripennis* (Coleoptera: Cerambycidae): effect of habitat and tree suitability.
1026 Insect Sci 23(2):288-296. https://doi.org/10.1111/1744-7917.12192
1027 Fallon DJ, Solter LF, Keena M et al (2004) Susceptibility of Asian longhorned beetle, *Anoplophora*
1028 *glabripennis (*Motchulsky) (Coleoptera: Cerambycidae) to entomopathogenic nematodes. Biol
1029 Control 30(2):430-438. https://doi.org/10.1016/j.biocontrol.2003.12.002
1030 Favaro R, Wichmann L, Ravn HP, Faccoli M (2015) Spatial spread and infestation risk assessment in
1031 the Asian longhorned beetle, *Anoplophora glabripennis*. Entomol Exp Appl 155(2):95-101.
1032 https://doi.org/10.1111/eea.12292
1033 Fisher JJ, Castrillo LA, Donzelli BG, Hajek AE (2017) Starvation and imidacloprid exposure
1034 influence immune response by *Anoplophora glabripennis* (Coleoptera: Cerambycidae) to a
1035 fungal pathogen. J Econ Entomol 110(4):1451-1459. https://doi.org/10.1093/jee/tox124
1036 Fisher JJ, Hajek AE (2014) Thermoregulatory behavior and fungal infection of *Anoplophora*
1037 *glabripennis* (Coleoptera: Cerambycidae). Environ Entomol 43(2):384-392.
1038 https://doi.org/10.1603/EN13267
1039 Fisher JJ, Hajek AE (2016) Influence of mating and age on susceptibility of the beetle *Anoplophora*
1040 *glabripennis* to the fungal pathogen *Metarhizium brunneum*. J Invertebr Pathol 136:142-148.
1041 https://doi.org/10.1016/j.jip.2016.04.004
1042 Fournier RE, Turgeon JJ (2017) Surveillance during monitoring phase of an eradication programme
1043 against *Anoplophora glabripennis* (Motschulsky) guided by a spatial decision support system.
1044 Biol Invasions 19(10):3013-3035. https://doi.org/10.1007/s10530-017-1505-2
1045 Fragnière Y, Forster B, Hölling D et al (2018) A local risk map using field observations of the Asian
1046 longhorned beetle to optimize monitoring activities. J Appl Entomol 142(6):578-588.
1047 https://doi.org/10.1111/jen.12491
1048 Goble TA, Rehner SA, Long SJ et al (2014) Comparing virulence of North American *Beauveria*
1049 *brongniartii* and commercial pathogenic fungi against Asian longhorned beetles. Biol Control
1050 72:91-97. https://doi.org/10.1016/j.biocontrol.2014.02.006
1051 Goble TA, Gardescu S, Fisher JJ et al (2016) Conidial production, persistence and pathogenicity of
1052 hydromulch formulations of *Metarhizium brunneum* F52 microsclerotia under forest
1053 conditions. Biol Control 95:83-93. https://doi.org/10.1016/j.biocontrol.2016.01.003
1054 Golec JR, Duan JJ, Aparicio E et al (2016) Life history, reproductive biology, and larval development
1055 of *Ontsira mellipes* (Hymenoptera: Braconidae), a newly associated parasitoid of the invasive

1056 Asian longhorned beetle (Coleoptera: Cerambycidae). J Econ Entomol 109(4):1545-1554.
1057 https://doi.org/10.1093/jee/tow122
1058 Golec JR, Li F, Cao L et al (2018) Mortality factors of *Anoplophora glabripennis* (Coleoptera:
1059 Cerambycidae) infesting *Salix* and *Populus* in central, northwest, and northeast China. Biol
1060 Control 126:198-208. https://doi.org/10.1016/j.biocontrol.2018.05.015
1061 Gould JR, Aflague B, Murphy TC et al (2018) Collecting nontarget wood-boring insects for host-
1062 specificity testing of natural enemies of cerambycids: a case study of *Dastarcus helophoroides*
1063 (Coleoptera: Bothrideridae), a parasitoid of the Asian longhorned beetle (Coleoptera:
1064 Cerambycidae). Environ Entomol 47(6):1440-1450. https://doi.org/10.1093/ee/nvy121
1065 Gugliuzzo A, Biedermann PHW, Carrillo D et al (2021) Recent advances toward the sustainable
1066 management of invasive *Xylosandrus ambrosia* beetles. J Pest Sci 94:615–637.
1067 https://doi.org/10.1007/s10340-021-01382-3
1068 Haack RA (2006) Exotic bark-and wood-boring Coleoptera in the United States: recent establishments
1069 and interceptions. Can J For Res 36(2):269-288. https://doi.org/10.1139/x05-249
1070 Haack RA, Hérard F, Sun J, Turgeon JJ (2010) Managing invasive populations of Asian longhorned
1071 beetle and citrus longhorned beetle: a worldwide perspective. Ann Rev Entomol 55:521-546.
1072 https://doi.org/10.1146/annurev-ento-112408-085427
1073 Haack RA, Britton KO, Brockerhoff EG et al. (2014) Effectiveness of the International Phytosanitary
1074 Standard ISPM No. 15 on reducing wood borer infestation rates in wood packaging material
1075 entering the United States. PLoS ONE 9(5):e96611.
1076 https://doi.org/10.1371/journal.pone.0096611
1077 Hansen L, Xu T, Wickham J et al (2015) Identification of a male-produced pheromone component of
1078 the Citrus longhorned beetle, *Anoplophora chinensis*. PLoS ONE 10(12):e0145355.
1079 https://doi.org/10.1371/journal.pone.0134358
1080 Hérard F, Cocquempot C, Lopez J et al (2005a) Field study to evaluate the egg parasitoid *Aprostocetus*
1081 *anoplophorae* sp. n. (Hymenoptera: Eulophidae) on two *Anoplophora* hosts. In: Gottschalk
1082 KW (ed) Proceedings, XV US Department of Agriculture interagency research forum on
1083 gypsy moth and other invasive species 2004; 2004 January 13-16; United States Department
1084 of Agriculture, Annapolis, MD, pp 40-42
1085 Hérard F, Bon M-C, Maspero M, et al (2005b) Survey and evaluation of potential natural enemies of
1086 *Anoplophora glabripennis* and *A. chinensis*. In: Gottschalk KW (ed) Proceedings, XV US
1087 Department of Agriculture interagency research forum on gypsy moth and other invasive
1088 species 2004; 2004 January 13-16; United States Department of Agriculture, Annapolis, MD,
1089 pp 34
1090 Hérard F, Ciampitti M, Maspero M et al (2006) *Anoplophora* species in Europe: infestations and
1091 management processes 1. EPPO Bull 36(3):470-474. https://doi.org/10.1111/j.1365-
1092 2338.2006.01046.x

1093 Hérard F, Maspero M, Ramualde N (2013) Potential candidates for biological control of the Asian
1094 longhorned beetle (*Anoplophora glabripennis*) and the citrus longhorned beetle (*Anoplophora*
1095 *chinensis*) in Italy. J Entomol Acarol Res 45(1s):22
1096 Hérard F, Maspero M, Bon MC (2017) Accidental introduction into Italy and establishment of
1097 *Aprostocetus fukutai* (Hymenoptera: Eulophidae) in Citrus Longhorned Beetle infestations. In:
1098 Mason PG, Gillespie DR, Vincent C (eds) Proceedings of the 5th International Symposium on
1099 Biological Control of Arthropods, September 11–15, Langkawi, Malaysia, pp 15–18
1100 Hérard F, Maspero M (2019) History of discoveries and management of the citrus longhorned beetle,
1101 *Anoplophora chinensis*, in Europe. J Pest Sci 92(1):117-130. https://doi.org/10.1007/s10340-
1102 018-1014-9
1103 Hoover K, Keena M, Nehme M et al (2014) Sex-specific trail pheromone mediates complex mate
1104 finding behavior in *Anoplophora glabripennis*. J Chem Ecol 40(2):169-180.
1105 https://doi.org/10.1007/s10886-014-0385-5
1106 Hoyer-Tomiczek U, Sauseng G (2013) Sniffer dogs to find *Anoplophora* spp. infested plants. J
1107 Entomol Acarol Res 45(1s):10-12
1108 Hoyer-Tomiczek U, Sauseng G, Hoch G (2016) Scent detection dogs for the Asian longhorn beetle,
1109 *Anoplophora glabripennis*. EPPO Bull 46(1):148-155. https://doi.org/10.1111/epp.12282
1110 Huang J, Qu B, Fang G et al (2020) The drivers of the Asian longhorned beetle disaster show
1111 significant spatial heterogeneity. Ecol Indic 117:106680.
1112 https://doi.org/10.1016/j.ecolind.2020.106680
1113 Hu J, Angeli S, Schuetz S et al (2009) Ecology and management of exotic and endemic Asian
1114 longhorned beetle *Anoplophora glabripennis*. Agric For Entomol 11(4):359-375.
1115 https://doi.org/10.1111/j.1461-9563.2009.00443.x
1116 Hull-Sanders H, Pepper E, Davis K, Trotter III RT (2017) Description of an establishment event by
1117 the invasive Asian longhorned beetle (*Anoplophora glabripennis*) in a suburban landscape in
1118 the northeastern United States. PLoS ONE 12(7):e0181655.
1119 https://doi.org/10.1371/journal.pone.0181655
1120 IPPC (International Plant Protection Convention) (2005) International Standards for Phytosanitary
1121 Measures: ISPM 23, Guidelines for inspection. Food and Agriculture Organization of the
1122 United Nations, Rome
1123 IPPC (International Plant Protection Convention) (2008) International Standards for Phytosanitary
1124 Measures: ISPM 31, Methodologies for Sampling of Consignments. Food and Agriculture
1125 Organization of the United Nations, Rome
1126 IPPC (International Plant Protection Convention) (2009) International Standards for Phytosanitary
1127 Measures: revision of ISPM No. 15, Regulation of wood packaging material in international
1128 trade. Food and Agriculture Organization of the United Nations, Rome

1129 Javal M, Lombaert E, Tsykun T et al (2019a) Deciphering the worldwide invasion of the Asian long-
1130 horned beetle: A recurrent invasion process from the native area together with a bridgehead
1131 effect. Mol Ecol 28(5):951-967. https://doi.org/10.1111/mec.15030
1132 Javal M, Roques A, Haran J et al (2019b) Complex invasion history of the Asian long-horned beetle:
1133 fifteen years after first detection in Europe. J Pest Sci 92:173-187.
1134 https://doi.org/10.1007/s10340-017-0917-1
1135 Javal M, Roques A, Roux G, Laparie M (2018a) Respiration-based monitoring of metabolic rate
1136 following cold-exposure in two invasive *Anoplophora* species depending on acclimation
1137 regime. Comp Biochem Physiol Part A Mol Integr Physiol 216:20-27.
1138 https://doi.org/10.1016/j.cbpa.2017.10.031
1139 Javal M, Roux G, Roques A, Sauvard D (2018b) Asian Long-horned Beetle dispersal potential
1140 estimated in computer-linked flight mills. J Appl Entomol 142(1-2):282-286.
1141 https://doi.org/10.1111/jen.12408
1142 Jucker C, Valentini M, Colombo M et al (2007) *Anoplophora chinensis*-Eradication programme in
1143 Lombardia (Italy).
1144 https://www.eppo.int/ACTIVITIES/plant_quarantine/shortnotes_qps/anoplophora_chinensis_e
1145 radication. Accessed 24 May 2020
1146 Kean JM, Suckling DM, Sullivan NJ et al (2015). Global eradication and response database
1147 (GERDA). http://b3.net.nz/gerda/index.php
1148 Keena MA (2018) Factors that influence flight propensity in *Anoplophora glabripennis* (Coleoptera:
1149 Cerambycidae). Environ Entomol 47(5):1233-1241. https://doi.org/10.1093/ee/nvy100
1150 Lee S, Lee Y, Lee S (2020) Population genetic structure of *Anoplophora glabripennis* in South Korea:
1151 Invasive populations in the native range? J Pest Sci 93:1181-96.
1152 https://doi.org/10.1007/s10340-020-01245-3
1153 Lesieur V, Lombaert E, Guillemaud T et al (2019) The rapid spread of *Leptoglossus occidentalis* in
1154 Europe: a bridgehead invasion. J Pest Sci 92: 189–200. https://doi.org/10.1007/s10340-018-
1155 0993-x
1156 Liebhold AM, Kean JM (2019) Eradication and containment of non-native forest insects: successes
1157 and failures. J Pest Sci 92:83-91. https://doi.org/10.1007/s10340-018-1056-z
1158 Li W (2004) Degradation and restoration of forest ecosystems in China. For Ecol Manag 201(1):33-
1159 41. https://doi.org/10.1016/j.foreco.2004.06.010
1160 Li G, Gao R, Smith MT, Kong L (2010) Study on dispersal of *Anoplophora glabripennis* (Motsch.)
1161 (Coleoptera: Cerambycidae) population. For Res, Beijing 23(5):678-684
1162 Li ML, Wang PX, Ma F, Yang ZQ (2007) Study on the parasitic efficiency of *Dastarcus*
1163 *helophoroides* on *Anoplophora glabripennis*. J Northwest A & F University (Natural Science
1164 Edition) 35(6):152-156

Li F, Zhang YL, Wang XY et al (2020) Discovery of parasitoids of *Anoplophora glabripennis* (Coleoptera: Cerambycidae) and their seasonal abundance in China using sentinel host eggs and larvae. J Econ Entomol 113(4):1656-65. https://doi.org/10.1093/jee/toaa068

Lingafelter SW, Hoebeke ER (2002) Revision of the genus *Anoplophora* (Coleoptera: Cerambycidae). The Entomological Society of Washington, Washington, DC

Lopez VM, Hoddle MS, Francese JA et al (2017) Assessing flight potential of the invasive Asian longhorned beetle (Coleoptera: Cerambycidae) with computerized flight mills. J Econ Entomol 110(3):1070-1077. https://doi.org/10.1093/jee/tox046

Lupi D, Favaro R, Jucker C et al (2017) Reproductive biology of *Sclerodermus brevicornis*, a European parasitoid developing on three species of invasive longhorn beetles. Biol Control 105:40-48. https://doi.org/10.1016/j.biocontrol.2016.11.008

Makihara H (2002) An exotic threat from China to North America, and this affair has affected to Japan: An Asian longhorned beetle, *Anoplophora glabripennis* (Motschulsky) (Coleoptera: Cerambycidae). Nat Insects 37(3):20-22

Mankin RW, Smith MT, Tropp JM et al (2008) Detection of *Anoplophora glabripennis* (Coleoptera: Cerambycidae) larvae in different host trees and tissues by automated analyses of sound-impulse frequency and temporal patterns. J Econ Entomol 101(3):838-849. https://doi.org/10.1093/jee/101.3.838

Maspero M (2015) Managing invasive populations of *Anoplophora chinensis* and *A. glabripennis* in Lombardy. Doctoral dissertation, ALMA DL, University of Bologna. https://doi.org/10.6092/unibo/amsdottorato/7184

Meng PS, Hoover K, Keena MA (2015) Asian longhorned beetle (Coleoptera: Cerambycidae), an introduced pest of maple and other hardwood trees in North America and Europe. J Integr Pest Manag 6(1):1-12. https://doi.org/10.1093/jipm/pmv003

Myers SW, Bailey SM (2011) Evaluation of a heat treatment schedule for the Asian longhorned beetle, *Anoplophora glabripennis* (Coleoptera: Cerambycidae). For Prod J 61(1):46-49. https://doi.org/10.13073/0015-7473-61.1.46

Nehme ME, Keena MA, Zhang A et al (2009) Attraction of *Anoplophora glabripennis* to male-produced pheromone and plant volatiles. Environ Entomol 38(6):1745-1755. https://doi.org/10.1603/022.038.0628

Nehme ME, Keena MA, Zhang A et al (2010) Evaluating the use of male-produced pheromone components and plant volatiles in two trap designs to monitor *Anoplophora glabripennis*. Environ Entomol 39(1):169-176. https://doi.org/10.1603/EN09177

Nehme ME, Trotter RT, Keena MA et al (2014) Development and evaluation of a trapping system for *Anoplophora glabripennis* (Coleoptera: Cerambycidae) in the United States. Environ Entomol 43(4):1034-1044. https://doi.org/10.1603/EN14049

1201 Nei M, Maruyama T, Chakraborty R (1975) The bottleneck effect and genetic variability in
1202 populations. Evolution 29(1):1-10. https://doi.org/10.2307/2407137
1203 Nowak DJ, Pasek JE, Sequeira RA et al. (2001) Potential effect of *Anoplophora glabripennis*
1204 (Coleoptera: Cerambycidae) on urban trees in the United States. J Econ Entomol 94(1):116-
1205 22. https://doi.org/10.1603/0022-0493-94.1.116
1206 Pan HY (2005) Review of the Asian longhorned beetle: research, biology, distribution and
1207 management in China. Food and Agriculture Organization, Forestry Department. Working
1208 Paper FBS/6E. FAO, Rome. http://www.fao.org/tempref/docrep/fao/012/j6355e/j6355e00.pdf
1209 Accessed 5 June 2020
1210 Pedlar JH, McKenney DW, Yemshanov D et al (2020) Potential economic impacts of the Asian
1211 longhorned beetle (Coleoptera: Cerambycidae) in Eastern Canada. J Econ Entomol
1212 113(2):839-50. https://doi.org/10.1093/jee/toz317
1213 Pelosi C, Bertrand C, Daniele G et al (2021) Residues of currently used pesticides in soils and
1214 earthworms: A silent threat? Agric Ecosyst Environ 305:107167.
1215 https://doi.org/10.1016/j.agee.2020.107167
1216 Peng J, Liu Y (1992) Iconography of forest insects in Hunan China. Hunan Forestry
1217 Department/Institute of Zoology, Acad Sin
1218 Pimentel D (2005) Environmental and economic costs of the application of pesticides primarily in the
1219 United States. In: Pimentel D, Peshin R (eds) Integrated Pest Management. Springer,
1220 Dordrecht. https://doi.org/10.1007/978-94-007-7796-5_2
1221 Sawyer AJ (2005) Annotated Categorization of ALB Host Trees [revised]. USDA APHIS-PPQ, Otis
1222 Pest Survey Detection and Exclusion Laboratory http://www.uvm.edu/albeetle/hosts.html
1223 Accessed 4 August 2020
1224 Sawyer AJ, Panagakos WS, Horner AE et al (2011) Asian longhorned beetle, over the river and
1225 through the woods: habitat-dependent population spread. In: McManus, Katherine A,
1226 Gottschalk KW (eds) Proceedings, 21st US Department of Agriculture interagency research
1227 forum on invasive species 2010; 2010 January 12-15; United States Department of
1228 Agriculture, Annapolis, MD, pp 52-54
1229 SFRL (Servizio Fitosanitario Regione Lombardia) (2020) Boll Ufficiale. Serie Ordinaria n. 7 - 12
1230 February 2020
1231 Sjöman H, Östberg J, Nilsson J (2014) Review of host trees for the wood-boring pests *Anoplophora*
1232 *glabripennis* and *Anoplophora chinensis*: an urban forest perspective. Arboric Urban For
1233 40(3):143-64
1234 Smith MT, Bancroft J, Li G et al (2001) Dispersal of *Anoplophora glabripennis* (Cerambycidae).
1235 Environ Entomol 30(6):1036-1040. https://doi.org/10.1603/0046-225X-30.6.1036

1236 Smith MT, Tobin PC, Bancroft J et al (2004) Dispersal and spatiotemporal dynamics of Asian
1237 longhorned beetle (Coleoptera: Cerambycidae) in China. Environ Entomol 33(2):435-442.
1238 https://doi.org/10.1603/0046-225X-33.2.435
1239 Stefan M, Markham C, Benjamin R, Coath J (2014) Case study. Invasive insects in plant biosecurity:
1240 the Asian longhorned beetle eradication program. In: Gordh G, McKirdy S (eds) The
1241 Handbook of Plant Biosecurity. Springer, Dordrecht, pp. 485-517.
1242 https://doi.org/10.1007/978-94-007-7365-3_16
1243 StMELF (Bayerisches Staatsministerium für Ernährung, Landwirtschaft und Forsten) (2020)
1244 https://www.lfl.bayern.de/ips/pflanzengesundheit/024167/index.php. Accessed 3rd January
1245 2020
1246 Strangi A, Binazzi F, Peverieri GS, Roversi PF (2017) The *Anoplophora chinensis* (Förster)
1247 (Coleoptera Cerambycidae Lamiinae) outbreaks in Italy: A possible geographical origin.
1248 Redia 100:175-179. https://doi.org/10.19263/REDIA-100.17.22
1249 Straw NA, Fielding NJ, Tilbury C et al (2016) History and development of an isolated outbreak of
1250 Asian longhorn beetle *Anoplophora glabripennis* (Coleoptera: Cerambycidae) in southern
1251 England. Agric For Entomol 18(3):280-293. https://doi.org/10.1111/afe.12160
1252 Sutin A, Yakubovskiy A, Salloum HR et al (2019) Towards an automated acoustic detection algorithm
1253 for wood-boring beetle larvae (Coleoptera: Cerambycidae and Buprestidae). J Econ Entomol
1254 112(3):1327-1336. https://doi.org/10.1093/jee/toz016
1255 Takahashi N, Ito M (2005) Detection and eradication of the Asian longhorned beetle in Yokohama,
1256 Japan. Res Bull Plant Prot Serv (Japan) 41:83-85
1257 Tobin PC, Kean JM, Suckling DM et al (2014) Determinants of successful arthropod eradication
1258 programs. Biol Invasions 16:401-414. https://doi.org/10.1007/s10530-013-0529-5
1259 Trotter RT, Hull-Sanders HM (2015) Quantifying dispersal of the Asian longhorned beetle
1260 (*Anoplophora glabripennis*, Coleoptera) with incomplete data and behavioral knowledge. Biol
1261 Invasions 17:3359-3369. https://doi.org/10.1007/s10530-015-0961-9
1262 Trotter RT, Keena MA (2016) A variable-instar climate-driven individual beetle-based phenology
1263 model for the invasive Asian longhorned beetle (Coleoptera: Cerambycidae). Environ
1264 Entomol 45(6):1360-1370. https://doi.org/10.1093/ee/nvw108
1265 Trotter RT, Pepper E, Davis K, Vazquez R (2019) Anisotropic dispersal by the Asian longhorned
1266 beetle (*Anoplophora glabripennis*): quantifying spatial risk and eradication effort with limited
1267 biological data. Biol Invasions 21:1179-1195. https://doi.org/10.1007/s10530-018-1894-x
1268 Tsykun T, Javal M, Hölling D et al (2019) Fine-scale invasion genetics of the quarantine pest,
1269 *Anoplophora glabripennis*, reconstructed in single outbreaks. Sci Rep 9:19436.
1270 https://doi.org/10.1038/s41598-019-55698-3

1271 Turgeon JJ, Pedlar J, De Groot P et al (2010) Density and location of simulated signs of injury affect
1272 efficacy of ground surveys for Asian longhorned beetle. Can Entomol 142(1):80-96.
1273 https://doi.org/10.4039/n09-049
1274 Turgeon JJ, Orr M, Grant C et al (2015) Decade-old satellite infestation of *Anoplophora glabripennis*
1275 Motschulsky (Coleoptera: Cerambycidae) found in Ontario, Canada outside regulated area of
1276 founder population. Coleopt Bull 69(4):674-678
1277 Turner RM, Plank MJ, Brockerhoff EG et al (2020) Considering unseen arrivals in predictions of
1278 establishment risk based on border biosecurity interceptions. Ecological Applications 30:
1279 e02194. https://doi.org/10.1002/eap.2194
1280 Turner RM, Brockerhoff EG, Bertelsmeier C et al (2021) Worldwide border interceptions provide a
1281 window into human-mediated global insect movement. Ecological Applications,
1282 https://doi.org/10.1002/eap.2412
1283 USDA-APHIS (United States Department of Agriculture- Animal and Plant Health Inspection
1284 Service) (2014) Asian longhorned beetle response guidelines.
1285 https://www.aphis.usda.gov/plant_health/plant_pest_info/asian_lhb/downloads/response-
1286 guidelines.pdf. Accessed 5 June 2020
1287 USDA-APHIS (United States Department of Agriculture- Animal and Plant Health Inspection
1288 Service) (2020a) U.S. Regulated Plant Pest Table.
1289 https://www.aphis.usda.gov/aphis/ourfocus/planthealth/import-information/rppl/rppl-table.
1290 Accessed 10 August 2020
1291 USDA-APHIS (United States Department of Agriculture- Animal and Plant Health Inspection
1292 Service) (2020b) ALB Survey Protocol.
1293 https://www.aphis.usda.gov/aphis/ourfocus/pfecollanthealth/plant-pest-and-disease-
1294 programs/pests-and-diseases/asian-longhorned-beetle/ct_survey. Accessed 5 June 2020
1295 USDA-APHIS (United States Department of Agriculture- Animal and Plant Health Inspection
1296 Service) (2021) Plant Protection Today - PPQ Scientists Evaluate Wasp's Ability to Detect
1297 and Attack the Asian Longhorned Beetle.
1298 https://www.aphis.usda.gov/aphis/ourfocus/planthealth/ppq-program-overview/plant-
1299 protection-today/articles/alb-biocontrol. Accessed 29 July 2021
1300 Van der Gaag DJ, Sinatra G, Roversi PF et al (2010) Evaluation of eradication measures against
1301 *Anoplophora chinensis* in early stage infestations in Europe. EPPO Bull 40(2):176-187.
1302 https://doi.org/10.1111/j.1365-2338.2010.02381.x
1303 Van der Gaag DJ, Loomans AJM (2014) Host plants of *Anoplophora glabripennis*, a review. EPPO
1304 Bull 44(3):518–528. https://doi.org/10.1111/epp.12151
1305 Yemshanov D, Haight RG, Koch FH et al (2019) Optimizing surveillance strategies for early detection
1306 of invasive alien species. Ecol Econ 162:87-99.
1307 https://doi.org/10.1016/j.ecolecon.2019.04.030

1308 Xu T, Hansen L, Cha DH et al (2020a) Identification of a female-produced pheromone in a destructive
1309 invasive species: Asian longhorn beetle, *Anoplophora glabripennis*. J Pest Sci 93:1321–1332.
1310 https://doi.org/10.1007/s10340-020-01229-3
1311 Xu T, Hansen L, Teale SA (2020b) Mating and adult feeding behaviour influence pheromone
1312 production in female Asian longhorn beetle *Anoplophora glabripennis* (Coleoptera:
1313 Cerambycidae). Agric For Entomol 23:276-286. https://doi.org/10.1111/afe.12430
1314 Xu T, Teale SA (2021) Chemical ecology of the Asian Longhorn Beetle, *Anoplophora glabripennis*. J
1315 Chem Ecol 47:489–503. https://doi.org/10.1007/s10886-021-01280-z
1316 Wang B, Gao R, Mastro VC, Reardon RC (2005) Toxicity of four systemic neonicotinoids to adults of
1317 *Anoplophora glabripennis* (Coleoptera: Cerambycidae). J Econ Entomol 98(6):2292-2300.
1318 https://doi.org/10.1093/jee/98.6.2292
1319 Wang X, Aparicio EM, Murphy TC et al (2019) Assessing the host range of the North American
1320 parasitoid *Ontsira mellipes*: potential for biological control of Asian longhorned beetle. Biol
1321 Control 137:104028. https://doi.org/10.1016/j.biocontrol.2019.104028
1322 Wang JH, Che SC, Qiu LF et al (2020) Efficacy of emamectin benzoate trunk injection against the
1323 Asian long-horned beetle [*Anoplophora glabripennis* (Coleoptera: Cerambycidae)]. J Econ
1324 Entomol 113(1):340-347. https://doi.org/10.1093/jee/toz299
1325 Wang X, Aparicio EM (2020) Reproductive traits of *Ontsira mellipes* (Hymenoptera: Braconidae), a
1326 North American parasitoid, as a novel biological control agent for exotic *Anoplophora*
1327 *glabripennis* (Coleoptera: Cerambycidae). J Econ Entomol 113(5):2112-2119.
1328 https://doi.org/10.1093/jee/toaa160
1329 Wang X, Aparicio EM, Duan JJ et al (2020) Optimizing parasitoid and host densities for efficient
1330 rearing of *Ontsira mellipes* (Hymenoptera: Braconidae) on Asian longhorned beetle
1331 (Coleoptera: Cerambycidae). Environ Entomol 49(5):1041-1048.
1332 https://doi.org/10.1093/ee/nvaa086
1333 Wang X, Wang XY, Kenis M et al (2021a) Exploring the potential for novel associations of generalist
1334 parasitoids for biological control of invasive woodboring beetles. BioControl 66(1):97-112.
1335 https://doi.org/10.1007/s10526-020-10039-6
1336 Wang X, Ramualde N, Aparicio EM et al (2021b) Optimal conditions for diapause survival of
1337 *Aprostocetus fukutai*, an egg parasitoid for biological control of *Anoplophora chinensis*.
1338 Insects 12(6):535. https://doi.org/10.3390/insects12060535
1339 Wickham JD, Xu Z, Teale SA (2012) Evidence for a female-produced, long range pheromone of
1340 *Anoplophora glabripennis* (Coleoptera: Cerambycidae). Insect Sci 19:355–371.
1341 https://doi.org/10.1111/j.1744-7917.2012.01504.x
1342 Williams DW, Lee HP, Kim IK (2004a) Distribution and abundance of *Anoplophora glabripennis*
1343 (Coleoptera: Cerambycidae) in natural *Acer* stands in South Korea. Environ Entomol
1344 33(3):540-545. https://doi.org/10.1603/0046-225X-33.3.540

Williams DW, Li G, Gao R (2004b) Tracking movements of individual *Anoplophora glabripennis* (Coleoptera: Cerambycidae) adults: application of harmonic radar. Environ Entomol 33(3):644-649. https://doi.org/10.1603/0046-225X-33.3.644

WITS (World Integrated Trade Solution) (2021). http://wits.worldbank.org. Accessed 16 July 2021

Wen J, Li Y, Xia N, Luo Y (1998) Study on dispersal pattern of *Anoplophora glabripennis* adults in poplars. Acta Ecol Sin 18(3):269-277

Yasui H (2009) Chemical communication in mate location and recognition in the white-spotted longicorn beetle, *Anoplophora malasiaca* (Coleoptera: Cerambycidae). App Entomol Zool 44:183–194. https://doi.org/10.1303/aez.2009.183

Yasui H, Yasuda T, Fukaya M et al (2007) Host plant chemicals serve intraspecific communication in the white-spotted longicorn beetle, *Anoplophora malasiaca* (Thomson) (Coleoptera: Cerambycidae). App Entomol Zool 42:255-268. https://doi.org/10.1303/aez.2007.255

Yasui H, Akino T, Fukaya M et al (2008) Sesquiterpene hydrocarbons: kairomones with a releaser effect in the sexual communication of the white-spotted longicorn beetle, *Anoplophora malasiaca* (Thomson) (Coleoptera: Cerambycidae). Chemoecology 18:233–242. https://doi.org/10.1007/s00049-008-0410-7

Zhang A, Oliver JE, Aldrich JR et al (2002) Stimulatory beetle volatiles for the Asian longhorned beetle, *Anoplophora glabripennis* (Motschulsky). Z Naturforsch C57(5-6):553-558. https://doi.org/10.1515/znc-2002-5-626

Zhang A, Oliver JE, Chauhan K et al (2003) Evidence for contact sex recognition pheromone of the Asian longhorned beetle, *Anoplophora glabripennis* (Coleoptera: Cerambycidae). Naturwissenschaften 90(9):410-413. https://doi.org/10.1007/s00114-003-0452-1

Zhao Z, Reddy GVP, Chen L et al (2020) The synergy between climate change and transportation activities drives the propagation of an invasive fruit fly. J Pest Sci 93: 615–625. https://doi.org/10.1007/s10340-019-01183-9

Zorović M, Čokl A (2015) Laser vibrometry as a diagnostic tool for detecting wood-boring beetle larvae. J Pest Sci 88:107-112. https://doi.org/10.1007/s10340-014-0567-5