



HAL
open science

The simple AMG model accurately simulates organic carbon storage in soils after repeated application of exogenous organic matter

Florent Levavasseur, Bruno Mary, Bent Christensen, Annie Duparque, Fabien Ferchaud, Thomas Kätterer, Hélène Lagrange, Denis Montenach, Camille Resseguier, Sabine Houot

► To cite this version:

Florent Levavasseur, Bruno Mary, Bent Christensen, Annie Duparque, Fabien Ferchaud, et al.. The simple AMG model accurately simulates organic carbon storage in soils after repeated application of exogenous organic matter. *Nutrient Cycling in Agroecosystems*, 2020, 117 (2), pp.215-229. 10.1007/s10705-020-10065-x . hal-02895345

HAL Id: hal-02895345

<https://hal.inrae.fr/hal-02895345v1>

Submitted on 25 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The simple AMG model accurately simulates organic carbon storage in soils after repeated application of exogenous organic matter

Florent Levavasseur^{1*}, Bruno Mary², Bent T Christensen³, Annie Duparque⁴, Fabien Ferchaud², Thomas Kätterer⁵, Hélène Lagrange⁶, Denis Montenach⁷, Camille Resseguier¹,
Sabine Houot¹

¹ UMR INRA AgroParisTech ECOSYS, Université Paris-Saclay, 78850 Thiverval-Grignon,
France

² BioEcoAgro Joint Research Unit, INRAE, Université de Liège, Université de Lille,
Université de Picardie Jules Verne, 02000 Barenton-Bugny, France

³ Department of Agroecology, AU Foulum, Aarhus University, Tjele, Denmark

⁴ Agro-Transfert Ressources et Territoires, 80200 Estrées-Mons, France

⁵ Department of Ecology, Swedish University of Agricultural Sciences, Uppsala, Sweden

⁶ ARVALIS-Institut du Végétal, 31450 Baziège, France

⁷ UE INRA 0871 Service d'expérimentation Agronomique et Viticole, Colmar, France

* Corresponding author: florent.levavasseur@inra.fr

Acknowledgments

This work was performed in partnership with the SAS PIVERT (www.institut-pivert.com). It was supported by the French Government (ANR-001-01) and the Genesys WP1 P13 Solebiom project.

The QualiAgro and Colmar field experiments are part of the SOERE-PRO (a network of long-term experiments dedicated to the study of impacts of organic residue recycling), certified by

24 ALLENI (Alliance Nationale de Recherche pour l'Environnement) and integrated as a
25 service of the "Investment in the Future" infrastructure AnaEE-France, overseen by the
26 French National Research Agency (ANR-11-INBS-0001).

27 The QualiAgro experiment was founded and is still supported by INRA and Veolia R&I.

28 The Danish contribution was financially supported by the Ministry of Environment and Food.

29 We thank the Lawes Agricultural Trust and Rothamsted Research for data from the e-RA
30 database. The Rothamsted Long-term Experiments National Capability (LTE-NC) is
31 supported by the UK BBSRC (Biotechnology and Biological Sciences Research Council,
32 BBS/E/C/000J0300) and the Lawes Agricultural Trust.

33 We thank the Station d'Expérimentation Rhône-Alpes Information Légumes for providing
34 data from the SERAIL experiment.

35

36 **Abstract**

37 Repeated application of exogenous organic matter (EOM) contributes to soil organic carbon
38 (SOC) stocks in cropped soils. Simple and robust models such as the AMG model are useful
39 tools for predicting the effects of various EOM practices on SOC. In AMG, EOM is
40 characterized by a single parameter: the isohumic coefficient K_1 , which represents the
41 proportion of carbon that is humified and incorporated into SOC. The AMG model has been
42 validated under various pedoclimatic conditions and cropping systems, but has not yet been
43 tested with data from long-term field experiments where EOM is regularly applied. The
44 calibration of the EOM parameter K_1 also remains an issue. In this study, AMG was used to
45 simulate SOC stocks in seven long-term field experiments with EOM application. AMG
46 simulated changes in SOC stocks with a mean *RMSE* of 3.0 t C ha⁻¹ (the difference in SOC
47 stocks between treatments with and without EOM). The optimized K_1 values were highly
48 correlated ($R^2=0.62$) with the indicator of remaining organic carbon (I_{ROC}), which is measured
49 by laboratory analysis. The present study (i) demonstrated the ability of the AMG model to
50 accurately simulate changes in SOC stocks in long-term field experiments with regular EOM
51 application and (ii) validated the parameterization of EOM in AMG using I_{ROC} , which is
52 routinely measured by commercial laboratories. Twenty-six different EOM types representing
53 a wide range of EOM sources were parameterized using more than 600 I_{ROC} values. The
54 AMG model could thus be used to predict the SOC increase following EOM addition with a
55 very simple calibration.

56 **Keywords:** organic amendment, organic fertilizer, EOM, soil organic carbon stock, model,
57 AMG.

58 **1 Introduction**

59 Exogenous organic matter (EOM) is organic matter that is not directly derived from crops but
60 applied to soil as an organic fertilizer or organic amendment to improve soil fertility. EOM

61 includes animal manures and other organic matter from urban or industrial activities, often
62 treated to produce compost and anaerobic digestates. Depending on its characteristics, the
63 use of EOM may increase the short-term and long-term nutrient supply (Gómez-Muñoz et al.,
64 2017), increase soil organic matter (SOM) stocks (Zavattaro et al., 2017) and improve soil
65 quality (Eden et al., 2017; Obriot et al., 2016). An increase in soil organic carbon (SOC)
66 stocks also contributes to reduced atmospheric CO₂ concentration (Maillard and Angers,
67 2014) provided that EOM had not been applied to the soil before (Powlson et al., 2011).
68 Repeated EOM application in field experiments increase SOM stocks, depending on EOM
69 type and characteristics, and the amount and frequency of EOM application (Bhogal et al.,
70 2018; Maltas et al., 2018). The physico-chemical characteristics of applied EOM, such as
71 nitrogen and lignin content, affect their initial rate of decomposition, whereas their influence
72 in the longer term is uncertain (Dignac et al., 2017). Additionally, specific site conditions (e.g.,
73 soil texture and climate) influence the potential soil carbon storage.

74 Models simulating soil carbon turnover are useful for predicting and comparing different EOM
75 application practices and their efficiency in increasing SOM stocks. The simpler these
76 models are, the more likely they are to be used by farming advisors. Several models have
77 been developed over the past decades, including, the Hénin-Dupuis (Hénin and Dupuis,
78 1945), Roth-C (Jenkinson and Rayner, 1977), Century (Parton et al., 1987), ICBM (Andrén
79 and Kätterer, 1997) and C-TOOL (Taghizadeh-Toosi et al., 2014) models. The AMG model
80 (Andriulo et al., 1999) was developed in the late 1990s to improve the Hénin-Dupuis model.
81 Compared to the Roth-C and Century models, AMG uses a simpler representation of organic
82 matter with only three pools: organic inputs (crop residues, roots or EOMs); an active SOM
83 pool, which is supplied by humified organic inputs and undergoes mineralization; and a
84 stable SOM pool, taken to be inert at the considered timescale (decades). AMG has been
85 parameterized and validated for a wide range of cropping systems and pedoclimatic
86 conditions (Bouthier et al., 2014; Clivot et al., 2019; Saffih-Hdadi and Mary, 2008). It is widely

87 used in France by farming advisors, researchers, teachers and students as it has a
88 dedicated online version (<http://www.simeos-amg.org/>). However, its ability to simulate the
89 effects of repeated EOM application on the long-term evolution of SOC stocks has not yet
90 been verified. An update of the EOM parameters is also needed due to the recent
91 development of the model (AMGv2). This version involves new formalisms and changes in
92 some parameter values (Clivot et al., 2019). Moreover, because of the required increase in
93 organic waste recycling in Europe (Directive 2018/851 of the European Parliament and of the
94 Council) and in anaerobic digestion of both urban and agricultural wastes, the use of certain
95 types of EOM is increasing together with the diversity of available EOM. Therefore,
96 parameters for a wider range of EOM types are needed to facilitate AMG use by both
97 researchers and stakeholders. The parameterization of EOM has previously been linked to
98 the Roth-C model using field data from long-term experiments (Dechow et al., 2019; Peltre et
99 al., 2012). However, long-term field trials are scarce and other methods based on laboratory
100 analysis are needed to parameterize AMG. The indicator of residual organic carbon (I_{ROC})
101 (Lashermes et al., 2009) is calculated from biochemical fractions of EOM (Van Soest and
102 Wine, 1967) and the proportion of EOM carbon mineralized during 3 days of incubation with
103 soil. I_{ROC} has been defined as a predictor of the EOM residual carbon after long-term
104 incubation of EOM with soil under controlled conditions. Peltre et al. (2012) used I_{ROC} to
105 predict the partition of EOM into the different carbon pools of the Roth-C model. Preliminary
106 results also indicate its potential use for parameterizing EOM in AMG (Bouthier et al., 2014).

107 The objectives of this study were (i) to evaluate the ability of the AMG model to accurately
108 simulate SOC stock evolution in long-term field experiments where EOM is regularly applied,
109 (ii) to validate a method for EOM parameterization in AMG from laboratory analysis, and (iii)
110 to use this method to parameterize a wide range of EOM types in AMG.

111 **2 Materials and Methods**

112 **2.1 Field data**

113 Data from seven long-term field experiments were used for this study: Askov K2 in Denmark
114 (Bruun et al., 2003; Christensen and Johnston, 1997), Colmar in eastern France (Obriot,
115 2016), QualiAgro in northern France (Obriot et al., 2016; Peltre et al., 2012), La Jaillière in
116 western France (Bouthier and Trochard, 2015), SERAIL in southern France (Peltre et al.,
117 2012), Broadbalk at Rothamsted Research in southern England (Jenkinson and Rayner,
118 1977; Perryman et al., 2018) and Ultuna in central Sweden (Gerzabek et al., 1997; Karhu et
119 al., 2012; Kätterer et al., 2011). These seven experiments cover a wide range of durations,
120 climates, soils, crop rotations and EOM types (Tables 1 and 2). More details about each
121 experiment can be found in the references given above.

122 Carbon inputs from plants were divided into aboveground and belowground inputs.
123 Belowground carbon inputs (roots and exudates) were computed for each treatment and
124 year according to the measured crop yields and allometric relationships (Clivot et al., 2019)
125 adapted from Bolinder et al. (2007). For aboveground carbon, allometric relationships were
126 also used for the Askov and SERAIL experiments, whereas carbon inputs were directly
127 estimated from biomass measurements in the field for the QualiAgro, Colmar, La Jaillière,
128 Ultuna and Rothamsted experiments.

129 The EOM carbon inputs were computed from the known amount of applied EOM and the dry
130 matter and carbon content of the EOM. When these variables were not determined every
131 year, mean values of the available data were used to estimate carbon inputs from EOM.

132 The SOC stocks were calculated by using SOC content and soil mass over the sampling
133 depth, by considering an equivalent soil mass (Ellert and Bettany, 1995). When the soil bulk
134 density changed over time (QualiAgro, Ultuna and Rothamsted experiments), SOC stocks
135 were estimated by adding a variable amount of subsoil. In the Askov, Colmar, La Jaillière

136 and SERAIL experiments, bulk density remained constant and SOC stocks were computed
137 with a constant soil depth. If any rock fragments were present (as at La Jaillière), the bulk
138 density of the fine soil was used, and the rock fragment fraction was subtracted from the soil
139 volume to compute the SOC stocks, as suggested by Poeplau et al. (2017).

140 **2.2 AMG model**

141 The AMG model originally developed by Andriulo et al. (1999) simulates the dynamics of
142 SOC stocks at an annual time step in response to climate and cropping systems. A full
143 description of the AMGv2 model version used in this study is given by Clivot et al. (2019)
144 along with its validation under various pedoclimatic and cropping conditions. Only the main
145 principles are described below.

146 AMG considers three carbon pools (Figure 1): (i) a pool including carbon inputs from crop
147 residues, roots and EOM, (ii) an active carbon pool and (iii) a stable carbon pool. A fixed
148 proportion, K_1 , of the carbon inputs is humified yearly and allocated to the active carbon pool.
149 The remaining $1-K_1$ is mineralized as CO_2 . Aboveground crop residues, roots, EOM are each
150 characterized by the specific K_1 parameter (also termed isohumic coefficient). The active
151 carbon pool decomposes according to first-order kinetics with a rate constant k affected by
152 mean annual water balance and air temperature, soil clay and carbonate contents, soil pH
153 and the C to N ratio of SOM (Clivot et al., 2017). The stable carbon pool is taken to be inert
154 during the simulated period. The initial proportion of stable carbon is preset by default to 65%
155 of the total SOC in long-term arable soils. The aboveground and belowground carbon inputs
156 from plants are estimated by allometric relationships (Clivot et al., 2019) adapted from
157 Bolinder et al. (2007) or directly as measured field data if available. The stocks are computed
158 for a soil depth given as input data. AMG can be described by the following set of equations
159 (Clivot et al., 2019):

$$160 \quad QC = QC_S + QC_A \quad (1)$$

$$161 \quad \frac{dQC_A}{dt} = \sum_i m_i K_{1i} - k QC_A \quad (2)$$

162 where QC is the total SOC stock (t ha^{-1}), QC_A and QC_S are the C stocks of the active and
163 stable C pools (t ha^{-1}), respectively, m_i is the annual C input from organic residue i (t ha^{-1}
164 yr^{-1}), K_{1i} is its isohumic coefficient (the fraction of C inputs which is incorporated in SOM after
165 1 year) and k is the mineralization rate constant of the active C pool (yr^{-1})

166 **2.3 Determination of the coefficient K_1 in long-term field experiments**

167 *2.3.1 EOM K_1 optimization*

168 The K_1 value of each EOM type was estimated by fitting the model to the time series of
169 differences in SOC stocks between the EOM treatments and the control treatments, i.e., the
170 additional SOC storage in the treatments with EOM, designated ΔSOC stocks. The
171 optimization was performed in R with the “optim” function and the “L-BFGS-B” method (Byrd
172 et al., 1995) by minimizing the sum of squared differences between the observed and
173 simulated ΔSOC stocks, with K_1 varying between 0 and 100%.

174 Using ΔSOC stocks instead of SOC stocks prevents the effects of a poor parameterization of
175 the initial proportion of stable soil organic carbon, as suggested by Peltre et al. (2012). It also
176 reduces the risk of bias, since any bias that occurred in simulating the SOC stocks of the
177 control treatment is expected to also occur when simulating the EOM treatment. Beyond
178 EOM inputs, we also considered the differences in crop residue input among treatments to
179 explain ΔSOC stocks, since the carbon input from crop residues was input data specific to
180 each treatment.

181 All other parameters of the model were kept as the default values to preserve the
182 consistency of the model and the general applicability of the optimized EOM K_1 values. This
183 included the K_1 values of aboveground crop residues and roots (Table S3 in Clivot et al.,

184 2019). Finally, for the field experiments in which soil pH and soil C to N ratio changed over
185 time, these changes were considered in the model input data.

186 2.3.2 Model evaluation

187 The model performance was first evaluated in control treatments without EOM (ASK_CT+N,
188 COL_CT+N, QUA_CT-N and QUA_CT+N, LAJA_CT+N, ROTH_CT_S0 and ROTH_CT_S1,
189 SER_CT_eqC and SER_CT_eqH, ULT_CT-N and ULT_CT+N). Indeed, because organic
190 carbon from EOM is incorporated in SOC, it was important to verify the ability of the model to
191 accurately simulate the evolution of SOC stocks under the various soil, climate and cropping
192 system conditions. The mean error (*ME*), root mean square error (*RMSE*), relative *RMSE*
193 (*RRMSE*), coefficient of determination (*R*²) and model efficiency (*EF*) were thus computed
194 using the observed and simulated SOC stocks in each control treatment (Equations 3 to 7).
195 The model performance for each treatment with EOM was then assessed by using the same
196 statistical criteria.

$$197 \quad ME = \frac{1}{n} \cdot \sum_{i=1}^n (O_i - S_i) \quad (3)$$

$$198 \quad RMSE = \sqrt{\frac{1}{n} \cdot \sum_{i=1}^n (O_i - S_i)^2} \quad (4)$$

$$199 \quad RRMSE = RMSE / \bar{O} \quad (5)$$

$$200 \quad R^2 = \left(\frac{\sum_{i=1}^n ((O_i - \bar{O}) \cdot (S_i - \bar{S}))}{\sqrt{\sum_{i=1}^n (O_i - \bar{O})^2} \cdot \sqrt{\sum_{i=1}^n (S_i - \bar{S})^2}} \right)^2 \quad (6)$$

$$201 \quad EF = 1 - \frac{\sum_{i=1}^n (O_i - S_i)^2}{\sum_{i=1}^n (O_i - \bar{O})^2} \quad (7)$$

202 where *O* and *S* are the observed and simulated values, respectively, and *n* is the number of
203 observations. \bar{O} and \bar{S} are the means of the observations and simulations, respectively.

204 **2.4 Using I_{ROC} to model SOC stocks**

205 The indicator of residual organic carbon (I_{ROC}) is computed from the biochemical fractions of
206 EOM (Van Soest and Wine, 1967) and from the proportion of organic carbon mineralized
207 during 3 days of soil incubation (Lashermes et al., 2009). I_{ROC} was defined as a predictor of
208 the EOM residual carbon after long-term incubation of EOM with soil under controlled
209 conditions.

210 The I_{ROC} values of the EOM types applied in the field experiments were available only for 4
211 out of the 7 experiments: Colmar, QualiAgro (one I_{ROC} value per EOM type and per
212 application year), Ultuna (three I_{ROC} values per EOM type corresponding to three pooled
213 samples of different years of application) and SERAIL (one or two I_{ROC} values per EOM type
214 corresponding to one or two application years). When several I_{ROC} values were available for
215 a given EOM type in a field experiment, the mean of the I_{ROC} values was used as the K_1
216 value in AMG. For the 4 experiments with measured I_{ROC} values, the relationship between
217 the optimized K_1 values based on field data (section 2.3) and the I_{ROC} values of the EOM was
218 investigated by calculating the coefficient of determination (R^2), mean error (ME) and root
219 mean square error ($RMSE$). The I_{ROC} values of the EOM were also directly used as K_1 input
220 values in AMG (without any rescaling). The model performance was assessed under the
221 same conditions as previously described (section 2.3.2) and compared to the model
222 performance with optimized K_1 .

223 **2.5 The I_{ROC} database**

224 The original database developed by Lashermes et al. (2009) provided K_1 values for EOM
225 types for which we did not have any field estimates. This database was expanded with
226 additional I_{ROC} values from various research projects. Thus, the database contained more
227 than 600 EOM types from 5 main groups: livestock liquid and solid manures (n=106),
228 anaerobic digestates (n=54), sewage sludge (n=71), composts (n=352) and others (n=69).
229 Most frequently used EOM types were present in the database; however, there were limited

230 number of values for cattle slurry (n=5) considering its relative abundance and for different
231 types of digestates (from 2 to 9 depending on the type of digestates).

232 **3 Results**

233 **3.1 Evaluation of the AMG model for the control treatments**

234 The performance of the AMG model for the control treatments without EOM application was
235 good for the Colmar, QualiAgro, La Jaillière, SERAIL and Ultuna experiments. In these
236 treatments, the *RMSE* was equal to or lower than 3 t C ha⁻¹ and the *RRMSE* was lower than
237 10% (electronic supplementary material, appendices A and B). The model thus succeeded in
238 representing the decrease in SOC stocks at La Jaillière (CT_N), Ultuna (CT_N and CT),
239 SERAIL (CT_EQC and CT_EQH), and QualiAgro (CT_N-), and the nearly constant SOC
240 stocks at Colmar (CT_N) and QualiAgro (CT_N+). For Askov and Rothamsted, the model
241 performance was worse, with an *RMSE* between 4 and 5 t C ha⁻¹ in all control treatments
242 (*RRMSE* from 19% to 30%). The model failed to simulate the increase in SOC stocks at
243 Askov (CT_N) and overestimated the decrease in SOC stocks at Rothamsted (CT_S0 and
244 CT_S1). If we consider all treatments and experiments, the simulation bias (*ME*) remained
245 low, smaller than 1 t C ha⁻¹, except for the Askov, Rothamsted, Ultuna (T0) and La Jaillière
246 experiments.

247 **3.2 Simulation of Δ SOC stocks with the optimized EOM K_1 value**

248 Using the optimized values of K_1 , AMG succeeded in simulating the dynamics of Δ SOC
249 stocks in all treatments and experiments, except in two treatments at QualiAgro (BIO_N+,
250 BIO_N-) and one at SERAIL (GW_EQH) which were underestimated (Figure 2). This was
251 confirmed by the low *ME* values (Table 3), except those for the underestimated treatments
252 (*ME* higher than 3.5 t C/ha). The *RMSE* remained low, with minimum, mean and maximum
253 values of 0.5, 3.0 and 8.2 t C ha⁻¹ respectively. Except for two treatments at La Jaillière

254 (POM, CPOM), which had very low variability in Δ SOC stocks, and the SERAIL experiment,
255 for which the dynamics of Δ SOC stocks were very noisy, the R^2 values were high (>0.6) to
256 very high, indicating a good simulation of the variability of Δ SOC stocks. For example, the
257 initial increase followed by stability in Δ SOC stocks at Rothamsted was well simulated, as
258 was the initial increase at La Jaillière followed by a decrease in Δ SOC stocks after the
259 cessation of EOM application. The model efficiency (EF) exhibited the same trends as R^2 .

260 **3.3 Determination of isohumic coefficient (K_1) values from long-term field** 261 **experiments**

262 The optimized K_1 values varied greatly among EOM types. For example, the lowest value
263 was 8% for the poultry manure at La Jaillière, while values of 100% were found for several
264 EOM types including green waste and sludge compost at QualiAgro (Table 3). The mean
265 optimized K_1 for all treatments was 63%. For a given type of EOM, a large variation also
266 existed. For example, the optimized K_1 value of cattle manure varied from 44% (FYM
267 treatment at Ultuna) to 99% (FYM_N+ at QualiAgro), with a mean value of 66%.

268 **3.4 Prediction of K_1 with I_{ROC}**

269 Based on the four experiments for which I_{ROC} values of EOM were available, the I_{ROC}
270 indicator appeared to be an acceptable predictor of K_1 (Figure 3). ME and $RMSE$ between K_1
271 and I_{ROC} were low for SERAIL and Ultuna (e.g., $RMSE$ equal to 11% for Ultuna) but higher for
272 Colmar and QualiAgro (Table 4). The I_{ROC} value systematically underestimated the optimized
273 K_1 ($ME=-30\%$) for QualiAgro. Within each experiment, the variation in K_1 was well explained
274 by the variability in I_{ROC} ($R^2 = 0.73 - 0.95$). Considering all experiments together, the I_{ROC}
275 explained slightly less of the K_1 variability ($R^2 = 0.62$).

276 **3.5 Simulations of C storage from EOM with K_1 values equal to I_{ROC}**

277 The I_{ROC} values were used as input K_1 values in AMG to simulate the Δ SOC stocks for the
278 four experiments for which I_{ROC} values were available. $RMSE$ logically increased in all

279 treatments when replacing optimized K_1 with I_{ROC} (Figure 4). The mean $RMSE$ increased
280 from 3.0 to 4.9 t C ha⁻¹, with a minimal increase of 0.01 t C ha⁻¹ for the FYM_EQH treatment
281 at SERAIL and a maximal increase of 5.4 t C ha⁻¹ for the treatment PEA+N at Ultuna. The
282 largest increases in $RMSE$ were associated with experiments with very high ΔSOC stocks
283 (more than 50 t C ha⁻¹ for PEA+N at Ultuna, for example), and thus the $RMSE$ increases
284 were relatively low.

285 **3.6 Definition of a database of reference K_1 values for AMG**

286 Using the I_{ROC} database (section 2.5), some marked differences among EOM types
287 appeared, (Figure 5, appendix D in electronic supplementary material). For example, the
288 median I_{ROC} values of chicken droppings, cattle manure and green waste compost were 16%,
289 67% and 82%, respectively. Variability was also found within a given type of EOM: for
290 example, the I_{ROC} values for cattle manure ranged from 21% to 79%. Some EOM types were
291 grouped when an insufficient amount of data existed and if the differences among these
292 EOMs were low. For example, all anaerobic digestates were grouped into one type. Another
293 group was sewage sludge from urban wastewater treatment plants. Finally, we proposed a
294 default K_1 value for 26 EOM types: 9 types of livestock manure, 1 type of digestate, 2 types
295 of sewage sludge, 13 types of compost and 1 type of other EOM.

296 **4 Discussion**

297 **4.1 The accurate simulation by AMG of SOC stock evolution after repeated** 298 **applications of EOM**

299 Many authors have observed an increase in SOC stocks following EOM addition, the
300 increase being dependent on the amount and type of EOM applied (Bhogal et al., 2018;
301 Maltas et al., 2018). The importance of the type of EOM in explaining differences in SOC
302 storage was confirmed in experiments where various EOM types applied similar carbon

303 inputs (Gerzabek et al., 1997). AMG succeeded in simulating these increases in SOC stocks
304 with various EOM types and under various pedoclimatic conditions and cropping systems
305 (Figure 2), for example, in the Ultuna and QualiAgro experiments. The ability of the AMG
306 model to simulate the additional SOC stocks from EOM addition (Δ SOC stocks) was
307 satisfactory. The mean *RMSE* value (3.0 t C ha^{-1}) was comparable to the mean standard
308 deviation of the SOC measurements in the different experiments (2.1 t C ha^{-1} in QualiAgro,
309 2.5 t C ha^{-1} for Colmar). This mean *RMSE* for Δ SOC stocks was also comparable to that
310 obtained by Peltre et al. (2012) with the Roth-C model. These authors found a mean *RMSE*
311 of 3.2 t C ha^{-1} in the Askov, QualiAgro (1998-2009 only), Ultuna and SERAIL experiments.
312 Even if EOM K_1 values were optimized to minimize the *RMSE* of Δ SOC stocks, the *RMSE* of
313 the simulated SOC stocks compared to the actual SOC stocks (electronic supplementary
314 material, appendix C) was also similar to that found in other modeling studies. Karhu et al.
315 (2012) reported *RMSE* values of SOC stocks between 2.5 and 10.3 t C ha^{-1} for different
316 treatments in the Ultuna experiment (from 2.1 to 6.6 t C ha^{-1} for the same treatments in our
317 study). Begum et al. (2017) obtained an *RRMSE* of 8.9% for the FYM treatment of the
318 Rothamsted experiment (FYM_S1), compared with 13.0% obtained in our study with AMG.
319 These simulations could have been further improved if we had optimized the size of the
320 stable SOC pool. However, this would not have affected the simulation of the Δ SOC stocks.

321 Finally, the mean *RMSE* that we obtained in simulating Δ SOC stocks for all EOM types
322 (3.0 t C ha^{-1}) was close to the *RMSE* obtained for SOC stocks in the control treatments
323 without EOM in our study (2.7 t C ha^{-1}). The mean *RMSE* from simulating Δ SOC stocks was
324 also close to the mean *RMSE* obtained by Clivot et al. (2019) with the AMG model on a
325 dataset of 60 treatments located at 20 sites in France (2.6 t C ha^{-1}). However, the mean
326 *RMSE* for SOC stocks with EOM application (4.3 t C ha^{-1} , electronic supplementary material,
327 appendix C) was higher. It is recalled that EOM K_1 values were not optimized to minimize the
328 *RMSE* for SOC stocks but rather to minimize the *RMSE* for Δ SOC stocks. Indeed, the poor

329 modeling performance for the control treatments at Rothamsted and Askov also impacted the
330 *RMSE* of SOC stocks for treatments with EOM (but not that of Δ SOC stocks). The poor
331 simulation of these control treatments could be related to inappropriate allometric coefficients
332 used for old crop varieties at Rothamsted and to the very low initial C content in the subsoil
333 used at the Askov experiment, for which the SOC mineralization function was not adapted.

334 **4.2 Variability and prediction of the K_1 isohumic coefficient for EOM**

335 The optimized K_1 values showed high variability among types of EOM, from 8% to 100%,
336 with a mean value of 63%. Most of the EOM K_1 values were higher than the K_1 values of
337 aboveground residues (from 22% to 32%) and roots (40%) (Clivot et al., 2019). This finding
338 confirms that certain EOM types make a greater contribution to SOC than crop residues, as
339 suggested by several authors (Kätterer et al., 2011; Kong et al., 2005), even if the
340 continuation of this greater contribution in the long term is debated (Maillard and Angers,
341 2014).

342 The variability in EOM K_1 values well reflected the difference in stability of different EOM
343 types, as approximated by their I_{ROC} value (Figure 3). Using the I_{ROC} indicator as a predictor
344 of K_1 only slightly decreased model performance (Figure 4). The largest increases in *RMSE*
345 were associated with experiments with very high Δ SOC stocks. The small *RMSE* values
346 confirmed the usefulness of I_{ROC} for parameterizing soil carbon models, as shown by Peltre
347 et al. (2012) with Roth-C. In comparison to the parameters used for this latter study, the use
348 of I_{ROC} was easier: I_{ROC} was directly used to derive K_1 values in AMG without any rescaling,
349 whereas several regressions were needed to predict the Roth-C EOM parameters. Some
350 studies used other laboratory measurements to calibrate EOM in soil carbon models.
351 Mondini et al. (2017) used EOM incubations to predict the partition coefficients and
352 mineralization rates of EOM in Roth-C. Pansu et al. (2017) used ^{13}C nuclear magnetic
353 resonance spectra to estimate the parameters of the TAO model. In comparison to these

354 studies, the advantages of I_{ROC} are that it is a standardized (XP U44-162 standard),
355 inexpensive method, and is already being conducted by several commercial laboratories.

356 **4.3 The need for further improvement of the AMG formalism for EOM?**

357 Even if the performance of AMG in simulating the effects of EOM on SOC stocks was
358 generally satisfying, AMG failed to simulate the substantial increase in SOC stocks observed
359 in some treatments of the QualiAgro experiment (+24 t C ha⁻¹ in 20 years for the BIO_N+
360 treatment), even with the maximal K_1 value ($K_1 = 100\%$). In AMG, organic carbon from EOM
361 is incorporated into the active SOC pool the year following EOM application and is then
362 mineralized at the same rate as the native active SOC pool. The result for QualiAgro
363 suggests that SOC derived from EOM could have a slower mineralization rate than the native
364 SOC. This hypothesis is supported by other modeling studies: tests done with the STICS
365 soil-crop model showed that a slower decomposition constant was needed to accurately
366 simulate the SOC stocks in QualiAgro (Levavasseur et al., in prep). Noiro-Cosson et al.
367 (2016) also calibrated a slower decomposition rate constant for EOM than for SOC to
368 simulate the QualiAgro experiment with the CERES-EGC model. Peltre et al. (2017)
369 observed a higher thermal stability of organic matter in EOM-amended soils. Yu et al. (2012)
370 indicated a lower specific mineralization rate in EOM-amended soils, but Liu et al. (2018)
371 found the opposite result.

372 Although the isohumic coefficient K_1 was generally well predicted by the I_{ROC} indicator,
373 significant residual variability remained. Some very different K_1 values were optimized for
374 similar EOM types in different experiments. For example, the optimized K_1 for cattle manure
375 varied greatly among the different experiments. Part of this variability is certainly due to the
376 true variability in OM stability, which could be related to differences in bedding materials and
377 storage duration (Helgason et al., 2005). More surprisingly, we observed an unexplained
378 variability in the K_1 of the green waste and sludge compost at QualiAgro and Colmar.
379 Although the compost often came from the same producer, the optimized K_1 values were

380 very different, at 100% and 41%, respectively. This unexplained variability in K_1 indicates that
381 some factors not considered in the AMG model affected the storage of carbon from EOM.
382 Zhang et al. (2018) showed that accounting for the effects of litter stoichiometry and soil N
383 availability may improve the simulation of SOM formation with the CENTURY model. Peltre
384 et al. (2012) suggested that changes in soil pH may explain some of the larger simulation
385 errors obtained with Roth-C. In our case, the evolution of pH was explicitly considered in the
386 AMG model (annual soil pH was a model input). Brillì et al. (2017) also indicated that poor
387 consideration of soil water conditions may lead to erroneous SOC modeling. The simple
388 annual water balance used in the AMG model to account for the effect of climate on SOC
389 dynamics might be a cause of some poor simulations. For example, in Colmar, the effect of
390 common thunderstorms in summer is not well represented by an annual water balance.

391 Finally, even if the AMG formalisms could be improved, AMG performed as well as other
392 models in the literature despite its simpler formalisms. The simple formalisms of the AMG
393 model make it possible for it to be a calibration-free model that is already easy to use outside
394 of academic research, especially thanks to the online tool (<http://www.simeos-amg.org/>).
395 Therefore, the need to modify the formalisms of AMG to more accurately simulate the effects
396 of EOM should be carefully investigated in other field experiments from both the modeling
397 and the soil chemistry points of view.

398 **5 Conclusion**

399 Using the AMG model, we simulated the evolution of SOC stocks in seven long-term field
400 experiments with repeated applications of different EOM types. The good model performance
401 confirmed its ability to adequately simulate long-term effects of EOM application on SOC
402 stocks. The only EOM parameter in AMG, the isohumic coefficient K_1 , was optimized for each
403 EOM to minimize the difference in SOC stocks between treatments with and without EOM.
404 These optimized K_1 values were well correlated with the organic matter stability of EOM,

405 approximated by the indicator of residual organic carbon in soil (I_{ROC}). This indicator is
406 routinely analyzed by commercial laboratories. The I_{ROC} value could be used directly as the
407 EOM K_f value in AMG while maintaining satisfactory model performance. A database of I_{ROC}
408 values was used to parameterize 26 types of EOM in AMG, which can thus be used by
409 stakeholders to model the effects of different EOM practices.

410 **6 References**

411 Andrén, O., Kätterer, T., 1997. Icbm: The Introductory Carbon Balance Model for Exploration
412 of Soil Carbon Balances. *Ecol. Appl.* 7, 1226–1236. [https://doi.org/10.1890/1051-](https://doi.org/10.1890/1051-0761(1997)007[1226:ITICBM]2.0.CO;2)
413 [0761\(1997\)007\[1226:ITICBM\]2.0.CO;2](https://doi.org/10.1890/1051-0761(1997)007[1226:ITICBM]2.0.CO;2)

414 Andriulo, A., Mary, B., Guerif, J., 1999. Modelling soil carbon dynamics with various cropping
415 sequences on the rolling pampas. *Agron. Sustain. Dev.* 19, 365–377.
416 <https://doi.org/10.1051/agro:19990504>

417 Begum, K., Kuhnert, M., Yeluripati, J., Glendining, M., Smith, P., 2017. Simulating soil carbon
418 sequestration from long term fertilizer and manure additions under continuous wheat using
419 the DailyDayCent model. *Nutr. Cycl. Agroecosystems* 109, 291–302.
420 <https://doi.org/10.1007/s10705-017-9888-0>

421 Bhogal, A., Nicholson, F.A., Rollett, A., Taylor, M., Litterick, A., Whittingham, M.J., Williams,
422 J.R., 2018. Improvements in the Quality of Agricultural Soils Following Organic Material
423 Additions Depend on Both the Quantity and Quality of the Materials Applied. *Front. Sustain.*
424 *Food Syst.* 2. <https://doi.org/10.3389/fsufs.2018.00009>

425 Bolinder, M.A., Janzen, H.H., Gregorich, E.G., Angers, D.A., VandenBygaart, A.J., 2007. An
426 approach for estimating net primary productivity and annual carbon inputs to soil for common
427 agricultural crops in Canada. *Agric. Ecosyst. Environ.* 118, 29–42.
428 <https://doi.org/10.1016/j.agee.2006.05.013>

- 429 Bouthier, A., Duparque, A., Mary, B., Sagot, S., Trochard, R., Levert, M., Houot, S., Damay,
430 N., Denoroy, P., Dinh, J.-L., Blin, B., Ganteil, F., 2014. Adaptation et mise en œuvre du
431 modèle de calcul de bilan humique à long terme AMG dans une large gamme de systèmes
432 de grandes cultures et de polyculture-élevage. *Innov. Agron.* 34, 125–139.
- 433 Bouthier, A., Trochard, R., 2015. Incidence d'apports répétés d'engrais de ferme sur la
434 fertilité physique, chimique et microbiologique des sols sous rotation de cultures annuelles et
435 sous prairie temporaire. *Fourrages* 223, 211–219.
- 436 Brilli, L., Bechini, L., Bindi, M., Carozzi, M., Cavalli, D., Conant, R., Dorich, C.D., Doro, L.,
437 Ehrhardt, F., Farina, R., Ferrise, R., Fitton, N., Francaviglia, R., Grace, P., Iocola, I., Klumpp,
438 K., Léonard, J., Martin, R., Massad, R.S., Recous, S., Seddaiu, G., Sharp, J., Smith, P.,
439 Smith, W.N., Soussana, J.-F., Bellocchi, G., 2017. Review and analysis of strengths and
440 weaknesses of agro-ecosystem models for simulating C and N fluxes. *Sci. Total Environ.*
441 598, 445–470. <https://doi.org/10.1016/j.scitotenv.2017.03.208>
- 442 Bruun, S., Christensen, B.T., Hansen, E.M., Magid, J., Jensen, L.S., 2003. Calibration and
443 validation of the soil organic matter dynamics of the Daisy model with data from the Askov
444 long-term experiments. *Soil Biol. Biochem.* 35, 67–76. [https://doi.org/10.1016/S0038-](https://doi.org/10.1016/S0038-0717(02)00237-7)
445 [0717\(02\)00237-7](https://doi.org/10.1016/S0038-0717(02)00237-7)
- 446 Byrd, R., Lu, P., Nocedal, J., Zhu, C., 1995. A Limited Memory Algorithm for Bound
447 Constrained Optimization. *SIAM J. Sci. Comput.* 16, 1190–1208.
448 <https://doi.org/10.1137/0916069>
- 449 Christensen, B.T., Johnston, A.E., 1997. Chapter 18 Soil organic matter and soil quality—
450 Lessons learned from long-term experiments at Askov and Rothamsted, in: Gregorich, E.G.,
451 Carter, M.R. (Eds.), *Developments in Soil Science, Soil Quality for Crop Production and*
452 *Ecosystem Health*. Elsevier, pp. 399–430. [https://doi.org/10.1016/S0166-2481\(97\)80045-1](https://doi.org/10.1016/S0166-2481(97)80045-1)

- 453 Clivot, H., Mary, B., Valé, M., Cohan, J.-P., Champolivier, L., Piraux, F., Laurent, F., Justes,
454 E., 2017. Quantifying in situ and modeling net nitrogen mineralization from soil organic
455 matter in arable cropping systems. *Soil Biol. Biochem.* 111, 44–59.
456 <https://doi.org/10.1016/j.soilbio.2017.03.010>
- 457 Clivot, H., Mouny, J.-C., Duparque, A., Dinh, J.-L., Denoroy, P., Houot, S., Vertès, F.,
458 Trochard, R., Bouthier, A., Sagot, S., Mary, B., 2019. Modeling soil organic carbon evolution
459 in long-term arable experiments with AMG model. *Environ. Model. Softw.* 118, 99–113.
460 <https://doi.org/10.1016/j.envsoft.2019.04.004>
- 461 Dechow, R., Franko, U., Kätterer, T., Kolbe, H., 2019. Evaluation of the RothC model as a
462 prognostic tool for the prediction of SOC trends in response to management practices on
463 arable land. *Geoderma* 337, 463–478. <https://doi.org/10.1016/j.geoderma.2018.10.001>
- 464 Dignac, M.-F., Derrien, D., Barré, P., Barot, S., Cécillon, L., Chenu, C., Chevallier, T.,
465 Freschet, G.T., Garnier, P., Guenet, B., Hedde, M., Klumpp, K., Lashermes, G., Maron, P.-
466 A., Nunan, N., Roumet, C., Basile-Doelsch, I., 2017. Increasing soil carbon storage:
467 mechanisms, effects of agricultural practices and proxies. A review. *Agron. Sustain. Dev.* 37,
468 14. <https://doi.org/10.1007/s13593-017-0421-2>
- 469 Eden, M., Gerke, H.H., Houot, S., 2017. Organic waste recycling in agriculture and related
470 effects on soil water retention and plant available water: a review. *Agron. Sustain. Dev.* 37,
471 11. <https://doi.org/10.1007/s13593-017-0419-9>
- 472 Gerzabek, M. h., Pichlmayer, F., Kirchmann, H., Haberhauer, G., 1997. The response of soil
473 organic matter to manure amendments in a long-term experiment at Ultuna, Sweden. *Eur. J.*
474 *Soil Sci.* 48, 273–282. <https://doi.org/10.1111/j.1365-2389.1997.tb00547.x>
- 475 Gómez-Muñoz, B., Magid, J., Jensen, L.S., 2017. Nitrogen turnover, crop use efficiency and
476 soil fertility in a long-term field experiment amended with different qualities of urban and

- 477 agricultural waste. *Agric. Ecosyst. Environ.* 240, 300–313.
478 <https://doi.org/10.1016/j.agee.2017.01.030>
- 479 Helgason, B.L., Larney, F.J., Janzen, H.H., 2005. Estimating carbon retention in soils
480 amended with composted beef cattle manure. *Can. J. Soil Sci.* 85, 39–46.
481 <https://doi.org/10.4141/S04-049>
- 482 Hénin, S., Dupuis, M., 1945. Essai de bilan de la matière organique du sol. *Ann. Agron.* 15,
483 17–29.
- 484 Jenkinson, D.S., Rayner, J.H., 1977. The Turnover of Soil Organic Matter in Some of the
485 Rothamsted Classical Experiments. *Soil Sci.* 123. [https://doi.org/10.1097/00010694-](https://doi.org/10.1097/00010694-200606001-00018)
486 [200606001-00018](https://doi.org/10.1097/00010694-200606001-00018)
- 487 Karhu, K., Gärdenäs, A.I., Heikkinen, J., Vanhala, P., Tuomi, M., Liski, J., 2012. Impacts of
488 organic amendments on carbon stocks of an agricultural soil — Comparison of model-
489 simulations to measurements. *Geoderma* 189–190, 606–616.
490 <https://doi.org/10.1016/j.geoderma.2012.06.007>
- 491 Kätterer, T., Bolinder, M.A., Andrén, O., Kirchmann, H., Menichetti, L., 2011. Roots
492 contribute more to refractory soil organic matter than above-ground crop residues, as
493 revealed by a long-term field experiment. *Agric. Ecosyst. Environ.* 141, 184–192.
494 <https://doi.org/10.1016/j.agee.2011.02.029>
- 495 Kong, A.Y.Y., Six, J., Bryant, D.C., Denison, R.F., van Kessel, C., 2005. The Relationship
496 between Carbon Input, Aggregation, and Soil Organic Carbon Stabilization in Sustainable
497 Cropping Systems. *Soil Sci. Soc. Am. J.* 69, 1078–1085.
498 <https://doi.org/10.2136/sssaj2004.0215>
- 499 Lashermes, G., Nicolardot, B., Parnaudeau, V., Thuriès, L., Chaussod, R., Guillotin, M.L.,
500 Linères, M., Mary, B., Metzger, L., Morvan, T., Tricaud, A., Vilette, C., Houot, S., 2009.

- 501 Indicator of potential residual carbon in soils after exogenous organic matter application. Eur.
502 J. Soil Sci. 60, 297–310. <https://doi.org/10.1111/j.1365-2389.2008.01110.x>
- 503 Liu, H., Zhang, J., Ai, Z., Wu, Y., Xu, H., Li, Q., Xue, S., Liu, G., 2018. 16-Year fertilization
504 changes the dynamics of soil oxidizable organic carbon fractions and the stability of soil
505 organic carbon in soybean-corn agroecosystem. Agric. Ecosyst. Environ. 265, 320–330.
506 <https://doi.org/10.1016/j.agee.2018.06.032>
- 507 Maillard, É., Angers, D.A., 2014. Animal manure application and soil organic carbon stocks: a
508 meta-analysis. Glob. Change Biol. 20, 666–679. <https://doi.org/10.1111/gcb.12438>
- 509 Maltas, A., Kebli, H., Oberholzer, H.R., Weisskopf, P., Sinaj, S., 2018. The effects of organic
510 and mineral fertilizers on carbon sequestration, soil properties, and crop yields from a long-
511 term field experiment under a Swiss conventional farming system. Land Degrad. Dev. 29,
512 926–938. <https://doi.org/10.1002/ldr.2913>
- 513 Mondini, C., Cayuela, M.L., Sinicco, T., Fornasier, F., Galvez, A., Sánchez-Monedero, M.A.,
514 2017. Modification of the RothC model to simulate soil C mineralization of exogenous organic
515 matter. Biogeosciences 14, 3253–3274. <https://doi.org/10.5194/bg-14-3253-2017>
- 516 Noirot-Cosson, P.E., Vaudour, E., Gilliot, J.M., Gabrielle, B., Houot, S., 2016. Modelling the
517 long-term effect of urban waste compost applications on carbon and nitrogen dynamics in
518 temperate cropland. Soil Biol. Biochem. 94, 138–153.
519 <https://doi.org/10.1016/j.soilbio.2015.11.014>
- 520 Obriot, F., 2016. Epannage de produits résiduels organiques et fonctionnement biologique
521 des sols: De la quantification des impacts sur les cycles carbone et azote à l'évaluation
522 multicritère de la pratique à l'échelle de la parcelle (PhD thesis). AgroParisTech.
- 523 Obriot, F., Stauffer, M., Goubard, Y., Cheviron, N., Peres, G., Eden, M., Revallier, A.,
524 Vieublé-Gonod, L., Houot, S., 2016. Multi-criteria indices to evaluate the effects of repeated

525 organic amendment applications on soil and crop quality. *Agric. Ecosyst. Environ.* 232, 165–
526 178. <https://doi.org/10.1016/j.agee.2016.08.004>

527 Pansu, M., Thuriès, L.J.-M., Soares, V.F., Simões, M.L., Neto, L.M., 2017. Modelling the
528 transformation of organic materials in soil with nuclear magnetic resonance spectra. *Eur. J.*
529 *Soil Sci.* 68, 90–104. <https://doi.org/10.1111/ejss.12405>

530 Parton, W.J., Schimel, D.S., Cole, C.V., Ojima, D.S., 1987. Analysis of Factors Controlling
531 Soil Organic Matter Levels in Great Plains Grasslands 1. *Soil Sci. Soc. Am. J.* 51, 1173–
532 1179. <https://doi.org/10.2136/sssaj1987.03615995005100050015x>

533 Peltre, C., Christensen, B.T., Dragon, S., Icard, C., Kätterer, T., Houot, S., 2012. RothC
534 simulation of carbon accumulation in soil after repeated application of widely different organic
535 amendments. *Soil Biol. Biochem.* 52, 49–60. <https://doi.org/10.1016/j.soilbio.2012.03.023>

536 Peltre, C., Gregorich, E.G., Bruun, S., Jensen, L.S., Magid, J., 2017. Repeated application of
537 organic waste affects soil organic matter composition: Evidence from thermal analysis, FTIR-
538 PAS, amino sugars and lignin biomarkers. *Soil Biol. Biochem.* 104, 117–127.
539 <https://doi.org/10.1016/j.soilbio.2016.10.016>

540 Perryman, S.A.M., Castells-Brooke, N.I.D., Glendining, M.J., Goulding, K.W.T., Hawkesford,
541 M.J., Macdonald, A.J., Ostler, R.J., Poulton, P.R., Rawlings, C.J., Scott, T., Verrier, P.J.,
542 2018. The electronic Rothamsted Archive (e-RA), an online resource for data from the
543 Rothamsted long-term experiments. *Sci. Data* 5, 180072.
544 <https://doi.org/10.1038/sdata.2018.72>

545 Poeplau, C., Vos, C., Don, A., 2017. Soil organic carbon stocks are systematically
546 overestimated by misuse of the parameters bulk density and rock fragment content. *SOIL* 3,
547 61–66. <https://doi.org/10.5194/soil-3-61-2017>

- 548 Powelson, D.S., Whitmore, A.P., Goulding, K.W.T., 2011. Soil carbon sequestration to
549 mitigate climate change: a critical re-examination to identify the true and the false. *Eur. J.*
550 *Soil Sci.* 62, 42–55. <https://doi.org/10.1111/j.1365-2389.2010.01342.x>
- 551 Saffih-Hdadi, K., Mary, B., 2008. Modeling consequences of straw residues export on soil
552 organic carbon. *Soil Biol. Biochem.* 40, 594–607.
553 <https://doi.org/10.1016/j.soilbio.2007.08.022>
- 554 Taghizadeh-Toosi, A., Christensen, B.T., Hutchings, N.J., Vejlin, J., Kätterer, T., Glendining,
555 M., Olesen, J.E., 2014. C-TOOL: A simple model for simulating whole-profile carbon storage
556 in temperate agricultural soils. *Ecol. Model.* 292, 11–25.
557 <https://doi.org/10.1016/j.ecolmodel.2014.08.016>
- 558 Van Soest, P.J., Wine, R.H., 1967. Use of detergents in the analysis of fibrous feeds. IV.
559 Determination of plant cell-wall constituents. *J. Assoc. Off. Anal. Chem.* 50, 50–55.
- 560 Yu, H., Ding, W., Luo, J., Geng, R., Ghani, A., Cai, Z., 2012. Effects of long-term compost
561 and fertilizer application on stability of aggregate-associated organic carbon in an intensively
562 cultivated sandy loam soil. *Biol. Fertil. Soils* 48, 325–336. [https://doi.org/10.1007/s00374-](https://doi.org/10.1007/s00374-011-0629-2)
563 [011-0629-2](https://doi.org/10.1007/s00374-011-0629-2)
- 564 Zavattaro, L., Bechini, L., Grignani, C., van Evert, F.K., Mallast, J., Spiegel, H., Sandén, T.,
565 Pecio, A., Giráldez Cervera, J.V., Guzmán, G., Vanderlinden, K., D’Hose, T., Ruyschaert,
566 G., ten Berge, H.F.M., 2017. Agronomic effects of bovine manure: A review of long-term
567 European field experiments. *Eur. J. Agron.* 90, 127–138.
568 <https://doi.org/10.1016/j.eja.2017.07.010>
- 569 Zhang, H., Goll, D.S., Manzoni, S., Ciais, P., Guenet, B., Huang, Y., 2018. Modeling the
570 effects of litter stoichiometry and soil mineral N availability on soil organic matter formation

571 using CENTURY-CUE (v1.0). Geosci. Model Dev. 11, 4779–4796.

572 <https://doi.org/10.5194/gmd-11-4779-2018>

573

574 **7 Tables**

575 Table 1 Main general characteristics of the experiments used in this study

Experiment	Location	Considered period	Soil type (FAO)	Sampling depth (cm)	Soil clay content (%)	Soil CaCO ₃ content (%)	Mean initial SOC stock (t C ha ⁻¹)	Mean annual temperature (°C)	Mean annual P-PET (mm)	Crop rotation
Askov (ASK)	Askov, Denmark (55°28' N, 9°55' E)	1956-1986 (31 years)	Dystric Arenosol	25	3	0	11	7.5	466	Silage maize – spring barley – fiber flax – winter wheat
Colmar (COL)	Colmar, France (48°06' N, 7°33' E)	2000-2013 (14 years)	Calcisol	28	18	12	45	11.4	-252	Grain maize – winter wheat – sugar beet - spring barley
QualiAgro (QUA)	Feucherolles, France (48°52' N, 1°57' E)	1998-2017 (20 years)	Luvisol	29	17	0	43	10.7	44	Grain maize – winter wheat
La Jaillièrè 2 (LAJA)	Loireauxence, France (47°45' N, 0°96' W)	1995-2009 (15 years)	Gleyic Cambisol	25	21	0	37	12.6	53	Silage maize – winter wheat
Rothamsted Broadbalk (ROTH)	Harpenden, UK (51°48' N, 0°22' W)	1843-2010 (168 years)	Chromic Luvisol	23	28	2	29	9.1	80	Winter wheat
SERAIL (SER)	Brindas, France (45°43' N, 4°42' E)	1995-2009 (15 years)	Luvisol	27	17	0	38	12.9	-135	Vegetables in rotation
Ultuna (ULT)	Uppsala, Sweden (59°82' N, 17°65' E)	1956-1991 (36 years)	Eutric Cambisol	20	37	0	43	5.4	-39	Several years of spring cereals (oat, wheat, barley) followed by one year of rape or mustard

576

577

578 Table 2 Description of the treatments in the different experiments used in this study

Experiment	Treatment	EOM type	Mean EOM amount (t C ha ⁻¹)	EOM return period (year)	Mineral fertilization or straw management differences
Askov (ASK)	FYM	Cattle farmyard manure	2.3	1	Mineral N fertilization
	PEA	White sphagnum peat	3.2	1	Mineral N fertilization
	SAW	Sawdust	3.2	1	Mineral N fertilization
	STR	Cereal straw	3.0	1	Mineral N fertilization
	CT+N	-	-	-	Mineral N fertilization
Colmar (COL)	GWS+N	Green waste and sewage sludge cocompost	2.2	2	Mineral N fertilization
	CT+N	-	-	-	Mineral N fertilization
QualiAgro (QUA)	BIO-N	Compost of green waste and home-sorted organic fraction of municipal solid waste	3.6	2	Low mineral N fertilization
	BIO+N	Compost of green waste and home-sorted organic fraction of municipal solid waste	3.6	2	Optimal mineral N fertilization
	GWS-N	Green waste and sewage sludge cocompost	4.0	2	Low mineral N fertilization
	GWS+N	Green waste and sewage sludge cocompost	4.0	2	Optimal mineral N fertilization
	MSW-N	Compost of mechanically separated organic fractions from residual waste after selective collection of dry and clean packaging	3.5	2	Low mineral N fertilization
			3.5	2	Optimal mineral N fertilization
	FYM-N	Cattle farmyard manure	3.7	2	Low mineral N fertilization
			3.7	2	Optimal mineral N fertilization
	CT-N	-	-	-	Low mineral N fertilization
	CT+N	-	-	-	Optimal mineral N fertilization
La Jaillièrè 2 (LAJA)	CM	Cattle farmyard manure	2.4	1 (until 2004)	Mineral P fertilization
	PIM	Pig farmyard manure	1.8	1 (until 2004)	Mineral P fertilization
	POM	Poultry farmyard manure	1.6	1 (until 2004)	Mineral P fertilization
	CCM	Composted cattle farmyard manure	1.9	1 (until 2004)	Mineral P fertilization
	CPIM	Composted pig farmyard manure	1.7	1 (until 2004)	Mineral P fertilization
	CPOM	Composted poultry farmyard manure	1.6	1 (until 2004)	Mineral P fertilization

Experiment	Treatment	EOM type	Mean EOM amount (t C ha ⁻¹)	EOM return period (year)	Mineral fertilization or straw management differences
	CT+N	-	-	-	Mineral NP fertilization
Rothamsted Broadbalk (ROTH)	FYM_S0	Cattle farmyard manure	3.0	1	Section 0: straw incorporated since 1986
	FYM_S1	Cattle farmyard manure	3.0	1	Section 1: straw removed
	CT_S0	-	-	-	Section 0: straw incorporated since 1986
	CT_S1	-	-	-	Section 1: straw removed
SERAIL (SER)	FUM_eqC	Cattle farmyard manure	2.6	1	P K Ca Mg
	FUM_eqH	Cattle farmyard manure	2.6	1	P K Ca Mg
	FUMT_eqC	Pelletized cattle farmyard manure	2.6	1	P K Ca Mg
	FUMT_eqH	Pelletized cattle farmyard manure	2.7	1	P K Ca Mg
	CDV_eqC	Green waste compost	2.6	1	P K Ca Mg
	CDV_eqH	Green waste compost	1.6	1	P K Ca Mg
	ALGO_eqC	Algoforestier: compost of bark, poultry manure, liquid manure and algae	2.6	1	P K Ca Mg
	ALGO_eqH	Algoforestier: compost of bark, poultry manure, liquid manure and algae	2.0	1	P K Ca Mg
	VEGET_eqC	Végethumus: compost of coffee pulp cakes (90%), sheep manure and wool waste	2.6	1	P K Ca Mg
	VEGET_eqH	Végethumus: compost of coffee pulp cakes (90%), sheep manure and wool waste	1.4	1	P K Ca Mg
	FUM_eqH	Cattle farmyard manure	2.6	1	P K Ca Mg
	CT_eqC	-	-	-	P K Ca Mg
	CT_eqH	-	-	-	P K Ca Mg
Ultuna (ULT)	FYM	Cattle farmyard manure	3.8	2	-
	FYM+P	Cattle farmyard manure	3.8	2	P
	GM	Green manure	3.5	2	-
	PEA	White sphagnum peat	3.9	2	-
	PEA+N	White sphagnum peat	3.9	2	Mineral N
	SAW	Sawdust	3.7	2	-
	SAW+N	Sawdust	3.7	2	Mineral N
	SLU	Anaerobically digested sewage sludge	3.7	2	-
	STR	Cereal straw	3.5	2	-
	STR+N	Cereal straw	3.5	2	Mineral N
	CT-N	-	-	-	-

Experiment	Treatment	EOM type	Mean EOM amount (t C ha ⁻¹)	EOM return period (year)	Mineral fertilization or straw management differences
	CT+N	-	-	-	Mineral N

579

580

581 Table 3 Model performance in simulating Δ SOC stocks in the treatments with EOM in the seven long-
582 term field experiments. The optimized value of the isohumic coefficient K_1 is given for each EOM type

Experiment	Treatment	EOM	ME (t C ha ⁻¹)	RMSE (t C ha ^{a-1})	RRMSE (%)	R ²	EF	K ₁ (%)
ASK	FYM	Cattle manure	0.09	1.32	11	0.96	0.96	63
ASK	PEA	Peat	0.80	4.91	20	0.89	0.88	86
ASK	SAW	Sawdust	-0.22	1.92	15	0.88	0.88	42
ASK	STR	Straw	0.19	1.47	15	0.94	0.93	40
COL	GWS_N+	Green waste and sludge compost	0.08	0.50	23	0.84	0.83	43
LAJA2	CM	Cattle manure	0.01	1.03	39	0.77	0.77	47
LAJA2	PIM	Pig manure	0.01	0.53	16	0.81	0.81	46
LAJA2	POM	Poultry manure	0.08	0.78	85	0.01	-0.08	8
LAJA2	CCM	Composted cattle manure	0.02	0.87	35	0.75	0.75	47
LAJA2	CPIM	Composted pig manure	0.06	0.91	54	0.71	0.71	50
LAJA2	CPOM	Composted poultry manure	0.00	0.65	-122	0.19	0.19	15
QUA	BIO_N-	Green waste and biowaste compost	-4.21	5.82	40	0.94	0.59	100
QUA	BIO_N+	Green waste and biowaste compost	-3.50	5.50	41	0.92	0.60	100
QUA	FYM_N-	Cattle manure	0.27	1.91	16	0.93	0.92	97
QUA	FYM_N+	Cattle manure	0.10	1.69	15	0.92	0.92	99
QUA	GWS_N-	Green waste and sludge compost	-1.20	3.09	21	0.94	0.90	100
QUA	GWS_N+	Green waste and sludge compost	-0.47	3.56	26	0.89	0.86	100
QUA	MSW_N-	Municipal solid waste compost	0.16	2.39	29	0.79	0.79	83
QUA	MSW_N+	Municipal solid waste compost	0.26	2.23	34	0.79	0.78	81
ROTH	FYM_S0	Cattle manure	0.08	6.59	17	0.75	0.75	74
ROTH	FYM_S1	Cattle manure	0.00	8.20	21	0.67	0.67	78
SER	ALGO_EQC	Algoforestier compost	-0.42	3.74	39	0.55	0.53	69
SER	ALGO_EQH	Algoforestier compost	-0.18	3.83	40	0.59	0.58	98
SER	FYM_EQC	Cattle manure	-0.44	3.10	64	0.57	0.54	56
SER	FYM_EQH	Cattle manure	-0.81	3.99	90	0.31	0.21	52
SER	GW_EQC	Green waste compost	-0.23	4.08	34	0.77	0.76	100
SER	GW_EQH	Green waste compost	-4.32	7.41	61	0.75	0.29	100
SER	PFYM_EQC	Pelletized cattle manure	0.11	2.01	47	0.71	0.71	40
SER	PFYM_EQH	Pelletized cattle manure	0.07	1.55	37	0.80	0.80	40
SER	VEGET_EQC	Végéthumus compost	-1.49	5.01	79	0.18	-0.20	60
SER	VEGET_EQH	Végéthumus compost	-2.40	4.83	70	0.30	0.04	100
ULT	FYM	Cattle manure	-0.38	2.61	15	0.86	0.83	44
ULT	FYM_P	Cattle manure	-0.32	2.42	13	0.89	0.87	46
ULT	GM	Green manure	-0.42	2.16	21	0.77	0.66	28
ULT	PEA	Peat	-0.31	4.16	14	0.88	0.88	78
ULT	PEA_N	Peat	0.07	4.52	13	0.90	0.90	86
ULT	SAW	Sawdust	-0.10	1.61	12	0.91	0.91	38
ULT	SAW_N	Sawdust	-0.06	1.62	11	0.91	0.91	37
ULT	SLU	Sludge	-0.86	5.46	19	0.73	0.66	66

Experiment	Treatment	EOM	ME (t C ha ⁻¹)	RMSE (t C ha ^{a-1})	RRMSE (%)	R ²	EF	K ₁ (%)
ULT	STR	Straw	-0.37	1.79	21	0.63	0.45	21
ULT	STR_N	Straw	-0.12	1.32	13	0.87	0.86	27
Min	-	-	-4.32	0.50	-122	0.01	-0.20	8
Mean	-	-	-0.50	3.00	28	0.74	0.67	63
Max	-	-	0.80	8.20	90	0.96	0.96	100

583

584 Table 4 Mean error (*ME*), root mean square error (*RMSE*) and coefficient of determination (*R*²)
 585 between the *I*_{ROC} indicator and the optimized *K*₁ for all treatments of experiments for which *I*_{ROC} values
 586 were available

Experiment	ME (%)	RMSE (%)	R ²
COL	24	24	-
QUA	-30	30	0.95
SER	-3	13	0.73
ULT	-2	11	0.73
All	-9	19	0.62

587

588

589 **8 Figure captions**


590 Figure 1 Conceptual diagram of the AMG model. The organic carbon inputs (m) from crop residues,
591 roots or EOM are either mineralized ($1-K_1$ fraction) or incorporated (K_1 fraction) into the active soil
592 organic carbon pool (Active C, QC_A). The active organic carbon pool mineralizes according to first-
593 order kinetics with the k decay constant. The stable organic carbon pool (Stable C, QC_S) is considered
594 completely inert (created with LibreOffice Draw)


595 Figure 2 Observed (dots) and simulated (lines) differences in SOC stocks (Δ SOC stocks) between the
596 treatments with EOM and without EOM (controls) for the seven long-term field experiments (created
597 with R)

598 Figure 3 Relationship between the optimized value of the isohumic coefficient K_1 and the I_{ROC} indicator
599 in the long-term field experiments of Colmar (COL), QualiAgro (QUA), SERAIL (SER) and Ultuna
600 (ULT). Error bars around the mean I_{ROC} values represent the standard deviations (when available)
601 (created with R)


602 Figure 4 Root mean square error ($RMSE$) between simulated and observed differences in SOC stocks
603 in treatments with and without EOM (Δ SOC stocks) in four long-term experiments. The isohumic K_1
604 coefficient was either optimized (black bars) or equal to I_{ROC} (gray bars) (created with R)

605 Figure 5 Distribution of I_{ROC} values for certain EOM types from the database and median values of I_{ROC}
606 proposed as reference K_1 values for AMG (in red). n represents the number of I_{ROC} values for a given
607 EOM. DIG=anaerobic digestate, SS=urban sewage sludge, CM=cattle manure, HM=horse manure,
608 PIS=pig slurry, CD=chicken droppings, GWC=green waste compost, CCM=composted cattle manure
609 (created with R)




ASK


FYM
PEA
SAW
STR

COL


GWS_N+

LAJA2


CCM
CPIM
CPOM
CM
PIM
POM

QUA


BIO_N+
BIO_N-
GWS_N+
GWS_N-
FYM_N+
FYM_N-
MSW_N+
MSW_N-

ROTH


FYM_S0
FYM_S1

SER


FYM_EQC
PFYM_EQC
GW_EQC
ALGO_EQC
VEGET_EQC
FYM_EQH
PFYM_EQH
GW_EQH
ALGO_EQH
VEGET_EQH

ULT

FYM
FYM_P
GM
PEA
PEA_N
SAW
SAW_N
SLU
STR
STR_N



RMSE Δ SOC stocks (t C/ha)



■ optimized h □ h=lROC

