

Data on the link between genomic integration of IS1548 and lineage of the strain obtained by bioinformatic analyses of sequenced genomes of Streptococcus agalactiae available at the National Center for Biotechnology Information database

Sarah Khazaal, Rim Al Safadi, Dani Osman, Aurélia Hiron, Philippe Gilot

▶ **To cite this version:**

Sarah Khazaal, Rim Al Safadi, Dani Osman, Aurélia Hiron, Philippe Gilot. Data on the link between genomic integration of IS1548 and lineage of the strain obtained by bioinformatic analyses of sequenced genomes of Streptococcus agalactiae available at the National Center for Biotechnology Information database. Data in Brief, 2020, 28, 11 p. 10.1016/j.dib.2019.105066 . hal-02905882

HAL Id: hal-02905882
https://hal.inrae.fr/hal-02905882

Submitted on 23 Jul 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Data in brief 28 (2020) 105066

Contents lists available at ScienceDirect

Data in brief

journal homepage: www.elsevier.com/locate/dib

Data Article

Data on the link between genomic integration of IS*1548* and lineage of the strain obtained by bioinformatic analyses of sequenced genomes of *Streptococcus agalactiae* available at the National Center for Biotechnology Information database

Sarah Khazaal [a, b], Rim Al Safadi [b], Dani Osman [b], Aurélia Hiron [a], Philippe Gilot [a, *]

[a] *ISP, Bactéries et Risque Materno-Foetal, Université de Tours, INRA, 37032, Tours, France*
[b] *Azm Center for Research in Biotechnology and its Applications, LBA3B, EDST, Lebanese University, Tripoli, 1300, Lebanon*

ARTICLE INFO

Article history:
Received 28 November 2019
Received in revised form 12 December 2019
Accepted 18 December 2019
Available online 31 December 2019

Keywords:
Mobile genetic element
IS*As1* family
Multi locus sequence typing
Sequence type
Clonal complex
Population structure

ABSTRACT

IS*1548*, a 1316-bp element of the IS*As1* family affects the expression of several genes of the opportunistic pathogen *Streptococcus agalactiae*. Furthermore, certain lineages of *S. agalactiae* are more frequently associated to particular diseases than other [1, 2]. We took advantage of the release of the genome sequences of a huge number of epidemiologically unrelated *S. agalactiae* strains of various origin to analyze the prevalence of IS*1548* among *S. agalactiae* strains. To this end, *S. agalactiae* genome available at the National Center for Biotechnology Information (NCBI) database were blasted with IS*1548* DNA sequences. A sequence type (ST), based on the allelic profile of seven housekeeping genes, was assigned to each strain possessing IS*1548*. These strains were then grouped into clonal complexes (CCs). The data obtained will give the opportunity to compare the sequenced genomes of *S. agalactiae* based on their lineage and/or possession of IS*1548*, and to select the corresponding strains for comparative experimental

* Corresponding author.
E-mail address: philippe.gilot@univ-tours.fr (P. Gilot).

https://doi.org/10.1016/j.dib.2019.105066
2352-3409/© 2019 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

studies. The data is related to the research article « Dual and divergent transcriptional impact of IS*1548* insertion upstream of the peptidoglycan biosynthesis *murB* gene of *Streptococcus agalactiae*" [2].

© 2019 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

Specifications Table

Subject	Microbiology
Specific subject area	Bacterial population structure and dynamics
Type of data	Raw: Figs. 1–9 Raw: Table 1
How data were acquired	BlastN program (https://blast.ncbi.nlm.nih.gov/) MLST program (http://pubmlst.org/sagalactiae/) goeBURST software (http://www.phyloviz.net/goeburst/)
Data format	Raw
Parameters for data collection	All the *Streptococcus agalactiae* genome sequences available as whole genome contigs or as complete genome sequences at the National Center for Biotechnology Information database on January 19, 2018 were analyzed
Description of data collection	Bioinformatic analyses of the above cited genomes by the BlastN program, the MLST program and the goeBURST software
Data source location	Institution: University of Tours City/Town/Region: Tours Country: France
Data accessibility	With the article
Related research article	Sarah Khazaal, Rim Al Safadi, Dani Osman, Aurelia Hiron, Philippe Gilot, Dual and divergent transcriptional impact of IS*1548* insertion upstream of the peptidoglycan biosynthesis *murB* gene of *Streptococcus agalactiae*, Gene, 720, https://doi.org/10.1016/j.gene.2019.144094

Value of the Data

- As IS*1548* plays a critical role in the evolution, the virulence and the host adaption ability of *Streptococus agalactiae* [1, 2], it is important to know which sequenced strains in the huge collection of the NCBI harbor (or not) this IS. It is also useful to know to which sequence type and clonal complex belong these strains as certain lineages of *S. agalactiae* are more frequently associated to particular diseases than other.
- Researchers involved in the study of: virulence and host adaptation mechanisms of *S. agalactiae*, of mobile genetic elements, of bacterial population structure and dynamics, and of *S. agalactiae* typing, can benefit from these data.
- The data gives the possibility to compare the sequenced genomes of *Streptococcus agalactiae* strains, available at the NCBI database, based on their lineage and/or possession of IS*1548*.
- The data allows the selection of sequenced strains based on their lineage and/or possession of IS*1548* for comparative experimental studies.

1. Data description

One hundred and twenty-one *S. agalactiae* strains with IS*1548* positive blastN similarity finding were identified among the 911 strains whose genomes where sequenced and available at the NCBI database on January 19, 2018 [2]. The dataset comprises the analysis of the Multi Locus Sequence type of these strains with the MLST program (Table 1) and their classification in clonal complex with the goeburst software (Figs. 1–9), respectively.

Table 1
Raw data of the Multi Locus Sequence Typing profiles of *Streptococcus agalactiae* strains possessing IS*1548*, obtained by submitting their sequenced genomes to the MLST database[a].

Strain	*adhP*	*pheS*	*atr*	*glnA*	*sdhA*	*glcK*	*tkt*	ST
351521	1	1	3	1	1	2	2	2
418136	1	1	3	2	2	2	2	19
446329	1	1	3	2	2	2	2	19
0506A_35_952	13	3	1	3	1	1	1	22
112469214_isolate1	13	3	1	3	1	1	1	22
2603V/R	1	1	3	2	2	2	9	110
357_SAGA	13	3	1	3	1	1	1	22
986_SAGA	156	1	3	2	2	2	2	853
B37VS	1	1	43	2	2	2	2	335
B41VS	116	1	4	1	3	3	2	652
B81VD	1	1	3	4	2	5	2	106
B848	1	1	3	2	2	2	2	19
B96V	2	1	1	2	1	1	1	17
BE-PW-051	1	1	3	2	2	2	2	19
BE-PW-095	1	1	3	2	2	2	2	19
BG-NI-012	1	1	3	2	2	2	2	19
BSU167	1	1	3	2	2	2	2	19
BSU174	10	1	12	1	3	2	2	41
BSU188	5	4	6	3	2	1	3	23
BSU442	13	3	1	3	1	1	1	22
BSU447	1	1	3	2	2	2	2	19
BV3L5	1	1	3	2	2	2	9	110
CCUG 19094	1	1	3	2	2	2	2	19
CCUG 24810	1	1	3	2	2	2	2	19
CCUG 37737	1	1	3	2	2	2	2	19
CCUG 37738	1	1	3	2	2	2	2	19
CCUG 37741	1	1	3	2	2	2	2	19
CCUG 37742	1	1	3	2	2	2	2	19
CCUG 44077	1	1	2	1	1	2	2	1
CCUG 44104	1	1	3	2	2	2	2	19
CCUG 45061	1	1	3	2	2	2	2	19
CZ-NI-002	1	1	3	2	2	2	2	19
CZ-NI-003	1	1	3	2	2	2	2	19
DE-NI-007	1	1	3	2	2	2	2	19
DE-NI-041	1	1	3	4	2	5	2	106
DE-NI-042	1	1	3	2	2	2	2	19
DK-NI-011	1	1	3	2	2	2	2	19
DK-PW-060	1	1	3	2	2	2	2	19
DK-PW-066	1	1	3	2	2	2	2	19
DK-PW-162	1	1	3	2	2	2	2	19
DML120817	1	1	3	2	2	2	2	19
ES-PW-008	1	1	3	2	2	2	2	19
ES-PW-033	1	1	3	2	2	2	2	19
ES-PW-063	1	1	3	2	2	2	2	19
ES-PW-085	1	1	3	2	2	2	2	19
ES-PW-118	1	1	3	2	2	2	2	19
ES-PW-156	1	1	3	2	2	2	2	19
FSL F2-338	13	3	1	3	1	1	1	22
FSL S3-003	1	1	3	2	2	2	2	19
FSL S3-005	13	3	1	3	1	1	1	22
FSL S3-268	13	3	1	3	1	1	1	22
FSL S3-337	1	1	3	2	2	2	2	19
GB00092	1	1	3	2	2	2	2	19
GB00174	13	3	1	3	1	1	1	22

(*continued on next page*)

Table 1 (*continued*)

Strain	*adhP*	*pheS*	*atr*	*glnA*	*sdhA*	*glcK*	*tkt*	ST
GB00264	9	1	4	1	3	3	2	10
GB00543	1	1	3	2	2	2	7	36
GB00561	1	1	3	2	2	2	2	19
GB00653	10	1	4	1	3	3	2	12
GB00663	1	1	3	2	2	2	2	19
GB00864	9	1	4	1	3	3	2	10
GB00865	1	1	3	2	2	2	2	19
GB00884	1	1	3	2	2	2	2	19
GB00904	1	1	3	2	2	2	2	19
GB00923	1	1	3	3	2	2	2	175
GB00929	1	1	3	2	2	2	2	19
GB00975	13	3	1	3	1	1	1	22
GB00984	1	1	3	2	2	2	2	19
GB01004	13	3	1	3	1	1	1	22
GB-NI-007	1	1	3	2	2	2	2	19
GB-NI-011	1	1	3	2	2	2	2	19
GB-NI-014	13	3	1	3	1	1	1	22
GB-NI-015	1	1	3	2	2	2	2	19
GB-PW-035	1	1	3	2	2	2	2	19
GB-PW-041	1	1	3	2	2	2	2	19
GB-PW-049	1	1	3	2	2	2	2	19
GB-PW-051	1	1	3	2	2	2	2	19
GB-PW-067	10	1	4	1	3	3	2	12
GB-PW-087	1	1	3	2	2	2	2	19
GB-PW-088	1	1	3	2	2	2	2	19
GBS1-NY	13	3	1	3	1	1	1	22
GBS2-NM	13	3	1	3	1	1	1	22
GBS6	13	3	1	3	1	1	1	22
GBS11	1	1	3	2	2	2	2	19
Gottschalk 1003A	1	1	3	2	2	2	2	19
H002	1	1	110	2	2	2	2	928
IMMI_409	1	1	3	2	2	2	2	19
IT-NI-036	1	1	5	4	1	4	6	26
IT-NI-037	1	1	5	4	1	4	6	26
LMG 15089	1	1	3	2	2	2	2	19
LMG 15093	1	1	3	2	2	2	9	110
Madagascar-IP-47	9	1	4	1	3	3	2	10
MC627	13	3	1	3	1	1	1	22
MC628	1	1	3	2	2	2	2	19
MC632	13	3	1	3	1	1	1	22
MC633	13	3	1	3	1	1	1	22
MC634	13	3	1	3	1	1	1	22
ML41151	1	1	4	2	2	3	2	328
Mother12	1	1	3	4	2	2	2	27
MRI Z1-022	32	1	3	2	2	2	2	121
PSS_7568	1	1	3	2	2	2	2	19
PSS_7632a	1	1	3	2	2	2	2	19
RBH01	1	1	3	2	2	2	2	19
RBH03	1	1	3	2	2	2	2	19
RBH04	1	1	3	2	2	2	2	19
RBH06	13	3	1	3	1	1	1	22
RBH11	1	1	3	2	2	2	2	19
RBH12	1	1	3	2	2	2	2	19
S10-201	1	1	3	2	2	2	2	19
Sag158	1	1	3	2	2	2	2	19
SG-M25	1	1	3	2	2	2	2	19
SH0334	1	1	3	2	2	2	2	19
SH0655	10	1	4	2	3	3	2	286
SH1370	1	1	2	1	1	2	2	1

Table 1 (*continued*)

Strain	*adhP*	*pheS*	*atr*	*glnA*	*sdhA*	*glcK*	*tkt*	ST
SH3601	1	1	3	2	2	2	2	19
ST 610	13	40	6	65	3	52	50	610
ST 612	111	40	6	65	3	52	51	612
ST 613	111	40	67	65	3	52	51	613
ST 615	13	40	6	65	3	52	51	615
ST 617	13	40	68	65	3	52	51	617
ST 618	111	40	69	65	3	52	51	618
USS-107	1	1	3	2	18	2	2	182

[a] Sequence type (ST), alcohol dehydrogenase gene (*adhP*), phenylalanine tRNA synthetase gene (*pheS*), amino acid transporter gene (*atr*), glutamine synthetase gene (*glnA*), serine dehydratase gene (*sdhA*), glucose kinase gene (*glcK*) and transketolase gene (*tkt*). Multi Locus Sequence Typing (MLST) database at https://pubmlst.org/sagalactiae/.

2. Experimental design, materials, and methods

The list of the S. *agalactiae* strains analyzed is available in Table S1 of reference [2]. Sequences of the genomes of these strains were available as whole genome contigs or as complete genome sequences at the NCBI database on January 19, 2018 (https://www.ncbi.nlm.nih.gov/genome/genomes/186?). Strains with an IS*1548* genomic insertion were identified by blasting these genomes with the IS*1548* DNA sequences described in Ref. [2]. To this end, the nucleotide collection (nr/nt) and the whole-genome shotgun contigs (wgs) databases of *S. agalactiae* (taxid:1311) were screened with the




Fig. 1. Raw data of the goeBURST analysis of the sequence types of strains possessing IS*1548*: Clonal Complex 1.




Fig. 2. Raw data of the goeBURST analysis of the sequence types of strains possessing IS*1548*: Clonal Complex 7.




Fig. 3. Raw data of the goeBURST analysis of the sequence types of strains possessing IS*1548*: Clonal Complex 10 and 12.




Fig. 4. Raw data of the goeBURST analysis of the sequence types of strains possessing IS*1548*: Clonal Complex 17.




Fig. 5. Raw data of the goeBURST analysis of the sequence types of strains possessing IS*1548*: Clonal Complex 19.

blastN program optimized for highly similar sequences (megablast). The algorithm parameters were defined as followed: maximum number of aligned sequences to display: 1000; parameters for short input sequences automatically adjusted; expect threshold: 10; word size: 28; maximum matches in a query range: 0; match/mismatch scores: 1,-2; gap costs: linear; filter for low complexity region activated; mask for lookup table only activated. *S. agalactiae* strains with an IS*1548* positive blastN similarity finding were typed by Multi Locus Sequence Typing (MLST). Seven housekeeping genes were analyzed: alcohol dehydrogenase gene (*adhP*), phenylalanine tRNA synthetase gene (*pheS*), amino acid transporter gene (*atr*), glutamine synthetase gene (*glnA*), serine dehydratase gene (*sdhA*), glucose kinase gene (*glcK*) and transketolase gene (*tkt*). A sequence type, based on the allelic profile of these housekeeping genes, was assigned to each strain by submitting the complete genome sequence or all of the contigs sequences of each strain to the *Streptococcus agalactiae* MLST database (http://pubmlst.org/sagalactiae/, [3], Table 1). Identified MLST types were then assigned to clonal complexes defined by the goeBURST algorithm (http://www.phyloviz.net/goeburst/, [4], Figs. 1–9)




Fig. 6. Raw data of the goeBURST analysis of the sequence types of strains possessing IS*1548*: Clonal Complex 22.




Fig. 7. Raw data of the goeBURST analysis of the sequence types of strains possessing IS*1548*: Clonal Complex 23.

with the ST1 to ST1193 MLST profiles available at the MLST database (https://pubmlst.org/bigsdb?db=pubmlst_sagalactiae_seqdef&page=downloadProfiles&scheme_id=1). A goeBURST clonal complex was defined as all allelic profiles sharing six identical alleles with at least one other member of the group.




Fig. 8. Raw data of the goeBURST analysis of the sequence types of strains possessing IS*1548*: Clonal Complex 27.




Fig. 9. Raw data of the goeBURST analysis of the sequence types of strains possessing IS*1548*: Clonal Complex 615.

Acknowledgments

SK was supported by PhD fellowships of the Lebanese University and AZM & SAADÉ and of the Lebanese Association for Scientific Research (LASeR). This publication made use of the PubMLST website (https://pubmlst.org/) developed by Keith Jolley and sited at the University of Oxford. The development of that website was funded by the Wellcome Trust.

Conflict of Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Appendix A. Supplementary data

Supplementary data to this article can be found online at https://doi.org/10.1016/j.dib.2019.105066.

References

[1] M. Fléchard, P. Gilot, Physiological impact of transposable elements encoding DDE transposases in the environmental adaptation of *Streptococcus agalactiae*, Microbiology 160 (2014) 1298−1315, https://doi.org/10.1099/mic.0.077628-0.
[2] S. Khazaal, R. Al Safadi, D. Osman, A. Hiron, P. Gilot, Dual and divergent transcriptional impact of IS*1548* insertion upstream of the peptidoglycan biosynthesis *murB* gene of *Streptococcus agalactiae*, Gene 720 (2019) 144094, https://doi.org/10.1016/j.gene.2019.144094.
[3] K. Jolley, J. Bray, M. Maiden, Open-access bacterial population genomics: BIGSdb software, the PubMLST.org website and their applications, Wellcome Open Res 3 (2018) 124, https://doi.org/10.12688/wellcomeopenres.14826.1.
[4] A.P. Francisco, M. Bugalho, M. Ramirez, J.A. Carrico, Global Optimal eBURST analysis of Multilocus typing data using a graphic matroid approach, BMC Bioinf. 10 (2009) 152, https://doi.org/10.1186/1471-2105-10-152.