



HAL
open science

Advances and prospects of environmental DNA in neotropical rainforests

Lucie Zinger, Julian Donald, Sébastien Brosse, Mailyn Adriana Gonzalez, Amaia Iribar, Céline Leroy, Jérôme Murienne, Jérôme Orivel, Heidy Schimann, Pierre Taberlet, et al.

► **To cite this version:**

Lucie Zinger, Julian Donald, Sébastien Brosse, Mailyn Adriana Gonzalez, Amaia Iribar, et al.. Advances and prospects of environmental DNA in neotropical rainforests. *Tropical Ecosystems in the 21st Century*, 62, Academic Press, pp.331-373, 2020, *Advances in Ecological Research*, 978-0-12-821134-2. 10.1016/bs.aecr.2020.01.001 . hal-02909735

HAL Id: hal-02909735

<https://hal.inrae.fr/hal-02909735v1>

Submitted on 25 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Running title: eDNA in Neotropical rainforests

Advances and prospects of environmental DNA in Neotropical rainforests

5 Lucie Zinger¹, Julian Donald², Sebastien Brosse², Maily A. Gonzalez³, Amaia Iribar²,
Céline Leroy^{4,5}, Jérôme Murienne², Jérôme Orivel⁵, Heidy Schimann⁵, Pierre
Taberlet^{6,7}, Carla Martins Lopes⁸

¹ Ecole Normale Supérieure, PSL Research University, CNRS, Inserm, Institut de
10 Biologie de l'Ecole Normale Supérieure (IBENS), F-75005, Paris France

² Université Toulouse 3 Paul Sabatier, CNRS, IRD, UMR 5174 EDB, F-31062
Toulouse France.

³ Programa Ciencias de la Biodiversidad, Instituto de Investigación de Recursos
Biológicos Alexander von Humboldt. Bogotá, Colombia.

15 ⁴ AMAP, Univ Montpellier, CIRAD, CNRS, INRAE, IRD, Montpellier, France

⁵ INRAE, CNRS, AgroParisTech, CIRAD, Université de Guyane, Université des
Antilles, UMR Ecologie des Forêts de Guyane (EcoFoG), Campus agronomique, BP
316, 97379, Kourou, France.

⁶ Laboratoire d'Ecologie Alpine (LECA), CNRS, Université Grenoble Alpes,
20 Grenoble, France

⁷ UiT – The Arctic University of Norway, Tromsø Museum, Tromsø, Norway

⁸ Departamento de Zoologia, Instituto de Biociências, Universidade Estadual
Paulista (UNESP). Rio Claro, SP, Brazil. CEP 13506-900.

25 **Abstract:** The rainforests of the Neotropics shelter a vast diversity of plant, animal
and microscopic species that provide critical ecosystem goods and services for both
local and worldwide populations. These environments face a major crisis due to
increased deforestation, pollution, and climate change, emphasizing the need for
more effective conservation efforts. The adequate monitoring of these ecosystems
30 has proven a complex and time consuming endeavour, which depends on ever
dwindling taxonomic expertise. To date, many species remain undiscovered, let
alone described, with otherwise limited information regarding known species
population distributions and densities. Overcoming these knowledge shortfalls and
practical limitations is becoming increasingly possible through techniques based on
35 environmental DNA (eDNA), i.e. DNA that can be obtained from environmental
samples (e.g. tissues, soil, sediment, water, etc.). When coupled with high-
throughput sequencing, these techniques now enable realistic, cost-effective, and
standardisable biodiversity assessments. This opens up enormous opportunities for
advancing our understanding of complex and species-rich tropical communities, but
40 also in facilitating large-scale biomonitoring programs in the neotropics. In this
review, we provide a brief introduction to eDNA methods, and an overview of their
current and potential uses in both terrestrial and aquatic ecosystems of neotropical
rainforests. We also discuss the limits and challenges of these methods for our
understanding and monitoring of biodiversity, as well as future research and applied
45 perspectives of these techniques in neotropical rainforests, and beyond.

Keywords: eDNA, DNA metabarcoding, environmental genomics, conservation
biology, biomonitoring, neotropics, rainforests.

1. Introduction

50 Faced with the current environmental crisis, there is an ever growing need to accurately assess existing policy and legislation which aims to protect ecosystems, such as the Paris Climate Agreement, the REDD+ framework, and the Aichi targets (Marques et al. 2014), as exemplified by the IPBES framework (Díaz et al. 2019). This is particularly true for the neotropical moist broadleaf forests, i.e. those
55 occurring from southern Mexico and Florida to Argentina (Olson et al. 2001; Morrone 2014). Of these forests, the rainforests occurring across Amazonia are the most substantial, covering 40% of the region, and representing the primary source of biodiversity across most taxa (Olson et al. 2001; Jenkins et al. 2013; Antonelli, Zizka, et al. 2018).

60 The biodiversity of neotropical rainforests provides critical ecosystem goods and services for both local and worldwide populations (Rice et al. 2018; Chaplin-Kramer et al. 2019), but these are threatened by increasing human pressures. The region has experienced a 10-fold increase in population densities over the past few decades (Tritsch & Le Tourneau 2016), coupled with a drastic increase in human
65 activities such as deforestation, agricultural expansion, mining and infrastructure construction (e.g. roads, dams; Castello et al. 2013; Rice et al. 2018). These unsustainable land transformations considerably modify abiotic conditions across habitats, and lead to species extinctions, resulting in altered ecosystem functioning and service provision (Rice et al. 2018; FAO 2019). In addition, current predictions
70 for the Amazon basin suggest that climate change will translate to increased droughts, forest-to-savanna transitions, carbon stock losses, and an alteration of the hydrologic and biogeochemical cycles which currently structure this ecosystem

(Nepstad et al. 2008; Davidson et al. 2012).

Assessing the fate of biodiversity with global change and the efficiency of
75 management policies relies largely on the measurement of biological variation at
genetic, population, community and ecosystem levels. Such measures, termed
“Essential Biodiversity Variables” (EBV, Pereira et al. 2013; Table 1) are most
effective when they can be measured in a standardized way that can be employed at
varying scales. Currently, these measurements are based on sampling and direct
80 observation of individuals and their description as species by taxonomists. However,
obtaining EBVs for neotropical forests is not straightforward. The majority of species
occurring in the Neotropics are rare and often exhibit a high level of cryptic diversity
(ter Steege et al. 2013; Antonelli, Ariza, et al. 2018; Zizka et al. 2018), making them
difficult to describe. The description of such hyperdiverse ecosystems thus relies on
85 considerable taxonomic expertise, yet these skills are in decline (Paknia et al. 2015).

Such a shortfall inherently affects our understanding of species spatial
distribution, abundance, evolutionary history, feeding and habitat preferences, as
well as functional properties (Hortal et al. 2015). Even when species are identifiable,
uncertainties surrounding their spatial distribution remain considerable for
90 neotropical rainforests, since biodiversity assessments are often spatially restricted
and biased towards a limited number of accessible areas. These issues pose major
limitations to characterising these ecosystems, to better anticipating their responses
to global change, and ultimately to implementing effective policies of biodiversity
conservation across the region.

95 Environmental DNA (eDNA) based methods (Figure 1) are now considered as
key tools to overcome the aforementioned challenges (Taberlet, Coissac,

Hajibabaei, et al. 2012; Deiner et al. 2017; Taberlet et al. 2018; Table 1), providing numerous advantages over classical inventory approaches. Firstly, DNA for taxonomic identification allows an objective analysis of sequence composition, as
100 opposed to more subjective determination using specimen morphology. Secondly, the sampling of the DNA released in the environment by organisms, or environmental DNA (eDNA) is straightforward, due to its prevalence almost everywhere.

In its narrowest sense, eDNA corresponds to the mixture of DNA that can be
105 found in any environmental matrix, whether consisting of soil, sediment or water. This DNA can belong to organisms that are present within the sample in an active or dormant stage (e.g. microbes, spores, pupae, or seeds). Alternatively, it can belong to organisms living in the sample vicinity, since organisms continuously expel DNA into the environment through excretion, secretion, decomposition, or sloughing of
110 tissues. An environmental sample therefore contains a “metagenome”, i.e. a pool of complete or partial genomes from many different species. This metagenome is made up of DNA that can be intracellular or extracellular, dissolved or adsorbed on organic or mineral particles (Nagler et al. 2018).

In its broadest sense, eDNA also corresponds to the DNA that can be
115 extracted from any biological material collected in natural systems, whether it corresponds to a single specimen or a whole community (e.g. bulk samples made of a mass trapping of arthropods or fish larvae). In both cases, the DNA recovered from such a sample does not only contain that of the specimens, but also encompasses the genes/genomes of the specimens symbionts, parasites, or more generally of
120 their microbiota, as well as of their prey (Taberlet et al. 2018; Hacquard et al. 2015).

Thus, the biodiversity retrieved from an eDNA sample is trans-kingdom and

multitrophic. Combined with high-throughput sequencing (HTS), environmental DNA-based methods (section 2, Figure 1) now make large-scale and multi-taxa surveys possible from material that is easy to collect, requiring minimal taxonomic expertise.

125 So far mostly used in temperate environments, such surveys could considerably speed up the acquisition of EBVs in general (Jetz et al. 2019), and in species rich and challenging ecosystems such as neotropical rainforests (Table 1).

First, eDNA can provide information on the occurrence of invasive species (Takahara et al. 2013; Valentin et al. 2018), human and agricultural pathogens or
130 pests (Lievens et al. 2006; Harwood et al. 2014; Bass et al. 2015), endangered species or populations (Harper et al. 2018; Tessler et al. 2018) and of wild species in general (Kirshtein et al. 2007; Scibetta et al. 2012). Likewise, it can be used to monitor species that indicate the health of ecosystems (i.e. bioindicators), in particular when these are microbes or invertebrates, of which identification requires
135 advanced and often rare taxonomic skills (Mächler et al. 2014; Pawlowski et al. 2014), especially in tropical ecosystems (Rousseau et al. 2013; Bowles & Courtney 2018 and references herein).

Second it can provide reliable information on the diversity and community composition of soil or aquatic microbes (e.g. Lauber et al. 2009; Zinger et al. 2011; Gilbert et al. 2012), as well as of invertebrates (Pansu et al. 2015; Bista et al. 2017; Zinger, Taberlet, et al. 2019), fish, amphibian, and mammalian communities (Boussarie et al. 2018; Schnell et al. 2018). eDNA can be further used as a standard impact assessment tool in both aquatic (Chariton et al. 2010; Li et al. 2018) and
140 terrestrial ecosystems (e.g. Drenovsky et al. 2010), or as an evaluation tool for the success of restoration and conservation strategies (Bohmann et al. 2014; Perring et al. 2015). Finally, eDNA can provide information for multiple taxon at the same time

(e.g. Li et al. 2018; Zinger, Taberlet, et al. 2019), and thus on biological interactions (Vacher et al. 2016). For example, using the eDNA retrieved from the faeces or gut content of a given species can reveal feeding habits (Pompanon et al. 2012), as well
150 as host microbiota and the occurrence of potential pathogens/parasites (Bass et al. 2015). This enables the study of full ecological networks across environmental or land disturbance gradients.

The objectives of this review are therefore (i) to provide a brief overview of eDNA-based methods, (ii) to assess their implementation to describe biodiversity in
155 both terrestrial and aquatic ecosystems of neotropical rainforests, (iii) to highlight the limits and challenges of these methods for providing reliable assessments of EBVs in these environments, and (iv) to propose several avenues for future research in this field.

[INSERT Table 1 here]

160 **2. Overview of eDNA methods**

The study of eDNA is made possible through the extraction of DNA from its environmental/biological matrix and its separation from any chemicals that can affect DNA amplification or sequencing reactions (e.g. humic substances, polyphenols, etc.). Once the DNA extract is obtained, four main methods are now routinely applied
165 depending on the final objective (Figure 1). They rely either on the amplification or enrichment of a target genomic region of the metagenome (i.e. species detection, DNA metabarcoding, or capture/enrichment), or on the direct - or “shotgun” - sequencing of the metagenome (i.e. metagenome skimming or metagenomics). We briefly describe each of these approaches below and in particular emphasize DNA
170 metabarcoding throughout this review, as this method is currently the most widely

used in the field, in particular in neotropical rainforests. For more detail regarding the molecular and bioinformatics procedures involved, we refer the reader to dedicated literature (e.g. de Bruijn 2011; Bálint et al. 2016; Deiner et al. 2017; Taberlet et al. 2018).

175 The “species detection” approach consists of detecting/quantifying the amount of a DNA marker that is specific to a single or a small set of species. This approach is most relevant when one aims to detect a species with a high level of sensitivity, including low density populations or dormant/juvenile life forms. The DNA markers used for this approach must correspond to a highly polymorphic locus, enabling the
180 design of primers which are highly species-specific. The approach currently preferred is a direct quantification of the number of copies of the target DNA marker through quantitative PCR (i.e. qPCR, sometimes referred to as real time PCR; Rees et al. 2014) or digital droplet PCR (ddPCR; Doi et al. 2015). These two quantitative methods can help to assess species population density or biomass in the studied
185 area (e.g. Pilliod et al. 2013). This approach is relatively cheap, since it does not rely on sequencing, and is therefore more suitable for large-scale or temporal studies, although is limited to focusing on only one or a reduced set of species.

 “DNA metabarcoding” (Taberlet, Coissac, Pompanon, et al. 2012) is the most popular approach to study eDNA (see Figure 1 for more detailed information). This
190 approach has also been referred to in the literature as “amplicon sequencing”, “ecometagenetics”, “metataxogenomics”, but should not be confused with “metagenomics”, which we define below. As with species detection, DNA metabarcoding relies on the amplification of a target DNA region by PCR. However in this case, the DNA region targeted is used as a “barcode” to discriminate the
195 species comprising the metagenome under study. A relatively large number of

samples processed with DNA metabarcoding can be sequenced in a single HTS run (Figure 1). The obtained sequencing reads are then processed bioinformatically to retrieve a list of species (or Operational Taxonomic Units, OTUs).

Enrichment capture on eDNA is very similar to DNA metabarcoding in that it consists of sequencing the same targeted regions. However, it differs in that the target DNA to be sequenced is not enriched through PCR amplification, but instead by capturing it with multiple, taxon-specific DNA probes bound to magnetic beads. This approach is often used for the analysis of ancient DNA of single species or simple species assemblages (e.g. Carpenter et al. 2013) and is increasingly used for the analysis of modern eDNA and complex communities (e.g. Shokralla et al. 2016; Wilcox et al. 2018), although the sensitivity and limitations of this approach are yet to be evaluated.

The last alternative relies on shotgun sequencing, i.e. random sequencing of DNA molecules from the environmental sample. "Metagenomics" is the most direct and comprehensive DNA-based technique, and consists of sequencing as much of the metagenome as possible so as to retrieve organisms taxonomic identity, their phylogenetic relationships, as well as to their metabolic properties. However, it is also the most challenging approach. First, much of the information contained within metagenomes remains undescribed. Second, a metagenome contains a huge diversity of genes and noncoding regions, of which a tiny fraction are highly repeated (e.g. ribosomal RNA genes), and a majority of which are rare. Fully describing this complexity therefore requires substantial sampling, in this case sequencing effort, which today remains costly. Finally, most environmental samples are dominated by microbial DNA, which reduces the probability of detecting larger organisms. Consequently, metagenomics is for now mostly used in environmental microbiology

(e.g. de Bruijn 2011) or for ancient DNA analyses (Thomsen & Willerslev 2015).

“Metagenome skimming” is a cheap version of metagenomics (Linard et al. 2015; Papadopoulou et al. 2015), albeit more expensive than methods targeting a particular DNA region. In this case, the metagenome is sampled at a shallow
225 sequencing depth so as to sequence only highly repeated DNA regions, i.e. the ribosomal RNA gene regions and the organelle genomes for eukaryotes. These regions can then be partially or fully reconstructed, and thus used to identify the species present but also their phylogenetic relationships.

[INSERT Figure 1 here, full width]

230 **3. Current use and challenges of eDNA applications in neotropical rainforests**

Current studies of biodiversity in neotropical rainforests that rely on eDNA based methods mainly describe community composition and diversity changes along environmental or disturbance gradients in order to identify patterns in diversity and
235 their drivers. These studies are reviewed below across ecosystems and focal organisms, and examined to determine what eDNA from different sources can reveal regarding ecological communities from neotropical rainforests and how sampling can be tailored to suit the ecological question. We will restrict our review to contemporary environments, as - to our knowledge - eDNA approaches *per se* have not been used
240 yet in the neotropical rainforests for palaeoecological purposes. We refer interested readers to dedicated reviews on this particular application (Rawlence et al. 2014; Thomsen & Willerslev 2015; Taberlet et al. 2018).

3.1. In terrestrial ecosystems

Microbial communities. These have been mostly analysed in the soil

245 environment, with the study of eDNA from soil samples having a relatively long
history in soil microbial ecology (Tiedje et al. 1999 for an early review). Available
studies for neotropical rainforests have shown that soil prokaryotic and
microeukaryotic communities vary across altitudes (Nottingham et al. 2018), soil
250 conditions, forest types and tree species composition (Ritter et al. 2019; Vasco-
Palacios et al. 2019). Numerous studies also report steep changes in composition
with increased drought (Waring & Hawkes 2015; Kivlin & Hawkes 2016a; Pajares et
al. 2018), deforestation and reconversion to different types of silviculture (Carney et
al. 2004; Ndaw et al. 2009; Kivlin & Hawkes 2016a; Kivlin & Hawkes 2016b), arable
farming (Rodrigues et al. 2013; Paula et al. 2014; Mendes et al. 2015; e.g. Franco et
255 al. 2019), and even as a result of pre-columbian activities (Kim et al. 2007;
Grossman et al. 2010; Navarrete et al. 2010). Likewise, soil microbial diversity differs
between old-growth and secondary forests (Araújo et al. 2014; McGee et al. 2019).
All these studies exemplify the utility of soil eDNA for providing microbial-derived
EBVs that are meaningful for monitoring the impact of climate change and land use
260 practices.

Invertebrates. Soil micro- and macro-invertebrates (i.e. nematodes,
earthworms, insects and springtails) have seldom been studied with eDNA from
neotropical rainforest soil samples (Wu et al. 2011; Zinger et al. 2016; Ritter et al.
2019), and in such cases, rather as part of the whole soil eukaryote diversity,
265 through the use of universal primers. A global-scale analysis suggests that
neotropical rainforests are dominated by arthropods and enriched in soil annelids
(Wu et al. 2011). Locally, soil micro- and mesofauna communities exhibit primarily
random spatial patterns that are more pronounced for the mesofauna as compared
to microscopic organisms, as shown at a forest site in French Guiana (Zinger,

270 Taberlet, et al. 2019).

The large majority of studies of soil or above-ground invertebrates have so far rather relied on eDNA extracted from bulk samples and analysed through DNA metabarcoding, which is a fast alternative to time consuming sorting and identification of hundreds to thousands of specimens that are difficult to identify.

275 Using this approach, Porazinska *et al.* (2012) were able to observe strong variation in soil nematodes communities across sites and habitats of Costa Rican rainforests. This approach has also enabled the description of above-ground terrestrial arthropods, such as sandflies occurring at several sites in French Guiana (Kocher, Gantier, et al. 2017), or arthropods from a forest canopy in Honduras (Creedy et al. 280 2019). The latter study also tested the effect of animal size on species detection, with results suggesting such effects are not visible when sequencing depth is sufficient. Enrichment capture has also been used to analyse bulk samples of arthropods sampled with malaise traps in a forest of Costa Rica (Shokralla et al. 2016). This method was found to be more accurate in describing biodiversity than 285 DNA metabarcoding on the same samples and classical observations.

Mammals. Using eDNA from bulk samples of faeces or hematophagous arthropods also seems particularly promising for sampling terrestrial vertebrate diversity as well. For example, DNA extracted from owl pellets in central Brazil provided meaningful information regarding the diversity of small mammals (Rocha et 290 al. 2015). Likewise, vertebrate communities are better described by the DNA contained in blood feeding arthropods collected with Malaise traps and pitfall traps than with classical or camera trap-based inventories, as shown for forests in Panama and Brazil (Rodgers et al. 2017; Lynggaard et al. 2019). This approach further revealed variation in vertebrate community composition, consistent with a gradient of

295 anthropogenic pressures in French Guiana, with a decline of diversity in the areas
experiencing the highest pressures (Kocher, de Thoisy, François Catzefflis, et al.
2017). Alternatively, water samples could also be used to study terrestrial mammals,
since water bodies should accumulate and transport material from the whole
catchment areas through erosion (Naiara Guimarães Sales et al. 2019).

300 **Plants.** Initial attempts to describe plant diversity with eDNA used bulk
samples of dried, fine roots isolated by hand from soil cores that were collected
following a grid or regular sampling scheme in the Barro Colorado Island in Panama
(Jones et al. 2011; Barberán et al. 2015). With this approach, one soil core exhibited
an average diversity of *ca.* 4 plant species and the DNA imprint of each tree
305 individual was detectable from 1 to *ca.* 20m from the stem. Similar figures can be
retrieved by directly using soil as starting material, as shown in a lowland rainforest
in French Guiana (Fig. 2A, see also Taberlet et al. 2018, (Yoccoz et al. 2012). Thus,
root and soil eDNA can offer new insights into plant root distribution in the soil and
their functional implications. The aboveground plant community might be better
310 assessed by targeting plant DNA markers on bulk samples of herbivorous
arthropods, but to our knowledge, this approach has not been tested yet.

Constraints and limits. The above shows that organisms from terrestrial
environments are either studied using environmental DNA extracted from soils,
which are noticeable reservoirs of both intra- and extracellular DNA and mostly
315 contain the signature of soil organisms, or using bulk samples of invertebrates. The
former material is probably the easiest to sample from a practical point of view, and
less biased/variable than different arthropod sampling techniques (Missa et al.
2009). However, one critical aspect when studying diversity using eDNA extracted
from soil is the heterogeneous and complex nature of soil substrates themselves, in

320 terms of physical, chemical and biological properties (Bardgett & van der Putten
2014). This can be an issue when comparing contrasting environments. Typically,
the amount of available extracellular DNA, useful for detecting non-microbial
organisms, is strongly constrained by soil chemical properties. For example, DNA
has a much stronger affinity to clay than sand (Levy-Booth et al. 2007), which thus
325 could introduce bias to comparisons between white-sand vs. *terra firme* forest soils.
This heterogeneity not only applies horizontally in space but also vertically, with clear
differences in prokaryotic and micro-eukaryotic communities between the organo-
mineral horizon and the litter layer from a taxonomic point of view (Figure 2B), and
most likely from a functional one (Fragoso & Lavelle 1992; Basset et al. 2015; Ritter
330 et al. 2019). This raises the question of if and how one should integrate this vertical
dimension.

The same applies for how much soil samples should be collected across
space, whatever the taxon targeted. For example, *terra firme* soils/litter and white
sand litter may require higher sampling effort than white sand soils to estimate the
335 plot-scale diversity, due to higher spatial heterogeneity (Figure 2C). Alternatively,
sampling effort could be reduced when comparing diversity or community turnover
across conditions by for example building composite samples at different sampling
points to capture local diversity while keeping down experimental costs. At the
sample scale, extracting eDNA from volumes of material that are larger than those
340 from most commercial soil DNA extraction kits (typically 250 µg) should best capture
local diversity, which is now possible for ≥10 grams of starting material, as shown for
neotropical rainforest soils (e.g. Zinger et al. 2016). Thus the required sampling effort
is likely to be highly system dependent, and further analyses across habitats will help
better define sampling standards for neotropical rainforests when using soil as a

345 starting material.

In any case, soil samples are unlikely to be the most relevant material for sampling the diversity of plants or aboveground animals at the plot scale because eDNA is poorly transported in soils and thus highly patchily distributed (Levy-Booth et al. 2007; Nagler et al. 2018). This patchiness most likely also results from the
350 reduced DNA persistence in tropical rainforest soils due to high demands of the living biomass for phosphorus, which is otherwise highly limiting (Dalling et al. 2016), thus reducing the probability of detecting large organisms. Accordingly, experiments show that dead root DNA is almost totally degraded after 15 days (Bithell et al. 2014). Likewise, microbes and soil fauna communities exhibit marked
355 seasonal and yearly dynamics (Fragoso & Lavelle 1992; Kivlin & Hawkes 2016b; Kivlin & Hawkes 2016a; Pajares et al. 2018), and so most likely does their DNA as compared to that of rooted plants, which continuously release DNA in soils (Figure 2D).

[INSERT Figure 2 here, full width]

360 **3.2. In aquatic ecosystems**

Microbial communities. Several studies using water eDNA have been conducted across different systems to study microbial communities. For example, Tessler *et al.* (2017) showed that bacterial communities from Brazilian floodplain lakes were highly distinct from other areas of the globe, while within Brazilian sites,
365 the composition was overall fairly similar. Other studies suggest the opposite for micro-eukaryotic plankton: Brazilian rivers seem to exhibit marked spatial patterns with relatively high community turnover, even within the same location (Lentendu et al. 2019). These discrepancies raise the question as to whether they arise from biological differences between microeukaryotes and bacteria or from methodological

370 inconsistencies, which emphasize the need for increased efforts in studying aquatic
microbial communities in these ecosystems.

Tank bromeliads (Bromeliaceae) accumulate rainwater at the base of each
leaf axil and thus represent freshwater islands in a terrestrial matrix. They harbour
various aquatic organisms ranging from prokaryotes to macroinvertebrates (Benzing
375 2000; Leroy et al. 2016). eDNA methods have provided insights into their community
structure through either metagenomics (Rodriguez-Nuñez et al. 2018) or DNA
metabarcoding (Louca et al. 2016; Louca et al. 2017), revealing bacterial
communities that are substantially different from freshwater lake sediments and soil,
but remarkably similar in functional structure due to an adaptation to oxygen-limited
380 conditions.

Invertebrates. The use of water/sediment eDNA for targeting aquatic
invertebrates (aquatic insects, crustaceans) has, to our knowledge, not yet been
applied to neotropical rainforest ecosystems. A recent study has shown its
usefulness for assessing macroinvertebrates community composition in the tropical
385 freshwaters of Singapore (Lim et al. 2016), suggesting that such an approach could
be relevant to neotropical rainforest ecosystems. As for terrestrial environments, the
use of bulk samples for aquatic systems is emerging, such as with the study of
Talaga et al. (2017), which details the development of DNA reference libraries for
Guianese mosquito larvae to distinguish species from bulk samples of freshwater
390 invertebrates. Still, eDNA studies of freshwater invertebrates in neotropical
rainforests are currently limited by knowledge deficits related to their taxonomy and
ecology and a lack of previously implemented studies . Although several
macroinvertebrate indices enabling the biological evaluation of freshwater
ecosystems are available (e.g. Couceiro et al. 2012; Dedieu et al. 2016), these are

395 seldom used because to our knowledge, there is currently no environmental law or
regulation relying on these in this ecoregion. One exception in that respect is French
Guiana, which must comply with the European Water Framework Directive.

Fishes. The potential of water/sediment eDNA has received comparatively
much more attention for studying fish communities. This has been particularly
400 stimulated by the strong limitations of traditional sampling methods, which provide
biased estimates and/or cause substantial fish mortalities. Indeed, gill nets provide
only partial inventories, and ichthyocides such as rotenone, which were widely
employed in the past, are increasingly banned. Electric fishing, which is often a good
sampling alternative in other environments can be inefficient in neotropical streams
405 because of the very low conductivity of the water (Allard et al. 2014). Hence, fish
eDNA has rapidly emerged as the most promising non invasive alternative to
traditional sampling for small streams, rivers, lakes and the sea. Cilleros *et al.* (2019)
compared eDNA and traditional sampling (nets and ichthyocides) both in small
streams and rivers across French Guiana. Not only did they find that species
410 assemblages were congruent between eDNA and traditional records, but also that
eDNA results were more efficient in distinguishing the fauna from different river
drainages. eDNA also enables the study of fish communities at cryptic life stages,
i.e. the ichthyoplankton. Nobile *et al.* (2019) used DNA metabarcoding on mock
communities built from fish eggs and larvae in the Grande River in Brazil, and
415 obtained an average detection rate higher than 95%, and a relatively good estimate
of larvae abundances. Likewise, capture enrichment on bulk samples for catfish
larvae from the Peruvian Amazon provided a good description of the community in
terms of both species and abundance (Maggia et al. 2017; Mariac et al. 2018).

Vertebrates. Several studies have focussed on vertebrates inhabiting aquatic

420 environments for at least a part of their life, such as amphibians. Comparing
traditional visual and audio survey techniques with DNA metabarcoding of water
samples showed that eDNA accurately reflects the conclusions of the other methods
whilst cutting the length of fieldwork required studying for frog communities in
freshwater streams in the Brazilian Atlantic forest (Lopes et al. 2017; Sasso et al.
425 2017). Likewise, a comparison of cost models suggests that eDNA-based surveys
are a cost-efficient alternative to traditional surveys in amphibian species rich areas
such as in the neotropical forest-savannah ecotones of Bolivia (Bálint et al. 2018). All
these studies further show that eDNA-methods circumvent biases of traditional
approaches linked with species abundance and life history traits. Indeed, they not
430 only allow for the detection of species closely associated with streams, but also of
frog species at cryptic life stages (e.g. tadpoles or eggs). These are often missed by
traditional surveys, but detectable with eDNA since they release DNA into the
environment irrespective of their life stage. Likewise, eDNA is also able to detect
endangered species in a non-destructive way, such as for the bromeliad inhabiting
435 Trinidad golden tree frog (Brozio et al. 2017). Beyond amphibians, Sales et al (2019)
also detected eDNA from both aquatic and terrestrial mammals when sampling water
in the Amazon's mainstream and tributaries, in addition to a river of the Brazilian
Atlantic forest. Comparing these results with camera trapping data confirmed the
congruence between the methods (Naiara Guimarães Sales et al. 2019).

440 Interestingly, some of the species detected using eDNA from water samples belong
to strictly terrestrial species such as bats or anteaters, which can be explained by the
fact that water conveys DNA from terrestrial to aquatic ecosystems. However, further
studies are needed to validate this protocol for capturing terrestrial vertebrate
diversity.

445 **Constraints and limits.** The above shows that eDNA for studying aquatic ecosystems can be extracted from either water or sediment samples, or bulk samples. For bulk samples, the trapping system is likely to be an important factor, as for traditional observations. For water or sediment samples, the interpretation of eDNA data from these two substrates remains unclear. Apart from microbial
450 communities that highly differ between these two environments due to contrasted oxygen nutrient availability (Zinger et al. 2011; Thompson et al. 2017), the discrepancy between the results obtained from water and sediments when targeting larger organisms has been highlighted by several studies. Some studies have shown that fish eDNA concentration in sediments is higher and detectable over longer
455 timescales than in water (Turner et al. 2015; Naiara G. Sales et al. 2019). However, other studies found that sediments were less effective than water samples, e.g. allowing the recovery of only 10% of the fish species in an oligotrophic lake in Mexico (Valdez-Moreno et al. 2019). Water remains to date the most commonly used substrate for eDNA studies in neotropical rainforests due to its ease of
460 collection. Sampling of eDNA is mainly conducted using filtration that is either directly performed in the field or subsequently in the laboratory (Lopes et al. 2017; Naiara G. Sales et al. 2019).

For both water and sediment samples, the concentration of eDNA in the environmental matrix strongly determines how much material should be collected to
465 appropriately sample freshwater diversity. For example, Cantera et al. (2019) sampled up to 340 L of water in streams and rivers in French Guiana to study the impact of sampling effort on fish detection. They showed that with a total filtration of 68 L, 91% of fish diversity could be detected in streams, and 74% in rivers. These results resonate with those obtained by Lopes et al. (2017), showing that filtering

470 larger quantities of water (from 20L to 60L) increases the detection probability for
amphibian species and thus covering local amphibian diversity in the Brazilian
Atlantic forest. Nevertheless, according to Cantera et al. (2019), filtering 34 L of
water is sufficient for the recovery of 64% of the local fish fauna in Guianese streams
and rivers, with a strong redundancy between eDNA replicates. Such a limited
475 sampling effort seems hence sufficient to distinguish fish communities between sites
and between ecosystem types (i.e. streams vs rivers).

The concentration of eDNA in freshwaters is a function of the local living
biomass, but also of the transport and degradation rate of eDNA in freshwater
ecosystems, which depends on environmental conditions (Barnes et al. 2014;
480 Barnes & Turner 2015). These processes require further investigation in both waters
and sediments of neotropical rainforest ecosystems, in order to best define the
sampling effort required to conduct reliable eDNA studies in these areas. It is now
well established that low pH conditions, high oxygen demand and primary
production, and high temperatures all accelerate the degradation of aquatic eDNA
485 (Barnes et al. 2014; Strickler et al. 2015), which is likely to strongly vary across
neotropical rainforest rivers and streams. As such, a study found an unexpected
higher mammal species richness in the Brazilian Atlantic forest compared to the
Amazon (Naiara Guimarães Sales et al. 2019), which is suspected to arise from a
higher degradation rate of eDNA due to the low pH of the Amazon waters ($\text{pH} \leq 4$).

490 Another important point, strongly linked to the degradation rate, is the
transportation of eDNA with water flow. Studies are ongoing on this aspect in
neotropical rivers, but Cantera et al. (2019) report that fish species detected from a
stream site were no longer detected in eDNA samples collected in a river site located
300m downstream from the confluence with the river. This suggests either a rapid

495 degradation and hence a relatively short distance of eDNA transportation in
neotropical waters, or more generally a high dilution downstream, which should
make eDNA detection more difficult at sites distant from where it has been released.
Finally, precipitations and stream size should also define local eDNA concentrations.
For example, Sales et al. (2019) reported noticeable compositional differences
500 between samples collected from the same location following a 3-week interval. While
this might be due to real variation in species composition, it is also possible that
variation in water volume linked to increased precipitation at the time of sampling
affected species recovery.

3.3. Common field, wet, and dry lab biases

505 Besides the clade- and environment-specific considerations mentioned above,
the processing of eDNA data typically consists of a series of methodological steps
(Figure 1) that are all subject to various biases (Dickie et al. 2018; Zinger, Bonin, et
al. 2019). We will briefly outline some of them and their associated solution when
crucial for applications in neotropical rainforests ecosystems, as these issues are
510 extensively addressed elsewhere (e.g. Deiner et al. 2017; Taberlet et al. 2018;
Alberdi et al. 2019). This discussion will be mostly focused on DNA metabarcoding,
as it is the approach the most widely used in eDNA research.

At the sampling step, the extent of the sampling area, sampling point
locations, number of biological replicates, sample conditioning and transport, *etc.* are
515 all important points to critically consider to avoid compromising the results (reviewed
in Dickie et al. 2018; Taberlet et al. 2018) and will inherently depend on the
particularities of the ecosystem and taxon under study (see above). Appropriate
sample conditioning is also critical in tropical climates, in which microbial growth and
DNA degradation is faster and more likely to occur during sample transport. Sample

520 cooling in ice can considerably slow down DNA degradation and microbial growth,
but this is seldom logistically feasible when working in remote and warm sites. To
circumvent this limitation, DNA extraction can be done directly in the field with
specified protocols requiring minimal infrastructure (e.g. Zinger et al. 2016; see
Taberlet et al. 2018 for a detailed protocol). Alternatively, the sample can be
525 dessicated with *silica gel* for soils or sediments, or more generally conserved with
preservation buffers. These are typically used for aquatic eDNA samples, either for
conserving water filters on which eDNA has been captured (Cilleros et al. 2019) or
for direct addition to water samples, although preservation buffers seems less
effective than sample cooling for eDNA recovery and taxon detection (Naiara G.
530 Sales et al. 2019).

After collection, the molecular processing of samples also has a variety of
biases that can reduce the detection or distort the abundance of the taxa retrieved,
an important limit for species population EBVs (Table 1). DNA extraction methods
are not equally efficient in extracting and purifying DNA, due to variable success of
535 cell lysis for microbes, and more generally to strong variations in the chemical
composition of the starting material, with some being noticeable PCR inhibitors (e.g.
humic acids). The methods employed for the extraction of DNA should be tailored to
the starting material and question, or it may miss or overrepresent certain taxa. Once
DNA is extracted, PCR amplification should be done with primers whose specificity-
540 to- and generality-within the clade of interest should have been verified following a
thorough literature review, preliminary tests, or the use of *in silico* PCR softwares
(e.g. Ficetola et al. 2010; Elbrecht & Leese 2017). Use of inappropriate primers will
both strongly bias the retrieved taxa abundances and in some cases, their detection
altogether.

545 Both PCR amplification and sequencing can also generate artifactual DNA
fragments/sequences, especially when the target DNA is rare (reviewed in Taberlet
et al., 2018). These artifacts are generally in low abundance and very similar to
genuine fragments (e.g. only one or a few different nucleotides). They are hence
difficult to identify and can artificially inflate taxonomic diversity estimates, this
550 attribute being a community composition EBV candidate (Table 1). Nevertheless,
such errors can be reduced by clustering DNA sequences at a certain sequence
identity level using supervised or unsupervised approaches (Figure 1). However, it
should be noted that the bioinformatics tools used, as well as their associated
parameters (e.g. clustering methods and thresholds, sequences distance indices)
555 are not all equally efficient in reducing this artifactual variability, and can even fail to
detect genuine biological variability (Coissac et al. 2012; Bálint et al. 2016; Zinger &
Philippe 2016; Deiner et al. 2017; Taberlet et al. 2018). The same applies when
using supervised approaches, as the taxonomic assignment quality of a sequence/
OTU inherently relies on the completeness and accuracy of the reference databases.
560 For example, using an incomplete reference database, i.e. without conspecific
sequences, can lead to an increase of 20% of erroneous taxonomic assignments as
compared to the use of a complete one, as shown for Amazonian mammals (Kocher,
de Thoisy, Francois Catzeflis, et al. 2017).

Diversity estimates can also be inflated through the presence of genuine DNA
565 fragments that are not initially present in the sample. The most obvious source of
such a problem is exogenous contamination, which can occur not only at the
sampling step, but also at the extraction, PCR, and sequencing steps because labs
and reagents all contain a number of contaminants (Salter et al. 2014). Beside this
problem, the multiplexing of samples within a single sequencing library or

570 sequencing lane also produces apparent cross-sample contamination. The exact
underlying mechanisms remain not well understood, but DNA fragments that are
multiplexed seem to exchange the small tags used to identify their sample of origin
(Figure 1), a bias often referred to as ‘tag-switches’, ‘tag-jumps’, or ‘cross-talks’ (e.g.
Schnell et al. 2015; Esling et al. 2015). Although this bias produces contaminants at
575 usually low abundances, it can have strong consequences if downstream analysis
relies on presence/absence and occurrences.

Given the different artifacts mentioned above, the reader should now be
aware that the inclusion of negative and positive controls at the sampling, extraction,
amplification and sequencing steps as well as technical replicates is critically
580 important to ensure not only data reliability but also to optimize the processing and
curation procedures of the obtained sequences through bioinformatics pipelines. The
problem of false positives can be reduced by using PCR-independent methods, such
as metagenomics/metagenome skimming, or capture enrichment. However both
approaches still require substantial developments and cost reductions to be
585 applicable in large-scale studies. In addition, these approaches are not error-free.
They still include tag-jumps or sequencing errors (Taberlet et al. 2018; Wilcox et al.
2018) that remain difficult to detect and filter out.

Artifactual signals can have dramatic effects on estimates and patterns of
alpha, and to a lesser extent beta diversity (Calderón–Sanou et al. 2019), as well as
590 on model parameters inference such as for Hubbell’s neutral model (Sommeria-Klein
et al. 2016). Since these artifacts are generally low in frequency, end-users should
also be careful when focusing on rare taxa. This corresponds to the majority of
species in neotropical rainforest ecosystems (ter Steege et al. 2013; Antonelli, Ariza,
et al. 2018; Zizka et al. 2018), which suggests that it is unlikely that current eDNA-

595 based approaches provide reliable estimates of species richness, i.e. the number of
species being present in the ecosystem studied. Nevertheless, these approaches
can still provide meaningful information on alpha or beta diversity patterns by using
diversity indices penalizing low-abundance OTUs or taxa such as those based on
Hill numbers, which includes well known indices such as Shannon or Simpson
600 diversity (Chao et al. 2014). These have been shown to provide more reliable
ecological inferences (Calderón–Sanou et al. 2019), and should be favored over
other indices where singletons (e.g. Chao, ACE, Fisher’s alpha indices) or rare
species have a strong weight (e.g. inferences based on species abundance
distribution or on presence-absence data). Nevertheless, new occupancy models
605 able to detect both false negative and false positives are currently emerging (Ficetola
et al. 2016; Guillera-Aroita et al. 2017), and their inclusion in current data curation
procedures will certainly allow overcoming the above-mentioned limitations.

3.4. Biological interpretation of eDNA

Beyond the methodological considerations raised above, eDNA has specific
610 intrinsic properties which must be considered when interpreting derived results. Even
if eDNA data resembles a traditional species abundance table, the abundances
correspond to sequencing read counts and species correspond to species, genera,
or to OTUs defined at a given level of sequence similarity. This difference can have
strong implications for the type of EBV that eDNA can actually measure (Table 1), as
615 well as on ecological inferences depending on the question addressed and types of
inference tools used, in particular when they involve theoretical frameworks and
models that rely heavily on species and abundances (e.g. niche or neutral models,
species abundance distributions).

A first uncertainty is on the extent to which sequences or OTUs can be used

620 as a proxy for species. In most eDNA studies, species or OTUs are defined by using
a threshold of 97% of sequence similarity. This threshold has been historically
defined for full-length barcode genes (e.g. Stackebrandt & Goebel 1994; Hebert et
al. 2003; Schoch et al. 2012). However, current eDNA studies target small regions
within these barcodes (Figure 1) in order to comply with both the sequencing limits of
625 current HTS instruments and, when applicable, with the fragmented nature of
extracellular DNA. This constraint inherently comes with a loss of taxonomic
resolution, which may have consequences for subsequent ecological inferences. The
“Amplicon-” or “Exact Sequence Variant” concept (ASV or ESV, Callahan et al. 2017)
has been recently proposed to, amongst other reasons, circumvent this problem, yet
630 this remains sensitive to some molecular artifacts. Sometimes interpreted as
intraspecific variability, which can be a desirable output of eDNA (Table 1), ASVs
may also yield ecological signals that differ from what one should expect when
considering species. Finally, eDNA markers do not have the same taxonomic
resolution across clades. For example, the fungal Internal Transcribed Spacers, or
635 the metazoan cytochrome oxidase subunit I (COI) can exhibit some intraspecific
variability for certain groups, and only genus to family level variability for others
(Schoch et al. 2012). Phylogenetic-based approaches can to a certain extent deal
with these limitations. However, while these can be employed with metagenomics or
metagenome skimming data (Andújar et al. 2015; Papadopoulou et al. 2015), the
640 short and hypervariable nature of most classical DNA markers used for DNA
metabarcoding do not enable making robust phylogenetic inferences, which limits
the use of such data to retrieve co-ancestry relationships (Table 1). For such data,
the phylogenetic diversity should be retrieved through phylogenetic placement
methods, provided that a robust backbone phylogenetic tree is available (e.g.

645 Matsen et al. 2010; Czech et al. 2019), which remains challenging for neotropical taxa (see section 4.1.).

The other uncertainty of eDNA data relates to the meaning of sequencing reads counts. As mentioned in section 3.3, a DNA extract is subjected to a suite of molecular manipulations that can distort the original distribution of DNA fragment
650 abundances. Adding spiked DNA of known composition and concentrations in environmental samples could allow for the retrieval of absolute values of eDNA molecules (e.g. Smets et al. 2015; Thomas et al. 2016). However, while the abundance (relative or not) of eDNA molecules has been found to correlate with organism biomass in simple experimental set ups (e.g. Nobile et al. 2019) or when
655 quantifying single species *in natura* with qPCR (reviewed in Taberlet et al. 2018), several factors can alter this relationship, and hence, assessment of population abundance (Table 1). First, eDNA persistence and transport in the environment makes it difficult to know whether this biomass is local and contemporary. This is likely to be especially true for soils or sediments as compared to water, the latter
660 being more exposed to high temperature and UV radiations, which favor DNA degradation (Barnes & Turner 2015; Nagler et al. 2018). Even if this bias is limited, relating eDNA abundance to population abundance *per se* remains challenging. Indeed, the number of DNA marker gene copies depends on the taxon, on the tissues from which eDNA is released, the biomass/size of the organisms, but also its
665 life stage (Maruyama et al. 2014). To our knowledge, there is no tool which can retrieve individual counts from sequencing reads or eDNA molecules at the scale of the biological community. These uncertainties has often led researchers to prefer presence-absence metrics over abundance-based ones. However, unless the representativeness of the data curation procedure can be proven, we advocate

670 against such reasoning due to the high error rate of PCR and sequencing based approaches (see section 3.3.).

Given the above-mentioned differences in the intrinsic nature of eDNA data as compared to traditional species abundance tables, this raises the question of whether one can draw ecological inferences with classical tools. Typically, it remains
675 largely uncertain whether inferring community diversity and related characteristics from eDNA-based species abundance distribution or using process models involving explicitly species and individuals is a correct approach. For example, adaptation of Hubbell's model to account for body size or biomass could be more appropriate (O'Dwyer et al. 2009; Sommeria-Klein et al. 2016). There is hence a need for
680 development of related tools and theories in ecology that would better comply with the nature of eDNA data.

4. Future directions and perspectives

The past decade has seen enormous advances in the development and extension of eDNA-based approaches, as well as a large number of potential
685 applications in various environments, including neotropical rainforests. However, these applications remain largely underused in this part of the world when compared with other far less diverse regions (this paper; Mulatu et al. 2017; Belle et al. 2019). This is because countries harboring lower diversity are in general more developed economically: infrastructure for molecular-based research is accessible, with
690 associated personnel now relatively well trained for eDNA data generation and analysis. On the other hand, the Nagoya Protocol on Access and Benefit Sharing restricts the access of genetic resources to the country where the sample has been collected, protecting local countries, which are often less economically developed,

from unethical practices by collectors outside of and within the scientific community.

695 We argue that current efforts to develop eDNA-based research in neotropical
countries should be encouraged and strengthened through international
collaborations between researchers from Neotropical countries and researchers from
countries that have already overcome issues relating to methodological application,
technical infrastructure and skill acquisition. Such efforts will enable the acquisition of
700 EBVs related to taxonomic diversity, but also beyond to provide information such as
species distributions or biotic interactions (Table 1), as well as associated underlying
processes. In this final section of the review, we will explore how eDNA can be better
used to improve research methods and their subsequent applications, and in doing
so ultimately contribute to improving conservation programs and management
705 strategies for these hyperdiverse ecosystems.

4.1. Making better sense of eDNA data with better reference databases

A key limit to current eDNA studies in neotropical rainforests is the provision
of relatively poor taxonomic information. This drawback arises in part from the
limitations of eDNA-based methods mentioned above, but is further exacerbated
710 when dealing with neotropical taxa in that they are largely underrepresented in
current DNA reference databases, and/or they have an unresolved taxonomy. This is
particularly true for micro-eukaryotes, for which a significant proportion of OTU and
sequencing reads remain unassigned to a taxon, even at the phylum level (Ritter et
al. 2019; Zinger, Taberlet, et al. 2019). The deficit in DNA references also applies to
715 less cryptic organisms. For example, only 58% of the São Paulo tree flora has
genetic records in international DNA reference databases (Lima et al. 2018). While
eDNA does facilitate the identification of challenging taxa at gross taxonomic levels,
it is therefore unlikely to provide a satisfactory solution for resolving the Linnean

shortfall and provide on its own information on EBVs related to species evolutionary
720 history and functional traits. We hence argue that the future of eDNA remains
inherently intertwined with the continued efforts of taxonomists and naturalists to
sample, identify and store physical specimens in order to complement DNA
reference databases, but also to describe their morphology, evolutionary history,
functional traits, and to solve taxonomic problems (Sheth & Thaker 2017; Dormontt
725 et al. 2018; Pinheiro et al. 2019).

Augmenting the completeness of DNA reference databases is crucial not only
to facilitate the assignment of unknown sequences. It is also essential to ensure, or
verify the plausibility of the retrieved signal, which can be extremely noisy as
discussed above. However, one of the difficulties in improving DNA reference
730 databases is the current lack of consensus when choosing the DNA regions to be
used across studies. Indeed, these may differ from the ones used in curated
reference databases linked to voucher specimens such as the BOLD system for
animals and plants (Ratnasingham & Hebert 2007) or databases dedicated to the
ribosomal clusters for microorganisms (e.g. UNITE, Abarenkov et al. 2010; SILVA,
735 Quast et al. 2013), which only contain gold standard barcoding genes (i.e. COI for
animals, *rbcL* or *matK* for plants, and ITS for fungi). This is because gold standard
barcodes are not necessarily compatible with all applications of eDNA, which often
require DNA primers that target broad taxonomic groups and DNA markers that are
short to suit existing sequencing technologies or the degraded state of eDNA.
740 Conserved priming sites across broad taxonomic groups are often absent within
these gold standard barcodes, an issue highlighted for animals (Bruce E. Deagle et
al. 2014) and plants (Hollingsworth et al. 2011). As a consequence, existing primer
sets targeting classical barcode subregions are often biased toward certain taxa or

on the contrary lack of specificity because they contain too many degenerate bases
745 (B. E. Deagle et al. 2014; Collins et al. 2019). Alternative DNA markers fulfilling
these conditions are often located in mitochondrial or chloroplastic introns or
ribosomal genes (Figure 1) which are better conserved. However these regions also
often exhibit lower taxonomic resolution and are much less referenced in DNA
databases. When choosing a DNA marker, the end-user must hence usually
750 compromise between more precise taxonomic information versus unbiased sampling
of biodiversity. These considerations go beyond the scope of this review and we
refer interested users to dedicated literature on the subject (Hollingsworth et al.
2011; Bruce E. Deagle et al. 2014; Taberlet et al. 2018).

As stated above, the choice of a given DNA marker strongly relies on the
755 biological question to be addressed, the starting material used and because current
reference databases have large deficits in neotropical organisms. Therefore, we
encourage the construction of custom reference databases for the targeted DNA
region from local taxa that are likely to be detected with the eDNA analysis, as done
for example in studies using the mt 12S rRNA gene of neotropical mammals
760 (Kocher, de Thoisy, Francois Catzeflis, et al. 2017) and of Guianese fishes (Cilleros
et al. 2019), or for the ITS1 region for the Basidiomycota of French Guiana (Jaouen
et al. 2019). Although often considered as a costly endeavour, it can be achieved at
relatively low expense (as low as ca. 5 \$USD / specimen) by using freshly collected
specimens, or herbarium/museum collections (e.g. Dormontt et al. 2018) and by
765 multiplexing thousands specimens in a single HTS run. Another promising alternative
that will alleviate the lack of standard DNA markers across studies lies in the building
of “marker-free” DNA reference databases. This is now possible with genome
skimming (Dodsworth 2015), which is similar to metagenome skimming but relies on

a single specimen. This approach produces sequences usable for both gold
770 standard and other barcodes as it generates sequences of the complete organelle
genomes and full nuclear ribosomal regions (Coissac et al. 2016). Although this
remains relatively expensive (as low as ca. 100 \$USD / specimen), it is likely to
become more affordable with continued decreases in sequencing costs.

In addition to compiling DNA information across species, reference databases
775 could complement taxonomic data with ecophysiological characteristics, such as
foliar, root or seed traits for plants, and morphological characteristics such as body
size for animals. Such information would be extremely valuable, allowing eDNA
studies to go beyond the simple description of taxonomic and phylogenetic diversity
of the studied system (Table 1). For example, inferring taxon function or gross
780 ecological traits from eDNA data is now possible for bacteria and fungi through
databases that compile both metabolic, life history traits, or broad lifestyle types (e.g.
PiCrust, Langille et al. 2013; FUNGuild, Nguyen et al. 2016/4). To our knowledge,
such tools are currently not directly available for macro-organisms, although several
databases compiling taxonomic and functional information in a number of groups
785 have been developed (e.g. FishBase, Froese & Pauly 2019; TRY, Kattge et al. 2019;
Atlantic Bird Traits, Rodrigues et al. 2019; or the Global Ants Database, Gibb et al.
2019). Their coupling with DNA reference databases would certainly help advance
the field of eDNA studies to include more process-based approaches.

4.2. Toward eDNA-based occurrence portals for the Neotropics?

790 The greatest strength of eDNA-based approaches is their relative ease of
implementation for both long-term and large-scale monitoring of complex
communities. Even if these data are not necessarily well resolved at the species
level, they still constitute invaluable occurrence data and thereby provide more

information on species distributions, another EBV (Table 1), that is currently largely
795 lacking for neotropical rainforest taxa (Antonelli, Ariza, et al. 2018).

To date, eDNA data and metadata reporting the location, time and exact
protocol of the sampling are disseminated individually using study specific web
repositories, as in data papers (e.g. Murienne et al. 2019) or more general
repositories (e.g. Dryad, datadryad.org , Zenodo, zenodo.org ; or the Short Read
800 Archives from Genbank, www.ncbi.nlm.nih.gov/sra). However, the construction of
dedicated portals compiling eDNA-based taxa occurrence can now be envisioned for
all neotropical rainforests and beyond following the examples of the occurrence
portal GBIF (www.gbif.org), the BOLD system (www.boldsystem.org) which
integrates DNA data with occurrence, or the EMP (www.earthmicrobiome.org) which
805 compiles occurrence and diversity of microbial taxa across the globe. The success of
such an endeavour depends on the adequate standardization of data, a challenge
given that ecological signal from eDNA data is influenced by the technique used, the
DNA region targeted, and the protocols of molecular biology and bioinformatics
chosen. While defining standards for such purposes will certainly facilitate the
810 integration of data across studies, it is also likely that this will be difficult to apply to
all desired situations, which may ultimately undermine scientific advances. Several
alternatives have been proposed to circumvent this issue. The first is to adopt
sequence taxonomy classification as a standard unit (Ramirez et al. 2018). As
highlighted above, such an approach heavily depends on taxonomic expertise and
815 enriched DNA reference databases to make the best use of eDNA data. The second
is the implementation of “eDNA biobanking”, i.e. the development of storage facilities
for eDNA samples that could be reused with different technologies (Jarman et al.
2018).

Although less precise than traditionally collected occurrence data, which are
820 limited in other ways, sections 3.1-2 demonstrate how eDNA-based studies can
unveil the abiotic determinants of neotropical diversity. Increasing eDNA-based
taxonomic inventories across environmental gradients will provide insights into taxa
environmental/physiological tolerances/preferences (Table 1), information which
remains scarce in neotropical rainforests. From a more applied perspective,
825 increasing eDNA sampling across land use gradients will enable the identification of
indicator taxa for environmental impacts or umbrella taxa that are specific to this
ecoregion. However, this application currently remains limited by the difficulty in
retrieving population size information from eDNA as discussed above. Without
significant developments for this particular aspect, eDNA-based approaches will
830 likely remain of limited utility when assessing the conservation status of neotropical
taxa.

4.3. Shedding new light on biotic interactions

The increasing use of eDNA will also certainly fill the current gap of
knowledge on species interactions (Table 1) by improving the description of complex
835 and multi-trophic communities for both well studied taxa and more elusive
organisms. Such assessments are urgently needed at a time where environmental
changes already cause direct species loss and cascading extinction via bottom-up or
top-down effects, especially in tropical ecosystems, including neotropical rainforests,
where biotic interactions are often expected to be highly specific (Barnes et al.
840 2017).

It is now possible to analyse the diet of a particular species by collecting
faeces, gut contents or even the DNA traces herbivores or pollinators leave on plants
([Koskinen et al. 2019](#); [Thomsen & Sigsgaard 2019](#)). These applications are routinely

used in temperate ecosystems (Bohmann et al. 2014; Taberlet et al. 2018; Alberdi et
845 al. 2019). By contrast, only few diet studies have been performed on neotropical
organisms, i.e. on tapirs from French Guiana (Hibert et al. 2013), on white-face
capuchins from Costa Rica (Mallott et al. 2018), on neotropical vampire bats
(Bohmann et al. 2018) and rodents (Lopes et al. 2015), and on particular arthropods
(Paula et al. 2016; Kocher, de Thoisy, François Catzeflis, et al. 2017; Rodgers et al.
850 2017). New protocols of diet assessment based on faeces or gut contents are now
available and optimized to reduce host DNA concentration in DNA extracts ([e.g.
Krehenwinkel et al. 2017](#)). Such improvements considerably reduce the costs
associated with molecular treatments and sequencing and hence allow for the
implementation of large-scale studies of full food-webs composed of understudied
855 and hyperdiverse taxa. This will certainly enable improved characterisation of trophic
niche and breadth for many neotropical taxa, thereby improving documentation of
feeding behaviour in relation to species functional traits and competitive interactions.

Likewise, eDNA can be used to unravel plant-pollinator networks. Pollinators
yield substantial amounts of pollen on their bodies, and conversely the surfaces of
860 leaves and flower petals also harbour traces of DNA belonging to visiting pollinators.
This material can be used to build reliable plant-pollinator insect interactions, as
shown in temperate ecosystems (Pornon et al. 2016; Thomsen & Sigsgaard 2019).
The applicability of the methods has, to our knowledge, not yet been tested in
neotropical rainforests and remains to be critically assessed due to the particular
865 climatic conditions, much greater richness, and also the greater amount of vertebrate
pollinators in these ecosystems, which can be more challenging to sample than
arthropods.

Similarly, improved understanding of host-microbiota interactions can have

important implications for threatened species conservation (West et al. 2019). This
870 can be done by studying microbial communities occurring at the surface or within
larger organisms in a more comprehensive way than before. So far, existing studies
have principally aimed to describe microbial communities and, in some cases, their
assembly mechanisms. This has been done mostly for leaf or root endophytes in
trees (Kembel et al. 2014; Bonfim et al. 2016; Schroeder et al. 2019; Donald et al.
875 2020), palms (Donald et al. 2019), grasses (Higgins et al. 2014) or fern species (Del
Olmo-Ruiz & Elizabeth Arnold 2017) and for the microbiota of frogs to assess its
potential role in the resistance to the chytrid fungus *Batrachochytrium dendrobatidis*
(Hughey et al. 2017; Catenazzi et al. 2018). To our knowledge the microbiota
associated with neotropical mammals as been only assessed for the endangered
880 Andean Bear (Borbón-García et al. 2017), and the same holds true for the microbiota
of invertebrates, which has been so far mostly studied on emblematic arthropods
such as ants (Sapountzis et al. 2015; Pringle & Moreau 2017). Although few studies
have shown experimentally that the plant microbiota can promote the growth and
survival of seedlings (Christian et al. 2017; Leroy et al. 2019), much remains to be
885 done to understand the functional contribution of the microbiota to host health, and
how this can affect community level distribution or diversity patterns (e.g. Janzen-
Connell effects accounting for the whole microbiota (Janzen 1970, Connell 1978)).

The approaches discussed above mostly enable reconstructing bi- or tri-
partite networks, but future applications are likely to span the whole ecological
890 network to advance our understanding of the resistance and resilience of biological
communities to disturbance. Indeed, eDNA can provide co-occurrence data for
multiple taxonomic, functional and trophic groups retrieved from soil, sediments or
water. While these co-occurrences do not represent biological interactions *per se*,

these can assist in the discovery of a large variety of interactions at larger
895 temporal/spatial scales, provided that these inferences are evaluated with *a priori*
knowledge of the system or statistical tools (Vacher et al. 2016).

4.4 Epidemiology and healthcare

Neotropical rainforests ecosystems harbours many emerging infectious
diseases, and use of eDNA for monitoring their agents or vectors has enormous
900 implications for human health. Most parasites and pathogens are usually only
detected when aggregating on or in their hosts and without eDNA, their detection
remains challenging in the environment (Bass et al. 2015). Recent results from
Sengupta *et al.* (2019) indicate that free living larval aquatic phases of *Schistosoma*
can be detected with eDNA from water samples, opening an avenue to the control of
905 this neglected tropical disease affecting >250 million people worldwide, mainly in
Africa, but with human infestations in several regions of South and Central America.
Although using eDNA as diagnostic evidence for pathogens or parasites requires
extensive validation before it is used in notification procedures or detection programs
(Bass et al. 2015), developments of such methods in the region would considerably
910 improve the monitoring and fight against agents of tropical diseases.

A number of human diseases require a vector, typically an insect, to transmit
the pathogen and surveillance programs usually rely on monitoring potential vector
populations. Such a task can prove daunting given a single night of trapping using a
standard CDC trap (Center for Disease and Control) could yield thousands of
915 mosquitoes / sandflies that need to be identified to species level. eDNA-based
approaches greatly reduce the time and costs related to these identifications
(Kocher, Gantier, et al. 2017; Talaga et al. 2017), and could be used for routine
monitoring of vector species and help in the control of vector-borne diseases.

Classical epidemiological monitoring programs largely focused on pathogens
920 or their vectors, yet it is increasingly recognised that the prediction of transmission
risk should include a better understanding of the ecosystem as a whole. This is
particularly true in a context of biodiversity erosion and habitat degradation, which
could be connected to the emergence of diseases as a result of trophic food-web
modifications. For example, deforestation has been suggested to lead to the
925 emergence of diseases such as malaria (Vittor et al. 2009) or Buruli ulcer (Morris et
al. 2016), through reductions in diversity and modifications to the species
composition of aquatic food-webs. Because eDNA-based methods can provide not
only rapid information on pathogens and vectors, but also a broad characterisation of
the whole ecological network, we believe they will strongly modify our approach to
930 epidemiology and understanding of disease emergence in the next few years.

4.5. Conservation and impact assessments in neotropical rainforests and beyond

Managing ecosystems and biodiversity requires efficient detection of the
species of interest, but also standard, cost- and time-effective protocols that can be
935 implemented repeatedly across large spatial scales and through time, with low, or
limited impact on organisms. Such protocols are currently not available for
monitoring neotropical rainforests and, more generally, neotropical ecosystems. This
review shows that eDNA-based methods fulfil these criteria while enabling
characterisation of the taxonomic composition of multiple trophic communities, and
940 could even constitute proxies of other EBVs. These methods complement remote
sensing tools since eDNA provides information at a much finer taxonomic resolution,
thereby better complying with some of the Aichi Targets that focus on endangered
and invasive species (Marques et al. 2014; Bush et al. 2017). Their use could hence
greatly facilitate the establishment of Rapid Biodiversity Assessment programs.

945 eDNA-based rapid biodiversity assessments hold great potential for the
evaluation of environmental impacts, in particular for the ever increasing
unsustainable use of land in neotropical rainforests, as exemplified with soil
organisms (e.g. Franco et al. 2019). Likewise, eDNA-based methods will be able to
help evaluate the success of different restoration and conservation strategies
950 (Fernandes et al. 2018). However, the use of eDNA for informing management and
political decisions will inherently require the development of quick and standardised
sampling protocols that work across varying environment types and can be easily
applied by practitioners. Beyond standardisation, which we show here to be a
challenging issue, such an application implies the development of biotic integrity
955 indices that are easily transferable to stakeholders, resource managers, and policy
makers, and eDNA research is still in its infancy on this particular matter (Cordier et
al. 2019). Nevertheless, we are confident that these limits can and will be overcome
in the near future.

Aside from rainforests, the Neotropics holds large areas of other biomes that
960 face threats that are not necessarily the same as for tropical rainforests but whose
diversity remains poorly described with both traditional and eDNA methods
(Antonelli, Ariza, et al. 2018). For example, eDNA could be particularly relevant to
describe and monitor white-sand ecosystems which harbour a unique flora and
fauna (Fine & Baraloto 2016), but which are currently threatened by increases in
965 cattle ranching, deforestation for firewood or mining for sand (Ferreira et al. 2013).
Likewise, it could be used for savanna and dry forest conservation, habitats which
currently experience greater pressures than other neotropical biomes, typically as
deforestation, localised human disturbance and increasing drought frequency and
intensity (Strassburg et al. 2017; González-M et al. 2018). For example, the revision

970 of Brazil's Forest Code in 2012, the Cerrado (Brazilian savanna) indirectly
encouraged Brazilian agribusiness to invest in this biome (Soares-Filho et al. 2014;
Strassburg et al. 2017). Estuaries, including mangrove forests, also represent
neglected and threatened habitats in the Neotropics, whilst harbouring rich
communities and serving as a nursery for many fish and crustaceans (Mumby et al.
975 2004). In these environments, the turbidity and strong water currents make species
inventories difficult, a limit that could be circumvented with eDNA (Stoeckle et al.
2017). A last example is the Pantanal biome, a savanna wetland which hosts a
unique diversity, supports essential ecosystem services, and is currently under
strong human pressure (Alho 2008). Descriptive and monitoring studies using eDNA
980 analysis in these neglected yet important ecosystems would therefore help to better
characterize their diversity and how they respond to various pressures.

From a more basic perspective, the possibility to implement comprehensive,
large-scale and long-term biodiversity observatories will certainly help to gain
insights into the origin and maintenance of neotropical biodiversity, and its singularity
985 in many ecosystems. Reconstructing past ecosystems from ancient DNA (Thomsen
& Willerslev 2015) would be extremely valuable in such a case, and would further
improve our understanding of the long-term dynamics of neotropical ecosystems,
and hence better predict their future. However, it remains unclear whether eDNA can
persist in the long term in tropical ecosystems and further studies are required in this
990 area. Nevertheless, long term dynamics can be assessed through monitoring
initiatives along transitions between different biomes. For example, savannas and
dry forests constitute transitory or alternative stable states of rainforests in response
to global changes (Nepstad et al. 2008; Dexter et al. 2018), and monitoring these
sites through eDNA should provide useful information on their dynamics, enabling

995 the identification of early warning markers of major ecological transitions. Acquisition
of such data will prove valuable for anticipating the status of these environments and
prioritizing corresponding conservation or restoration actions to mitigate such
transitions.

1000

Acknowledgements

Some of the research presented in this paper received funding from the French Agence Nationale de la Recherche (ANR) (METABAR: ANR-11-BSV7-0020; GlobNets: ANR-16-CE02-0009; DEBIT: ANR-17-CE02-0007-01), from

1005 'Investissement d'Avenir' grants managed by the ANR (CEBA: ANR-10-LABX-25-01; TULIP: ANR-10-LABX-0041), and from São Paulo Research Foundation (FAPESP #2013/50741-7).

References

- 1010 Abarenkov, K. et al., 2010. The UNITE database for molecular identification of fungi - recent updates and future perspectives. *New phytologist*, 186(2), pp.281–285.
- Alberdi, A. et al., 2019. Promises and pitfalls of using high-throughput sequencing for diet analysis. *Molecular Ecology Resources*, 19(2), pp.327–348.
- 1015 Alho, C.J.R., 2008. Biodiversity of the Pantanal: response to seasonal flooding regime and to environmental degradation. *Brazilian Journal of Biology*, 68(4 Suppl), pp.957–966.
- Allard, L. et al., 2014. Electrofishing efficiency in low conductivity neotropical streams: towards a non-destructive fish sampling method. *Fisheries Management and Ecology*, 21(3), pp.234–243.
- 1020 Andújar, C. et al., 2015. Phylogenetic community ecology of soil biodiversity using mitochondrial metagenomics. *Molecular Ecology*, 24(14), pp.3603–3617.
- Antonelli, A., Zizka, A., et al., 2018. Amazonia is the primary source of Neotropical biodiversity. *Proceedings of the National Academy of Sciences of the United States of America*, 115(23), pp.6034–6039.
- 1025 Antonelli, A., Ariza, M., et al., 2018. Conceptual and empirical advances in Neotropical biodiversity research. *PeerJ*, 6, p.e5644.
- Araújo, A.S.F. et al., 2014. Soil bacterial diversity in degraded and restored lands of Northeast Brazil. *Antonie van Leeuwenhoek*, 106(5), pp.891–899.
- 1030 Bálint, M. et al., 2018. Accuracy, limitations and cost efficiency of eDNA-based community survey in tropical frogs. *Molecular Ecology Resources*, 18(6), pp.1415–1426.
- Bálint, M. et al., 2016. Millions of reads, thousands of taxa: microbial community

structure and associations analyzed via marker genes. *FEMS Microbiology Reviews*, 40(5), pp.686–700.

- 1035 Barberán, A. et al., 2015. Relating belowground microbial composition to the taxonomic, phylogenetic, and functional trait distributions of trees in a tropical forest. *Ecology Letters*, 18(12), pp.1397–1405.
- Bardgett, R.D. & van der Putten, W.H., 2014. Belowground biodiversity and ecosystem functioning. *Nature*, 515(7528), pp.505–511.
- 1040 Barnes, A.D. et al., 2017. Direct and cascading impacts of tropical land-use change on multi-trophic biodiversity. *Nature Ecology & Evolution*, 1(10), pp.1511–1519.
- Barnes, M.A. et al., 2014. Environmental conditions influence eDNA persistence in aquatic systems. *Environmental Science & Technology*, 48(3), pp.1819–1827.
- Barnes, M.A. & Turner, C.R., 2015. The ecology of environmental DNA and implications for conservation genetics. *Conservation Genetics*, pp.1–17.
- 1045 Bass, D. et al., 2015. Diverse Applications of Environmental DNA Methods in Parasitology. *Trends in Parasitology*, 31(10), pp.499–513.
- Basset, Y. et al., 2015. Arthropod distribution in a tropical rainforest: tackling a four dimensional puzzle. *PloS One*, 10(12), p.e0144110.
- 1050 Belle, C.C., Stoeckle, B.C. & Geist, J., 2019. Taxonomic and geographical representation of freshwater environmental DNA research in aquatic conservation. *Aquatic conservation: marine and freshwater ecosystems*. Available at: [https://doi.org/10.1002/aqc.3208@10.1002/\(ISSN\)1099-0755.Using-environmental-DNA-in-freshwater-conservation](https://doi.org/10.1002/aqc.3208@10.1002/(ISSN)1099-0755.Using-environmental-DNA-in-freshwater-conservation).
- 1055 Benzing, D.H., 2000. *Bromeliaceae: Profile of an Adaptive Radiation*, Cambridge University Press.
- Bista, I. et al., 2017. Annual time-series analysis of aqueous eDNA reveals ecologically relevant dynamics of lake ecosystem biodiversity. *Nature Communications*, 8, p.14087.
- 1060 Bithell, S.L. et al., 2014. DNA analysis of soil extracts can be used to investigate fine root depth distribution of trees. *AoB Plants*, 7, p.lu091.
- Bohmann, K. et al., 2014. Environmental DNA for wildlife biology and biodiversity monitoring. *Trends in Ecology & Evolution*, 29, pp.358–367.
- 1065 Bohmann, K. et al., 2018. Using DNA metabarcoding for simultaneous inference of common vampire bat diet and population structure. *Molecular Ecology Resources*, 18(5), pp.1050–1063.
- Bonfim, J.A. et al., 2016. Dark septate endophytic fungi of native plants along an altitudinal gradient in the Brazilian Atlantic forest. *Fungal Ecology*, 20, pp.202–210.

- 1070 Borbón-García, A. et al., 2017. Captivity shapes the gut microbiota of Andean bears: insights into health surveillance. *Frontiers in Microbiology*, 8, p.1316.
- Boussarie, G. et al., 2018. Environmental DNA illuminates the dark diversity of sharks. *Science Advances*, 4(5), p.eaap9661.
- Bowles, D.E. & Courtney, G.W., 2018. Advances in aquatic insect systematics and biodiversity in the Neotropics: introduction. *Aquatic Insects*, 39(2-3), pp.89–93.
- 1075 Brozio, S. et al., 2017. Development and Application of an eDNA Method to Detect the Critically Endangered Trinidad Golden Tree Frog (*Phytotriades auratus*) in Bromeliad Phytotelmata. *PLOS ONE*, 12(2), p.e0170619. Available at: <http://dx.doi.org/10.1371/journal.pone.0170619>.
- 1080 de Bruijn, F.J. ed., 2011. *Handbook of Molecular Microbial Ecology I: Metagenomics and Complementary Approaches*, John Wiley & Sons.
- Bush, A. et al., 2017. Connecting Earth observation to high-throughput biodiversity data. *Nature Ecology & Evolution*, 1(7), p.176.
- 1085 Calderón–Sanou, I. et al., 2019. From environmental DNA sequences to ecological conclusions: How strong is the influence of methodological choices? *Journal of biogeography*. Available at: <https://onlinelibrary.wiley.com/doi/abs/10.1111/jbi.13681>.
- Callahan, B.J., McMurdie, P.J. & Holmes, S.P., 2017. Exact sequence variants should replace operational taxonomic units in marker-gene data analysis. *The ISME Journal*, 11(12), pp.2639–2643.
- 1090 Cantera, I. et al., 2019. Optimizing environmental DNA sampling effort for fish inventories in tropical streams and rivers. *Scientific Reports*, 9(1), p.3085.
- Carney, K.M., Matson, P.A. & Bohannon, B.J.M., 2004. Diversity and composition of tropical soil nitrifiers across a plant diversity gradient and among land-use types. *Ecology Letters*, 7(8), pp.684–694.
- 1095 Carpenter, M.L. et al., 2013. Pulling out the 1%: whole-genome capture for the targeted enrichment of ancient DNA sequencing libraries. *American Journal of Human Genetics*, 93, pp.852–864.
- Castello, L. et al., 2013. The vulnerability of Amazon freshwater ecosystems. *Conservation Letters*, 6(4), pp.217–229.
- 1100 Catenazzi, A. et al., 2018. Widespread elevational occurrence of antifungal bacteria in Andean amphibians decimated by disease: a complex role for skin symbionts in defense against Chytridiomycosis. *Frontiers in Microbiology*, 9, p.465.
- 1105 Chao, A., Chiu, C.-H. & Jost, L., 2014. Unifying species diversity, phylogenetic diversity, functional diversity, and related similarity and differentiation measures through Hill numbers. *Annual Reviews of Ecology, Evolution and Systematics*, 45, pp.297–324.

- Chaplin-Kramer, R. et al., 2019. Global modeling of nature's contributions to people. *Science*, 366(6462), pp.255–258.
- 1110 Chariton, A.A. et al., 2010. Ecological assessment of estuarine sediments by pyrosequencing eukaryotic ribosomal DNA. *Frontiers in Ecology and the Environment*, 8(5), pp.233–238.
- Christian, N. et al., 2017. Exposure to the leaf litter microbiome of healthy adults protects seedlings from pathogen damage. *Proceedings of the Royal Society B: Biological Sciences*, 284(1858), p.20170641.
- 1115 Cilleros, K. et al., 2019. Unlocking biodiversity and conservation studies in high-diversity environments using environmental DNA (eDNA): A test with Guianese freshwater fishes. *Molecular Ecology Resources*, 19(1), pp.27–46.
- Coissac, E. et al., 2016. From barcodes to genomes: extending the concept of DNA barcoding. *Molecular Ecology*, 25(7), pp.1423–1428.
- 1120 Coissac, E., Riaz, T. & Puillandre, N., 2012. Bioinformatic challenges for DNA metabarcoding of plants and animals. *Molecular Ecology*, 21, pp.1834–1847.
- Collins, R.A. et al., 2019. Non-specific amplification compromises environmental DNA metabarcoding with COI D. Yu, ed. *Methods in ecology and evolution / British Ecological Society*, 10(11), pp.1985–2001.
- 1125 Cordier, T. et al., 2019. Embracing environmental genomics and machine learning for routine biomonitoring. *Trends in Microbiology*, 27(5), pp.387–397.
- Couceiro, S.R.M. et al., 2012. A macroinvertebrate multimetric index to evaluate the biological condition of streams in the Central Amazon region of Brazil. *Ecological indicators*, 18, pp.118–125.
- 1130 Creedy, T.J., Ng, W.S. & Vogler, A.P., 2019. Toward accurate species-level metabarcoding of arthropod communities from the tropical forest canopy. *Ecology and Evolution*, 9(6), pp.3105–3116.
- 1135 Czech, L., Barbera, P. & Stamatakis, A., 2019. Methods for automatic reference trees and multilevel phylogenetic placement. *Bioinformatics*, 35(7), pp.1151–1158.
- Dalling, J.W. et al., 2016. Nutrient availability in tropical rain forests: the paradigm of phosphorus limitation. In *Tropical tree physiology*. Cham: Springer, pp. 261–273.
- 1140 Davidson, E.A. et al., 2012. The Amazon basin in transition. *Nature*, 481(7381), pp.321–328.
- Deagle, B.E. et al., 2014. DNA metabarcoding and the COI marker: not a perfect match. *Biology letters*, 10, p.UNSP 20140562.
- Deagle, B.E. et al., 2014. DNA metabarcoding and the cytochrome c oxidase subunit I marker: not a perfect match. *Biology letters*, 10(9). Available at:

- 1145 <http://dx.doi.org/10.1098/rsbl.2014.0562>.
- Dedieu, N. et al., 2016. A multimetric macroinvertebrate index for the implementation of the European Water Framework Directive in French Guiana, East Amazonia. *River research and applications*, 32(3), pp.501–515.
- 1150 Deiner, K. et al., 2017. Environmental DNA metabarcoding: Transforming how we survey animal and plant communities. *Molecular Ecology*, 26(21), pp.5872–5895.
- Del Olmo-Ruiz, M. & Elizabeth Arnold, A., 2017. Community structure of fern-affiliated endophytes in three neotropical forests. *Journal of tropical ecology*, 33(1), pp.60–73.
- 1155 Dexter, K.G. et al., 2018. Inserting tropical dry forests into the discussion on biome transitions in the tropics. *Front. Ecol. Evol.*, 6, p.104.
- Díaz, S. et al., 2019. Pervasive human-driven decline of life on Earth points to the need for transformative change. *Science*, 366(6471). Available at: <http://dx.doi.org/10.1126/science.aax3100>.
- 1160 Dickie, I.A., Boyer, S. & Buckley, H.L., 2018. Towards robust and repeatable sampling methods in eDNA-based studies. *Molecular Ecology*, 18(5), pp.940–952.
- Dodsworth, S., 2015. Genome skimming for next-generation biodiversity analysis. *Trends in Plant Science*, 20(9), pp.525–527.
- 1165 Doi, H. et al., 2015. Droplet digital polymerase chain reaction (PCR) outperforms real-time PCR in the detection of environmental DNA from an invasive fish species. *Environmental Science & Technology*, 49(9), pp.5601–5608.
- Donald, J. et al., 2020. A test of community assembly rules using foliar endophytes from a tropical forest canopy. *Journal of Ecology*. Available at:
- 1170 <http://dx.doi.org/10.1111/1365-2745.13344>.
- Donald, J. et al., 2019. Tropical Palm Endophytes Exhibit Low Competitive Structuring When Assessed Using Co-occurrence and Antipathogen Activity Analysis. *Frontiers in Forests and Global Change*, 2, p.86.
- 1175 Dormontt, E.E. et al., 2018. Advancing DNA barcoding and metabarcoding applications for plants requires systematic analysis of herbarium collections-an Australian perspective. *Frontiers in Ecology and Evolution*, 6, p.134.
- Drenovsky, R.E. et al., 2010. Land use and climatic factors structure regional patterns in soil microbial communities. *Global Ecology and Biogeography*, 19(1), pp.27–39.
- 1180 Elbrecht, V. & Leese, F., 2017. PrimerMiner : an r package for development and in silico validation of DNA metabarcoding primers. *Methods in Ecology and Evolution*, 8(5), pp.622–626.

- 1185 Esling, P., Lejzerowicz, F. & Pawlowski, J., 2015. Accurate multiplexing and filtering for high-throughput amplicon-sequencing. *Nucleic Acids Research*, 43(5), pp.2513–2524.
- FAO, 2019. *The State of the World's Biodiversity for Food and Agriculture* J. Bélanger & D. Pilling, eds., Rome: FAO Commission on Genetic Resources for Food and Agriculture Assessments.
- 1190 Fernandes, K. et al., 2018. DNA metabarcoding-a new approach to fauna monitoring in mine site restoration. *Restoration Ecology*, 26(6), pp.1098–1107.
- Ferreira, L.V. et al., 2013. A extração ilegal de areia como causa do desaparecimento de campinas e campinaranas no Estado do Pará, Brasil. *Pesquisas (Botânica)*, 64, pp.157–173.
- 1195 Ficetola, G.F. et al., 2010. An in silico approach for the evaluation of DNA barcodes. *BMC Genomics*, 11, p.434.
- Ficetola, G.F., Taberlet, P. & Coissac, E., 2016. How to limit false positives in environmental DNA and metabarcoding? *Molecular Ecology Resources*, 16, pp.604–607.
- 1200 Fine, P.V.A. & Baraloto, C., 2016. Habitat Endemism in White-sand Forests: Insights into the Mechanisms of Lineage Diversification and Community Assembly of the Neotropical Flora. *Biotropica*, 48(1), pp.24–33.
- Fragoso, C. & Lavelle, P., 1992. Earthworm communities of tropical rain forests. *Soil Biology & Biochemistry*, 24(12), pp.1397–1408.
- 1205 Franco, A.L.C. et al., 2019. Amazonian deforestation and soil biodiversity. *Conservation Biology*, 33(3), pp.590–600.
- Froese, R. & Pauly, D., 2019. FishBase. *World Wide Web electronic publication*. Available at: www.fishbase.org.
- Gibb, H. et al., 2019. The Global Ants Database. Available at: <http://globalants.org/>.
- 1210 Gilbert, J.A. et al., 2012. Defining seasonal marine microbial community dynamics. *The ISME journal*, 6(2), pp.298–308.
- González-M, R. et al., 2018. Disentangling the environmental heterogeneity, floristic distinctiveness and current threats of tropical dry forests in Colombia. *Environmental Research Letters*, 13(4), p.045007.
- 1215 Grossman, J.M. et al., 2010. Amazonian anthrosols support similar microbial communities that differ distinctly from those extant in adjacent, unmodified soils of the same mineralogy. *Microbial Ecology*, 60(1), pp.192–205.
- Guillera-Aroita, G. et al., 2017. Dealing with false-positive and false-negative errors about species occurrence at multiple levels. *Methods in ecology and evolution / British Ecological Society*, 8(9), pp.1081–1091.

- 1220 Hacquard, S. et al., 2015. Microbiota and Host Nutrition across Plant and Animal Kingdoms. *Cell Host & Microbe*, 17(5), pp.603–616.
- Harper, L.R. et al., 2018. Needle in a haystack? A comparison of eDNA metabarcoding and targeted qPCR for detection of the great crested newt (*Triturus cristatus*). *Ecology and Evolution*, 8(12), pp.6330–6341.
- 1225 Harwood, V.J. et al., 2014. Microbial source tracking markers for detection of fecal contamination in environmental waters: relationships between pathogens and human health outcomes. *FEMS Microbiology Reviews*, 38(1), pp.1–40.
- Hebert, P.D.N. et al., 2003. Biological identification through DNA barcodes. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 270, pp.313–321.
- 1230 Hibert, F. et al., 2013. Unveiling the diet of elusive rainforest herbivores in next generation sequencing era? The tapir as a case study. *PLoS One*, 8, p.e60799.
- Higgins, K.L. et al., 2014. Communities of fungal endophytes in tropical forest grasses: highly diverse host- and habitat generalists characterized by strong spatial structure. *Fungal Ecology*, 8, pp.1–11.
- 1235 Hollingsworth, P.M., Graham, S.W. & Little, D.P., 2011. Choosing and using a plant DNA barcode. *PLoS One*, 6(5), p.e19254.
- Hortal, J. et al., 2015. Seven shortfalls that beset large-scale knowledge of biodiversity. *Annual Review of Ecology, Evolution, and Systematics*, 46(1), pp.523–549.
- 1240 Hughey, M.C. et al., 2017. Skin bacterial microbiome of a generalist Puerto Rican frog varies along elevation and land use gradients. *PeerJ*, 5, p.e3688.
- Jaouen, G. et al., 2019. Fungi of French Guiana gathered in a taxonomic, environmental and molecular dataset. *Scientific data*, 6(1), p.206.
- 1245 Jarman, S.N., Berry, O. & Bunce, M., 2018. The value of environmental DNA biobanking for long-term biomonitoring. *Nature Ecology & Evolution*, 2(8), pp.1192–1193.
- Jenkins, C.N., Pimm, S.L. & Joppa, L.N., 2013. Global patterns of terrestrial vertebrate diversity and conservation. *Proceedings of the National Academy of Sciences of the United States of America*, 110(28), pp.E2602–10.
- 1250 Jetz, W. et al., 2019. Essential biodiversity variables for mapping and monitoring species populations. *Nature ecology & evolution*, 3(4), pp.539–551.
- Jones, F.A. et al., 2011. The roots of diversity: below ground species richness and rooting distributions in a tropical forest revealed by DNA barcodes and inverse modeling. *PLoS One*, 6, p.e24506.
- 1255 Kattge, J. et al., 2019. TRY plant trait database - enhanced coverage and open access. *Global change biology*. Available at:

<http://dx.doi.org/10.1111/gcb.14904>.

- 1260 Kembel, S.W. et al., 2014. Relationships between phyllosphere bacterial communities and plant functional traits in a neotropical forest. *Proceedings of the National Academy of Sciences of the United States of America*, 111(38), pp.13715–13720.
- Kim, J.-S. et al., 2007. Bacterial diversity of terra preta and pristine forest soil from the Western Amazon. *Soil Biology & Biochemistry*, 39(2), pp.684–690.
- 1265 Kirshtein, J.D. et al., 2007. Quantitative PCR detection of *Batrachochytrium dendrobatidis* DNA from sediments and water. *Diseases of Aquatic Organisms*, 77(1), pp.11–15.
- 1270 Kivlin, S.N. & Hawkes, C.V., 2016a. Temporal and Spatial Variation of Soil Bacteria Richness, Composition, and Function in a Neotropical Rainforest. *PloS One*, 11(7), p.e0159131.
- Kivlin, S.N. & Hawkes, C.V., 2016b. Tree species, spatial heterogeneity, and seasonality drive soil fungal abundance, richness, and composition in Neotropical rainforests. *Environmental Microbiology*, 18(12), pp.4662–4673.
- 1275 Kocher, A., de Thoisy, B., Catzeflis, F., et al., 2017. Evaluation of short mitochondrial metabarcodes for the identification of Amazonian mammals. *Methods in Ecology and Evolution*, 8(10), pp.1276–1283.
- Kocher, A., de Thoisy, B., Catzeflis, F., et al., 2017. iDNA screening: Disease vectors as vertebrate samplers. *Molecular Ecology*, 26(22), pp.6478–6486.
- 1280 Kocher, A., Gantier, J.-C., et al., 2017. Vector soup: high-throughput identification of Neotropical phlebotomine sand flies using metabarcoding. *Molecular Ecology Resources*, 17(2), pp.172–182.
- Langille, M.G.I. et al., 2013. Predictive functional profiling of microbial communities using 16S rRNA marker gene sequences. *Nature Biotechnology*, 31(9), pp.814–821.
- 1285 Lauber, C.L. et al., 2009. Pyrosequencing-based assessment of soil pH as a predictor of soil bacterial community structure at the continental scale. *Applied and Environmental Microbiology*, 75, pp.5111–5120.
- 1290 Lentendu, G. et al., 2019. Protist Biodiversity and Biogeography in Lakes From Four Brazilian River--Floodplain Systems. *Journal of Eukaryotic Microbiology*, 66(4), pp.592–599.
- Leroy, C. et al., 2019. How significant are endophytic fungi in bromeliad seeds and seedlings? Effects on germination, survival and performance of two epiphytic plant species. *Fungal Ecology*, 39, pp.296–306.
- 1295 Leroy, C. et al., 2016. The contribution of microorganisms and metazoans to mineral nutrition in bromeliads. *Journal of Plant Ecology*, 9(3), pp.241–255.

- Levy-Booth, D.J. et al., 2007. Cycling of extracellular DNA in the soil environment. *Soil Biology & Biochemistry*, 39(12), pp.2977–2991.
- 1300 Lievens, B. et al., 2006. Real-time PCR for detection and quantification of fungal and oomycete tomato pathogens in plant and soil samples. *Plant Science*, 171(1), pp.155–165.
- Li, F. et al., 2018. Application of Environmental DNA Metabarcoding for Predicting Anthropogenic Pollution in Rivers. *Environmental Science & Technology*, 52(20), pp.11708–11719.
- 1305 Lima, R.A.F. de et al., 2018. Can plant DNA barcoding be implemented in species-rich tropical regions? A perspective from São Paulo State, Brazil. *Genetics and Molecular Biology*, 41(3), pp.661–670.
- Lim, N.K.M. et al., 2016. Next-generation freshwater bioassessment: eDNA metabarcoding with a conserved metazoan primer reveals species-rich and reservoir-specific communities. *Royal Society open science*, 3(11), p.160635.
- 1310 Linard, B. et al., 2015. Metagenome skimming of insect specimen pools: potential for comparative genomics. *Genome Biology and Evolution*, 7(6), pp.1474–1489.
- Lopes, C.M. et al., 2015. DNA metabarcoding diet analysis for species with parapatric vs sympatric distribution: a case study on subterranean rodents. *Heredity*, 114, pp.525–536.
- 1315 Lopes, C.M. et al., 2017. eDNA metabarcoding: a promising method for anuran surveys in highly diverse tropical forests. *Molecular Ecology Resources*, 17(5), pp.904–914.
- 1320 Louca, S. et al., 2017. Functional structure of the bromeliad tank microbiome is strongly shaped by local geochemical conditions. *Environmental Microbiology*, 19(8), pp.3132–3151.
- Louca, S. et al., 2016. High taxonomic variability despite stable functional structure across microbial communities. *Nature Ecology & Evolution*, 1(1), p.15.
- Lynggaard, C. et al., 2019. Vertebrate diversity revealed by metabarcoding of bulk arthropod samples from tropical forests. *Environmental DNA*, 56, p.1637.
- 1325 Mächler, E. et al., 2014. Utility of environmental DNA for monitoring rare and indicator macroinvertebrate species. *Freshwater Science*, 33(4), pp.1174–1183.
- Maggia, M.E. et al., 2017. DNA Metabarcoding of Amazonian Ichthyoplankton Swarms. *PloS One*, 12(1), p.e0170009.
- 1330 Mallott, E.K., Garber, P.A. & Malhi, R.S., 2018. trnL outperforms rbcL as a DNA metabarcoding marker when compared with the observed plant component of the diet of wild white-faced capuchins (*Cebus capucinus*, Primates). *PloS One*, 13(6), p.e0199556.
- Mariac, C. et al., 2018. Metabarcoding by capture using a single COI probe (MCSP)

- 1335 to identify and quantify fish species in ichthyoplankton swarms. *PloS one*, 13(9), p.e0202976.
- Marques, A. et al., 2014. A framework to identify enabling and urgent actions for the 2020 Aichi Targets. *Basic and Applied Ecology*, 15(8), pp.633–638.
- Maruyama, A. et al., 2014. The release rate of environmental DNA from juvenile and adult fish. *PLoS One*, 9, p.e114639.
- 1340 Matsen, F.A., Kodner, R.B. & Armbrust, E.V., 2010. pplacer: linear time maximum-likelihood and Bayesian phylogenetic placement of sequences onto a fixed reference tree. *BMC Bioinformatics*, 11, p.538.
- McGee, K.M. et al., 2019. Determinants of soil bacterial and fungal community composition toward carbon-use efficiency across primary and secondary forests in a Costa Rican conservation area. *Microbial Ecology*, 77(1), pp.148–167.
- 1345 Mendes, L.W. et al., 2015. Land-use system shapes soil bacterial communities in Southeastern Amazon region. *Applied Soil Ecology*, 95, pp.151–160.
- Missa, O. et al., 2009. Monitoring arthropods in a tropical landscape: relative effects of sampling methods and habitat types on trap catches. *Journal of insect conservation*, 13(1), p.103.
- 1350 Morris, A.L. et al., 2016. Deforestation-driven food-web collapse linked to emerging tropical infectious disease, *Mycobacterium ulcerans*. *Science advances*, 2(12), p.e1600387.
- Morrone, J.J., 2014. Biogeographical regionalisation of the Neotropical region. *Zootaxa*, 3782, pp.1–110.
- 1355 Mulatu, K.A. et al., 2017. Biodiversity monitoring in changing tropical forests: a review of approaches and new opportunities. *Remote Sensing*, 9(10), p.1059.
- Mumby, P.J. et al., 2004. Mangroves enhance the biomass of coral reef fish communities in the Caribbean. *Nature*, 427(6974), pp.533–536.
- 1360 Murienne, J. et al., 2019. Aquatic eDNA for monitoring French Guiana biodiversity. *Biodiversity data journal*, 7, p.e37518.
- Nagler, M. et al., 2018. Extracellular DNA in natural environments: features, relevance and applications. *Applied Microbiology and Biotechnology*, 102(15), pp.6343–6356.
- 1365 Navarrete, A.A. et al., 2010. A molecular survey of the diversity of microbial communities in different Amazonian agricultural model systems. *Diversity*, 2(5), pp.787–809.
- 1370 Ndaw, S.M. et al., 2009. Relationships between bacterial diversity, microbial biomass, and litter quality in soils under different plant covers in northern Rio de Janeiro State, Brazil. *Canadian Journal of Microbiology*, 55(9), pp.1089–1095.

- Nepstad, D.C. et al., 2008. Interactions among Amazon land use, forests and climate: prospects for a near-term forest tipping point. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363(1498), pp.1737–1746.
- 1375 Nguyen, N.H. et al., 2016/4. FUNGuild: An open annotation tool for parsing fungal community datasets by ecological guild. *Fungal Ecology*, 20, pp.241–248.
- Nobile, A.B. et al., 2019. DNA metabarcoding of Neotropical ichthyoplankton: Enabling high accuracy with lower cost. *Metabarcoding and Metagenomics*, 3, p.e35060.
- 1380 Nottingham, A.T. et al., 2018. Microbes follow Humboldt: temperature drives plant and soil microbial diversity patterns from the Amazon to the Andes. *Ecology*, 99(11), pp.2455–2466.
- O'Dwyer, J.P. et al., 2009. An integrative framework for stochastic, size-structured community assembly. *Proceedings of the National Academy of Sciences of the United States of America*, 106(15), pp.6170–6175.
- 1385
- Olson, D.M. et al., 2001. Terrestrial Ecoregions of the World: A New Map of Life on Earth. *Bioscience*, 51(11), p.933.
- Pajares, S. et al., 2018. Environmental controls on soil microbial communities in a seasonally dry tropical forest. *Applied and Environmental Microbiology*, 84(17), pp.e00342–18.
- 1390
- Paknia, O., Rajaei Sh., H. & Koch, A., 2015. Lack of well-maintained natural history collections and taxonomists in megadiverse developing countries hampers global biodiversity exploration. *Organisms, Diversity & Evolution*, 15(3), pp.619–629.
- 1395 Pansu, J. et al., 2015. Landscape-scale distribution patterns of earthworms inferred from soil DNA. *Soil Biology & Biochemistry*, 83(0), pp.100–105.
- Papadopoulou, A., Taberlet, P. & Zinger, L., 2015. Metagenome skimming for phylogenetic community ecology: a new era in biodiversity research. *Molecular Ecology*, 24(14), pp.3515–3517.
- 1400 Paula, D.P. et al., 2016. Uncovering trophic interactions in arthropod predators through DNA shotgun-sequencing of gut contents. *PloS One*, 11(9), p.e0161841.
- Paula, F.S. et al., 2014. Land use change alters functional gene diversity, composition and abundance in Amazon forest soil microbial communities. *Molecular Ecology*, 23(12), pp.2988–2999.
- 1405
- Pawlowski, J. et al., 2014. Environmental monitoring through protist next-generation sequencing metabarcoding: assessing the impact of fish farming on benthic foraminifera communities. *Molecular Ecology Resources*, 14, pp.1129–1140.
- Pereira, H.M. et al., 2013. Ecology. Essential biodiversity variables. *Science*,

- 1410 339(6117), pp.277–278.
- Perring, M.P. et al., 2015. Advances in restoration ecology: rising to the challenges of the coming decades. *Ecosphere*, 6(8), pp.1–25.
- Pilliod, D.S. et al., 2013. Estimating occupancy and abundance of stream amphibians using environmental DNA from filtered water samples. *Canadian Journal of Fisheries and Aquatic Sciences*, 70, pp.1123–1130.
- 1415
- Pinheiro, H.T. et al., 2019. Will DNA barcoding meet taxonomic needs? *Science*, 365(6456), pp.873–874.
- Pompanon, F. et al., 2012. Who is eating what: diet assessment using next generation sequencing. *Molecular Ecology*, 21, pp.1931–1950.
- 1420 Porazinska, D.L. et al., 2012. Nematode spatial and ecological patterns from tropical and temperate rainforests. *PloS One*, 7(9), p.e44641.
- Pornon, A. et al., 2016. Using metabarcoding to reveal and quantify plant-pollinator interactions. *Scientific Reports*, 6, p.27282.
- Pringle, E.G. & Moreau, C.S., 2017. Community analysis of microbial sharing and specialization in a Costa Rican ant–plant–hemipteran symbiosis. *Proceedings of the Royal Society B: Biological Sciences*, 284(1850), p.20162770.
- 1425
- Quast, C. et al., 2013. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Research*, 41(Database issue), pp.D590–6.
- 1430 Ramirez, K.S. et al., 2018. Detecting macroecological patterns in bacterial communities across independent studies of global soils. *Nature Microbiology*, 3(2), pp.189–196.
- Ratnasingham, S. & Hebert, P.D.N., 2007. bold: The Barcode of Life Data System (<http://www.barcodinglife.org>). *Molecular Ecology Notes*, 7(3), pp.355–364.
- 1435 Rawlence, N.J. et al., 2014. Using palaeoenvironmental DNA to reconstruct past environments: progress and prospects. *Journal of Quaternary Science*, 29, pp.610–626.
- Rees, H.C. et al., 2014. The detection of aquatic animal species using environmental DNA - a review of eDNA as a survey tool in ecology. *Journal of Applied Ecology*, 51(5), pp.1450–1459.
- 1440
- Rice, J. et al., 2018. Summary for policymakers of the regional assessment report on biodiversity and ecosystem services for the Americas of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. *IPBES book*. Available at: <https://www.fs.usda.gov/treesearch/pubs/56826>.
- 1445 Ritter, C.D. et al., 2019. Locality or habitat? Exploring predictors of biodiversity in Amazonia. *Ecography*, 42(2), pp.321–333.

- Rocha, R.G. et al., 2015. DNA from owl pellet bones uncovers hidden biodiversity. *Systematics and Biodiversity*, 13(4), pp.403–412.
- 1450 Rodgers, T.W., Xu, C.C.Y. & Giacalone, J., 2017. Carrion fly-derived DNA metabarcoding is an effective tool for mammal surveys: Evidence from a known tropical mammal community. *Molecular Ecology*, 17(6), pp.e133–e145.
- 1455 Rodrigues, J.L.M. et al., 2013. Conversion of the Amazon rainforest to agriculture results in biotic homogenization of soil bacterial communities. *Proceedings of the National Academy of Sciences of the United States of America*, 110(3), pp.988–993.
- Rodrigues, R.C. et al., 2019. ATLANTIC BIRD TRAITS: a data set of bird morphological traits from the Atlantic forests of South America. *Ecology*, 100(6), p.e02647.
- 1460 Rodriguez-Nuñez, K.M., Rullan-Cardec, J.M. & Rios-Velazquez, C., 2018. The metagenome of bromeliads phytotelma in Puerto Rico. *Data in Brief*, 16, pp.19–22.
- Rousseau, L. et al., 2013. Soil macrofauna as indicators of soil quality and land use impacts in smallholder agroecosystems of western Nicaragua. *Ecological Indicators*, 27, pp.71–82.
- 1465 Sales, N.G. et al., 2019. Assessing the potential of environmental DNA metabarcoding for monitoring Neotropical mammals: a case study in the Amazon and Atlantic Forest, Brazil. *bioRxiv*, p.750414. Available at: <https://www.biorxiv.org/content/biorxiv/early/2019/08/31/750414.full.pdf>.
- 1470 Sales, N.G. et al., 2019. Influence of preservation methods, sample medium and sampling time on eDNA recovery in a neotropical river. *Environmental DNA*, 1(2), pp.119–130.
- Salter, S.J. et al., 2014. Reagent and laboratory contamination can critically impact sequence-based microbiome analyses. *BMC Biology*, 12, p.87.
- 1475 Sapountzis, P. et al., 2015. Acromyrmex leaf-cutting ants have simple gut microbiota with nitrogen-fixing potential. *Applied and Environmental Microbiology*, 81(16), pp.5527–5537.
- 1480 Sasso, T. et al., 2017. Environmental DNA characterization of amphibian communities in the Brazilian Atlantic forest: Potential application for conservation of a rich and threatened fauna. *Biological Conservation*, 215, pp.225–232.
- Schnell, I.B. et al., 2018. Debugging diversity - a pan-continental exploration of the potential of terrestrial blood-feeding leeches as a vertebrate monitoring tool. *Molecular Ecology Resources*, 18(6), pp.1282–1298.
- 1485 Schnell, I.B., Bohmann, K. & Gilbert, M.T.P., 2015. Tag jumps illuminated - reducing sequence-to-sample misidentifications in metabarcoding studies. *Molecular Ecology Resources*, 15(6), pp.1289–1303.

- Schoch, C.L. et al., 2012. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proceedings of the National Academy of Sciences of the United States of America*, 109(16), pp.6241–6246.
- 1490 Schroeder, J.W. et al., 2019. Host plant phylogeny and abundance predict root-associated fungal community composition and diversity of mutualists and pathogens R. Shefferson, ed. *Journal of Ecology*, 107(4), pp.1557–1566.
- Scibetta, S. et al., 2012. A molecular method to assess *Phytophthora* diversity in environmental samples. *Journal of Microbiological Methods*, 88(3), pp.356–368.
- 1495 Sengupta, M.E. et al., 2019. Environmental DNA for improved detection and environmental surveillance of schistosomiasis. *Proceedings of the National Academy of Sciences of the United States of America*, 116(18), pp.8931–8940.
- Sheth, B.P. & Thaker, V.S., 2017. DNA barcoding and traditional taxonomy: an integrated approach for biodiversity conservation. *Genome*, 60(7), pp.618–628.
- 1500 Shokralla, S. et al., 2016. Environmental DNA Barcode Sequence Capture: Targeted, PCR-free Sequence Capture for Biodiversity Analysis from Bulk Environmental Samples. *bioRxiv*. Available at: <http://dx.doi.org/10.1101/087437>.
- Sigsgaard, E.E. et al., 2016. Population characteristics of a large whale shark aggregation inferred from seawater environmental DNA. *Nature Ecology & Evolution*, 1, p.0004.
- 1505 Smets, W. et al., 2015. A method for simultaneous measurement of soil bacterial abundances and community composition via 16S rRNA gene sequencing. *PeerJ Preprints*, 3(e1622), p.e1622.
- Soares-Filho, B. et al., 2014. Land use. Cracking Brazil's Forest Code. *Science*, 344(6182), pp.363–364.
- 1510 Sommeria-Klein, G. et al., 2016. Inferring neutral biodiversity parameters using environmental DNA data sets. *Scientific Reports*, 6, p.35644.
- Stackebrandt, E. & Goebel, B.M., 1994. Taxonomic note: a place for DNA-DNA reassociation and 16S rRNA sequence analysis in the present species definition in bacteriology. *International Journal of Systematic and Evolutionary Microbiology*, 44(4), pp.846–849.
- 1515 ter Steege, H. et al., 2013. Hyperdominance in the Amazonian tree flora. *Science*, 342(6156), p.1243092.
- Strassburg, B.B.N. et al., 2017. Moment of truth for the Cerrado hotspot. *Nature ecology & evolution*, 1(4), p.99.
- 1520 Strickler, K.M., Fremier, A.K. & Goldberg, C.S., 2015. Quantifying effects of UV-B, temperature, and pH on eDNA degradation in aquatic microcosms. *Biological Conservation*, 183, pp.85–92.
- Taberlet, P., Coissac, E., Hajibabaei, M., et al., 2012. Environmental DNA. *Molecular*

- 1525 *Ecology*, 21(8), pp.1789–1793.
- Taberlet, P. et al., 2018. *Environmental DNA: For Biodiversity Research and Monitoring*, Oxford University Press.
- 1530 Taberlet, P., Coissac, E., Pompanon, F., et al., 2012. Towards next-generation biodiversity assessment using DNA metabarcoding. *Molecular Ecology*, 21(8), pp.2045–2050.
- Takahara, T., Minamoto, T. & Doi, H., 2013. Using environmental DNA to estimate the distribution of an invasive fish species in ponds. *PLoS One*, 8, p.e56584.
- Talaga, S. et al., 2017. DNA reference libraries of French Guianese mosquitoes for barcoding and metabarcoding. *PLoS one*, 12(6), p.e0176993.
- 1535 Tessler, M. et al., 2017. A Global eDNA Comparison of Freshwater Bacterioplankton Assemblages Focusing on Large-River Floodplain Lakes of Brazil. *Microbial Ecology*, 73(1), pp.61–74.
- Tessler, M. et al., 2018. Bloodlines: mammals, leeches, and conservation in southern Asia. *Systematics and Biodiversity*, 16(5), pp.488–496.
- 1540 Thomas, A.C. et al., 2016. Quantitative DNA metabarcoding: improved estimates of species proportional biomass using correction factors derived from control material. *Molecular Ecology Resources*, 16(3), pp.714–726.
- Thompson, L.R. et al., 2017. A communal catalogue reveals Earth’s multiscale microbial diversity. *Nature*, 551(7681), pp.457–463.
- 1545 Thomsen, P.F. & Sigsgaard, E.E., 2019. Environmental DNA metabarcoding of wild flowers reveals diverse communities of terrestrial arthropods. *Ecology and Evolution*, 9(4), pp.1665–1679.
- 1550 Thomsen, P.F. & Willerslev, E., 2015. Environmental DNA – An emerging tool in conservation for monitoring past and present biodiversity. *Biological Conservation*, 183, pp.4–18.
- Tiedje, J.M. et al., 1999. Opening the black box of soil microbial diversity. *Applied Soil Ecology*, 13(2), pp.109–122.
- 1555 Tritsch, I. & Le Tourneau, F.-M., 2016. Population densities and deforestation in the Brazilian Amazon: New insights on the current human settlement patterns. *Applied Geography*, 76, pp.163–172.
- Turner, C.R., Uy, K.L. & Everhart, R.C., 2015. Fish environmental DNA is more concentrated in aquatic sediments than surface water. *Biological Conservation*, 183, pp.93–102.
- 1560 Vacher, C. et al., 2016. Learning ecological networks from next-generation sequencing data. *Advances in Ecological Research*, 54, pp.1–39.
- Valdez-Moreno, M. et al., 2019. Using eDNA to biomonitor the fish community in a

- tropical oligotrophic lake. *PloS One*, 14(4), p.e0215505.
- 1565 Valentin, R.E. et al., 2018. Early detection of invasive exotic insect infestations using eDNA from crop surfaces. *Frontiers in Ecology and the Environment*, 16(5), pp.265–270.
- Vasco-Palacios, A.M. et al., 2019. Carbon content and pH as important drivers of fungal community structure in three Amazon forests. *Plant and soil*. Available at: <https://doi.org/10.1007/s11104-019-04218-3>.
- 1570 Vittor, A.Y. et al., 2009. Linking deforestation to malaria in the Amazon: characterization of the breeding habitat of the principal malaria vector, *Anopheles darlingi*. *The American journal of tropical medicine and hygiene*, 81(1), pp.5–12.
- 1575 Waring, B.G. & Hawkes, C.V., 2015. Short-term precipitation exclusion alters microbial responses to soil moisture in a wet tropical forest. *Microbial Ecology*, 69(4), pp.843–854.
- West, A.G. et al., 2019. The microbiome in threatened species conservation. *Biological Conservation*, 229, pp.85–98.
- Wilcox, T.M. et al., 2018. Capture enrichment of aquatic environmental DNA: A first proof of concept. *Molecular Ecology Resources*, 18(6), pp.1392–1401.
- 1580 Wu, T.H. et al., 2011. Molecular study of worldwide distribution and diversity of soil animals. *Proceedings of the National Academy of Sciences of the United States of America*, 108, pp.17720–17725.
- Yoccoz, N.G. et al., 2012. DNA from soil mirrors plant functional and structural diversity. *Molecular Ecology*, 21.
- 1585 Zinger, L., Taberlet, P., et al., 2019. Body size determines soil community assembly in a tropical forest. *Molecular Ecology*, 28(3), pp.528–543.
- Zinger, L., Bonin, A., et al., 2019. DNA metabarcoding—Need for robust experimental designs to draw sound ecological conclusions. *Molecular Ecology*, 28(8), pp.1857–1862.
- 1590 Zinger, L. et al., 2016. Extracellular DNA extraction is a fast, cheap and reliable alternative for multi-taxa surveys based on soil DNA. *Soil Biology & Biochemistry*, 96, pp.16–19.
- Zinger, L. et al., 2011. Global patterns of bacterial beta-diversity in seafloor and seawater ecosystems. *PLoS One*, 6, p.e24570.
- 1595 Zinger, L. & Philippe, H., 2016. Coalescing molecular evolution and DNA barcoding. *Molecular Ecology*, 25(9), pp.1908–1910.
- Zizka, A. et al., 2018. Finding needles in the haystack: where to look for rare species in the American tropics. *Ecography*, 41(2), pp.321–330.

1600 **Tables**

1605 **Table 1: Essential Biodiversity Variables (EBVs) and potential utility of eDNA-based methods to measure them in neotropical rainforests.** EBVs are as defined by Pereira et al. (2013). Sections of this review or reference paper discussing such applications, or associated limitations are also indicated. NA: no documentation available yet. Usefulness levels are attributed depending on the biases of eDNA for each EBV candidate, the potential costs, as well as the extent to which eDNA information has to be complemented by other sources (e.g. species functional traits).

| EBV Class | EBV Candidate | Utility of eDNA | Sections or references |
|-----------------------|--|------------------------|-------------------------------|
| Genetic composition | Co-ancestry | Fairly useful | 2, 3.4 |
| | Allelic diversity and population genetic differentiation | Fairly useful | (Sigsgaard et al. 2016) |
| | Breed and variety diversity | Unknown | NA |
| Species populations | Species distribution | Very useful | 1, 2, 3.1, 3.2, 4.3, 4.4 |
| | Population abundance | Poorly useful | 2, 3.3., 3.4 |
| | Population structure by age/size class | Useless | NA |
| Species traits | Phenology | Fairly useful | 3.1, 4.1 |
| | Morphology and Reproduction | Useless | NA |
| | Physiology and movement | Fairly useful | 4.1, 4.2 |
| Community composition | Taxonomic diversity | Very useful | 1, 2, 3 |
| | Species interactions | Very useful | 1, 4.3 |
| Ecosystem function | Net lary or llary productivity | Poorly useful | NA |
| | Nutrient retention | Useless | NA |
| | Disturbance regime | Fairly useful | 3.1, 3.2, 4.5 |
| Ecosystem structure | Habitat structure | Fairly useful | 3.1, 3.2, 4.5 |
| | Ecosystem extent and fragmentation | Fairly useful | 3.1, 3.2, 4.5 |
| | Ecosystem composition by functional type | Useful | 4.1, 4.3 |

Figures caption

1610 **Figure 1: Overview of the main eDNA-based methods with a focus on DNA**
metabarcoding applied to fish diversity assessment. The broad information that
can be retrieved through each of these methods is depicted in white boxes. Step 1
corresponds to DNA sampling and extraction, which is common to all eDNA-based
methods (black boxes). Each step of DNA metabarcoding is then described: Step 2
1615 depicts the DNA amplification step and which DNA regions are generally used. It
also shows how multiple samples can be sequenced in parallel: by adding a small
sample-specific nucleotidic label in the 5' region of each primer (here corresponding
to sample A) prior to or after DNA amplification. Step 3 illustrates a multiplex of
samples that has been sequenced in a single sequencing run. Between ca. 500-
1620 1000 samples can be multiplexed on Illumina sequencers depending on the sample
diversity. The sequencing step can be seen as a sampling process; the more diverse
the pool of amplicons (i.e. containing different barcodes), the more sequencing reads
are required to appropriately describe the sample diversity and composition. The
dashed sequence in sample B illustrates a tag-jump event. Step 4 broadly
1625 summarizes the bioinformatic procedures used to curate/annotate the sequencing
data and ultimately retrieve a site by OTU/species table.

Figure 2: Examples of soil/litter eDNA signals in 1 ha forest plots of French
Guiana. A) Comparison of the eDNA imprints of different tree species in soil
samples collected every 5 m across a 1 ha plot in the Nourague Reserve, and in the
1630 top 10 cm of the soil layer. The colour gradient represents the log₁₀ relative
abundance of sequencing reads from each species. Black stars correspond to the
location of tree stems with diameter at breast height ≥10 cm. The two left panels
show signatures that are consistent with the locations of conspecific stems. The two
right panels show inconsistent trends, where “false absences” (i.e. absence of DNA
1635 when a stem is present) is likely due to deep rooting systems and “false presences”
to roots of small trees not included in the botanical inventory. It is unlikely that they
correspond to pollen, seeds or litter, because such material should be present
around the other conspecific stems. **B-C)** DNA metabarcoding based analyses of
bacterial and eukaryotic communities from soils and litter samples (ca. 10 g and 0.5
1640 m³ each respectively) collected in 1 ha plots of a *terra firme* forest (Nouragues
Reserve) and white-sand forest (Mana). The plots show differences **B)** in community
composition as measured with the Bray-Curtis index on hellinger-transformed data,
summarized with a principal coordinate analysis and **C)** in plot-scale diversity and
spatial heterogeneity, as depicted with species accumulation curves. **D)** Seasonal
1645 variations in bacterial, eukaryotic and plant community composition in the same plot
as in A) and retrieved with soil eDNA. The figure has been produced using the same
indices and techniques as in B).

1650

Species detection

quantification of a specific marker with qPCR or ddPCR


Species presence/
biomass

DNA metabarcoding

2- Amplification & labelling of a standard DNA region (i.e. barcode *lato sensu*)


conserved sites across taxa

variable region




3 - Sample multiplexing and High-throughput Sequencing

Sequencing reads




1 - Sampling and DNA extraction



4 - DNA sequence curation & classification


Unsupervised classification: pairwise similarities between sequences

Supervised classification: similarities against reference sequences



Taxonomic diversity

% reads



Capture/enrichment

enrichment of the DNA extract with DNA barcodes & HTS

Taxonomic diversity

Metagenome skimming

DNA shearing and shallow shotgun sequencing with HTS


Taxonomic and phylogenetic diversity

Metagenomics


DNA shearing and deep shotgun sequencing with HTS

Taxonomic, phylogenetic, and functional diversity


A



B




C



Sample type ● T-F I litter ● T-F I soil ● W-S I litter ● W-S I soil

D



Dear Alex,

Thank you very much to you Alex for these recommendations as well as to the referees for their comments. We have followed most of these suggestions, and explain below why when we did not. Overall, we took care to reduce the length and restructured the whole manuscript to make it more fluid and accessible. We also provide a roadmap paragraph at the end of the introduction to facilitate the reading. We believe that all these comments greatly helped us to improve the manuscript, and we hope that this new version will be satisfying and of help for the readers.

Best,

Lucie, on behalf of co-authors.

#####

REFEREE 1:

R1.1: Authors have reviewed many aspects of the use of eDNA in neotropical rainforest. Based on the structure of the review, reader can easily get lost in all the sections/text and lose track of the reason of why is currently reading this review. Lots of information are repetitions of other reviews that could be removed from the main text and included as table/box or figure, authors even suggest other reviews for specific topics (mainly introduction and current applications). When the reader finally get through those sections, challenges and perspectives are actually providing new information with very good point for better sampling design and how eDNA could be applied in the region.

>>> We understand the referee's point and strongly reduced the length of the manuscript accordingly.

R1.2: In general, the manuscript is rarely interrupt with a figure or table (not present at all) and when present figures do not add additional information than the text (see specific comment on figure 3).

>>> We have included an additional table that we refer to at several instances in the MS. We also removed the former figure 3 and modified figures 1 and 2.

R1.3: The other problem of this review is that not include any information on other techniques that are not sequencing even if one of their main point is that eDNA is not yet a quantitative measure. At the moment all most all the studies that have tried to link eDNA to biomass are based on qPCR results. As the estimation of biomass from eDNA is big problem and authors make it clear in their review it also important adding information on studies that used qPCR in eDNA. eDNA used in combination of qPCR for detecting threatened species or invasive is one of the most advance area of research especially for the standardisation of the results across different species/labs (see for example "Reporting the limits of detection and quantification for environmental DNA assays" or "Combining ddPCR and environmental DNA to improve detection capabilities of a critically endangered freshwater invertebrate") and have practical application in conservation efforts.

>>> We added such considerations briefly, but did not extend the text on this aspect due to the manuscript length limit and the very limited number of such studies in neotropical ecosystems. In addition, we respectfully disagree that this approach is the most advanced area of research on eDNA, as, to our knowledge, the amount of proof of concept papers equals, or is even lower than for e.g. DNA metabarcoding applications. Therefore, we prefer to refer only to dedicated reviews here.

R1.4: Another point that should be revised is how much importance should be given to

microbial ecology studies. Microbial studies are really interesting and are considered the base on which eDNA is developing as the former has a longer tradition as research field. Based on existing literature, studies and reviews have made a clear separation between the microbial ecology studies and eDNA studies, with some overlapping possible within the microeukaryotes communities. In this review, a big part of text is devoted to pure microbial studies (e.g. terrestrial ecosystems and aquatic ecosystems sections) and should be reduced.

>>> We understand the referee's point and took care to strongly reduce the text dealing with microbial ecology. However, we have still let some references to it given the increasing overlap between classical and microbial ecology that are now possible such as with eDNA techniques, as well as with the increasing interest of classical ecologists to host-associated microbes.

R1.5: Keywords: should include one or two words relative to the rainforest or geographical area used for the review

>>> Done

R1.6: 1.2 Environmental DNA definitions: Line 124 Authors use the word "metagenome" to indicate all the environmental DNA that is present in the sample. As the target reader would include non-molecular expert the word "metagenome" can create confusion and been associated with shotgun sequencing. See for example line 260 and following, "Metagenomics is the most direct ...". I would suggest using a different word to avoid any confusion.

>>> We disagree with the referee on this point, because metagenome is the exact term that one should use in this case. However, to avoid any confusion, we provide a clear definition of this word l.110ff.

R1.7: Line 130 Authors state that they are going to use the eDNA in the broadest sense including also faeces or gut content in the eDNA starting material. Authors should add a phrase to explain what is included in eDNA in the narrowest term, again as many readers are not familiar with terminology.

>>> We now provide an explicit definition of eDNA in its narrowest and broadest sense l.104ff and 114ff

R1.8: Lines 153-155 and 158-160: Authors suggest that the amount of eDNA is roughly positive correlated with biomass, this is still a matter of debate and an active research area. Authors should point out this uncertainty, the reader can get the wrong notion that biomass can be estimated based on eDNA. The same comment is for the following phrases about spatio-temporal information. The proximity of the source of eDNA is clearly a matter of debate especially in water system in which eDNA can travel up to 200 km from the source.

>>> We have removed this consideration and now discuss more about this particular aspects throughout section 3, in particular section 3.3. and 3.4.

R1.9: Line 161 "*Extracellular DNA can also persist in the environment from several hours...*" *this is also true for intracellular DNA and therefore I suggest changing extracellular DNA to eDNA.*

>>> We removed this part of the introduction to comply with the referee's comment R1.1

R1.10: 1.3 Methods for processing eDNA: As said in general comment, this section can easily cut it down or convert in tables/figure. Authors correctly pointed out that several reviews are already present (line 188-189) to cover most aspects of this section and repetition should be avoided.

>>> We now present this in a dedicated section (section 2) and strongly reduced the text to better rely on Figure 1. We still believe that minimum information has to be provided for readers that are not familiar with these technologies (and often use incorrect terminology when talking about eDNA-based methods).

R1.11: Figure 1 In general it is a very clear figure with detailed information for all the aspect that's should cover. Sample B is not too clear, from what I understood is a sample in which an error has occurred, and the wrong specie has been assigned to the wrong fish, is it correct? Authors should explain in the description.

>>> We thank the referee and have now better explained what is new or artifactual in sample B. We also modified the figure to make it more informative for the whole section 2.

R1.12: 2.1 Terrestrial ecosystems: Lines 399-431 Authors dedicate a large section of this section to microbial ecology, which is really interesting, but it is off topic in this context. Authors should consider removing it or reduce it drastically. The following part even if still include microbiology linked with eukaryotes and provide an excellent review on how to sample soil core and what has been done in neotropics bioregion.

>>> We thank the referee for this advice and have amended this section (now section 3.1) accordingly. As mentioned above, we still kept some references on microbial communities, which could be of interest for some of the potential readers of this review.

R1.13: Lines 433-435 Bulk samples and soil eDNA studies are compared but authors have stated in line 130 that they will use eDNA in the broadest sense which include bulk samples therefore they should specify better what they mean with soil eDNA. The use of eDNA in its broadest and narrowest sense through the manuscript is a weak point as this led to inevitable confusion in non-expert reader.

>>> In the revised manuscript, we made sure to clearly specify whether the examples/limitations we discuss apply to soil/sediment/water eDNA, or to eDNA obtained from bulk samples throughout the section 3 when this information is important to mention.

R1.14: Figure 3. Figure is not adding/explaining any particular information that cannot be retrieved by the text. Figure should be removed or changed to add meaningful information compared to the text

>>> We have removed this figure.

R1.15: 2.2 Aquatic ecosystems: Line 636-652 As stated in general comments a good section of this paragraph is about more microbial ecology than eDNA. All microbial ecology part should be shortened.

>>> Done

R1.16: 2.5 Host associated microbiota: This paragraph mostly belongs to microbiology research field. Endophyte communities, root associated microbiota, microbial symbiosis with eukaryotes are all classic area of microbiology research. The gut microbiota could be incorporated with the diet analysis in one section.

>>> We have considerably revised this section and that focusing on diet, which we moved to the perspectives section, as this is probably the application of eDNA that has least been covered in neotropical rainforests. In this new structure, we also considerably reduced the length of the discussion on host-microbiota. l.865ff

R1.17: 3.1 Field and, wet and dry lab biases: In general, the paragraph is well written with several example specific based on neotropical area (lines 831-837) while other parts are mainly repetition present in several reviews that authors cite and invite the readers for

further reading (lines 838-859). The second part should be cut or presented in a different format and only the relevant point should be present in the text.

>>> We agree and have shortened the text and make better use of Figure 1 here.

R1.18: Lines 874-875 Authors should include a mention to occupancy model that would likely become routinely used to detect false positive/negative.

>>> We now present occupancy models as a promising way of improving eDNA data l.601

R1.19: 3.2 Biological interpretations of eDNA: Line 971-973 eDNA molecules are able to persist in the environment for long period if trapped in soil or sediment, but in water the persistence is reduced, and this should be explicitly said otherwise the reader can get a wrong notion.

>>> We now state this explicitly l.655ff

R1.20: Authors make a clear point in this section on how at the moment eDNA is not quantitative measure for many reasons however in all the manuscript they never mention qPCR techniques even if they include citation as Maruyama et al 2014 (line 991) to explain why eDNA is still a not quantitative measure. Most of the research to make eDNA a quantitative measure is based on qPCR and information on this topic have to be included in the manuscript.

>>> We did not extend the MS on this particular aspect due to length constraints, but now mention this point more explicitly l.649ff.

R1.21: 4.1 Increasing the breath of reference database: Authors are pointing out that at the moment there is a lack of consensus on which marker gene should be considered as gold standard in eDNA studies, I will suggest adding which one they would suggest using to someone new in the field. For example, if one of the readers would like to start an eDNA project in neotropical rainforest authors would suggest using COI or other markers?

>>> While we agree that such discussion is extremely useful for end-users, we do not believe that our review should focus on this aspect, for which guidelines depend on the taxon and ecosystem studied, as well as on the initial question addressed. Providing meaningful advice on this matter would actually require a whole separate review, so we prefer to refer the readers to more specific literature l.746ff

R1.22: I do not completely agree with the sentences in lines 1085-1087 as it will be not so easily achieved. Tropical ecosystems harbour huge biodiversity and as the authors said later in their conclusion usually they lack the facilities to carry out molecular work and the transport of specimen outside the countries are not usually an easy path to follow.

>>> While we certainly agree that improving DNA reference databases can represent a non negligible certain burden (but see our discussions l.760ff), we do believe that this effort has to be mentioned and encouraged through collaboration between northern and southern countries. The same concern (on costs) holds true for use of eDNA in general, which is why we emphasize on the need for international collaborations in the introduction of section 4.

R1.23: Line 1145 and following. Several web repositories for eDNA metadata sharing are available

>>> We could not retrieve to what the referee was referring to, as the line number does not match the referee comment. However, we have added information on existing facilities to share eDNA data and metadata l.793.

R1.24: 4.3 Shedding new light on tropical ecological networks: Line 1236-1254 Authors are highlight how eDNA can be used to test several hypotheses as the Janzen-Connell however

then they expand their discussion without any additional information for the reader excluding the final recommendation that should be moved further up in the paragraph.

>>> We totally suppressed this paragraph to shorten the manuscript and in particular the part related to microorganisms, as recommended by both referees.

REFEREE 2:

R2.1: The manuscript “Advances and prospects of environmental DNA in Neotropical rainforests” reviews the existing literature on eDNA use in the neotropics. It introduces eDNA, reviews its current applications in the tropical biomes of the neotropics, discusses challenges of eDNA methods, and looks at future uses of eDNA.

The manuscript presents an extensive review of the literature and could be of relevance to researchers in the neotropics, but has a few serious problems that need to be resolved prior to publication. The main problem is the length of the ms, which is 8,000 words over the suggested word count for this journal. The second major issue is the lack of cohesion throughout the ms, there is no clear focus or goal, and even the exact topic seems to fluctuate between different sections. Lastly, the review would benefit from a more thorough synthesis and interpretation of the literature, rather than mainly providing summaries of previous research.

I recommend major revisions prior to a decision of acceptance. As it stands, the ms is not yet suitable for a high-impact journal such as *Advances in Ecological Research*. Revisions to the length, cohesiveness, and interpretation of the literature would greatly improve the suitability of this ms for publication in *Adv Ecol Res*.

>>> We thank the referee for his/her assessment and suggestions. We overall agree with his/her criticisms and in this revision, we took special care to reduce the length of the MS, homogenize the different parts, improve cohesiveness, and clarify the objectives. We hope the revised text will be clearer.

R2.2: General comments: A big issue with the ms is its length. At 22,755 words (16,186 excluding refs), or 77 pages, it is too long (it exceeds the journal’s suggested word limit by more than 8,000 words). The considerable effort that undoubtedly went in this ms is drowned out by unnecessary details and repetitiveness, turning this potentially very useful resource into an unwieldy ms that is unlikely to be used by many people. The word count could easily be cut by at least a third, or even half.

>>> We agree and have considerably reduced the length of the MS thanks to both referees comments.

R2.3: Another issue is the lack of a clear cohesive structure in the bulk of the review. The ms reads as if the authors each wrote their section, but there was limited post-writing effort to streamline the manuscript and make it into a cohesive paper. This unfortunately translates in a bulky ms that is in times hard to follow, with considerable repetition and varying quality of English language.

>>> We did our best to improve the manuscript cohesiveness both conceptually and with the language.

R2.4: Maybe this is because no clear goal of the review? The different chapters vary considerably in their focus (“neotropical rainforests” vs. “tropical biomes of the Neotropics” vs. “Neotropics” vs. “Neotropical ecosystems” vs. “Tropical ecosystems of the Central and South Americas”). Setting up a clear focus and goal of the review in the introduction would help guide readers (and it seems the authors as well).

>>> We agree with the referee and now better make clear that our focus are neotropical rainforests throughout the MS, although we extend the scope of the review in the last section as a perspective (4.5).

R2.5: Sections (particularly Part 2) of the review read as a very traditional literature review (listing summaries of research papers) with limited synthesis of literature. While such reviews can be useful to researchers in a narrow field, it excludes a wider scientific audience and reduces the value for a high impact ecology journal.

>>> We have considerably revised all parts and in particular former sections 2 and 3 (now section 3) in order to better reveal what are the known and unknown for each of these systems from an eDNA perspective.

R2.6: While the conclusion raises very important points, most of the points raised are not mentioned anywhere in the review. This hiatus makes the conclusion seem like an afterthought to rationalise the review, rather than an integral part of the review.

>>> We agree and now consider the problems related to costs, training and Nagoya protocols in the introduction of the perspective section (section 4), as they apply if one is to apply these techniques more routinely in neotropical rainforests and beyond.

R2.7: Throughout the ms there is a tendency for “fuzzy language” to describe quantities (“fairly good”, “some”, “non-negligible”, etc.), which should be avoided in scientific writing. There are some minor language issues (grammar, word order)

>>> In this revised manuscript, we have taken care of avoiding any vague writing and revised the english.

R2.8: I don't really understand the use of all the specific “shortfalls” (Darwinian, Linnean, Wallacean,...) in the text. As they seem to be explained each time, just leaving the explanation would make more sense than including more unnecessary terminology in an already jargon-heavy manuscript (except for referencing Hortal et al. 2015)

>>> We have excluded this jargon in the revised MS.

R2.9: Detailed comments: Title: The title does not seem to reflect the content of the review, which seems to be about more than only rainforests.

>>> We now better make the point on what ecosystem we focus on throughout the MS (neotropical rainforests).

R2.10: Abstract: l28: “wholly dependant” seems like rather strong statement; l29-30: incorrect use of the term “let alone”, consider switching to: “many species remain undiscovered, let alone described”

>>> Corrected.

R2.11: 40-44: It would be useful to explain this in detail in the ms intro

>>> We now provide a roadmap paragraph in the intro l.150ff.

R2.12: 41: “this ecoregion”: which one? Not introduced in abstract

>>> We have now replaced this by the actual system to which we refer to.

R2.13: Part 1: Intro: Intro does not clearly lay out what the aim of the review is, which makes for a confusing read for the rest of the ms: Is the aim to give an update of knowledge on neotropical rainforests? Explain value of eDNA in neotropical forests? Give a list and summary of each eDNA study in this bioregion? Etc..

>>> We now provide a roadmap paragraph in the intro I.150ff.

R2.14: First two paragraphs in intro are very long, suggest splitting for increased readability (potentially lines 61 and 93)

>>> Thanks for the suggestion. We reduced the whole introduction.

R2.14: I.53: Really? I did not know this, larger than the central Africa, Southeast Asia, etc. combined? Are there references available for this statement?

>>> Corrected.

R2.15:

- I.83: remove “now”
- I.84: not clear what the meaning is of “in general”
- I.98: consider replacing “for” by “even”
- I.109: “reduced” or “limited”?
- I.111: not clear of solving the aforementioned limitations will mitigate global change
- I.116-118: reference would be useful
- I.132-133: not relevant

>>> Done/Clarified/Modified

R2.16: 158-160: I do not agree with this statement, an extremely rare species depositing faeces where a sample is collected would (incorrectly) suggest a very high biomass. Suggest altering statement or providing clear references

>>> We agree and, as indicated for R1.8, the whole paragraph has been revised. These considerations are now only mentioned in section 3.

R2.17: 178: would be useful to refer sooner to figure 1 in this paragraph

>>> Done.

R2.18:

- 321: “extremely” is an exaggeration, consider removing
- 328-332: Repetitive section

>>> We modified the paragraph to avoid repetition.

R2.19: 332-334: The references used here are not risk assessments, but impact assessments. The two are entirely different management tools and cannot be interchanged

>>> We corrected the text accordingly.

R2.20: 340: “bioindication” is jargon and should be defined

>>> We now provide its definition.

R2.21: Part 2: Current applications; The different sections of part 2 are all differently structured. I would expect at least some parallels when comparing how eDNA has been used in different ecosystems / substrates

>>> We thoroughly revised the text in this part to make it more homogeneous and structured.

R2.22: Terrestrial ecosystems: No clear structure + too much detail in describing results of other studies, a review should synthesize more, rather than merely summarise.

Additionally, the paragraphs on different biota in this section seem to have different goals, which is quite confusing

- Section (407-440): Microbial DNA: very detailed description of results older studies
- Section (441-479): Invertebrates: no detailed info, but synthesis and suggestions for uses
- Section (480-502): Larger animals/plants: most detailed info on methods, high level detail on results, interpretation of those results (not done in any other sections)
- Section (503-527): what is the topic? Vertical heterogeneity AND microbial and plant seasonal variation AND eDNA persistence in soils?
- Section (528-551): This section goes back to animals, but now on other, better sampling methods than soil?

>>> We modified the whole structure of that paragraph and of that on aquatic ecosystems so that to go beyond a simple listing of existing papers and harmonized the structure of both paragraphs. In particular, we provide paragraph headings to better reflect this new structure, which follows discussion on each taxon and then general concerns/challenges particular to each of these environments.

R2.23: Aquatic ecosystems: some language mistakes (grammar, incorrect word use), could do with language edit. Follows different structure (again) from previous section. This section focuses more on method use, as I expected the entire review to do. Would suggest to try to follow format/order of terrestrial section though (e.g. microbial – invert – large species).

>>> See reply to R2.22.

R2.24:

- Diet analyses: interesting section, just some language errors
- Bulk samples: some minor language mistakes

>>> These paragraphs have been integrated either in terrestrial/aquatic ecosystems, or in perspectives (diet, section 4.3)

R2.25: Host associated microbiota: First sentence states this is done using faeces, tissues, or bulk samples, which places it in the previous two sections of Part 2. It is not clear why this is a separate section, especially since parts of this have already been discussed on 699 – 713. Consider integrating this part in the other sections, or re-writing this (and previous) sections so it's clear why this deserves separate section. Too much detail in describing results of other studies.

>>> This section has been strongly reduced and moved in to section 4.5 which discusses the perspective research on biological interactions.

R2.26:

- 370: “most” is too vague, reword
- 372: “certain” is too vague, reword
- 374: not clear what is meant by “achievements”: studies, results, conclusions?

>>> Corrected

R2.27: 386-392: repetitive already discussed in intro

>>> We modified the introduction and this paragraph in response to another comment we had, so this comment does not apply anymore.

R2.28: 393: “non-negligible” is too vague, reword

>>> Vague wording has been excluded throughout the MS.

R2.29:

- 394-395: consecutive sentences starting with “however”
- 397: “second”: where was “first”?
- 419-420: irrelevant
- 445-447: references needed
- 490-491: change word order: “by directly using soil as”

>>> Corrected/Clarified.

R2.30: 495-498: Commenting on the methodological details of this single study, but not on any other study seems strangely out of place in this section

>>> We removed this part of the text, and only discuss the results briefly in the Figure 2 caption.

R2.31: 506: “horizontal heterogeneity” was not mentioned above, or anywhere else in the ms

>>> We have clarified this point I.323

R2.32: 513-516: not clear how seasonality fits into a paragraph on vertical heterogeneity

>>> We respectfully disagree with the referee. Both deal with variations in abiotic conditions, either on a temporal axis or on a spatial one. Nevertheless, we separated these two ideas in the revised MS to comply with other comments.

R2.33: 513-528: entire paragraph does not seem to fit into this section

>>> We removed this discussion.

R2.33:

- 529: previous paragraph already started with “finally”
- 562: Remove “the”
- 568: Remove “typically”
- 578: Is this sentence about soil or water?
- 579: does “it” refer to water or eDNA?
- 580: what does “particles” refer to? Water, eDNA, other?
- 581: incorrect use of “therefore”
- 607 + 612: suggest using “toxins” or “ichthyocides” instead of “toxicants”
- 691: “some” is too vague, reword
- 692: “their diet”
- 700: “also now” is grammatically incorrect
- 706: change word order to: “by directly targeting”
- 732: “fairly good” is too vague, reword
- 749: remove “indeed”
- 754-755: faeces and bulk tissues already addressed in the previous section
- 792: “exploding” is not suitable scientific language

>>> Thanks, all these points are now corrected/modified in the revised text.

R2.34: Part 3: Challenges; This part has very limited linking back to topic of the review (neotropical rainforests), I could find only three small examples in 10 pages. While it might be more difficult in a technical section, clearly relating back to the topic of the review would greatly increase relevance to readers. Word length should be reduced, removing repetitive

sections and unnecessary fillers could cut word length by almost half.

>>> We have considerably reduced this section accordingly (now section 3.3 and 3.4). However, we feel that it is important to provide a reminder for these limitations. In our experience as readers, referees and editors ourselves, we still read many manuscripts that poorly account for (or even totally omit) these considerations. We hence believe that they will be useful for naive users/readers.

R2.35: 820 – 834 (intro): the same content could be written in a quarter of the word length

>>> See our reply to R2.34.

R2.36:

- 824: remove “indeed”
- 829: change to “prevent” (grammar: it the particularities that prevent, not the eDNA)

>>> Do not apply anymore.

R2.37:

- 836-845: another intro after the intro?
- 846-935: reduce length, too much circular reasoning and unnecessary filler

>>> See our reply to R2.34.

R2.37:

- 862: Change “beyond” to “After”
- 864: “Usually” or “unusually” low abundances?

>>> Corrected

R2.38: 1025: Not clear what reasoning authors argue against: the use of presence/absence metrics? Or the use of abundance metrics? If PA is considered unreliable, what is favoured instead? Abundance of reads does not reliably relate to real abundance, would be useful to explain more clearly what is meant.

>>> The whole paragraph has been modified and clarified.

R2.38: Part 4: Perspectives Too long, often repetitive, and includes irrelevant information

>>> We have also considerably revised and removed the redundancies in this section. We did our best to remove potential irrelevant information, although the referee did not single out any in particular.

R2.39:

- 1054: “has come of age” is a big statement, which may or may not be entirely correct
- 1061-1066: seems like a rather lengthy way to say “reference databases are poorly resolved”

>>> Corrected

R2.40: 1084-1099:

- It would be useful to introduce/explain what this list is meant to achieve, there is no clear lead to the start of it.

- It is not entirely clear to me how the first and second point differ: they both seem to argue for more complete, high quality reference databases

>>> We have considerably shortened this paragraph to remove potential redundancies

R2.41:

- 1088: “non-negligible” is vague, non-scientific wording
- 1084-1086: Both sentences start with “first” and “firstly”, but there is no “secondly” to follow up

>>> Corrected

R2.42: 1131 – 1134: Trait databases are available for some marine taxa (e.g. FishBase), but not necessarily linked to DNA reference databases

>>> We added examples of such databases l.782.

R2.43:

- 1147: different format of the term “terra-firme” to previous use (402: “terra firme”), same issue with “white sand” (1341), “white sands” (402), “white-sand” (1145)
- 1142: remove “nonetheless”

>>> Corrected.

R2.44: 1136-1172: savannas, tropical dry forest, mangroves, deep sea environments, etc are not rainforests, so not clear why they have been included here.

>>> We excluded deep sea environments to remain focused on terrestrial habitats, but still propose an opening to other ecosystems. Even if the review is focused on Neotropical rainforests, we make clear in section 4.5 (and also in the initial version) that going beyond rainforests in the Neotropics is relevant to better understand the fate of rainforests with global change. More generally, we did not find many studies on other ecosystems, which is also one of the reasons that led us to focus the review on rainforests.

R2.45: 1231-1246: While interesting, the threats to rainforest are well known and have been described in the intro, so most of this section could be removed

>>> These considerations are now partly excluded from the intro and the targeted paragraph has been reformulated.

R2.46:

- 1246: “non-negligible” is vague, reword
- 1262: “a number of empirical evidence”: incorrect grammar
- 1287-1291: needs references
- 1301: what are EBVs?
- 1323: “remains”

>>> Corrected

R2.47: 1310 - 1318: I thought the review was about rainforest and not savannas?

>>> See our reply to R2.44.

R2.48: 1321-1336: this paragraph on sanitation/parasites/healthcare does not belong in a section on “conservation biology and ecological risk assessments”

>>> We created a dedicated section to the topic accordingly l.894

R2.49: Conclusion: 1364: Is this review supposed to be about the Neotropics or neotropical rainforests?

>>> See our reply to R2.44.

R2.50: 1365-1366: This review did not compare research eDNA effort in other, less diverse regions. So this statement should either be removed or backed with references.

>>> Corrected

R2.51: 1365-1376: none of this was discussed in the review, so unclear why it is used as a conclusion

>>> We agreed and discuss these aspects much earlier in the manuscript now l.685ff

R2.52: Figures: Figure 1: needs a more detailed legend to make it easy to interpret

>>> Now provided.

R2.53: Figure 3: not possible to distinguish between the different shades of grey, so cannot interpret this figure

>>> Figure 3 has been removed.