



**HAL**  
open science

## Metabolic flux in cellulose batch and cellulose-fed continuous cultures of *Clostridium cellulolyticum* in response to acidic environment

Mickaël Desvaux, Emmanuel Guedon, Henri Petitdemange

► **To cite this version:**

Mickaël Desvaux, Emmanuel Guedon, Henri Petitdemange. Metabolic flux in cellulose batch and cellulose-fed continuous cultures of *Clostridium cellulolyticum* in response to acidic environment. *Microbiology*, 2001, 147 (6), pp.1461-1471. 10.1099/00221287-147-6-1461 . hal-02910800

**HAL Id: hal-02910800**

**<https://hal.inrae.fr/hal-02910800>**

Submitted on 3 Aug 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Metabolic flux in cellulose batch and cellulose-fed continuous cultures of *Clostridium cellulolyticum* in response to acidic environment

Mickaël Desvaux, Emmanuel Guedon and Henri Petitdemange

Author for correspondence: Henri Petitdemange. Tel: +33 3 83 91 20 53. Fax: +33 3 83 91 25 50.  
e-mail: hpetitde@lcb.uhp-nancy.fr

Laboratoire de Biochimie  
des Bactéries Gram +,  
Domaine Scientifique  
Victor Grignard, Université  
Henri Poincaré, Faculté des  
Sciences, BP 239, 54506  
Vandœuvre-lès-Nancy  
Cédex, France

***Clostridium cellulolyticum*, a nonruminant cellulolytic mesophilic bacterium, was grown in batch and continuous cultures on cellulose using a chemically defined medium. In batch culture with unregulated pH, less cellulose degradation and higher accumulation of soluble glucides were obtained compared to a culture with the pH controlled at 7.2. The gain in cellulose degradation achieved with pH control was offset by catabolite production rather than soluble sugar accumulation. The pH-controlled condition improved biomass, ethanol and acetate production, whereas maximum lactate and extracellular pyruvate concentrations were lower than in the non-pH-controlled condition. In a cellulose-fed chemostat at constant dilution rate and pH values ranging from 7.4 to 6.2, maximum cell density was obtained at pH 7.0. Environmental acidification chiefly influenced biomass formation, since at pH 6.4 the dry weight of cells was more than fourfold lower compared to that at pH 7.0, whereas the specific rate of cellulose assimilation decreased only from 11.74 to 10.13 milliequivalents of carbon (g cells)<sup>-1</sup> h<sup>-1</sup>. The molar growth yield and the energetic growth yield did not decline as pH was lowered, and an abrupt transition to washout was observed. Decreasing the pH induced a shift from an acetate-ethanol fermentation to a lactate-ethanol fermentation. The acetate/ethanol ratio decreased as the pH declined, reaching close to 1 at pH 6.4. Whatever the pH conditions, lactate dehydrogenase was always greatly in excess. As pH decreased, both the biosynthesis and the catabolic efficiency of the pyruvate-ferredoxin oxidoreductase declined, as indicated by the ratio of the specific enzyme activity to the specific metabolic rate, which fell from 9.8 to 1.8. Thus a change of only 1 pH unit induced considerable metabolic change and ended by washout at around pH 6.2. *C. cellulolyticum* appeared to be similar to rumen cellulolytic bacteria in its sensitivity to acidic conditions. Apparently, the cellulolytic anaerobes studied thus far do not thrive when the pH drops below 6.0, suggesting that they evolved in environments where acid tolerance was not required for successful competition with other microbes.**

Keywords: cellulolytic bacteria, flux analysis, environmental pH, cellulose degradation, chemostat

## INTRODUCTION

Biological degradation of cellulosic materials is signifi-

**Abbreviations:** AADH, acetaldehyde dehydrogenase; AK, acetate kinase; ADH, alcohol dehydrogenase; ATP-Eff, efficiency of ATP generation; Fd, ferredoxin; G1P, glucose 1-phosphate; G6P, glucose 6-phosphate; LDH, lactate dehydrogenase; meq C, milliequivalent of carbon; PFO, pyruvate-ferredoxin oxidoreductase; PTA, phosphotransacetylase; R, ratio of specific enzyme activity to metabolic flux.

cant in swamps, marshes, sediments, composts and anaerobic waste treatment, and in the intestinal tracts of herbivorous animals and insects (Ljungdahl & Eriksson, 1985). Within the biosphere, cellulolytic clostridia participate significantly in this process (Bayer & Lamed, 1992; Leschine, 1995; Tomme *et al.*, 1995), which is strongly linked to the global carbon cycle (Wolin & Miller, 1987).

High concentrations of fermentation acids and, as a

result, low pH conditions are often found in anaerobic habitats (Ljungdahl & Eriksson, 1985; Goodwin & Zeikus, 1987). Yet due to their particular pattern of intracellular pH regulation (Huang *et al.*, 1985) compared with other low-G + C Gram-positive anaerobes, mainly the so-called lactic acid bacteria, the clostridial-type bacteria are generally considered as restricted to less acidic ecological niches (Russell *et al.*, 1996).

Cellulolytic clostridia digest cellulose through extracellular multienzyme complexes (Béguin & Lemaire, 1996; Bayer *et al.*, 1998). These cellulosomes are found at the surface of the bacteria and allow both cell adhesion to cellulose fibres (Bayer *et al.*, 1996) and very efficient degradative activity against crystalline cellulose due to a high synergism of the different cellulase components (Boisset *et al.*, 1999).

*Clostridium cellulolyticum* is a low-G + C Gram-positive nonruminant cellulolytic mesophilic bacterium belonging to the clostridial group III, and also classified in family 4, genus 2, of a new proposed hierarchical structure for clostridia (Collins *et al.*, 1994). Using cellobiose, a soluble substrate, several advances in understanding of the metabolism of this bacterium have been made, such as (i) a better control of catabolism in a mineral salt-based medium (Payot *et al.*, 1998; Guedon *et al.*, 1999b), (ii) recognition of major differences in regulatory responses in cellobiose-limited and cellobiose-saturated chemostat cultures (Guedon *et al.*, 2000b), and (iii) the importance of glucose 6-phosphate (G6P) and glucose 1-phosphate (G1P) nodes in regulation of metabolic fluxes (Guedon *et al.*, 2000a). Earlier investigations using cellulose have been mainly devoted to cellulolytic performance and bacterial behaviour towards insoluble substrates (Giallo *et al.*, 1985; Gelhaye *et al.*, 1993a, b). A few studies, however, have focused on the metabolism of this bacterium on cellulose: recent investigation of cellulose fermentation performed in batch culture (Desvaux *et al.*, 2000) indicated (i) variation of metabolite yields as a function of initial cellulose concentration, and (ii) an early growth inhibition related to pyruvate overflow as with cellobiose (Guedon *et al.*, 1999a). A study in a cellulose-limited chemostat indicated that bacterial metabolism was not as distorted as with cellobiose and *C. cellulolyticum* appeared well adapted to a cellulolytic lifestyle (Desvaux *et al.*, 2001).

Whereas the effects of acidic conditions on the growth of cellulolytic rumen bacteria have been the subject of considerable research (Russell & Dombrowski, 1980; Kalachniuk *et al.*, 1994; Russell & Wilson, 1996; Russell & Diez-Gonzalez, 1998), little is known of how these conditions affect the metabolism of cellulolytic clostridia (Duong *et al.*, 1983; Mitchell, 1998). The aim of the present work was to investigate the cellulose degradation and metabolic changes of *C. cellulolyticum* on insoluble cellulose caused by environmental pH conditions. Since conditions in natural environments most likely resemble those somewhere between a closed batch culture and an open continuous culture system (Kovárová-Kovar & Egli, 1998), both types of culture

systems were used to study the effects of pH on growth and metabolism of *C. cellulolyticum*.

## METHODS

**Organism and medium.** *Clostridium cellulolyticum* ATCC 35319 was isolated from decayed grass by Petitdemange *et al.* (1984). Germination of stocks of spores and anaerobic cell culture were performed as described by Desvaux *et al.* (2000). The defined medium used in all experiments was a modified CM3 medium as described by Guedon *et al.* (1999a) containing various amounts of cellulose MN301 (Macherey-Nagel) as specified in Results.

**Growth conditions.** *Clostridium cellulolyticum* was grown either in batch or in continuous culture with cellulose as sole carbon and energy source. All experiments were performed in a 1.5 l working volume fermenter (LSL Biolafite) as previously described (Guedon *et al.*, 1999a, b); the temperature was maintained at 34 °C and the pH was controlled by automatic addition of 3 M NaOH or 1 M HCl as specified in Results. The inoculum was 10% (v/v) from an exponentially growing culture.

Batch cultures were prepared as previously described (Desvaux *et al.*, 2000). The chemostat system used was a segmented gas-liquid continuous culture device as described by Weimer *et al.* (1991b) with some modifications (Desvaux *et al.*, 2001). The cultures were maintained for a period of eight to nine residence times (Desvaux *et al.*, 2001); for each condition the data were the mean of at least three samples.


**Analytical procedures.** Biomass, cellulose concentration, gases, extracellular proteins, amino acids, glucose, soluble celloextrins, glycogen, acetate, ethanol and lactate extracellular pyruvate were assayed as described by Desvaux *et al.* (2001). Pyridine nucleotides, coenzyme A (CoA), acetyl-CoA, G1P and G6P were extracted and fluorimetric determination performed as previously described (Desvaux *et al.*, 2001). ATP and ADP were measured using the luciferin-luciferase luminescence system (Microbial Biomass Test Kit, Celsis Lumac) (Guedon *et al.*, 2000a).

**Enzyme assays.** Cell extracts were prepared and enzyme assays performed as previously described (Guedon *et al.*, 2000a). Pyruvate-ferredoxin (Fd) oxidoreductase (PFO) (EC 1.2.7.1), lactate dehydrogenase (LDH) (EC 1.1.1.27), phosphotransacetylase (PTA) (EC 2.3.1.8), acetate kinase (AK) (EC 2.7.2.1), acetaldehyde dehydrogenase (AADH) (EC 1.2.1.10) and alcohol dehydrogenase (ADH) (EC 1.1.1.1) were assayed as described by Desvaux *et al.* (2001).

**Calculations and carbon flux analysis.** The metabolic pathways and equations of cellulose fermentation by *C. cellulolyticum*, expressed as *n* hexose equivalents (hexose eq) corresponding to *n* glucose residues of the cellulose chain, were previously reported (Desvaux *et al.*, 2000).

The  $q_{\text{cellulose}}$  is the specific rate of hexose residue fermented in  $\text{mmol (g cells)}^{-1} \text{ h}^{-1}$ .  $q_{\text{acetate}}$ ,  $q_{\text{ethanol}}$  and  $q_{\text{lactate}}$  are the specific rates of product formation in  $\text{mmol (g cells)}^{-1} \text{ h}^{-1}$ .  $q_{\text{extracellular pyruvate}}$  is the specific rate of extracellular pyruvate formation in  $\mu\text{mol (g cells)}^{-1} \text{ h}^{-1}$ .  $q_{\text{NADH produced}}$  and  $q_{\text{NADH consumed}}$  are the specific rates of NADH production and NADH consumption respectively in  $\text{mmol (g cells)}^{-1} \text{ h}^{-1}$  and were calculated as follows:  $q_{\text{NADH produced}} = q_{\text{pyruvate}}$  and  $q_{\text{NADH consumed}} = 2q_{\text{ethanol}} + q_{\text{lactate}}$ . The specific rate of acid production (O'Sullivan & Condon, 1999), was calculated as follows:  $q_{\text{H}^+} = q_{\text{acetate}} + q_{\text{lactate}} + q_{\text{extracellular pyruvate}}$ .

The molar growth yield ( $Y_{x/s}$ ) is expressed in g cells (mol hexose eq fermented) $^{-1}$ . The energetic yield of biomass ( $Y_{\text{ATP}}$ )



**Fig. 1.** Carbon flux distribution within the central metabolic pathways of *C. cellulolyticum* when grown on cellulosic substrate ( $n$  is the number of hexose residues inside the biopolymer). Carbon flow corresponds to the equations given in Methods.

was (Desvaux *et al.*, 2000):  $Y_{\text{ATP}} = \text{concn}_{\text{biomass}} / (1.94 \text{ concn}_{\text{acetate}} + 0.94 \text{ concn}_{\text{ethanol}} + 0.94 \text{ concn}_{\text{lactate}} + 0.94 \text{ concn}_{\text{extracellular pyruvate}})$ .  $Y_{\text{ATP}}$  is expressed in g cells (mole ATP produced) $^{-1}$ .  $q_{\text{ATP}}$  is the specific rate of ATP generation in  $\text{mmol (g cells)}^{-1} \text{ h}^{-1}$ , calculated by the following equation (Desvaux *et al.*, 2000):  $q_{\text{ATP}} = 1.94 q_{\text{acetate}} + 0.94 q_{\text{ethanol}} + 0.94 q_{\text{lactate}} + 0.94 q_{\text{extracellular pyruvate}}$ . The energetic efficiency (ATP-Eff) corresponding to the ATP generation in cellulose catabolism is given by the ratio of  $q_{\text{ATP}}$  to  $q_{\text{cellulose}}$  (Miyagi *et al.*, 1994).

Distribution of the carbon flow by stoichiometric flux analysis (Papoutsakis, 1984; Desai *et al.*, 1999a, b) was determined by adapting the model developed by Holms (1996) to *C. cellulolyticum* metabolism (Desvaux *et al.*, 2001).

At steady state, the carbon flux through each enzyme of the known metabolic pathway, as indicated in Fig. 1, was calculated in milliequivalents of carbon ( $\text{meq C (g cells)}^{-1} \text{ h}^{-1}$ ), as follows:

$$q_{\text{G1P}} = 0.63 q_{\text{cellulose}}$$

$$q_{\text{glucose}} = 0.37 q_{\text{cellulose}}$$

$$q_{\text{biosynthesis}} = q_{\text{biomass}} - q_{\text{glycogen}} + q_{\text{amino acid}} + q_{\text{extracellular protein}}$$

$$q_{\text{G6P}} = q_{\text{biosynthesis}} + q_{\text{pyruvate}}$$

$$q_{\text{phosphoglucomutase}} = q_{\text{G6P}} - q_{\text{glucose}}$$

$$q_{\text{exopolysaccharide}} = q_{\text{G1P}} - q_{\text{phosphoglucomutase}} - q_{\text{glycogen}}$$

$$q_{\text{acetyl-CoA}} = q_{\text{acetate}} + q_{\text{ethanol}}$$

$$q_{\text{carbon dioxide}} = (q_{\text{acetate}} + q_{\text{ethanol}}) / 2$$


$q_{\text{pyruvate}} = q_{\text{acetate}} + q_{\text{ethanol}} + q_{\text{lactate}} + q_{\text{extracellular pyruvate}} + q_{\text{carbon dioxide}}$   
The turnover of a pool ( $\text{h}^{-1}$ ) was calculated from specific rate and pool size expressed in moles or in carbon equivalents (Holms, 1996). The ratio  $R$  corresponded to the ratio of specific enzyme activity to metabolic flux (Holms, 1996; Desvaux *et al.*, 2001).

## RESULTS

### Kinetic profile in batch cellulose fermentation

*C. cellulolyticum* was grown in a bioreactor on defined medium with  $6.7 \text{ g cellulose l}^{-1}$  under non-pH-controlled and pH-controlled conditions (Fig. 2). In the non-pH-controlled run, the pH dropped to 5.3 after 6 d culture (Fig. 2Ia). Compared to the batch culture controlled at pH 7.2, the final biomass attained in the non-pH-controlled condition was lower, as was the percentage of cellulose degradation, which reached 67%, as against 86% in the pH-controlled fermentation (Figs 2Ib and 2IIf). The maximum specific growth rate, however, was similar in non-pH-controlled and pH-controlled batch culture, i.e.  $0.053$  and  $0.056 \text{ h}^{-1}$  respectively.

The metabolite production profiles were clearly different in the two fermentation modes (Fig. 2Ic and 2If). With pH regulation, higher final levels of acetate and ethanol were reached, whereas the lactate and extracellular



**Fig. 2.** pH variation (a), growth and residual cellulose concentration (b), metabolite concentrations (c), and soluble glucide accumulation (d) during cellulose batch fermentation without (I) and with (II) pH control. ▼, pH; ○, cellulose; ●, biomass; □, acetate; ▲, ethanol; △, lactate; ■, extracellular pyruvate; ▽, glucose; ◆, cellobiose; ◇, cellotriose.

pyruvate concentrations attained in the course of fermentation were higher in a non-pH-controlled run. Moreover, with the pH controlled at 7.2, pyruvate was reconsumed in the course of the fermentation more rapidly than without pH control. Thus once growth stopped, biocatabolic activity of cells ceased in the non-pH-controlled culture whereas in the pH-controlled condition the degraded cellulose was further metabolized by resting cells.

The kinetic profile for individual soluble glucide accumulation indicated that it was a non-growth-associated event (Figs 2Id and 2IId). A higher level of soluble cello-oligosaccharides was achieved in fermentation without pH control: up to 1.56 mM cellobiose was detected as well as 0.36 mM cellotriose. With pH control, the sugar level in the broth medium was very limited since only 0.16 mM cellobiose was found and no cellotriose could be detected. In both culture conditions,

longer celldextrins could not be detected either by HPLC or by TLC techniques. Inasmuch as cellulose was degraded to a greater extent in the pH-controlled culture, the difference in accumulation of sugars may reflect a difference in the cellulose substrate, e.g. its surface area (Weimer *et al.*, 1990, 1991a; Fields & Russell, 2000), at the time cultures entered stationary phase, and/or differences in metabolism of stationary-phase cells.

#### Effect of pH on the growth and metabolite production of *C. cellulolyticum* in cellulose continuous culture


Growth parameters, notably end products measured at each steady state, as a function of the pH value are compiled in Table 1. *C. cellulolyticum* was grown in continuous culture under cellulose-limited conditions and at a constant dilution rate of 0.053 h<sup>-1</sup> with pH

**Table 1.** Fermentation parameters from continuous steady-state cultures of *C. cellulolyticum* as a function of environmental pH at  $D = 0.053 \text{ h}^{-1}$

The cellulose input was 0.37% (w/v). Values are the means of samples at steady state and are shown  $\pm$  standard deviation where appropriate. Where no standard deviation is given, individual values did not vary from the mean by more than 10%.


Parameter	Results obtained at pH value of				
	7.4	7.2	7.0	6.8	6.4
Biomass ( $\text{g l}^{-1}$ )	$0.129 \pm 0.011$	$0.207 \pm 0.018$	$0.234 \pm 0.021$	$0.164 \pm 0.014$	$0.056 \pm 0.005$
Cellulose consumed (mM hexose eq)	$4.75 \pm 0.22$	$7.46 \pm 0.38$	$8.14 \pm 0.41$	$5.31 \pm 0.25$	$1.79 \pm 0.09$
$q_{\text{cellulose}}$ [ $\text{mmol (g cells)}^{-1} \text{ h}^{-1}$ ]	1.95	1.91	1.84	1.72	1.69
$q_{\text{pyruvate}}$ [ $\text{mmol (g cells)}^{-1} \text{ h}^{-1}$ ]	2.76	2.63	2.46	2.31	2.18
Product yield (%)*					
Acetate	70.3	68.11	67.2	62.3	47.7
Ethanol	29.3	30.4	30.8	33.4	44.8
Lactate	0.5	1.5	2.0	4.4	7.5
$q_{\text{extracellular pyruvate}}$ [ $\mu\text{mol (g cells)}^{-1} \text{ h}^{-1}$ ]	3.91	9.47	7.29	13.90	32.42
Glycogen [ $\text{mg (g cells)}^{-1}$ ]	$91.5 \pm 2.6$	$96.8 \pm 2.9$	$107.3 \pm 4.0$	$78.8 \pm 2.3$	$62.7 \pm 1.7$
Extracellular proteins ( $\text{mg l}^{-1}$ )	$11.3 \pm 0.7$	$18.5 \pm 0.9$	$18.2 \pm 0.8$	$15.4 \pm 0.6$	$4.0 \pm 0.3$
Free amino acids ( $\text{mg l}^{-1}$ )	$33.4 \pm 1.6$	$63.3 \pm 3.6$	$62.9 \pm 4.0$	$47.1 \pm 2.1$	$13.9 \pm 0.8$
Carbon recovery (%)	94.7	94.2	92.3	95.3	91.4

\* The product yields were expressed as percentages of  $q_{\text{pyruvate}}$ .



**Fig. 3.** Dry weight of cells, percentage cellulose degradation, and molar and energetic growth yields during growth of *C. cellulolyticum* in continuous culture ( $D = 0.053 \text{ h}^{-1}$ ) under various pH conditions and with cellulose as sole carbon and energy source. ●, Biomass; ○, percentage cellulose degradation; □,  $Y_{X/S}$ ; ■,  $Y_{ATP}$ .

values ranging from 7.4 to 6.4. The primary metabolic end products of the cellulose fermentation were acetate, ethanol, lactate,  $\text{H}_2$  and  $\text{CO}_2$ . In addition to carbon conversion into biomass, amino acids and extracellular proteins were also detected in the supernatant (Table 1). Exopolysaccharides were readily observable by micro-



**Fig. 4.** Specific production rate (a) and product ratio (b) during growth of *C. cellulolyticum* in cellulose-fed continuous culture ( $D = 0.053 \text{ h}^{-1}$ ) under various pH conditions. □,  $q_{\text{acetate}}$ ; ▲,  $q_{\text{ethanol}}$ ; ▾,  $q_{\text{lactate}}$ ; ●,  $q_{\text{H}^+}$ ; ○,  $\text{H}_2/\text{CO}_2$ ; ■, ethanol/acetate ratio.

**Table 2.** Redox and energetic balance of *C. cellulolyticum* cells at steady state

Parameter	Results obtained at pH value of				
	7.4	7.2	7.0	6.8	6.4
NADH [ $\mu\text{mol (g cells)}^{-1}$ ]	4.75 $\pm$ 0.85	4.68 $\pm$ 0.90	4.90 $\pm$ 0.97	5.26 $\pm$ 1.02	5.93 $\pm$ 1.14
NAD <sup>+</sup> [ $\mu\text{mol (g cells)}^{-1}$ ]	11.31 $\pm$ 2.22	12.01 $\pm$ 2.36	11.40 $\pm$ 2.25	11.22 $\pm$ 2.17	11.69 $\pm$ 2.33
NADH/NAD <sup>+</sup> ratio	0.42	0.39	0.43	0.47	0.51
$q_{\text{NADH produced}}$ [ $\text{mmol (g cells)}^{-1} \text{ h}^{-1}$ ]	2.76	2.63	2.46	2.31	2.18
$q_{\text{NADH used}}$ [ $\text{mmol (g cells)}^{-1} \text{ h}^{-1}$ ]	1.63	1.64	1.56	1.64	2.11
$q_{\text{NADH produced}}/q_{\text{NADH used}}$ ratio	1.69	1.61	1.57	1.41	1.03
Pool ATP+ADP [ $\mu\text{mol (g cells)}^{-1}$ ]	5.13	4.79	5.39	5.92	6.11
ATP/ADP ratio	0.50	0.46	0.43	0.39	0.34
$q_{\text{ATP}}$ [ $\text{mmol (g cells)}^{-1} \text{ h}^{-1}$ ]	4.53	4.26	3.97	3.61	3.08
ATP-Eff*	2.32	2.23	2.15	2.10	1.82

\* ATP-Eff is the ATP generation efficiency.

**Table 3.** Estimation of carbon flow based on the steady-state values from a cellulose chemostat of *C. cellulolyticum* at  $D = 0.053 \text{ h}^{-1}$  as a function of environmental pH

Carbon flow was calculated as specified in Methods. See Fig. 1 for the metabolic pathways. The values are expressed both as  $\text{meq C (g cells)}^{-1} \text{ h}^{-1}$  and, in parentheses, as a percentage of the specific rate of cellulose consumed ( $q_{\text{cellulose}}$ ).


Carbon flow	Results obtained at pH value of				
	7.4	7.2	7.0	6.8	6.4
$q_{\text{cellulose}}$	11.74 (100.0)	11.46 (100.0)	11.04 (100.0)	10.56 (100.0)	10.13 (100.0)
$q_{\text{G1P}}$	7.39 (63.0)	7.22 (63.0)	6.95 (63.0)	6.65 (63.0)	6.38 (63.0)
$q_{\text{glycogen}}$	0.18 (1.5)	0.19 (1.7)	0.21 (1.9)	0.16 (1.5)	0.12 (1.2)
$q_{\text{exopolysaccharide}}$	0.41 (3.5)	0.42 (3.7)	0.60 (5.5)	0.51 (4.8)	0.63 (6.2)
$q_{\text{glucose}}$	4.35 (37.0)	4.24 (37.0)	4.09 (37.0)	3.91 (37.0)	3.75 (37.0)
$q_{\text{phosphoglucomutase}}$	6.80 (57.9)	6.60 (57.6)	6.14 (55.6)	5.99 (56.7)	5.63 (55.5)
$q_{\text{G6P}}$	11.15 (94.9)	10.85 (94.7)	10.23 (92.6)	9.90 (93.7)	9.38 (92.6)
$q_{\text{extracellular pyruvate}}$	0.01 (0.1)	0.03 (0.2)	0.02 (0.2)	0.04 (0.4)	0.10 (1.0)
$q_{\text{biosynthesis}}$	2.84 (24.2)	2.94 (25.7)	2.84 (25.7)	2.92 (27.6)	2.78 (27.4)
$q_{\text{pyruvate}}$	8.31 (70.7)	7.90 (69.0)	7.39 (66.9)	6.98 (66.1)	6.60 (65.2)
$q_{\text{acetyl-CoA}}$	5.50 (46.9)	5.17 (45.1)	4.81 (43.6)	4.43 (41.9)	4.01 (39.6)
$q_{\text{lactate}}$	0.04 (0.3)	0.12 (1.0)	0.15 (1.4)	0.30 (2.9)	0.49 (4.8)
$q_{\text{CO}_2}$	2.75 (23.4)	2.59 (22.6)	2.41 (21.8)	2.21 (21.0)	2.01 (19.8)
$q_{\text{ethanol}}$	1.62 (13.8)	1.60 (13.9)	1.51 (13.7)	1.54 (14.6)	1.94 (19.2)
$q_{\text{acetate}}$	3.88 (33.1)	3.58 (31.2)	3.30 (29.9)	2.88 (27.3)	2.07 (20.4)

scopic examination but could not be measured as previously described (Payot *et al.*, 1998) due to the significant interference with cellulose fibres leading to erroneous estimation of their concentration. Taking into account amino acids, proteins, fermentative end products and biomass concentration, the carbon balance ranged between 91.4 and 95.3% (Table 1).

As the pH was lowered from 7.4 to 7.0, the dry weight of cells increased (Fig. 3); with a further pH decline, however, the cell density decreased and a steady state of the culture could not be established at pH 6.2 since washout occurred. Whatever the pH conditions, *C. cellulolyticum* always left some cellulose undigested

(Fig. 3). When the pH value was decreased from 7.4 to 7.0, the percentage of cellulose degradation increased from 20.6 to 34.2% and then dropped to reach 7.3% at pH 6.4. Whereas *C. cellulolyticum* showed depressed dry weight of cells at pH values higher than 7.0, the observed cell yields ( $Y_{X/S}$ ) did not decline, even at a pH value close to washout (Fig. 3);  $Y_{X/S}$  increased from 27.1 to 31.3 g cells (mol ATP)<sup>-1</sup> between pH 7.4 and 6.4. The energetic yield of biomass ( $Y_{\text{ATP}}$ ) changed in the same way as  $Y_{X/S}$ ; such a result means that approximately the same amount of ATP was used for the process implicated in cell growth regardless of the pH value.

In all runs, acetate was the main fermentative end



**Fig. 5.** Effect of pH on the pools (a, c) and ratio (b, d) of G1P and G6P (a, b), and of acetyl-CoA and CoA (c, d), when *C. cellulolyticum* is grown on cellulose in continuous culture at  $D = 0.053 \text{ h}^{-1}$ . ○, G6P; ●, G1P; △, G6P/G1P; ■, acetyl-CoA; □, CoA; ▲, acetyl-CoA/CoA.

product: it represented between 70.3 and 47.7% of the carbon directed towards catabolites (Table 1). With lowering of pH, the specific production rate of ethanol increased while that of acetate declined, the latter always remaining the higher (Fig. 4a). The ethanol-to-acetate ratio increased with decreasing pH and reached 0.94 at pH 6.4 (Fig. 4b). In contrast, the  $\text{H}_2/\text{CO}_2$  ratio followed a downward trend with decreasing pH (Fig. 4b). Lactate was also produced at all pH values, but whereas the specific production rate ( $q_{\text{lactate}}$ ) was low at pH 7.4, i.e.  $0.01 \text{ mmol (g cells)}^{-1} \text{ h}^{-1}$ , it increased as the pH decreased, reaching  $0.16 \text{ mmol (g cells)}^{-1} \text{ h}^{-1}$  at pH 6.4 (Fig. 4a). As the pH was lowered, the specific rate of acid production ( $q_{\text{H}^+}$ ), calculated from the spectrum of acid end products formed in cellulose chemostat culture

grown at the different pH values, continuously decreased from 1.95 to  $1.23 \text{ mmol (g cells)}^{-1} \text{ h}^{-1}$  (Fig. 4a).

### Redox and energetic balance in a cellulose-fed chemostat at various pH values

The NADH balance calculated from the known metabolic pathways producing and consuming reducing equivalents, i.e.  $q_{\text{NADH produced}}/q_{\text{NADH used}}$ , continuously decreased with pH from its value of 1.69 at pH 7.4 and was approximately 1 at pH 6.4 (Table 2). Whereas  $\text{NAD}^+$  molecules are reduced by glyceraldehyde 3-phosphate dehydrogenase during ethanol, lactate or acetate formation, only the ethanol and lactate production pathways allow the regeneration of the  $\text{NAD}^+$  pool via dehydrogenase activities. Despite the apparent imbalance at pH values higher than 6.4, the intracellular  $\text{NADH}/\text{NAD}^+$  ratio was always lower than 1 (Table 2). At pH values higher than 6.4,  $q_{\text{lactate}}$  and  $q_{\text{ethanol}}$  were then not sufficient to regenerate  $\text{NADH}$  to  $\text{NAD}^+$ . This result correlated with the  $\text{H}_2/\text{CO}_2$  ratios, which were always higher than 1 (Fig. 4b), suggesting that low lactate and ethanol production was balanced by  $\text{H}_2$  gas formation via  $\text{NADH-Fd}$  reductase and hydrogenase activities.

The stoichiometry of ATP generated over hexose eq fermented (ATP-Eff) decreased from 2.32 to 1.82 as the pH was lowered from 7.4 to 6.4;  $q_{\text{ATP}}$  also declined, from  $4.53$  to  $3.08 \text{ mmol (g cells)}^{-1} \text{ h}^{-1}$  (Table 2). The ATP + ADP pool fluctuated between 4.79 and  $6.11 \text{ μmol (g cells)}^{-1}$ , while the ATP/ADP ratio decreased from 0.50 to 0.34 with environmental acidification (Table 2). This variation was probably correlated with the change of the end products, namely the decrease of acetate biosynthesis.

### Metabolic flux analysis of cells grown on cellulose in an acid environment

The metabolism of *C. cellulolyticum* when grown on cellulose is depicted in Fig. 1. With acidification of the growth medium from pH 7.4 to 6.4, the rate of cellulose consumption declined from 11.74 to  $10.13 \text{ meq C (g cells)}^{-1} \text{ h}^{-1}$  (Table 3). The proportion of the cellulose converted into biomass, free amino acids and extracellular proteins, i.e.  $q_{\text{biosynthesis}}$ , varied from 24.2 to 27.4% with this pH decline (Table 3); considering each, the biomass formation increased from 17.9 to 20.7% of the original carbon while cellulose conversion into amino acid and extracellular protein accounted for around 5.3 and 1.6% of the carbon uptake respectively. Another part of the carbon flow was directed towards metabolite fermentation, i.e. acetate, ethanol,  $\text{CO}_2$ , extracellular pyruvate and lactate, which as a whole decreased from 70.7 to 65.2% of the cellulose consumed (Table 3). Metabolism distributed carbon differently over known catabolic routes as the pH declined. Carbon flow through the  $\text{CO}_2$  and acetate formation pathways declined from 23.4 to 19.8% and from 33.1 to 20.4% respectively. However, ethanol formation increased from 13.8 to 19.2% of the carbon consumed; lactate



**Table 4.** Specific enzymic activity and flux relative to available enzyme activity in *C. cellulolyticum* cell extract at steady state

Enzyme*	Results obtained at pH value of									
	7.4		7.2		7.0		6.8		6.4	
	SEA†	R‡	SEA	R	SEA	R	SEA	R	SEA	R
PTA	0.83 ± 0.08	19.1	0.97 ± 0.07	24.0	0.71 ± 0.08	19.2	0.61 ± 0.05	18.8	0.46 ± 0.06	19.5
AK	0.88 ± 0.09	20.6	0.83 ± 0.10	20.1	0.64 ± 0.06	17.2	0.49 ± 0.05	15.2	0.30 ± 0.03	12.9
AADH	0.12 ± 0.01	6.3	0.18 ± 0.02	10.1	0.21 ± 0.01	12.4	0.19 ± 0.01	11.3	0.22 ± 0.03	9.8
ADH	0.29 ± 0.02	15.7	0.28 ± 0.03	15.6	0.25 ± 0.02	14.9	0.23 ± 0.01	13.3	0.27 ± 0.03	12.3
LDH	0.15 ± 0.01	529.5	0.19 ± 0.03	215.3	0.18 ± 0.01	155.4	0.14 ± 0.01	62.9	0.17 ± 0.02	46.1
PFO	0.61 ± 0.05	9.8	0.66 ± 0.03	11.2	0.34 ± 0.04	6.3	0.13 ± 0.02	2.5	0.08 ± 0.01	1.8

\* PTA, phosphotransacetylase; AK, acetate kinase; AADH, acetaldehyde dehydrogenase; ADH, alcohol dehydrogenase; LDH, L-lactate dehydrogenase; PFO, pyruvate-Fd oxidoreductase.

† SEA is the specific activity of enzyme expressed in  $\mu\text{mol min}^{-1} (\text{mg protein})^{-1}$ .

‡ R is the ratio of specific enzymic activity to metabolic flux through the considered metabolic pathway. Flux was previously expressed as  $\mu\text{mol} (\text{mg protein})^{-1} \text{min}^{-1}$ .

formation also increased, from only 0.3% at pH 7.4 to 4.8% at pH 6.4. At the same time, the proportion of carbon flowing towards extracellular pyruvate rose from 0.1 to 1% with lower environmental pH.

As the pH decreased, the G6P pool slightly increased (Fig. 5a); in terms of turn-over, this pool varied from 116.4 to 86.5  $\text{h}^{-1}$  from pH 7.4 to 6.4. This result was correlated with the concomitant decrease of the carbon flow through glycolysis on the one hand and the increase of the carbon flow through the biosynthesis pathway on the other, as the environmental pH fell. The  $q_{\text{G6P}}$  decreased from 11.15 to 9.38  $\text{meq C (g cells)}^{-1} \text{h}^{-1}$  with acidification, which represented 94.9–92.6% of the original carbon metabolized (Table 3). As for the G1P pool, it increased with decreasing pH (Fig. 5a) which was expressed as a decrease of the turn-over of this pool from 91.3 to 42.7  $\text{h}^{-1}$  from pH 7.4 to 6.4. As a result the G6P/G1P ratio ranged from 1.19 to 0.73 with decreasing environmental pH (Fig. 5b). G1P is an important node in cellulose metabolism since carbon could either flow down glycolysis via phosphoglucose mutase or stored as glycogen, or be converted into exopolysaccharide.  $q_{\text{phosphoglucose mutase}}$  decreased from 6.80 to 5.63  $\text{meq C (g cells)}^{-1} \text{h}^{-1}$  with lowering of pH; when expressed as a percentage of  $q_{\text{cellulose}}$ , this flux ranged from 57.9 to 55.5% (Table 3). The proportion of carbon through  $q_{\text{glycogen}}$  varied from 1.2 to 1.9%, while  $q_{\text{exopolysaccharide}}$  represented 3.5% of the carbon uptake at pH 7.4 to reach 6.2% at pH 6.4 (Table 3).

The CoA pool was fuelled by phosphotransacetylase and acetaldehyde dehydrogenase, whereas acetyl-CoA was formed via pyruvate-Fd oxidoreductase activity (Fig. 1). With lowering of the pH, both pools increased (Fig. 5c), however the acetyl-CoA/CoA ratio decreased from 4.46 to 3.06 (Fig. 5d). Such results were correlated with (i) the rerouting of carbon flow towards biosynthesis, which increased from 24.2 to 27.4% at the

expense of  $q_{\text{pyruvate}}$ , which decreased from 70.7 to 65.2  $\text{meq C (g cells)}^{-1} \text{h}^{-1}$  (Table 3) and (ii) the decrease of carbon flowing through acetate production from 33.1 to 20.4% whereas it increased only from 13.8 to 19.2% through the ethanol pathway as pH declined (Table 3).

#### Enzymic activities as a function of environmental pH

The influence of environmental pH on the specific activities of the enzymes studied is compiled in Table 4. *In vitro*, PFO, PTA, AK and AADH activities were higher under conditions giving higher *in vivo* specific production rates (Table 4). When the carbon flow was expressed as  $\mu\text{mol} (\text{mg protein})^{-1} \text{h}^{-1}$  from previously calculated values (Table 3), R (the ratio of the specific enzyme activity to metabolic rate) could be calculated (Holms, 1996). In the metabolic branch leading to acetate production through PTA and AK, R fluctuated between 18.8 and 24.0, and between 20.6 and 12.9, respectively, as pH decreased (Table 4). R for the enzymes of the ethanol pathway varied between 6.3 and 12.4, and between 15.7 and 12.3 for AADH and ADH, respectively (Table 4). At each step in the central metabolic pathways, the intracellular concentration of substrates, products, cofactors or effector molecules as well as intracellular ionic strength, redox potential or pH can influence the partition and regulation of the carbon flux (Holms, 1986). Nevertheless, the fact that fluxes were much less than the available enzyme activity indicated that the carbon flows were determined by the concentration of substrate available rather than the enzyme activity (Holms, 1996). Despite the variation of enzyme biosynthesis, the amount of these enzymes was always sufficient to catabolize the flowing carbon since R was much higher than 1. At pH 7.4, for the lactate formation pathway, R was very high, i.e. 529.5. This indicated that although LDH was readily available, little carbon was catabolized by this metabolic pathway. As

the pH decreased, however,  $R$  declined to 46.1, indicating that the LDH was more and more implicated in carbon conversion (Table 4). As for the metabolic route through PFO,  $R$  was in the same range as PTA, AK, AADH or ADH for environmental pH values between 7.4 and 7.0, i.e.  $R$  between 6.3 and 11.2 (Table 4), indicating that the fluxes were much less than the available enzyme activity. Yet for a pH lower than 7.0,  $R$  markedly decreased and reached 1.8 at pH 6.4 (Table 4). Then both biosynthesis and catalytic efficiency of PFO declined with pH since the specific enzyme activity and  $q_{\text{acetyl-CoA}}$  decreased (Tables 3 and 4).

## DISCUSSION

Contrary to what was first observed by Giallo *et al.* (1983), pH control during batch culture fermentation of *C. cellulolyticum* greatly influenced cell growth and metabolism. In fact, maintaining the pH at 7.2, increased cell density, enhanced ethanol and acetate production and raised the extent of cellulose hydrolysis, but did not increase the amount of soluble glucides formed, contrary to what was observed in non-pH-controlled fermentation. As previously observed with increasing the concentration of cellulose (Desvaux *et al.*, 2000), the maximum rate of cellulose degradation observed in pH-controlled cultivation reflects the higher cell mass compared to a non-pH-controlled culture. Cellulolysis continued after the cessation of growth and a high level of soluble glucide accumulation was only observed in a non-pH-controlled culture. The gain in cellulose degradation achieved under pH control was offset by catabolite production rather than soluble sugar accumulation.

In continuous culture, maximum cell density was obtained at pH 7.0, but as the pH declined from 7.0 to 6.4 at constant  $D$ , biomass was lowered more than fourfold. At the same time, the specific rate of cellulose consumption, however, decreased only from 1.84 to 1.69 meq C (g cells)<sup>-1</sup> h<sup>-1</sup>. Thus environmental acidification influenced chiefly the biomass formation rather than cellulose degradation and assimilation. *C. cellulolyticum* did not show depressed yields and the transition to wash-out appeared abrupt. This result would be more consistent with a direct effect on a cellular constituent, such as the negative effect of acid on an enzyme or transport protein (Russell & Dombrowski, 1980; Russell & Diez-Gonzalez, 1998).

During cellulose catabolism by *C. cellulolyticum*, soluble  $\beta$ -glucans are first converted into G1P and G6P (Desvaux *et al.*, 2000). With environmental acidification, the G1P pool increased, since the proportion of carbon flowing via phosphoglucomutase varied between 57.9 and 55.5%. The remaining G1P was directed towards exopolysaccharides (up to 9.9% of the G1P) rather than glycogen synthesis (3.0% maximum of the G1P), both allowing dissipation of carbon surplus (Guedon *et al.*, 2000b). As the culture pH was lowered, the flow through glycolysis decreased while carbon directed to biosynthesis increased; as a result, the G6P pool was between 15.9 and 18.1  $\mu\text{mol}$  (g cells)<sup>-1</sup>. Compared with

conditions of uncoupling between catabolism and anabolism encountered during ammonium-limited chemostat performed with cellobiose (Guedon *et al.*, 2000a), the excess of carbon at the G1P–G6P branch point was here limited; in fact, exopolysaccharides and glycogen could represent up to 16.0 and 21.4% respectively of the cellobiose consumed and cellotriose was detected extracellularly (Guedon *et al.*, 2000a).

The increase of the acetyl-CoA pool was corroborated by the analysis of carbon flux; the proportion of cellulose consumed flowing through PFO diminished as the pH declined, as did the ratio of specific enzymic activity to metabolic flux ( $R$ ), and the flux was rerouted away from acetate production. The acetyl-CoA/CoA ratio decrease was paralleled by decreases in  $\text{H}_2/\text{CO}_2$  and  $q_{\text{NADH produced}}/q_{\text{NADH used}}$ . Despite the variation of the NADH balance, calculated from catabolic pathways producing and consuming reducing equivalents, the intracellular NADH/NAD<sup>+</sup> ratio was well regulated. Such a result is in good agreement with the model of Decker *et al.* (1976), where the NADH-Fd reductase is activated by the acetyl-CoA and inhibited by CoA and which underlines that the fates of NADH and acetyl-CoA regulation are intertwined. From acetyl-CoA, acetate was mainly formed but the flux split differently as the environment was acidified, favouring ethanol production. In addition, as pH declined, the level of lactate production rose and coincided with the pyruvate leak, indicating that PFO could no longer support carbon flowing from glycolysis,  $R$  decreasing to 1.8 at pH 6.4. Whatever the pH, LDH was always biosynthesized. This enzyme operated mainly as the pH declined but always remained in excess since even at pH 6.4,  $R$  was 46.1. In these experimental conditions, LDH allowed draining off part of the pyruvate surplus. At high pH values,  $\text{H}_2/\text{CO}_2$  ratios higher than 1 suggested that  $\text{H}_2$  was produced via NADH-Fd reductase and hydrogenase activities in addition to pyruvate-Fd oxidoreductase and hydrogenase activities. With lower pH values, this ratio decreased and was compensated by the increase of ethanol production until washout occurred.

Reinvestigation of cellulose degradation by *C. cellulolyticum* (Desvaux *et al.*, 2000) showed marked differences in the catabolism of this bacterium as compared with the first investigations carried out (Giallo *et al.*, 1985). The present paper demonstrates that the inhibition of growth first observed with batch culture performed in penicillin flasks sealed with butyl rubber stoppers and without shaking of the medium (Giallo *et al.*, 1983, 1985) is mainly the result of low pH due to acid production in the course of fermentation. The range of pH allowing maximum cell density is restricted; strict control of pH is therefore necessary to obtain the optimum cellulolytic performance in biotechnological processes using *C. cellulolyticum*. Cellulolytic bacteria so far investigated cannot grow at pH values significantly less than 6.0 (Stewart, 1977; Russell & Dombrowski, 1980; Russell & Diez-Gonzalez, 1998). However, it is well established that in anaerobic habitats, particularly in the natural environment, high

fermentation acid concentrations and, as a result, low pH values are often encountered (Ljungdahl & Eriksson, 1985; Goodwin & Zeikus, 1987). Since these bacteria have not developed resistance to low pH environments, this implies that they have evolved in an ecological niche where competition for efficient metabolism in acidic conditions is not crucially important. In the same way that growth of *C. cellulolyticum* under an excess of nutrients (Guedon *et al.*, 1999b) or with an easily available carbon source, such as soluble glucides, appeared as aberrations considering the natural bacterial ecosystem (Desvaux *et al.*, 2000, 2001), cultures without pH control have been shown to be detrimental for optimum growth of this bacterium. These data from monospecies laboratory culture must be extrapolated to microbial ecosystems to explain the maintenance of *C. cellulolyticum* in natural environments. Clearly much remains to be learned about the complex interactions in which this bacterium takes part in microbiota (Kuznetsov *et al.*, 1979; Ljungdahl & Eriksson, 1985; Leschine, 1995; Costerton *et al.*, 1995).

## ACKNOWLEDGEMENTS

This work was supported by the Commission of European Communities FAIR programme (contract no. CT95-0191 [DG 12 SSMA]) and by the programme Agrice (Contract no. 9701041).

The authors thank G. Raval for technical assistance and E. McRae for correcting the English and for critical reading of the manuscript.

## REFERENCES

- Bayer, E. A. & Lamed, R. (1992). The cellulose paradox: pollutant par excellence and/or a reclaimable natural resource? *Bio-degradation* **3**, 171–188.
- Bayer, E. A., Shoham, Y., Tormo, J. & Lamed, R. (1996). The cellulosome: a cell surface organelle for the adhesion to and degradation of cellulose. In *Bacterial Adhesion: Molecular and Ecological Diversity*, pp. 155–182. Edited by M. Fletcher. New York: Wiley-Liss.
- Bayer, E. A., Chanzy, H., Lamed, R. & Shoham, Y. (1998). Cellulose, cellulases and cellulosomes. *Curr Opin Struct Biol* **8**, 548–557.
- Béguin, P. & Lemaire, M. (1996). The cellulosome: an exocellular, multiprotein complex specialized in cellulose degradation. *Crit Rev Biochem Mol Biol* **31**, 201–236.
- Boisset, C., Chanzy, H., Henrissat, B., Lamed, R., Shoham, Y. & Bayer, E. A. (1999). Digestion of crystalline cellulose substrates by *Clostridium thermocellum* cellulosome: structural and morphological aspects. *Biochem J* **340**, 829–835.
- Collins, M. D., Lawson, P. A., Willems, A., Cordoba, J. J., Fernandez-Garayzabal, J., Garcia, P., Cai, J., Hippe, H. & Farrow, J. A. E. (1994). The phylogeny of the genus *Clostridium*: proposal of five new genera and eleven species combinations. *Int J Syst Bacteriol* **44**, 812–826.
- Costerton, J. W., Lewandowski, Z., Caldwell, D. E., Korber, D. R. & Lappin-Scott, H. M. (1995). Microbial biofilms. *Annu Rev Microbiol* **49**, 711–745.
- Decker, K., Rössle, M. & Kreuzsch, J. (1976). The role of nucleotides in the regulation of the energy metabolism of *Clostridium kluyveri*. In *Microbial Production and Utilization of Gases*, pp. 75–83. Edited by H. G. Schlegel, G. Gottschalk & N. Pfennig. Göttingen: Akademie der Wissenschaften zu Göttingen.
- Desai, R. P., Harris, L. M., Welker, N. E. & Papoutsakis, E. T. (1999a). Metabolic flux analysis elucidates the importance of acid-formation pathways in regulating solvent production by *Clostridium acetobutylicum*. *Metab Eng* **1**, 206–213.
- Desai, R. P., Nielsen, L. K. & Papoutsakis, E. T. (1999b). Metabolic flux analysis elucidates the importance of acid-formation pathways in regulating solvent production by *Clostridium acetobutylicum* fermentations with non-linear constraints. *J Biotechnol* **71**, 191–205.
- Desvaux, M., Guedon, E. & Petitdemange, H. (2000). Cellulose catabolism by *Clostridium cellulolyticum* growing in batch culture on defined medium. *Appl Environ Microbiol* **66**, 2461–2470.
- Desvaux, M., Guedon, E. & Petitdemange, H. (2001). Carbon flux distribution and kinetics of cellulose fermentation in steady-state continuous cultures of *Clostridium cellulolyticum* on a chemically defined medium. *J Bacteriol* **183**, 119–130.
- Duong, T. V. C., Johnson, E. A. & Demain, A. L. (1983). Thermophilic, anaerobic and cellulolytic bacteria. In *Topics in Enzyme and Fermentation Biotechnology*, pp. 156–195. Edited by A. Weisman. New York: Wiley.
- Fields, M. W. & Russell, J. B. (2000). *Fibrobacter succinogenes* S85 ferments ball-milled cellulose as fast as cellobiose until cellulose surface area is limiting. *Appl Microbiol Biotechnol* **54**, 570–574.
- Gelhay, E., Gehin, A. & Petitdemange, H. (1993a). Colonization of crystalline cellulose by *Clostridium cellulolyticum* ATCC 35319. *Appl Environ Microbiol* **59**, 3154–3156.
- Gelhay, E., Petitdemange, H. & Gay, R. (1993b). Adhesion and growth rate of *Clostridium cellulolyticum* ATCC 35319 on crystalline cellulose. *J Bacteriol* **175**, 3452–3458.
- Giallo, J., Gaudin, C., Belaich, J. P., Petitdemange, E. & Caillet-Mangin, F. (1983). Metabolism of glucose and cellobiose by cellulolytic mesophilic *Clostridium* sp. strain H10. *Appl Environ Microbiol* **45**, 843–849.
- Giallo, J., Gaudin, C. & Belaich, J. P. (1985). Metabolism and solubilization of cellulose by *Clostridium cellulolyticum* H10. *Appl Environ Microbiol* **49**, 1216–1221.
- Goodwin, S. & Zeikus, J. G. (1987). Ecophysiological adaptations of anaerobic bacteria to low pH: analysis of anaerobic digestion in acidic bog sediments. *Appl Environ Microbiol* **53**, 57–64.
- Guedon, E., Desvaux, M., Payot, S. & Petitdemange, H. (1999a). Growth inhibition of *Clostridium cellulolyticum* by an inefficiently regulated carbon flow. *Microbiology* **145**, 1831–1838.
- Guedon, E., Payot, S., Desvaux, M. & Petitdemange, H. (1999b). Carbon and electron flow in *Clostridium cellulolyticum* grown in chemostat culture on synthetic medium. *J Bacteriol* **181**, 3262–3269.
- Guedon, E., Desvaux, M. & Petitdemange, H. (2000a). Kinetic analysis of *Clostridium cellulolyticum* carbohydrate metabolism: importance of glucose 1-phosphate and glucose 6-phosphate branch points for distribution of carbon fluxes inside and outside cells as revealed by steady-state continuous culture. *J Bacteriol* **182**, 2010–2017.
- Guedon, E., Payot, S., Desvaux, M. & Petitdemange, H. (2000b). Relationships between cellobiose catabolism, enzyme levels and metabolic intermediates in *Clostridium cellulolyticum* grown in a synthetic medium. *Biotechnol Bioeng* **67**, 327–335.
- Holms, H. (1986). The central metabolic pathways of *Escherichia coli*: relationship between flux and control at a branch point,

- efficiency of conversion to biomass, and excretion of acetate. *Curr Top Cell Regul* **28**, 69–105.
- Holms, H. (1996).** Flux analysis and control of the central metabolic pathways in *Escherichia coli*. *FEMS Microbiol Rev* **19**, 85–116.
- Huang, L., Gibbins, L. N. & Forsberg, C. W. (1985).** Transmembrane pH gradient and membrane potential in *Clostridium acetobutylicum* during growth under acetogenic and solventogenic conditions. *Appl Environ Microbiol* **50**, 1043–1047.
- Kalachniuk, H. I., Marounek, M., Kalachniuk, L. H. & Savka, O. H. (1994).** Rumen bacterial metabolism as affected by extracellular redox potential. *Ukr Biokhim Zh* **66**, 30–40.
- Kovárová-Kovar, K. & Egli, T. (1998).** Growth kinetics of suspended microbial cells: from single-substrate-controlled growth to mixed-substrate kinetics. *Microbiol Mol Biol Rev* **62**, 646–666.
- Kuznetsov, S. I., Dubinina, G. A. & Lapteva, N. A. (1979).** Biology of oligotrophic bacteria. *Annu Rev Microbiol* **33**, 377–387.
- Leschine, S. B. (1995).** Cellulose degradation in anaerobic environments. *Annu Rev Microbiol* **8**, 237–299.
- Ljungdahl, L. G. & Eriksson, K. E. (1985).** Ecology of microbial cellulose degradation. *Adv Microb Ecol* **8**, 237–299.
- Mitchell, W. J. (1998).** Physiology of carbohydrate to solvent conversion by clostridia. *Adv Microb Physiol* **39**, 31–130.
- Miyagi, A., Ohta, H., Kodama, T., Fukui, K., Kato, K. & Shimono, T. (1994).** Metabolic and energetic aspects of the growth response of *Streptococcus rattus* to environmental acidification in anaerobic continuous culture. *Microbiology* **140**, 1945–1952.
- Mokrasch, L. C. (1967).** Use of 2,4,6-trinitrobenzenesulfonic acid for the coestimation of amines, amino acids, and proteins in mixtures. *Anal Biochem* **18**, 64–71.
- O'Sullivan, E. & Condon, S. (1999).** Relationship between acid tolerance, cytoplasmic pH, and ATP and H<sup>+</sup>-ATPase levels in chemostat cultures of *Lactococcus lactis*. *Appl Environ Microbiol* **65**, 2287–2293.
- Papoutsakis, E. T. (1984).** Equations and calculations for fermentations of butyric acid bacteria. *Biotechnol Bioeng* **26**, 174–187.
- Payot, S., Guedon, E., Cailliez, C., Gelhaye, E. & Petitdemange, H. (1998).** Metabolism of cellobiose by *Clostridium cellulolyticum* growing in continuous culture: evidence for decreased NADH reoxidation as a factor limiting growth. *Microbiology* **144**, 375–384.
- Petitdemange, E., Caillet, F., Giallo, J. & Gaudin, C. (1984).** *Clostridium cellulolyticum* sp. nov., a cellulolytic mesophilic species from decayed grass. *Int J Syst Bacteriol* **34**, 155–159.
- Russell, J. B. & Diez-Gonzalez, F. (1998).** The effects of fermentation acids on bacterial growth. *Adv Microb Physiol* **39**, 205–234.
- Russell, J. B. & Dombrowski, D. B. (1980).** Effect of pH on the efficiency of growth by pure cultures of rumen bacteria in continuous culture. *Appl Environ Microbiol* **39**, 604–610.
- Russell, J. B. & Wilson, D. B. (1996).** Why are ruminal cellulolytic bacteria unable to digest cellulose at low pH? *J Dairy Sci* **79**, 1503–1509.
- Russell, J. B., Bond, D. R. & Cook, G. M. (1996).** The fructose diphosphate/phosphate regulation of carbohydrate metabolism in low G+C Gram positive anaerobes. *Res Microbiol* **147**, 528–534.
- Stewart, C. S. (1977).** Factors affecting the cellulolytic activity of rumen contents. *Appl Environ Microbiol* **33**, 497–502.
- Tomme, P., Warren, R. A. J. & Gilkes, N. R. (1995).** Cellulose hydrolysis by bacteria and fungi. *Adv Microb Physiol* **37**, 1–81.
- Weimer, P. J., Lopez-Guisa, J. M. & French, A. D. (1990).** Effect of cellulose fine structure on kinetics of its digestion by mixed ruminal microorganisms in vitro. *Appl Environ Microbiol* **56**, 2421–2429.
- Weimer, P. J., French, A. D. & Calamari, T. A. (1991a).** Differential fermentation of cellulose allomorphs by ruminal cellulolytic bacteria. *Appl Environ Microbiol* **57**, 3101–3106.
- Weimer, P. J., Shi, Y. & Odt, C. L. (1991b).** A segmented gas/liquid delivery system for continuous culture of microorganisms on insoluble substrates and its use for growth of *Ruminococcus flavefaciens* on cellulose. *Appl Microbiol Biotechnol* **36**, 178–183.
- Wolin, M. J. & Miller, T. L. (1987).** Bioconversion of organic carbon to CH<sub>4</sub> and CO<sub>2</sub>. *Geomicrobiol J* **5**, 239–259.

Received 6 November 2000; revised 29 January 2001; accepted 12 February 2001.