



HAL
open science

Kinetics and Metabolism of Cellulose Degradation at High Substrate Concentrations in Steady-State Continuous Cultures of *Clostridium cellulolyticum* on a Chemically Defined Medium

Mickaël Desvaux, Emmanuel Guedon, Henri Petitdemange

► **To cite this version:**

Mickaël Desvaux, Emmanuel Guedon, Henri Petitdemange. Kinetics and Metabolism of Cellulose Degradation at High Substrate Concentrations in Steady-State Continuous Cultures of *Clostridium cellulolyticum* on a Chemically Defined Medium. *Applied and Environmental Microbiology*, 2001, 67 (9), pp.3837 - 3845. 10.1128/AEM.67.9.3837-3845.2001 . hal-02910803

HAL Id: hal-02910803

<https://hal.inrae.fr/hal-02910803>

Submitted on 3 Aug 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Kinetics and Metabolism of Cellulose Degradation at High Substrate Concentrations in Steady-State Continuous Cultures of *Clostridium cellulolyticum* on a Chemically Defined Medium

MICKAËL DESVAUX, EMMANUEL GUEDON, AND HENRI PETITDEMANGE*

Laboratoire de Biochimie des Bactéries Gram +, Domaine Scientifique Victor Grignard, Université Henri Poincaré, Faculté des Sciences, 54506 Vandœuvre-lès-Nancy Cédex, France

Received 2 March 2001/Accepted 31 May 2001

The hydrolysis and fermentation of insoluble cellulose were investigated using continuous cultures of *Clostridium cellulolyticum* with increasing amounts of carbon substrate. At a dilution rate (D) of 0.048 h^{-1} , biomass formation increased proportionately to the cellulose concentration provided by the feed reservoir, but at and above $7.6 \text{ g of cellulose liter}^{-1}$ the cell density at steady state leveled off. The percentage of cellulose degradation declined from 32.3 to 8.3 with 1.9 and $27.0 \text{ g of cellulose liter}^{-1}$, respectively, while cellodextrin accumulation rose and represented up to 4.0% of the original carbon consumed. The shift from cellulose-limited to cellulose-sufficient conditions was accompanied by an increase of both the acetate/ethanol ratio and lactate biosynthesis. A kinetics study of *C. cellulolyticum* metabolism in cellulose saturation was performed by varying D with $18.1 \text{ g of cellulose liter}^{-1}$. Compared to cellulose limitation (M. Desvaux, E. Guedon, and H. Petitdemange, J. Bacteriol. 183:119–130, 2001), in cellulose-sufficient continuous culture (i) the ATP/ADP, NADH/NAD⁺, and $q_{\text{NADH produced}}/q_{\text{NADH used}}$ ratios were higher and were related to a more active catabolism, (ii) the acetate/ethanol ratio increased while the lactate production decreased as D rose, and (iii) the maximum growth yield ($Y_{X/S}^{\text{max}}$) ($40.6 \text{ g of biomass per mol of hexose equivalent}$) and the maximum energetic yield ($Y_{\text{ATP}}^{\text{max}}$) ($19.4 \text{ g of biomass per mol of ATP}$) were lowered. *C. cellulolyticum* was then able to regulate and optimize carbon metabolism under cellulose-saturated conditions. However, the facts that some catabolized hexose and hence ATP were no longer associated with biomass production with a cellulose excess and that concomitantly lactate production and pyruvate leakage rose suggest the accumulation of an intracellular inhibitory compound(s), which could further explain the establishment of steady-state continuous cultures under conditions of excesses of all nutrients. The following differences were found between growth on cellulose in this study and growth under cellobiose-sufficient conditions (E. Guedon, S. Payot, M. Desvaux, and H. Petitdemange, Biotechnol. Bioeng. 67:327–335, 2000): (i) while with cellobiose, a carbon flow into the cell of as high as $5.14 \text{ mmol of hexose equivalent g of cells}^{-1} \text{ h}^{-1}$ could be reached, the maximum entering carbon flow obtained here on cellulose was $2.91 \text{ mmol of hexose equivalent g of cells}^{-1} \text{ h}^{-1}$; (ii) while the NADH/NAD⁺ ratio could reach 1.51 on cellobiose, it was always lower than 1 on cellulose; and (iii) while a high proportion of cellobiose was directed towards exopolysaccharide, extracellular protein, and free amino acid excretions, these overflows were more limited under cellulose-excess conditions. Such differences were related to the carbon consumption rate, which was higher on cellobiose than on cellulose.

Cellulose is the most abundantly produced biopolymer on earth (5, 28). Due to its recalcitrant, durable nature, cellulose accumulates in terrestrial environments, where a variety of cellulolytic microorganism, existing in virtually every niche and climate, decompose it (4, 28, 29). Around 5 to 10% of cellulosic materials are degraded anaerobically, and the final products released during fermentation are methane and carbon dioxide (28, 51); among cellulolytic bacteria, clostridia play an important role in such processes (28).

Clostridium cellulolyticum, a nonruminant, strictly anaerobic, cellulolytic bacterium (45), digests cellulose through the cellulosome (43). This extracellular multienzymatic complex is composed of a variety of cellulases organized around a scaffolding protein called CipC (6, 41, 42). The cellulosomes are found at

the surface of the cells and allow both adhesion and efficient degradative activity against the cellulose fibers (3, 8).

Using cellobiose, which is one of the soluble cellodextrins released during cellulolysis, major differences in the regulation of the carbon flow between carbon-limited and carbon-sufficient continuous cultures have been reported (19, 20). As the dilution rate increases, in cellobiose-limited chemostats the metabolic pathways towards ethanol and lactate contribute to balance the reducing equivalents supplied by acetate formation (19), while under cellobiose-saturated conditions (20) the redox balance is essentially maintained by NADH-ferredoxin (NADH-Fd) reductase-hydrogenase and ethanol dehydrogenase activities and the carbon flow is equilibrated by three overflows, i.e., exopolysaccharide, extracellular protein, and amino acid excretions.

Using a substrate more closely related to the natural ecosystem of the bacterium, recent investigations with cellulose-limited chemostats (11) have indicated that there is neither a shift from an acetate-ethanol fermentation to a lactate-ethanol fermentation nor pyruvate overflow at high catabolic rates as

* Corresponding author. Mailing address: Laboratoire de Biochimie des Bactéries Gram +, Domaine Scientifique Victor Grignard, Université Henri Poincaré, Faculté des Sciences, BP 239, 54506 Vandœuvre-lès-Nancy Cédex, France. Phone: 33 3 83 91 20 53. Fax: 33 3 83 91 25 50. E-mail: hpetitde@lcb.uhp-nancy.fr.




FIG. 1. Carbon flow within the central metabolic pathways of *C. cellulolyticum* grown in cellulose excess. n , number of hexose residues inside the biopolymer. The specific consumption and production rates (q) correspond to the equations given in Table 1.

TABLE 1. Calculations for flux analysis during cellulose-excess fermentation by *C. cellulolyticum* in chemostat culture

Carbon flow ^a	Equation ^b
$q_{\text{cellulose}}$	$(C_{\text{cellulose}}/X) \times D$
$q_{\text{cellobiose}}$	$(C_{\text{cellobiose}}/X) \times D$
$q_{\text{cellotriose}}$	$(C_{\text{cellotriose}}/X) \times D$
q_{lactate}	$(C_{\text{lactate}}/X) \times D$
q_{acetate}	$(C_{\text{acetate}}/X) \times D$
q_{ethanol}	$(C_{\text{ethanol}}/X) \times D$
$q_{\text{extracellular pyruvate}}$	$(C_{\text{extracellular pyruvate}}/X) \times D$
$q_{\text{biosynthesis}}$	$[(C_{\text{biomass}} - C_{\text{glycogen}} + C_{\text{amino acid}} + C_{\text{protein}})/X] \times D$
$q_{\text{carbon dioxide}}$	$1/2 \times (q_{\text{acetate}} + q_{\text{ethanol}})$
q_{pyruvate}	$q_{\text{acetate}} + q_{\text{ethanol}} + q_{\text{lactate}} + q_{\text{extracellular pyruvate}} + q_{\text{carbon dioxide}}$
$q_{\text{acetyl-CoA}}$	$q_{\text{acetate}} + q_{\text{ethanol}}$
q_{G1P}	$0.63 \times q_{\text{cellulose}}$
$q_{\text{G1P towards cellobiose}}$	$1/2 \times q_{\text{cellobiose}}$
$q_{\text{G1P towards cellotriose}}$	$2/3 \times q_{\text{cellotriose}}$
$q_{\text{G1P towards celloedextrin}}$	$q_{\text{G1P towards cellobiose}} + q_{\text{G1P towards cellotriose}}$
q_{glucose}	$0.37 \times q_{\text{cellulose}}$
$q_{\text{celloedextrin}}$	$q_{\text{cellobiose}} + q_{\text{cellotriose}}$
$q_{\beta\text{-glucan towards cellobiose}}$	$1/2 \times q_{\text{cellobiose}}$
$q_{\beta\text{-glucan towards cellotriose}}$	$1/3 \times q_{\text{cellotriose}}$
$q_{\beta\text{-glucan towards celloedextrin}}$	$q_{\beta\text{-glucan towards cellobiose}} + q_{\beta\text{-glucan towards cellotriose}}$
q_{G6P}	$q_{\text{biosynthesis}} + q_{\text{pyruvate}}$
$q_{\text{phosphoglucomutase}}$	$q_{\text{G6P}} - q_{\text{glucose}}$
q_{glycogen}	$(C_{\text{glycogen}}/X) \times D$
$q_{\text{exopolysaccharide}}$	$q_{\text{G1P}} - q_{\text{phosphoglucomutase}} - q_{\text{glycogen}} - q_{\text{celloedextrin}}$

^a Carbon flows are diagrammed in Fig. 1.

^b C is the concentration in meqC liter⁻¹ of compound produced or consumed; X is the biomass concentration expressed as grams liter⁻¹; D is the dilution rate in hour⁻¹; and q is the specific metabolic rate in meqC gram of cells⁻¹ hour⁻¹.

previously observed on cellobiose (18, 19). Thus, with this culture condition, *C. cellulolyticum* appeared well adapted and even restricted to a cellulolytic lifestyle (11, 12). In its natural biota, however, growth under carbon-sufficient conditions is undoubtedly experienced by bacteria and probably more frequently than carbon limitation, since cellulose accumulates in environments (4, 5).

The aim of the present study, then, was to investigate kinetically the *C. cellulolyticum* fermentation under carbon-saturated conditions by using continuous culture analysis with cel-

lulose and a mineral salt-based medium, which are more closely related to the natural ecosystem of the bacterium (11, 19).

MATERIALS AND METHODS

Organism and growth conditions. *C. cellulolyticum* ATCC 35319 (45) was cultured as previously reported (10) on a defined medium (19) containing cellulose MN301 (Macherey-Nagel, Düren, Germany) at various concentrations as specified in Results. All experiments in segmented gas-liquid continuous culture (49) were performed in a 1.5-liter-working-volume fermentor (LSL Biolaftite, St.

TABLE 2. Fermentation parameters from continuous culture of *C. cellulolyticum* with increasing concentrations of cellulose at a D value of 0.048 h⁻¹

Parameter (unit)	Results ^a obtained with a cellulose concn (g liter ⁻¹) of:				
	1.9	3.8	7.6	14.4	27.0
Biomass (g liter ⁻¹)	0.095 ± 0.008	0.195 ± 0.017	0.212 ± 0.023	0.297 ± 0.031	0.294 ± 0.033
Consumed cellulose (mM hexose eq)	3.7 ± 0.2	7.7 ± 0.3	9.7 ± 0.5	14.3 ± 0.8	13.9 ± 0.7
$q_{\text{cellulose}}$ (mmol g of cells ⁻¹ h ⁻¹)	1.87	1.89	2.21	2.31	2.26
Acetate (mM)	3.50 ± 0.13	7.59 ± 0.21	10.59 ± 0.37	17.03 ± 0.55	16.04 ± 0.57
Ethanol (mM)	1.73 ± 0.07	3.34 ± 0.11	3.64 ± 0.14	4.29 ± 0.17	4.15 ± 0.13
Lactate (mM)	0.05 ± 0.01	0.13 ± 0.01	0.62 ± 0.03	0.56 ± 0.02	0.75 ± 0.03
Extracellular pyruvate (μM)	11.4 ± 0.7	33.3 ± 1.8	161.6 ± 9.1	179.2 ± 8.5	194.6 ± 10.2
Glycogen (mg g of cells ⁻¹)	99.1 ± 3.7	102.3 ± 3.1	55.6 ± 1.9	61.4 ± 2.5	62.7 ± 2.2
Cellobiose (mM)	ND ^b	ND	0.08 ± 0.02	0.15 ± 0.02	0.17 ± 0.05
Cellotriose (mM)	ND	ND	0.04 ± 0.01	0.06 ± 0.02	0.07 ± 0.01
Extracellular proteins (mg liter ⁻¹)	8.4 ± 0.5	14.1 ± 0.7	9.9 ± 0.4	6.1 ± 0.3	7.6 ± 0.3
Free amino acids (mg liter ⁻¹)	33.0 ± 1.7	62.4 ± 3.9	77.7 ± 4.2	118.1 ± 5.7	112.4 ± 6.1
$Y_{X/S}$ (g of cells mol of hexose eq ⁻¹)	25.6	25.4	21.8	20.8	21.2
Y_{ATP} (g of cells mol of ATP eq ⁻¹)	11.2	10.8	8.6	7.9	8.2
q_{ATP} (mmol g of cells ⁻¹ h ⁻¹)	4.27	4.43	5.56	6.08	5.84
Carbon recovery (%)	95.3	95.5	97.4	96.8	96.1

^a Values are the averages for samples at steady state ± standard deviations. Values without standard deviations were determined with an average accuracy of ±10%.

^b ND, not detectable.

TABLE 3. Fermentation parameters from continuous steady-state cultures^a of *C. cellulolyticum* under cellulose-sufficient conditions

Parameter (unit)	Results ^b obtained at a <i>D</i> value (h ⁻¹) of:			
	0.026	0.051	0.061	0.080
Biomass (g liter ⁻¹)	0.367 ± 0.034	0.266 ± 0.023	0.205 ± 0.019	0.161 ± 0.012
Consumed cellulose (mM hexose eq)	23.3 ± 1.1	12.2 ± 0.6	8.5 ± 0.4	5.95 ± 0.3
<i>q</i> _{cellulose} (mmol g of cells ⁻¹ h ⁻¹)	1.65	2.33	2.53	2.91
Product yield (%) of <i>q</i> _{pyruvate}				
Acetate	70.8	75.5	76.6	78.0
Ethanol	24.1	21.2	20.8	20.5
Lactate	5.2	3.3	2.6	1.5
Extracellular pyruvate	0.8	0.5	0.4	0.3
<i>q</i> _{pyruvate} (mmol g of cells ⁻¹ h ⁻¹)	2.68	3.55	3.67	4.02
Glycogen (mg g of cells ⁻¹)	52.3 ± 1.7	63.6 ± 2.3	95.4 ± 3.5	81.2 ± 2.8
Cellobiose (mM)	0.24 ± 0.09	0.10 ± 0.04	0.07 ± 0.02	ND ^c
Celotriose (mM)	0.11 ± 0.05	0.05 ± 0.02	0.03 ± 0.01	ND
Extracellular proteins (mg liter ⁻¹)	13.9 ± 0.7	6.2 ± 0.4	4.4 ± 0.3	2.2 ± 0.1
Free amino acids (mg liter ⁻¹)	190.6 ± 8.5	104.9 ± 5.9	62.9 ± 3.7	47.1 ± 2.1
<i>Y</i> _{X/S} (g of cells mol of hexose eq ⁻¹)	15.8	21.9	24.1	27.5
Carbon recovery (%)	97.4	96.7	93.5	92.9

^a The cellulose input was 1.81% (wt/vol), and ammonium was at 15.13 mM.

^b Values are the averages for samples at steady state ± standard deviations. Values without standard deviations were determined with an average accuracy of ±10%.

^c ND, not detectable.

Germain en Laye, France) at 34°C and pH 7.2 and monitored as previously indicated (11).

Analytical procedures. Biomass, cellulose concentration, gas analysis, extracellular protein, amino acid, glucose, soluble cellodextrins, glycogen, acetate, ethanol, lactate, and extracellular pyruvate were determined as described previously (10, 11, 17).

The percentage of cells that were nonadherent to cellulose fibers was determined by vacuum filtration through 3-μm-pore-size polycarbonate membrane (Millipore, Molsheim, France) as described by Wells et al. (50).

The intracellular compounds NAD⁺, NADH, ATP, ADP, AMP, glucose-1-phosphate (G1P), and glucose-6-phosphate (G6P) and the enzymes glyceraldehyde-3-phosphate dehydrogenase (GAPDH) (EC 1.2.1.12), pyruvate-Fd oxidoreductase (PFO) (EC 1.2.7.1), lactate dehydrogenase (LDH) (EC 1.1.1.27), acetate kinase (AK) (EC 2.7.2.1), and alcohol dehydrogenase (ADH) (EC 1.1.1.1) were extracted and assayed as reported previously (11, 17).

Calculations. The metabolic pathways and equations for cellulose fermentation by *C. cellulolyticum* (expressed as *n* hexose equivalents [hexose eq], corresponding to *n* glucose residues of the cellulose chain) were reported previously (10, 11).

The specific rate of hexose residue fermentation (*q*_{cellulose}) and the specific rates of product formation (*q*_{acetate}, *q*_{ethanol}, *q*_{extracellular pyruvate}, *q*_{lactate}, and *q*_{pyruvate}) are expressed in millimoles per gram of cells per hour and were calculated as indicated previously (11). *q*_{NADH produced} and *q*_{NADH used} are the specific rates of NADH production and NADH consumption, respectively, in millimoles per gram of cells per hour and were calculated as follows: *q*_{NADH produced} = *q*_{pyruvate} and *q*_{NADH used} = 2 *q*_{ethanol} + *q*_{lactate}. *q*_{NADH-Fd} was the specific rate of H₂ production via the NADH-Fd-H₂ path and corresponded to *q*_{NADH produced} - *q*_{NADH used}.

The molar growth yield (*Y*_{X/S}) was expressed in grams of cells per mole of hexose eq fermented. The energetic yield of biomass (*Y*_{ATP}) was expressed in grams of cells per mole of ATP produced and calculated as described previously (11): *Y*_{ATP} = concentration_{biomass} / (1.94 concentration_{acetate} + 0.94 concentration_{ethanol} + 0.94 concentration_{lactate} + 0.94 concentration_{extracellular pyruvate}). The specific rate of ATP generation (*q*_{ATP}) was expressed in millimoles per gram of cells per hour and calculated by the following equation (11): *q*_{ATP} = 1.94 *q*_{acetate} + 0.94 *q*_{ethanol} + 0.94 *q*_{lactate} + 0.94 *q*_{extracellular pyruvate}. The energetic efficiency (ATP-Eff) corresponding to the ATP generation in cellulose catabolism is given by the ratio of *q*_{ATP} to *q*_{cellulose} (11).

A Pirt plot was used for the determination of the maximum yield (*Y*^{max}) and the maintenance coefficient (*m*) (46). The energetic charge and oxidation/reduction index (O/R) were calculated as described by Gottschalk (22). The first-order rate constant of cellulose removal was determined with the equation established by Pavlostathis et al. (44) as described previously (11).

Determination of the distribution of the carbon flow by stoichiometric flux analysis (9) was done by adapting the model developed by Holms (26) to *C. cellulolyticum* metabolism as depicted in Fig. 1. For further direct calculation of

the carbon flow at steady state through each enzyme of the known metabolic pathways, the fluxes were expressed in milliequivalents of carbon (meqC) per gram of cells per hour and calculated as indicated in Table 1.

The turnover of a pool (hours⁻¹) corresponded to the rate of input or output divided by the pool size, which is then the number of times that the pool turns over every hour (27). *R* is the ratio of the specific enzyme activity to metabolic flux (11, 27).

RESULTS

***C. cellulolyticum* continuous culture with increasing concentrations of cellulose.** *C. cellulolyticum* was grown on cellulose in independent runs using a segmented gas-liquid continuous culture device at a dilution rate of 0.048 h⁻¹ with substrate concentrations ranging from 1.9 to 27.0 g liter⁻¹ (Table 2). With increasing amounts of substrate, the concentration of consumed cellulose rose, but at above 7.6 g liter⁻¹ (i.e., 47.0 mM hexose eq) the concentration of consumed cellulose stagnated at 13.9 to 14.3 mM hexose eq (Table 2). The biomass concentration at each steady state increased with the cellulose concentration in the feed medium reservoir, but at above 7.6 g liter⁻¹ it remained quite constant at around 0.296 g liter⁻¹ (Table 2), and thus the production of biomass paralleled the amount of digested cellulose. Microscopic examination indicated that at low cellulose concentrations unattached cells were observable and that almost all of the cellulose fibers were colonized by bacteria. All of these results indicated that at above 7.6 g of cellulose liter⁻¹, continuous cultures were carried out under cellulose-sufficient conditions.

Shifting from cellulose-limited to cellulose-excess conditions was accompanied by a drop of both *Y*_{X/S} and *Y*_{ATP}, whereas *q*_{ATP} increased (Table 2), showing that an uncoupling growth phenomenon occurred. The shift from cellulose limitation to cellulose saturation was accompanied by an increase of lactate biosynthesis (Table 2) as well as the acetate/ethanol ratio, which increased from 2.02 to 3.86 with 1.9 and 27.0 g of cellulose liter⁻¹, respectively. As lactate production rose, extracellular pyruvate production increased as well (Table 2). The decrease in ethanol production in favor of acetate production

was associated with additional ATP, explaining the fact that the q_{ATP} increased during the shift to cellulose saturation (Table 2).

On a synthetic medium, cellulose was converted into cell mass, fermentative catabolites, extracellular amino acids, and proteins (Table 2). Exopolysaccharides were observable by microscopic examination but could not be measured due to significant interference, as already explained (11). While cello-dextrins were not present in cellulose limitation, cellobiose and celotriose were detected in the supernatant under cellulose-excess conditions (Table 2). However, neither glucose nor celodextrins with longer chains than celotriose could be assayed by enzymatic, high-pressure liquid chromatography, and thin-layer chromatography techniques. Taking these compounds into account, the global carbon balance was found to be in the range of 95.3 to 97.4% (Table 2).

Cellulose degradation in continuous culture at high substrate concentrations. *C. cellulolyticum* was cultivated in cellulose excess with 18.1 g of cellulose liter⁻¹ at different D values, which ranged from 0.026 to 0.080 h⁻¹ (Table 3). From the lowest to the highest D value tested, the cell density at steady state decreased while the observed cell yield ($Y_{X/S}$) increased (Table 3). The Pirt plot of these data ($r^2 = 0.992$) permitted determination of a $Y_{X/S}^{\text{max}}$ of 40.6 g of biomass mol of hexose eq consumed⁻¹ and a maintenance coefficient (m) of 1.0 mmol of hexose eq g of cells⁻¹ h⁻¹. The percentage of nonadherent cells remained very low, ranging from 7.2 to 2.7% (Fig. 2a). In a cellulose-limited chemostat (i.e., 3.7 g of cellulose liter⁻¹), however, the percentage of planktonic cells decreased from 59.6 to 21.0% as D increased from 0.027 to 0.083 h⁻¹ and was always much higher than in cellulose saturation (Fig. 2a). The proportion of undegraded cellulose was much lower in cellulose limitation than under cellulose-sufficient conditions; with increasing D it rose from 49.6 to 79.0% and from 78.6 to 94.7%, respectively (Fig. 2b). With this culture condition, *C. cellulolyticum* always left undigested cellulose. Plots of S_R/S_0 versus t_R ($t_R = 1/D$) were linear, with a first-order rate constant of 0.008 h⁻¹ determined from linear regression of the data ($r^2 = 0.996$) (Fig. 2b).

Acetate was always the predominant fermentation end product (Table 3), with the ratio of acetate to ethanol increasing from 2.94 to 3.80. Lactate was also significantly produced, while extracellular pyruvate did not exceed 0.8% of the q_{pyruvate} . Another part of the carbon was oriented towards amino acid, protein, and biomass (Table 3). These compounds were taken into account in addition to fermentative end products and celodextrins for calculation of carbon recovery, which then ranged between 92.9 and 97.4%.

Kinetics analysis of microbial cellulose conversion under cellulose-sufficient conditions. The carbon flow in the central metabolic pathway of *C. cellulolyticum* (Fig. 1) grown in cellulose-sufficient continuous culture is compiled in Table 4. With increasing D , the proportion of carbon flowing down the catabolite declined from 81.2 to 69.1%, while it was enhanced through biosynthesis pathways from 16.3 to 23.8%. In parallel, q_{G6P} gradually rose, but this increase represented a decreasing proportion of the original carbon. As a result, the G6P pool slowly declined (Fig. 3) as D rose, and when expressed in term of turnover this pool increased from 22.1 to 41.7 h⁻¹. The proportion of carbon directed towards exopolysaccharide and




FIG. 2. Percentage of nonadherent cells (a) and proportion of undigested cellulose (b) in cellulose-limited (i.e., 3.7 g liter⁻¹) (○, □) and in cellulose-excess (i.e., 18.1 g liter⁻¹) (●, ■) continuous culture of *C. cellulolyticum*. Error bars indicated standard deviations. Inset, correlation between S_R/S_0 and t_R under cellulose-sufficient conditions.

glycogen was low at a D of 0.026 h⁻¹ and reached 5.7 and 1.4%, respectively, with the highest D tested (Table 4). In contrast, the q_{G1P} towards celodextrin declined from 2.0% to nil at a D of 0.080 h⁻¹, since no celodextrin could then be detected (Table 4). The G1P flux through phosphoglucomutase varied from 60.4 to 55.9% (Table 4). G1P then accumulated with increasing D (Fig. 3), which resulted in the turnover decreasing from 46.4 to 15.4 h⁻¹. The proportion of the carbon flux towards phosphoglucomutase declined, and the G1P, which was directed towards celodextrin at low D values, was rerouted towards glycogen and exopolysaccharide at higher D values. The percentage of carbon directed towards the fermentative end products declined as D rose (Table 4). One part of the flux was converted to acetyl coenzyme A (acetyl-CoA), i.e., from 50.9 to 45.2%. In the same time, q_{acetate} and q_{ethanol} increased, but when expressed as a percentage of $q_{\text{cellulose}}$, the two fluxes declined (Table 4). Another part of the carbon flowing down glycolysis was oriented towards the lactate production pathway. As D was enhanced, lactate production decreased, as did the pyruvate leak.

TABLE 4. Carbon fluxes under cellulose-saturated conditions with *C. cellulolyticum*

Carbon flow ^a	Results obtained at a <i>D</i> value (h ⁻¹) of:							
	0.026		0.051		0.061		0.080	
	meqC g of cells ⁻¹ h ⁻¹	%	meqC g of cells ⁻¹ h ⁻¹	%	meqC g of cells ⁻¹ h ⁻¹	%	meqC g of cells ⁻¹ h ⁻¹	%
<i>q</i> _{cellulose}	9.91	100.0	13.98	100.0	15.20	100.0	17.44	100.0
<i>q</i> _{G1P}	6.24	63.0	8.80	63.0	9.57	63.0	10.98	63.0
<i>q</i> _{celloextrin}	0.34	3.5	0.41	3.0	0.37	2.6	ND ^b	ND
<i>q</i> _{G1P towards celloextrin}	0.20	2.0	0.24	1.7	0.21	1.5	ND	ND
<i>q</i> _{β-glucan towards celloextrin}	0.15	1.5	0.18	1.3	0.16	1.1	ND	ND
<i>q</i> _{glycogen}	0.05	0.5	0.12	0.9	0.22	1.4	0.24	1.4
<i>q</i> _{exopolysaccharide}	0.01	0.1	0.10	0.7	0.55	3.6	0.99	5.7
<i>q</i> _{glucose}	3.67	37.0	5.18	37.0	5.63	37.0	6.46	37.0
<i>q</i> _{phosphoglucomutase}	5.99	60.4	8.34	59.7	8.59	56.5	9.75	55.9
<i>q</i> _{G6P}	9.66	97.5	13.52	96.7	14.22	93.5	16.21	92.9
<i>q</i> _{biosynthesis}	1.61	16.3	2.88	20.6	3.22	21.2	4.16	23.8
<i>q</i> _{pyruvate}	8.04	81.2	10.64	76.1	11.00	72.3	12.05	69.1
<i>q</i> _{acetyl-CoA}	5.05	50.9	6.82	48.8	7.11	46.8	7.89	45.2
<i>q</i> _{lactate}	0.41	4.2	0.35	2.5	0.29	1.9	0.18	1.0
<i>q</i> _{extracellular pyruvate}	0.06	0.6	0.05	0.4	0.05	0.3	0.04	0.2
<i>q</i> _{carbon dioxide}	2.52	25.5	3.41	24.4	3.56	23.4	3.95	22.6
<i>q</i> _{ethanol}	1.28	12.9	1.50	10.7	1.52	10.0	1.64	9.4
<i>q</i> _{acetate}	3.77	38.0	5.33	38.1	5.59	36.8	6.25	35.8

^a Carbon flows were calculated as specified in Materials and Methods and are diagrammed in Fig. 1.

^b ND, not determined.

Relationships between carbon flow and enzymatic activities, energetic balance, and redox balance. In vitro GAPDH, PFO, ADH, and AK activities were higher under growth conditions giving higher in vivo specific production rates (Table 5). For LDH, however, the specific enzyme activities decreased with *D*, which was correlated with the in vivo lactate production rate. A ratio of specific enzyme activity to metabolic flux (*R*) (11, 27) was then calculated; *R* was higher than 1 for all enzymes tested. Therefore, these enzymes were not limiting with respect to the carbon flow, and thus fluxes were determined more by the concentration of substrate available than by the enzyme activity (12, 26).

As *D* increased, *q*_{ATP} was enhanced while the stoichiometry




FIG. 3. G1P (○), G6P (●), and G6P/G1P ratio (□) as a function of dilution rate in cellulose-sufficient continuous culture of *C. cellulolyticum*.

of ATP generated over fermented cellulose, i.e., ATP-Eff, declined from 2.66 to 2.37 (Table 6). Thus, the acetate production could not compensate for the ATP loss per hexose eq fermented due to the decrease of both ethanol and lactate production. A mean value of 0.77 was obtained for the adenylate energy charge (Table 6). The apparent energetic yield increased with *D* (Table 6), and from a Pirt plot of the data ($r^2 = 0.988$) a $Y_{\text{ATP}}^{\text{max}}$ of 19.4 g of cells mol of ATP⁻¹ and an m_{ATP} of 3.1 mmol of ATP g of cells⁻¹ h⁻¹ were determined.

Calculating the coenzyme balance, it could first be observed that both $q_{\text{NADH produced}}$ and $q_{\text{NADH used}}$ increased with *D*, as did the $q_{\text{NADH produced}}/q_{\text{NADH used}}$ ratio (Table 7). This excess of produced NADH correlated with increases of both the H₂/CO₂ ratio, which was always higher than 1, and the $q_{\text{NADH-Fd}}$. These results suggested that the intracellular NADH/NAD⁺ ratio was maintained by the NADH-Fd reductase and hydrogenase, since these interconnected enzymatic activities can oxidize NADH via H₂ production (18–20). The O/R, determined from the gas production ratio and fermentative end product concentration, was very close to 1 and indicated an efficient reoxidation of NADH via H₂ production in addition to carbon fermentative pathways (11).

DISCUSSION

A continuous culture system in which the feed rate is set externally is generally regarded as a chemostat if cell growth is also limited by a selected nutrient(s) (21, 52). Upon increasing the carbon substrate concentration in the feed reservoir, cellulose was no longer the growth-limiting nutrient at and above 7.6 g of cellulose liter⁻¹, and all other nutrients appeared in excess (10, 20). Growth under substrate excess generally results in oscillations and hysteresis (23, 24, 37, 52), but such phenomena did not occur, since steady states of both residual cellulose concentration and biomass monitored during cell culture could

TABLE 5. Specific enzymatic activities in *C. cellulolyticum* cell extract at steady-state growth under cellulose-excess conditions

Enzyme	Results obtained at D value (h^{-1}) of:							
	0.026		0.051		0.061		0.080	
	Sp act ^a	R^b	Sp act	R	Sp act	R	Sp act	R
GAPDH (EC 1.2.1.12)	1.47 ± 0.22		2.87 ± 0.38		3.24 ± 0.41		3.91 ± 0.47	
LDH (EC 1.1.1.27)	0.25 ± 0.06	80.8	0.22 ± 0.03	82.8	0.16 ± 0.04	74.6	0.11 ± 0.05	81.9
PFO (EC 1.2.7.1)	0.62 ± 0.07	10.9	0.72 ± 0.06	9.4	0.89 ± 0.09	11.1	0.95 ± 0.11	10.7
AK (EC 2.7.2.1)	0.31 ± 0.04	7.3	0.66 ± 0.08	11.0	1.03 ± 0.12	16.4	1.33 ± 0.19	18.9
ADH (EC 1.1.1.1)	0.23 ± 0.03	16.0	0.32 ± 0.04	19.0	0.39 ± 0.03	22.8	0.45 ± 0.06	24.4

^a Expressed in micromoles minute⁻¹ milligram of protein⁻¹. Results are means and standard deviations.

^b R , ratio of specific enzymatic activity to metabolic flux through the considered metabolic path; flux was expressed as micromoles milligram of protein⁻¹ minute⁻¹.

be maintained (39). Then, as previously observed with cellobiose (20), a stable carbon excess continuous culture could be imposed on *C. cellulolyticum*. From studies of growth of *C. cellulolyticum* in cellobiose-fed continuous culture in a stepwise fashion (18) and in cellulose batch culture with a reinoculation mode (10), it was demonstrated that growth arrest was not associated with the production of extracellular toxic compounds, and thus it is unlikely that the steady state of the present continuous culture could be maintained by such growth inhibition. Yet, comparing the maximum growth yields and maximum energetic yield obtained under cellulose-sufficient conditions (i.e., $Y_{X/S}^{\max} = 40.6$ g of biomass mol of hexose eq consumed⁻¹ and $Y_{\text{ATP}}^{\max} = 19.4$ g of cells mol of ATP⁻¹) to those resulting from cellulose limitation (i.e., $Y_{X/S}^{\max} = 50.5$ g of biomass mol of hexose eq consumed⁻¹ and $Y_{\text{ATP}}^{\max} = 30.3$ g of cells mol of ATP⁻¹ [11]), it could be observed that both maximum yields were lowered in the presence of a cellulose excess. The decline of these yields indicated that an uncoupling growth phenomenon had occurred; it also took place with the rise in lactate production accompanied by a pyruvate leak. Thus, the growth stagnation was certainly related to an accumulation of an intracellular inhibitory compound(s) (18); intracellular inhibition could furthermore explain the establishment of a steady state under the condition of an excess of all nutrients (52).

The understanding of microbial cellulose metabolism is of both ecological and biotechnological interest. Cellulose degradation plays a key role in the global carbon cycle (28, 29, 51) and is a promising strategy in consolidated bioprocessing for the production of biochemical compounds (25, 31–34). So far, however, very few studies have been devoted to cellulose digestion by cellulolytic bacteria under substrate-saturated conditions (38, 40, 48). Under cellulose-sufficient culture condi-

tions, cellulose digestion always follows first-order kinetics, where k (0.008 h⁻¹) was much lower than under cellulose-limited conditions (0.046 h⁻¹) (11). Once cellulose-saturated conditions were attained, approximately the same amount of cellulose was digested, since the biomass concentration at steady state stagnated. Therefore, k will vary for each cellulose concentration used with this growth condition, since cellulose degradation will follow first-order kinetics with respect to the remaining cellulose concentration (44). Opposite to what was observed with cellulose limitation (11), the lactate-ethanol production was lowered as D rose and had to be balanced by dihydrogen production via the NADH-Fd reductase, which led to an increased H₂/CO₂ ratio. The acetate/ethanol ratio was always higher than 1, but in contrast to the case with cellulose limitation, it increased with D (11). The specific lactate production rate as well as the pyruvate leak decreased with increasing D but always remained higher than in a cellulose-limited chemostat (11). The R values for LDH were around 80.0 and were not as high as with cellulose limitation, where R could reach 434.2 (11). From the G1P metabolic node, cello-dextrins were produced and represented up to 3.5% of the carbon uptake at the lowest D value tested. G1P was rerouted towards exopolysaccharide and glycogen as D rose; as in cellulose limitation, these biosyntheses could be adjusted as a function of carbon flux. Such glycogen turnover was recently observed with *Fibrobacter succinogenes* on cellulose (7, 14). A concomitant decrease of the percentage of carbon flowing through $q_{\text{phosphoglucumutase}}$, q_{G6P} , and q_{pyruvate} in favor of biosynthesis pathways could explain the relative drop in lactate production as D increased. Compared to that in cellulose-limited chemostats, the proportion of unattached cells was low. With cellulose limitation the available surface area is saturated by bacteria, while under cellulose-sufficient conditions the cel-

TABLE 6. Energetic balance in cellulose-saturated continuous culture of *C. cellulolyticum*

Parameter (unit)	Results ^a obtained at a D value (h^{-1}) of:			
	0.026	0.051	0.061	0.080
ATP ($\mu\text{mol g of cells}^{-1}$)	2.84 ± 0.15	2.31 ± 0.11	2.17 ± 0.19	1.89 ± 0.10
ADP ($\mu\text{mol g of cells}^{-1}$)	1.41 ± 0.08	1.21 ± 0.06	1.29 ± 0.09	1.25 ± 0.05
AMP ($\mu\text{mol g of cells}^{-1}$)	0.32 ± 0.05	0.27 ± 0.03	0.11 ± 0.01	0.25 ± 0.02
Energetic charge	0.78	0.77	0.79	0.74
q_{ATP} (mmol g of cells ⁻¹ h ⁻¹)	4.39	5.98	6.23	6.89
ATP-Eff	2.66	2.57	2.46	2.37
Y_{ATP} (g of cells mol of ATP ⁻¹)	5.9	8.5	9.8	11.6

^a Values are the averages for samples at steady-state ± standard deviations. Values without standard deviations were determined with an average accuracy of ±10%.

TABLE 7. Redox balance of *C. cellulolyticum* at steady state under cellulose-sufficient conditions

Parameter (unit)	Results ^a obtained at a <i>D</i> value (h ⁻¹) of:			
	0.026	0.051	0.061	0.080
NADH (μmol g of cells ⁻¹)	6.67 ± 1.36	7.56 ± 1.59	5.16 ± 1.09	8.12 ± 1.62
NAD ⁺ (μmol g of cells ⁻¹)	11.31 ± 2.33	12.01 ± 2.43	11.40 ± 2.25	12.32 ± 2.44
NADH/NAD ⁺ ratio	0.59	0.63	0.45	0.66
<i>q</i> _{NADH produced} (mmol g of cells ⁻¹ h ⁻¹)	2.68	3.55	3.67	4.02
<i>q</i> _{NADH used} (mmol g of cells ⁻¹ h ⁻¹)	1.42	1.61	1.61	1.70
<i>q</i> _{NADH-Fd} (mmol g of cells ⁻¹ h ⁻¹)	1.26	1.93	2.05	2.32
<i>q</i> _{NADH produced} / <i>q</i> _{NADH used} ratio	1.89	2.20	2.27	2.36
H ₂ /CO ₂ ratio	1.43	1.55	1.61	1.67
O/R	1.03	1.01	0.98	0.96

^a Values are the averages for samples at steady state ± standard deviations. Values without standard deviations were determined with an average accuracy of ±10%.

lulose surface area is largely accessible for bacterial adhesion (10, 13). This was correlated with a higher cellulose digestion rate reflected by higher *q*_{cellulose} under substrate excess conditions, since most of the cells adhered to cellulose fibers and thus participated directly in cellulose digestion. While cello-dextrin was undetectable with cellulose limitation (11), cellobiose and cellotriose were detected here in the supernatant; such a finding was certainly related to a reversible phosphorylase reaction (1, 2, 30, 35, 36, 47, 50). Under cellobiose-sufficient conditions (17, 20), only cellotriose was detected, but the present results suggest that cellobiose could also be synthesized de novo during cell growth on cellobiose.

Previous reports on experiments with cellobiose stated that the adhesion-colonization phase of the process of cellulose digestion by *C. cellulolyticum* (15, 16) corresponded to a carbon-sufficient period (20). It was thus argued that a carbon flow of as high as 5.14 mmol of hexose eq g of cells⁻¹ could be attained with cellulose as a substrate (20). With cellulose saturation, however, the entering carbon flow remained lower than expected, i.e., 2.91 mmol of hexose eq g of cells⁻¹ h⁻¹. The NADH/NAD⁺ ratio was always lower than 1 on cellulose, whereas a ratio of as high as 1.51 was obtained with cellobiose excess (20); this result was most probably related to a higher carbon consumption rate which led to rate-limiting fluxes through ethanol and dihydrogen production pathways on cellobiose. Thus, the proper NADH/NAD⁺ ratio was maintained only when ethanol and lactate production complemented the path towards H₂ production via NADH-Fd reductase activity. With a carbon excess, free amino acid could account for 15.4% of the cellobiose fermented (20), against a maximum of 5.8% on cellulose, while exopolysaccharide represented up to 38.1% of the cellobiose consumed (20) and there was a maximum of only 5.7% with cellulose as the substrate. It thus appeared that some of the general metabolic trends associated with carbon-sufficient conditions, such as (i) the ATP/ADP ratio always being higher than 1, (ii) the elevated production of lactate at a low *D*, and (iii) the concomitant increase of *q*_{ethanol}, *q*_{NADH-Fd}, *q*_{NADH produced}/*q*_{NADH used}, NADH/NAD⁺, and H₂/CO₂ as *D* rose and some other regulations of bacterial metabolism, were not observed on cellulose compared to cellobiose. Even if cellulose degradation must be considered as a microbial process rather than a purely enzymatic event, the strong influence of the cellulosome on the entering carbon flow must be taken into account (10, 11). The study of *C. cellulolyticum* catabolism with soluble glucide allowed the demonstration of bacterial

metabolic limitation, but this response should be interpreted as deregulation of the metabolism. All of these results demonstrate that *C. cellulolyticum* was able to correctly regulate and optimize carbon metabolism in limited and saturated conditions using a substrate more representative of its natural environment, i.e., cellulose (12).

ACKNOWLEDGMENTS

This work was supported by the Commission of European Communities FAIR program (contract CT950191 [DG12SSMA]) and by the program Agrice (contract 9701041).

We thank Anne-Cécile Aubry and Guy Raval for excellent technical assistance and Edward McRae for correcting the English and for critical reading of the manuscript.

REFERENCES

- Alexander, J. K. 1972. Cellobiose phosphorylase from *Clostridium thermo-cellum*. *Methods Enzymol.* **28**:944–948.
- Alexander, J. K. 1972. Cellodextrin phosphorylase from *Clostridium thermo-cellum*. *Methods Enzymol.* **28**:948–953.
- Bayer, E. A., H. Chanzy, R. Lamed, and Y. Shoham. 1998. Cellulose, cellulases and cellulosomes. *Curr. Opin. Struct. Biol.* **8**:548–557.
- Bayer, E. A., and R. Lamed. 1992. The cellulose paradox: pollutant par excellence and/or a reclaimable natural resource? *Biodegradation* **3**:171–188.
- Béguin, P., and J. P. Aubert. 1996. The biological degradation of cellulose. *FEMS Microbiol. Rev.* **13**:25–58.
- Belaich, J. P., C. Tardiff, A. Belaich, and C. Gaudin. 1997. The cellulolytic system of *Clostridium cellulolyticum*. *J. Biotechnol.* **57**:3–14.
- Bibollet, X., N. Bosc, M. Matulova, A. M. Delort, G. Gaudet, and E. Forano. 2000. ¹³C and ¹H NMR study of cellulose metabolism by *Fibrobacter succinogenes* S85. *J. Biotechnol.* **77**:37–47.
- Boisset, C., H. Chanzy, B. Henrissat, R. Lamed, Y. Shoham, and E. A. Bayer. 1999. Digestion of crystalline cellulose substrates by *Clostridium thermocellum* cellulosome: structural and morphological aspects. *Biochem. J.* **340**:829–835.
- Desai, R. P., L. K. Nielsen, and E. T. Papoutsakis. 1999. Stoichiometric modeling of *Clostridium acetobutylicum* fermentations with non-linear constraints. *J. Biotechnol.* **71**:191–205.
- Desvaux, M., E. Guedon, and H. Petitdemange. 2000. Cellulose catabolism by *Clostridium cellulolyticum* growing in batch culture on defined medium. *Appl. Environ. Microbiol.* **66**:2461–2470.
- Desvaux, M., E. Guedon, and H. Petitdemange. 2001. Carbon flux distribution and kinetics of cellulose fermentation in steady-state continuous cultures of *Clostridium cellulolyticum* on a chemically defined medium. *J. Bacteriol.* **183**:119–130.
- Desvaux, M., and H. Petitdemange. 2001. Flux analysis of the metabolism of *Clostridium cellulolyticum* grown in cellulose-fed continuous culture on a chemically defined medium under ammonium-limited conditions. *Appl. Environ. Microbiol.* **67**:3846–3851.
- Fields, M. W., and J. B. Russell. 2000. *Fibrobacter succinogenes* S85 ferments ball-milled cellulose as fast as cellobiose until cellulose surface area is limiting. *Appl. Microbiol. Biotechnol.* **54**:570–574.
- Gaudet, G., E. Forano, G. Dauphin, and A. M. Delort. 1992. Futile cycling of glycogen in *Fibrobacter succinogenes* as shown by in situ ¹H-NMR and ¹³C-NMR investigation. *Eur. J. Biochem.* **207**:155–162.
- Gelhaye, E., A. Gehin, and H. Petitdemange. 1993. Colonization of crystal-

- line cellulose by *Clostridium cellulolyticum* ATCC 35319. Appl. Environ. Microbiol. **59**:3154–3156.
16. Gelhaye, E., H. Petitdemange, and R. Gay. 1993. Adhesion and growth rate of *Clostridium cellulolyticum* ATCC 35319 on crystalline cellulose. J. Bacteriol. **175**:3452–3458.
 17. Guedon, E., M. Desvaux, and H. Petitdemange. 2000. Kinetic analysis of *Clostridium cellulolyticum* carbohydrate metabolism: importance of glucose 1-phosphate and glucose 6-phosphate branch points for distribution of carbon fluxes inside and outside cells as revealed by steady-state continuous culture. J. Bacteriol. **182**:2010–2017.
 18. Guedon, E., M. Desvaux, S. Payot, and H. Petitdemange. 1999. Growth inhibition of *Clostridium cellulolyticum* by an inefficiently regulated carbon flow. Microbiology **145**:1831–1838.
 19. Guedon, E., S. Payot, M. Desvaux, and H. Petitdemange. 1999. Carbon and electron flow in *Clostridium cellulolyticum* grown in chemostat culture on synthetic medium. J. Bacteriol. **181**:3262–3269.
 20. Guedon, E., S. Payot, M. Desvaux, and H. Petitdemange. 2000. Relationships between cellobiose catabolism, enzyme levels and metabolic intermediates in *Clostridium cellulolyticum* grown in a synthetic medium. Biotechnol. Bioeng. **67**:327–335.
 21. Gottschal, J. C. 1992. Continuous culture, p. 559–572. In J. Lederberg (ed.), Encyclopedia of microbiology, vol. 1. Academic Press, New York, N.Y.
 22. Gottschalk, G. 1985. Bacterial metabolism. Springer-Verlag, New York, N.Y.
 23. Harrison, D. E. F., and H. H. Topiwala. 1974. Transient and oscillatory states of continuous culture. p. 168–219. In T. H. Ghose and A. Fiechter (ed.), Advanced biochemical engineering, vol. 3. Springer-Verlag, Berlin, Germany.
 24. Hjortso, M. A., and J. Nielsen. 1994. A conceptual model of autonomous oscillations in microbial cultures. Chem. Eng. Sci. **49**:1083–1095.
 25. Hogsett, D. A., H. J. Alm, T. D. Bernardez, C. R. South, and L. R. Lynd. 1992. Direct microbial conversion: prospects, progress, and obstacles. Appl. Biochem. Biotechnol. **34–35**:527–541.
 26. Holms, H. 1986. The central metabolic pathways of *Escherichia coli*: relationship between flux and control at a branch point, efficiency of conversion to biomass, and excretion of acetate. Curr. Top. Cell. Regul. **28**:69–105.
 27. Holms, H. 1996. Flux analysis and control of the central metabolic pathways in *Escherichia coli*. FEMS Microbiol. Rev. **19**:85–116.
 28. Leschine, S. B. 1995. Cellulose degradation in anaerobic environments. Annu. Rev. Microbiol. **49**:399–426.
 29. Ljungdahl, L. G., and K. E. Eriksson. 1985. Ecology of microbial cellulose degradation. Adv. Microb. Ecol. **8**:237–299.
 30. Lou, J., K. A. Dawson, and H. J. Strobel. 1997. Cellobiose and cellodextrin metabolism by the ruminal bacterium *Ruminococcus albus*. Curr. Microbiol. **35**:221–227.
 31. Lynd, L. R. 1996. Overview and evaluation of fuel ethanol from cellulosic biomass: technology, economics, the environments, and policy. Annu. Rev. Energy Environ. **21**:403–465.
 32. Lynd, L. R., H. E. Grethlein, and R. H. Wolkin. 1989. Fermentation of cellulosic substrates in batch and continuous culture of *Clostridium thermocellum*. Appl. Environ. Microbiol. **55**:3131–3139.
 33. Lynd, L. R., J. H. Cushman, R. J. Nichols, and C. E. Wyman. 1991. Fuel ethanol from cellulosic biomass. Science **251**:1318–1323.
 34. Lynd, L. R., C. E. Wyman, and T. U. Gerngross. 1999. Biocommodity engineering. Biotechnol. Prog. **15**:777–793.
 35. Matheron, C., A. M. Delort, G. Gaudet, and E. Forano. 1996. Simultaneous but differential metabolism of glucose and cellobiose in *Fibrobacter succinogenes* cells, studied by in vivo ¹³C NMR. Can. J. Microbiol. **42**:1091–1099.
 36. Matheron, C., A. M. Delort, G. Gaudet, and E. Forano. 1998. In vivo ¹³C NMR study of glucose and cellobiose metabolism by four cellulolytic strains of the genus *Fibrobacter*. Biodegradation **9**:451–461.
 37. Menzel, K., A. P. Zeng, H. Biebl, and W. D. Deckwer. 1996. Kinetic, dynamic and pathway studies of glycerol metabolism by *Klebsiella pneumoniae* in anaerobic continuous culture. 1. The phenomena and characterization of oscillation and hysteresis. Biotechnol. Bioeng. **52**:549–560.
 38. Mitchell, W. J. 1998. Physiology of carbohydrate to solvent conversion by clostridia. Adv. Microb. Physiol. **39**:31–130.
 39. Monod, J. 1950. La technique de culture continue théorie et applications. Ann. Inst. Pasteur **79**:390–410.
 40. Ohmiya, K., K. Nokura, and S. Shimizu. 1983. Enhancement of cellulose degradation by *Ruminococcus albus* at high cellulose concentration. J. Ferment. Technol. **61**:25–30.
 41. Pages, S., A. Belaich, H. P. Fierobe, C. Tardif, C. Gaudin, and J. P. Belaich. 1999. Sequence analysis of scaffolding protein CipC and ORFXp, a new cohesin-containing protein in *Clostridium cellulolyticum*: comparison of various cohesin domains and subcellular localization of ORFXp. J. Bacteriol. **181**:1801–1810.
 42. Pages, S., A. Belaich, C. Tardif, C. Reverbel-Leroy, C. Gaudin, and J. P. Belaich. 1996. Interaction between the endoglucanase CelA and the scaffolding protein CipC of the *Clostridium cellulolyticum* cellulosome. J. Bacteriol. **178**:2279–2286.
 43. Pages, S., L. Gal, A. Belaich, C. Gaudin, C. Tardif, and J. P. Belaich. 1997. Role of scaffolding protein CipC of *Clostridium cellulolyticum* in cellulose degradation. J. Bacteriol. **179**:2810–2816.
 44. Pavlostathis, S. G., T. L. Miller, and M. J. Wolin. 1988. Kinetics of insoluble cellulose fermentation by continuous cultures of *Ruminococcus albus*. Appl. Environ. Microbiol. **54**:2660–2663.
 45. Petitdemange, E., F. Caillet, J. Giallo, and C. Gaudin. 1984. *Clostridium cellulolyticum* sp. nov., a cellulolytic mesophilic species from decayed grass. Int. J. Syst. Bacteriol. **34**:155–159.
 46. Pirt, S. J. 1975. Principles of microbe and cell cultivation. Blackwell Scientific Publishers, Oxford, United Kingdom.
 47. Russell, J. B. 1985. Fermentation of cellodextrins by cellulolytic and non-cellulolytic rumen bacteria. Appl. Environ. Microbiol. **49**:572–576.
 48. Russell, J. B. 1998. Strategies that ruminal bacteria use to handle excess carbohydrate. J. Anim. Sci. **76**:1955–1963.
 49. Weimer, P. J., Y. Shi, and C. L. Odt. 1991. A segmented gas/liquid delivery system for continuous culture of microorganisms on insoluble substrates and its use for growth of *Ruminococcus flavefaciens* on cellulose. Appl. Microbiol. Biotechnol. **36**:178–183.
 50. Wells, J. E., J. B. Russell, Y. Shi, and P. J. Weimer. 1995. Cellodextrin efflux by the cellulolytic ruminal bacterium *Fibrobacter succinogenes* and its potential role in the growth of nonadherent bacteria. Appl. Environ. Microbiol. **61**:1757–1762.
 51. Wolin, M. J., and T. L. Miller. 1987. Bioconversion of organic carbon to CH₄ and CO₂. Geomicrobiol. J. **5**:239–259.
 52. Zeng, A. P. 1999. Continuous culture, p. 151–164. In A. L. Demain and J. E. Davies (ed.), Manual of industrial microbiology and biotechnology. ASM Press, Washington, D.C.