



HAL
open science

Digital soil mapping of key GlobalSoilMap properties in Northern Karnataka Plateau

S. Dharumarajan, B. Kalaiselvi, Amar Suputhra, M. Lalitha, Rajendra Hegde, S.K. Singh, Philippe Lagacherie

► To cite this version:

S. Dharumarajan, B. Kalaiselvi, Amar Suputhra, M. Lalitha, Rajendra Hegde, et al.. Digital soil mapping of key GlobalSoilMap properties in Northern Karnataka Plateau. *Geoderma Régional*, 2020, 20, pp.e00250. 10.1016/j.geodrs.2019.e00250 . hal-02913799

HAL Id: hal-02913799

<https://hal.inrae.fr/hal-02913799>

Submitted on 23 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

26 **Key words:** Digital soil mapping, Vertisols, Alfisols, Quantile Regression Forest, soil
27 properties, Field capacity, Permanent wilting point

28

29 **1. Introduction**

30 Soil properties are assessed through resource inventorisation with the main objective
31 to delineate areas which need uniform management practices and provide users with
32 information on soil properties. Assessment of spatial distribution of soil properties for each
33 location is important for site-specific land management, land evaluation and land suitability
34 analysis (Gessler et al., 2000; McBratney et al., 2003). Although several spatial soil databases
35 are developed throughout the world, they are neither exhaustive nor precise enough for
36 ensuring enlightened decisions. For example, though digitized soil maps are available for
37 most of the world (Grunwald et al., 2011), those information are at very small scale (1:1
38 million or coarser) for many areas and do not adequately represent soil variability in a format
39 that is useful for a non-pedologists (Sanchez et al., 2009). Digital soil mapping (DSM)
40 represents a ground-breaking solution compared to conventional soil survey by its ability to
41 exploit large sets of spatial data, to produce uncertainty estimates associated with soil
42 predictions and can be revised once new data are collected (Lagacherie and McBratney,
43 2007). Soil database generated through field sampling and laboratory analysis are used to
44 feed a DSM model that predicts soil properties in areas not sampled. Digital soil maps also
45 provide the uncertainties associated with such predictions. The overall uncertainty of the
46 prediction is estimated by combining uncertainties of input data, spatial inference, and soil
47 functions (Dharumarajan et al., 2019a). Uncertainties are essential for understanding and
48 dealing with risk in decision-making.

49 DSM has moved from a largely academic towards an operational activity through
50 GlobalSoilMap project (<http://www.globalsoilmap.net/>). The project aims to map the several

51 key soil properties of globe onto a three-dimensional grid at fine spatial resolution with local
52 uncertainty estimates (Arrouays et al., 2014). The first versions of GlobalSoilMap products
53 have already been produced in various countries (Mulder et al., 2016; Grundy et al., 2015;
54 Adhikari et al., 2014; Poggio & Gimona., 2017) with spatial inference functions using
55 globally available landscape parameters such as Digital Elevation Models, multispectral
56 remote sensing, geology maps, and legacy soil maps as inputs.

57 In India, ICAR-National Bureau of Soil Survey and Land Use Planning
58 (ICAR-NBSS&LUP), Nagpur has recently launched an ambitious program called
59 "IndianSoilGrids" with the objective to develop soil properties map as per GlobalSoilMap
60 Specifications. In recent past, effort has been made to compile the legacy soil data in the form
61 of harmonized databases and stored in NBSS&LUP Geoportal. Besides pursuing the storage
62 effort, IndianSoilGrids project paved the ways to exploit legacy soil data through DSM
63 models. In this context, the present exploratory study was carried out to produce a fine
64 resolution map of major GlobalSoilMap soil properties such as organic carbon, pH, CEC,
65 clay, sand, silt, field capacity and permanent wilting point in part of Northern Karnataka
66 Plateau region representing semi arid tropics of south India using Quantile Regression Forest
67 Model techniques.

68

69 **2. Materials and methods**

70 **2.1 Study area**

71 The present study was carried out in part of Koppal and Gadag districts of Northern
72 Karnataka Plateau (Fig.1). The study area is located in 14° 56' to 15° 37' N latitude and
73 75°23' to 76° 25' longitude with an area of 3655 km². The study region represents hot-semi
74 arid climate with rainfall range of 600-750 mm and potential evapo-transpiration (PET) of

75 1600-1700 mm. The average annual rainfall is 672 mm. This area includes mountainous,
76 expansive plateau with substantial area is underlined by basalts with continuation of Deccan
77 trap of Maharashtra. The major area comes under rainfed cultivation with crops like
78 Sorghum, Pigeon pea and Pearl millet. The major soils represented by shallow to deep
79 vertisols, alfisols and inceptisols.

80 ***2.2 Sampling methodology***


81 The profiles studied under Sujala III (Karnataka Watershed Development Project II)
82 project were used for mapping of soil properties. Sixty soil profiles were studied upto 2m or
83 hard rock based on variability in landform and land use. The soil horizons were demarcated
84 and from the representative soil horizons, soil samples were collected for laboratory analysis.
85 Collected soil samples were air dried in shade and passed through 2 mm sieve by gently
86 ground with a wooden mallet. The samples were analysed for particle-size distribution
87 following International Pipette method (Richards, 1954), pH and electrical conductivity (EC)
88 in 1:2.5 soil:water suspension (Jackson, 1962). Organic carbon was estimated by Walkley
89 and Black (1934) method. The cation exchange capacity (CEC) and exchangeable cations
90 were determined as described by Jackson (1973). Field capacity (FC) and permanent wilting
91 point (PWP) were estimated using pressure plate apparatus (Richards, 1956). The profile soil
92 properties were pre-processed by harmonization of soil depth interval (GlobalSoilMap depth
93 specification) predictions using equal-area spline functions (Bishop et al., 1999).

94

95 ***2.3 Environmental covariates and models used***

96 A Digital elevation model (DEM) with 30 m resolution was obtained from SRTM and
97 processed using ArcGIS10 data management tool box. The primary and secondary derivatives

106 of DEM like elevation, slope, aspect, curvatures (plan and profile), topographic wetness
107 Index (TWI) and topographic position index (TPI), LS factor, Multi-resolution Ridge Top
108 Flatness (MrRTF) and Multi-resolution Index of Valley Bottom Flatness (MrVBF) were
109 derived by using Saga-GIS 6.3.0 version. Along with DEM attributes, all the bands of
110 Sentinel- 2 imagery (13 bands), Normalized Difference Vegetation Index (NDVI) and
111 Enhanced vegetation index (EVI) (MOD13Q1) were used as covariates for prediction of soil
112 properties (Table.1). The environmental variables were intersected for all the sampling points
113 for prediction of soil properties.



106

107

Fig.1. Study area with profile locations

108

109

110

Table.1. Different covariates used in the model

111

112

113

114

115

116

117

118

119

Predictor	Source	Resolution
Elevation (m)	SRTM DEM	30 m
Slope (%)	SRTM DEM	30 m
Aspect	SRTM DEM	30 m
TPI	SRTM DEM	30 m
TWI	SRTM DEM	30 m
Plan curvature	SRTM DEM	30 m
Profile curvature	SRTM DEM	30 m
MrVBF	SRTM DEM	30 m
MrRTF	SRTM DEM	30 m
NDVI	MOD13Q1(2011-2015)	250m_16 days
EVI	MOD13Q1(2011-2015)	
Sentinel-2	13 bands of Sentinel 2 data	10-60m

120

121

122

123

124

125

126

127

128

129

130

131

132

Quantile regression forest (QRF) model was used for prediction of soil properties and uncertainty estimates in the study area. QRF is an extension of Random forest model and the advantage of QRF over Random Forest model (RFM) is for each node in each tree, RFM keeps only the mean of the observations that fall into this node and neglects all other information whereas QRF keeps the value of all observations in this node, and assesses the conditional distribution based on the information (Meinshausen, 2006; Vaysse & Lagacherie, 2017; Dharumarajan et al., 2019a). For the present study, ranger package was used for running the QRF algorithm in R environment. Ranger package helps to identify the best RF properties for running the model. Ten folds cross validation techniques with 20 times repetition was used to evaluate the performance of QRF model. The performance of QRF was evaluated using indicators such as Coefficient of determination (R^2), Root Mean Square Error (RMSE), mean error (ME). Prediction interval coverage percentage (PICP) was used to evaluate the uncertainty of prediction.

133 3. Results and Discussion

134 3.1 Summary statistics of soil properties

135 Summary of the soil properties are presented in Table. 2. The soil pH ranged from 4.7
136 to 9.9 with a mean and standard deviation of 8.2 and 1.1, respectively. The organic carbon
137 content ranged between 0.11 and 1.16% with mean of 0.5 % and standard deviation of 0.23%.
138 The organic carbon skewed positively whereas pH skewed negatively showed that
139 asymmetrical distribution. The higher variability in pH is mainly attributed to soil
140 pedological factors and land management. The soil hydraulic properties such as field capacity
141 and permanent wilting point were ranged from 6 to 60% and 1.5 to 43.7% with mean and
142 standard deviation of 29.2, 18.5 and 12.2, 10.5 % respectively. Cation exchange capacity of
143 the soil varied from 2.0 to 80.9 cmol(+) kg⁻¹ with mean and SD of 32.3 and 20.5 cmol(+) kg⁻¹
144 respectively. Except pH and silt content, all other soil properties had registered negative
145 kurtosis. Similar way except, clay content and pH, all other properties showed positive
146 skewness. The correlation analysis showed that field capacity and permanent wilting point
147 has significant positive correlation with clay and silt and negative correlation with sand
148 content.

149 **Table2. Statistical results of soil properties**

Properties	Mean	Min	Max	Std dev.	Skewness	Kurtosis
pH	8.2	4.7	9.9	1.1	-1.2	0.9
OC (%)	0.5	0.11	1.16	0.23	0.5	-0.35
Clay(%)	42.6	4.1	75.8	18.2	-0.1	-0.8
Sand(%)	41.3	8.7	87.6	22	0.3	-1
Silt(%)	16.1	4.7	40.7	6.9	0.6	0.3
CEC (C mol(+) kg ⁻¹)	32.3	2	80.9	20.5	0.3	-1.1
FC(%)	29.2	6	60	12.2	0.3	-0.6
PWP(%)	18.5	1.5	43.7	10.5	0.5	-0.6

150

151

152

153 ***3.2 Performance of Quantile Regression Forest Model in predicting soil properties***

154 The performance of Quantile Regression Forest model was evaluated by calculating
155 statistical indicators viz., Coefficient of determination (R^2), Mean error (ME) and Root Mean
156 Square Error (RMSE). The cross validation results (Table 3 and Fig.2a-h) showed that the
157 combination of different covariates explained the variability's of predicted soil properties
158 viz., pH, organic carbon, CEC, clay, sand, silt, FC and PWP. The model could capture low to
159 medium variability ($R^2=0-56\%$) while predicting pH, Organic carbon and CEC for different
160 depth ranges. Among these soil properties, CEC prediction was good compared to pH and
161 Organic carbon. The present model explained 31-56 % of variation for prediction of CEC in
162 different depth intervals. Similar results were observed by different researchers (Gallo et al.,
163 2018, $R^2=40\%$; Chagas et al., 2018, $R^2=47\%$; Ghaemi et al., 2013, $R^2=45-65\%$). In case of
164 pH, only 8-23 % of variability was captured by the model. The poor prediction may be
165 attributed to more variability in pH influenced by soil intrinsic (pedogenic) and extrinsic
166 (land management) factors. Like, pH, the performance of the model for prediction of
167 organic carbon is also very low ($R^2=0-27\%$). The poor performance may be related to the low
168 levels of soil organic carbon compared to soils having high organic carbon (Lo seen et al.,
169 2010; Carvalho junior et al., 2014; Gastaldi et al., 2012; Dharumarajan et al., 2017;
170 Dharumarajan et al., 2019a). The prediction of particle size quantities viz., clay, sand and silt
171 content were fairly good. Prediction accuracy for sand is 41-49 % with RMSE of 15.4-
172 17.9%. R^2 of silt varied from 29 to 49 % for different depth intervals. Similar results were
173 observed by Akpa et al. (2014) who recorded R^2 value of 16-56 % for prediction of particle
174 size fractions in Nigeria using RFM whereas Santra et al. (2017) found only 21-28 % of
175 variation in sand content captured by Random forest algorithm.

176

177

178 **Table. 3 Performance of Quantile Regression Forest model for prediction of soil**
 179 **properties**

		Mean error	RMSE	R²(%)	PICP
pH	0-5 cm	-0.19± 0.02	0.96 ± 0.02	10± 5	88.7± 1.3
	5-15 cm	-0.18± 0.02	0.94± 0.03	13± 6	89.0 ± 0.78
	15-30 cm	-0.18± 0.02	0.95± 0.03	8± 9	90.0± 1.1
	30-60cm	-0.14± 0.03	1.00± 0.03	9± 6	88.3 ± 2.0
	60-100 cm	-0.3± 0.02	1.02± 0.02	23±3	89.3 ±2.1
	100-200cm	-0.23± 0.01	0.88± 0.03	4±8	86.4 ± 3.2
OC (%)	0-5 cm	-0.02 ± 0.0	0.22 ± 0.01	08 ± 6	87.5 ± 1.7
	5-15 cm	-0.02± 0.01	0.22± 0.01	07 ± 5	86.2 ± 1.9
	15-30 cm	0.0 ± 0.00	0.21 ± 0.01	10 ± 4	86.8 ± 2.3
	30-60cm	0.02 ± 0.01	0.20± 0.01	27± 4	88.6 ± 1.5
	60-100 cm	0.03± 0.01	0.19± 0.00	0± 2	89.4 ± 1.7
	100-200cm	0.03± 0.00	0.20± 0.00	5± 3	84.8 ± 2.1
Clay (%)	0-5 cm	0.19 ± 0.2	6.10 ± 0.29	37 ± 6	88.4 ± 2.3
	5-15 cm	2 0.22 ± 0.17	6.04 ± 0.22	39 ± 5	88.3 ± 2.0
	15-30 cm	0.59 ± 0.16	6.09 ± 0.20	39 ± 4	88.2 ± 1.2
	30-60cm	-0.04 ± 0.51	12.39 ± 0.42	43 ± 4	87.2 ± 2.3
	60-100 cm	0.0 ± 0.09	4.95 ± 0.1	18 ± 3	88.3 ±2.1
	100-200cm	0.05 ± 0.21	5.2 ± 0.22	0 ± 8	83.2 ± 1.6
Sand (%)	0-5 cm	1.54 ± 0.56	17.24 ± 0.59	48 ± 4	86.6 ± 1.7
	5-15 cm	2.25 ± 0.56	16.92 ± 0.72	49 ±4	87.6 ± 2.0
	15-30 cm	1.37 ± 0.44	16.10 ± 0.34	45 ± 2	85.3 ± 1.4
	30-60cm	40.04 ± 0.45	17.86 ± 0.43	42 ± 3	92.2 ± 1.2
	60-100 cm	-0.24 ± 0.52	15.43 ± 1.18	45 ± 9	93.6 ± 1.3
	100-200cm	0.1 ± 0.92	16.66 ± 0.94	41± 7	88 ± 0.0
Silt (%)	0-5 cm	-0.20 ± 0.39	13.47 ± 0.5	49 ± 4	91.8 ± 1.5
	5-15 cm	0.05 ± 0.32	13.90 ± 0.44	45 ± 4	90.8 ± 1.6
	15-30 cm	-0.08 ± 0.31	13.45 ± 0.40	40 ± 4	89.5 ± 1.2
	30-60cm	0.57± 0.17	6.30 ± 0.21	29 ± 5	84.0 ± 1.8
	60-100 cm	0.53 ± 0.49	11.78 ± 0.58	52± 5	90.8 ± 1.3
	100-200cm	0.57± 0.17	6.30 ± 0.21	29 ± 5	84.0 ± 1.8
CEC (C mol kg ⁻¹)	0-5 cm	1.26 ± 0.27	13.48 ± 0.66	51 ± 5	87.6 ± 1.2
	5-15 cm	1.19 ± 0.51	13.47 ± 0.42	51 ± 3	86.9 ± 1.5
	15-30 cm	1.03 ± 0.38	12.27 ± 0.37	56 ± 3	87.6 ± 1.9
	30-60cm	1.19 ± 0.67	14.88 ± 0.55	43 ± 4	87.1 ± 2.3
	60-100 cm	1.52 ± 0.61	15.98 ± 0.66	40 ± 5	88.8 ± 2.3
	100-200cm	1.55 ± 0.69	17.1 ± 0.71	31 ±6	87.1 ± 2.3
FC (%)	0-5 cm	0.03 ± 0.25	8.42 ± 0.29	36 ± 4	88 ± 2.9
	5-15 cm	0.12 ± 0.28	8.69 ± 0.21	30 ± 3	88.0 ± 2.5

	15-30 cm	-0.04 ± 0.2	7.79 ± 0.15	37 ± 3	86.0 ± 1.6
	30-60cm	0.26 ± 0.16	8.49 ± 0.15	38 ± 2	85.7 ± 2.1
	60-100 cm	0.66 ± 0.37	10.1 ± 0.41	38 ± 5	89.6 ± 1.5
	100-200cm	-0.29 ± 0.31	10.7 ± 0.41	40 ± 5	84.6 ± 2.4
PWP (%)	0-5 cm	0.26 ± 0.21	6.8 ± 0.2	41 ± 3	92.0 ± 1.5
	5-15 cm	0.3 ± 0.21	6.74 ± 0.25	41 ± 4	91.9 ± 1.4
	15-30 cm	0.3 ± 0.16	6.31 ± 0.13	43 ± 2	89.9 ± 1.7
	30-60cm	0.50 ± 0.17	6.91 ± 0.16	47 ± 3	91.6 ± 2.1
	60-100 cm	0.79 ± 0.19	8.62 ± 0.39	42 ± 5	90.8 ± 1.8
	100-200cm	0.84 ± 0.37	8.89 ± 0.53	49 ± 6	90.6 ± 2.7

180

181 Soil hydraulic properties are important for irrigation scheduling and proper landuse
182 planning (Dharumarajan et al., 2019b). Soil hydraulic properties such as field capacity and
183 permanent wilting point determines the availability and retention of the water for crop
184 growth. Field capacity and permanent wilting point were well predicted by QRF model.
185 Compared to field capacity ($R^2=30-38\%$), permanent wilting point was predicted with high
186 accuracy ($R^2=41-49\%$). Hong et al. (2013) recorded digital soil mapping approach for
187 prediction of soil hydraulic property with maximum accuracy ($R^2=61\%$) whereas Román
188 Dobarco (2019) reported prediction accuracy (R^2) of FC and PWP were 21 and 29 %
189 respectively.

190 Prediction interval coverage probability (PICP) is an indication of efficiency of
191 uncertainty measurements. The present predictions found that the PICP values ranged from
192 83.2 to 92.2 %. Overall, the prediction performance of this model was high for soil hydraulic
193 properties. Higher sample density is required for better results in tropical countries where
194 soil pattern is complex due to the geological uplift than other regions (Carvalho junior et al.,
195 2014).

196

197

198 **3.3. Importance of predictor variables for predicting soil properties**

199 RFM model estimates the importance of covariates based on how best or worse the prediction
 200 would be if one or more variable is removed and also it protects elimination of good predictor
 201 variables which are important for the model (Prasad et al. 2006). Figures 3a-h shows the
 202 variable importance rankings of Random Forest model for pH, OC, clay, sand, silt, CEC, FC
 203 and PWP. Elevation is emerged as top predictor for prediction of clay and organic carbon.
 204 MRVBF and TWI are ranked as most important predictor for prediction of pH and PWP.
 205 Different bands of Sentinel -2 imagery occupies in the top position for prediction of majority
 206 of soil properties. Different researchers recorded usefulness of Sentinel-2 imageries in
 207 prediction of different soil properties (Castaldi et al., 2019; Gholizadeh et al., 2018; Vaudour
 208 et al., 2019). Recently, Gomez et al. (2019) showed good discrimination ability of time series
 209 Sentinel-2 images in identifying different texture class and associated uncertainty.

211 **4. Spatial prediction of soil properties**

212 **Table. 4. Summary statistics of predicted soil properties**

		Mean	Min	Max	stdev	kurtosis	skewness
pH	0-5 cm	8.1	5.5	9.2	0.6	2.2	-1.5
	5-15 cm	8.2	5.9	9.2	0.4	1.8	-0.8
	15-30 cm	8.2	6.0	9.2	0.3	5.7	-1.6
	30-60cm	8.2	6.4	9.3	0.4	3.8	-0.9
	60-100 cm	8.5	6.5	9.2	0.3	5.3	-1.5
	100-200cm	8.6	8.0	9.1	0.2	1.7	-1.4
OC (%)	0-5 cm	0.61	0.35	0.83	0.08	0.9	1.3
	5-15 cm	0.60	0.40	0.84	0.07	3.0	1.7
	15-30 cm	0.59	0.37	0.83	0.10	-0.5	0.5
	30-60cm	0.59	0.28	0.80	0.11	-0.7	0.0
	60-100 cm	0.4	0.3	0.6	0.1	-0.9	-0.1
	100-200cm	0.4	0.3	0.6	0.0	9.9	0.9
Clay (%)	0-5 cm	16.2	5.5	32.6	5.1	-1.0	0.0
	5-15 cm	36.6	7.7	62.1	11.3	-0.7	-0.2

	15-30 cm	39.9	7.3	64.1	8.6	0.0	0.2
	30-60cm	45.7	8.5	66.5	8.6	0.1	0.6
	60-100 cm	15.2	8.0	19.0	2.4	-1.3	-0.2
	100-200cm	17.2	12.1	21.7	1.7	1.6	-0.5
Sand (%)	0-5 cm	45.5	17.8	82.6	15.4	-0.7	0.7
	5-15 cm	45.3	17.7	82.9	15.0	-0.8	0.7
	15-30 cm	41.7	15.5	76.5	13.5	-1.0	0.5
	30-60cm	36.1	13.4	80.5	11.9	-0.6	0.3
	60-100 cm	38.4	12.4	57.8	13.9	-1.3	-0.3
	100-200cm	38.0	11.2	62.6	15.6	-1.4	-0.4
Silt (%)	0-5 cm	36.5	7.7	61.9	12.0	-0.8	-0.2
	5-15 cm	16.7	6.2	32.6	5.6	-0.7	0.2
	15-30 cm	17.5	7.1	35.6	6.4	-1.0	0.3
	30-60cm	15.8	8.7	27.5	4.2	-0.4	0.3
	60-100 cm	46.9	20.2	69.0	10.7	-1.2	0.4
	100-200cm	45.0	24.1	71.4	12.9	-1.5	0.3
CEC (C mol kg ⁻¹)	0-5 cm	27.5	4.7	62.3	14.0	-1.3	0.1
	5-15 cm	27.5	5.2	62.3	14.5	-1.3	0.2
	15-30 cm	29.9	5.0	64.0	14.6	-1.3	0.2
	30-60cm	32.9	7.0	62.3	13.3	-1.2	0.1
	60-100 cm	32.4	10.6	65.3	14.9	-1.6	0.1
	100-200cm	31.0	13.9	54.5	14.7	-1.5	0.3
FC (%)	0-5 cm	24.9	9.0	41.0	6.8	-0.5	-0.7
	5-15 cm	25.6	10.1	39.0	6.3	-0.4	-0.6
	15-30 cm	26.7	12.1	39.9	5.7	-0.8	-0.2
	30-60cm	29.3	10.4	46.8	5.9	-1.1	0.4
	60-100 cm	32.2	18.8	45.7	7.9	-1.5	-0.2
	100-200cm	30.5	15.2	50.4	8.3	-1.4	0.3
PWP (%)	0-5 cm	14.6	3.2	29.5	5.7	-0.9	0.1
	5-15 cm	15.1	3.5	27.3	5.2	-0.9	0.3
	15-30 cm	16.3	4.8	27.9	4.7	-1.1	0.3
	30-60cm	18.5	2.9	36.1	5.7	-1.0	0.3
	60-100 cm	19.8	9.0	37.5	7.0	-1.3	0.1
	100-200cm	19.8	7.8	39.1	8.0	-1.5	-0.1

213

214 Mapping of soil properties is a preliminary step due its variability for decision making
215 such as the delineation of suitable crop growing areas or identification of degraded areas.
216 Summary statistics of predicted soil properties are presented in Table.4. Predicted maps of
217 sand, silt, clay, CEC, FC and PWP in the surface (0-5 cm) along with uncertainty using
218 Quantile Regression Forest are presented in Fig.4&5. The predicted sand content in 0-5 cm
219 varied from 17.8-82.6%. The predicted silt and clay content varied from 7.7-61.9 % and 5.5-

220 32.6 % respectively. High sand content recorded in North-eastern part of study area and high
221 clay and silt content recorded in north-western part. The high sand content of surface soils in
222 North-eastern part might be due to severity of the erosion where finer particles are moved
223 into the low lying areas. The predicted cation exchange capacity varied from 4.7 – 62.3 %
224 and recorded low CEC in north-eastern part. Predicted hydraulic properties viz., field
225 capacity and permanent wilting point ranged from 9.1-41 % and 3.2-29.5% respectively. The
226 spatial prediction of soil properties suggested that distribution of soil properties on the
227 surface are highly variable due to variations in environmental factors, land management and
228 land use. The spatial resolution of the maps helps to assess and monitor the soil health and
229 preparation of proper land use plan.

230 **4. Conclusion**

231 The prediction of soil properties and uncertainty by QRF model was reasonable and
232 varied from 8-51% for surface and 0-56% for subsoil. Except pH and OC, the present model
233 predicted better for most of the soil properties compared to previous studies. Weak variations
234 in soil properties, mixed lithologic occurrence and sparse sample density are linked with
235 performance of the model. The data augmentation certainly helps in reducing the uncertainty
236 and over fitting and to improve model accuracy further. The prediction can also be
237 improvised by increasing the environment covariates such as geology map and climatic
238 datasets.

239 **Acknowledgements**

240 The funding received from the project ATCHA, ANR-16-CE03-0006 to attend and present
241 this paper in the 2019 digital soil mapping and GlobalSoilMap workshop held in The
242 University of Chile, Santiago is highly acknowledged. The authors also acknowledge Dr.

243 Laurent Ruiz, Indo-French Cell for Water Sciences, Bangalore for his guidance in Indian
244 digital soil mapping programme.

245 **Reference**

246 Adhikari, K., Minasny, B., Greve, M.B., Greve, M.H., 2014. Constructing a soil class map of
247 Denmark based on the FAO legend using digital techniques. *Geoderma* 214–215, 101–113.

248 Akpa, S.I.C., Odeh, I.O.A., Bishop, T.F.A., Hartemink, A.E., 2014. Digital mapping of soil
249 particle-size fractions for Nigeria. *Soil Sci Soc Am J.* 78, 1953–1966.

250 Arrouays, D., Grundy, M. G., Hartemink, A. E., Hempel, J. W., Heuvelink, G.B.M.,
251 Hong, S.Y., Lagacherie, P., Lelyk, G., McBratney, A. B., McKenzie, N. J.,
252 Mendonça-Santos, M.D., Minasny, B., Montanarella, L., Odeh, I. O. A., Sanchez, P. A.,
253 Thompson, J. A., Zhang, G. L., 2014. GlobalSoilMap: towards a fine-resolution global grid
254 of soil properties. *Advances in Agronomy* 125, 93-134.

255 Bishop, T.F.A., McBratney, A.B., Laslett, G.M., 1999. Modelling soil attribute depth
256 functions with equal-area quadratic smoothing splines. A comparison of prediction methods
257 for the creation of field-extent soil property maps. *Geoderma* 91, 27-45.

258 Gallo, B.C., José, A. M., Demattê, Rodnei Rizzo, José L. Safanelli, Wanderson de S. Mendes,
259 Igo F. Lepsch, Marcus V. Sato, Danilo J. Romero, Marilusa P. C. Lacerda, 2018. Multi-
260 Temporal Satellite Images on Topsoil Attribute Quantification and the Relationship with Soil
261 Classes and Geology. *Remote Sens.* 10, 1571. doi:10.3390/rs10101571

262 Carvalho Junior W de, Philippe Lagacherie , Cesar da Silva Chagas , BrazCalderano Filho,
263 Silvio Barge Bhering, 2014. A regional-scale assessment of digital mapping of soil attributes
264 in a tropical hillslope environment. *Geoderma* 232–234, 479–486.

265 Castaldi, F., Hueni, A., Chabrillat, S., Ward, K., Buttafuoco, G., Bomans, B., Vreys, K.,
266 Brell, M., van Wesemael, B., 2019. Evaluating the capability of the Sentinel 2 data for soil
267 organic carbon prediction in croplands. *ISPRS J. Photogramm. Remote Sens.* 147, 267–282.

268 Chagas, C.S., Carvalho Júnior, W., Pinheiro, H.S.K., Xavier, P.A.M., Bhering, S.B., Pereira,
269 N.R., Calderano Filho, B., 2018. Mapping soil cation exchange capacity in a semiarid region
270 through predictive models and covariates from remote sensing data. *Rev Bras Cienc Solo.*
271 2018;42:e0170183. <https://dx.doi.org/10.1590/18069657rbc20170183>

272 Dharumarajan, S., Rajendra Hegde, Janani, N., Singh, S.K., 2019a. The need for digital soil
273 mapping in India, *Geoderma Regional* 15. <https://doi.org/10.1016/j.geodrs.2019.e00204>.

274 Dharumarajan, S., Rajendra Hegde, Lalitha, M., Kalaiselvi, B., Singh, S. K., 2019b.
275 Pedotransfer functions for predicting soil hydraulic properties in semi-arid regions of
276 Karnataka Plateau, India. *Current Science* 116(7),1237-1246.

277 Dharumarajan, S., Rajendra Hegde, Singh, S.K., 2017. Spatial prediction of major soil
278 properties using Random Forest techniques - A case study in semi-arid tropics of South
279 India. *Geoderma Regional* 10C, 154-162. doi: 10.1016/j.geodrs.2017.07.005

280 Gastaldi, G., Minasny, B., McBratney, A.B., 2012. Mapping the occurrence and thickness of
281 soil horizons within soil profiles. In: Minasny, B., Malone, B.P., McBratney, A.B. (Eds.),
282 *Digital Soil Assessments and Beyond*. CRC Press/Balkema, London, pp. 145–148.

283 Gessler, P.E., Chadwick, O.A., Chamran, F., Althouse, L., Holmes, K., 2000. Modeling soil-
284 landscape and ecosystem properties using terrain attributes. *Soil Science Society of America*
285 *Journal* 64, 2046-2056.

286 Ghaemi. M., Astaraei. A.R., Sanaeinejad, S.H., Zare, H., 2013. Using satellite data for soil
287 cation exchange capacity studies. *Int. Agrophys.* 27, 409-417. doi: 10.2478/intag-2013-0011.

288 Gholizadeh, A., Daniel Žižala, Mohammadmehdi Saberioon, LubošBorůvka, 2018. Soil
289 organic carbon and texture retrieving and mapping using proximal, airborne and Sentinel-2
290 spectral imaging. *Remote Sensing of Environment* 218, 89–103.

291 Gomez, C., Dharumarajan, S., Féret, J.B., Lagacherie, P., Ruiz, L., Sekhar, M., 2019. Use of
292 Sentinel-2 Time-Series Images for Classification and Uncertainty Analysis of Inherent
293 Biophysical Property: Case of Soil Texture Mapping. *Remote Sens.* 11, 565.

294 Grundy, M.J., Viscarra Rossel, R.A., Searle, R.D., Wilson, P.L., Chen, C., Gregory, L.J.,
295 2015. Soil and landscape grid of Australia. *Soil Res.* 53, 835–844.

296 Grunwald, S., Thompson, J. A., Boettinger, J. L., 2011. Digital soil mapping and modeling at
297 continental scales: Finding solutions for global issues. *Soil Sci. Soc. Am. J.* 75, 1201–1213.

298 Hong, S.Y., Budiman Minasny, Kyung Hwa Han, Yihyun Kim and Kyungdo Lee, 2013.
299 Predicting and mapping soil available water capacity in Korea. *Peer J.* 1,e71; DOI
300 10.7717/peerj.71

301 Jackson, M.L.,1973. *Soil Chemical Analysis*. Prentice Hall of India Pvt. Ltd. New Delhi.

302 Jackson, M. L., 1962. *Soil chemical analysis*. Prentice Hall. Englewood Cliffs, New York,
303 U.S.A

304 Lagacherie, P., McBratney, A.B., 2007. Spatial soil information systems and spatial Soil
305 inference systems: perspectives for digital soil mapping. In: Lagacherie, P., McBratney, A.B.,
306 McBratney, A.B., Mendonça-Santos, M.L., Minasny, B., 2003. On digital soil mapping.
307 *Geoderma* 117,3–52. doi:10.1016/S0016-7061(03)00223-4.

308 Lo Seen, D., Ramesh, B. R., Nair, K. M., Martin, M., Arrouays, D., Bourgeon, G.,2010. Soil
309 carbon stocks, deforestation and land-cover changes in the Western Ghats biodiversity
310 hotspot (India). *Global Change Biology* 16, 1777-1792. doi:10.1111/j.1365-
311 2486.2009.02127.x

312 Meinshausen, N., 2006. Quantile regression forests. *J. Mach. Learn. Res.* 7, 983–999.

313 Mulder, V.L., Lacoste, M., Richer de Forges, A.C., Martin, M.P., Arrouays, D., 2016.
314 National versus global modelling the 3D distribution of soil organic carbon in mainland
315 France. *Geoderma* 263, 13-34.

316 Poggio, L., Gimona, A., 2017. 3D mapping of soil texture in Scotland, *Geoderma Regional*
317 9, 5-16. <https://doi.org/10.1016/j.geodrs.2016.11.003>, 2017.

318 Prasad, A. M., Iverson, L. R., Liaw. A., 2006. Newer classification and regression tree
319 techniques: bagging and random forests for ecological prediction. *Ecosystems* 9, 181–199.

320 Richards, L.A., 1954. Diagnosis and improvement of saline and alkali soils. USDA, Hand
321 Book No, 60:101.

322 Richards, L.A., Gardner, W.R., Gen Ogata, 1956. Physical processes determining water loss
323 from soil. *Soil Sci. Soc. Am. Proc.* 20,310-314.

324 Román Dobarco, M., Bourennane, H., Arrouays, D., Saby, N. P.A, Cousin, I., Martin, M.
325 P.,2019. Uncertainty assessment of GlobalSoilMap soil available water capacity products: A
326 French case study. *Geoderma* 344, 14-30. doi: 10.1016/j.geoderma.2019.02.036

327 Sanchez, P.A., Ahamed, S., Carr´e, F., Hartemink, A.E., Hempel, J., Huising, J., Lagacherie,
328 P., McBratney, A.B., McKenzie, N.J., De Lourdes Mendonc, Santos, M., Minasny, B.,
329 Montanarella, L., Okoth, P., Palm, C.A., Sachs, J.D., Shepherd, K.D., Vagen, T.G.,

330 Vanlauwe, B., Walsh, M.G., Winowiecki, L.A., Zhang, G.L., 2009. Digital soil map of the
331 world. *Science* 325,680–681. DOI 10.1126/science.1175084.

332 Santra, P., Kumar, M., Panwar, N., 2017. Digital soil mapping of sand content in arid western
333 India through geostatistical approaches. *Geoderma Reg.* 9, 56–72.

334 Vaudour, E., Gomez, C., Fouad, Y., Lagacherie, P., 2019. Sentinel-2 image capacities to
335 predict common topsoil properties of temperate and Mediterranean agroecosystems. *Remote*
336 *Sens. Environ.* 223, 21–33.

337 Vaysse, K., Lagacherie, P., 2017. Using quantile regression forest to estimate uncertainty of
338 digital soil mapping products. *Geoderma* 291, 55–64.
339 <https://doi.org/10.1016/j.geoderma.2016.12.017>

340 Walkley, A., Black, I.A., 1934. An estimation of the method for determining soil organic
341 matter and a proposed modification of the chromic acid titration method. *Soil science* 37, 29-
342 38.

343

344




Fig. 2(a-h). Observed soil properties Vs Predicted soil properties in 0-5 cm depth




Fig.3(a-h). Variation importance ranking of Random forest model in prediction of different soil properties




Fig.4. Predicted sand, silt, clay and CEC content in 0-5 cm depth




Fig.5. Predicted Organic carbon content, Field capacity and permanent wilting point in 0-5 cm depth