



Early life solid food ingestion remodels the gut microbiota composition and increases the production of protective bacterial metabolites

Martin Beaumont, Charlotte Paes, Eloïse Mussard, Laurent Cauquil, Patrick Aymard, Céline Barily, Olivier Bouchez, Cécile Canlet, Sylvie Combes

► To cite this version:

Martin Beaumont, Charlotte Paes, Eloïse Mussard, Laurent Cauquil, Patrick Aymard, et al.. Early life solid food ingestion remodels the gut microbiota composition and increases the production of protective bacterial metabolites. digestive disease week 2019, May 2019, San diego, United States.
hal-02914647

HAL Id: hal-02914647

<https://hal.inrae.fr/hal-02914647>

Submitted on 12 Aug 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EARLY LIFE SOLID FOOD INGESTION REMODELS THE GUT MICROBIOTA COMPOSITION AND INCREASES THE PRODUCTION OF PROTECTIVE BACTERIAL METABOLITES

Martin Beaumont¹, Charlotte Paës^{1,2}, Eloïse Mussard¹, Laurent Cauquil¹, Patrick Aymard¹, Céline Barily¹, Olivier Bouchez³, Cécile Canlet⁴, Sylvie Combes¹.



1 - GenPhySE, Université de Toulouse, INRA, INPT, ENVT, Castanet-Tolosan, France; 2 - GEC consortium CCPA, Eivalis, Inzo, MixScience, Techna; 3 - GeT-PlaGe, Genotoul, INRA, Castanet-Tolosan, France; 4 - Toxalim (Research Centre in Food Toxicology), Université de Toulouse, INRA, ENVT, INP-Purpan, UPS, Toulouse, France.

Background




Methods


Neonatal rabbit model



1- Microbiota composition




The microbiota composition shifted after the onset of solid food ingestion




The relative abundance of families including pathobionts decreased after the onset of solid food ingestion while the abundance of a butyrate producing family increased

2- Bacterial metabolites




Short chain fatty acids concentrations increased after the onset of solid food





3-phenylpropionate concentration increased after the onset of solid food while trimethylamine concentration decreased

3- Epithelial barrier *in vitro*



The mixture of metabolites produced by the microbiota after the onset of solid food promoted epithelial barrier formation



Butyrate and 3-phenylpropionate promoted epithelial barrier formation while trimethylamine had no effect

Conclusion

- The microbiota composition and its metabolic activity shifted after the onset of solid food ingestion
- Since milk intake was still predominant at 25 days, our results suggest that the introduction of solid food (*versus* weaning) is sufficient to induce microbiota maturation
- Metabolites produced by the gut microbiota after the onset of solid food ingestion might contribute to the postnatal maturation of the epithelial barrier