

Combining partial cutting and direct seeding to overcome regeneration failures in dune forests

Arthur Guignabert, Laurent Augusto, Florian Delerue, Francis Maugard, Céline Gire, Clément Magnin, Sylvie Niollet, Maya Gonzalez

▶ To cite this version:

Arthur Guignabert, Laurent Augusto, Florian Delerue, Francis Maugard, Céline Gire, et al.. Combining partial cutting and direct seeding to overcome regeneration failures in dune forests. Forest Ecology and Management, 2020, 476, 10.1016/j.foreco.2020.118466 . hal-02915058

HAL Id: hal-02915058 https://hal.inrae.fr/hal-02915058

Submitted on 22 Aug2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Version of Record: https://www.sciencedirect.com/science/article/pii/S0378112720312354 Manuscript_04afd504b79388fcbb610f52480d59ca

1 Title:

2 Combining partial cutting and direct seeding to overcome regeneration failures in dune forests.

3

4 Authors:

5 Arthur Guignabert^{a*}, Laurent Augusto^a, Florian Delerue^{b,c}, Francis Maugard^d, Céline Gire^a, Clément

- 6 Magnin^d, Sylvie Niollet^a, Maya Gonzalez^a.
- 7
- ^a UMR 1391 ISPA, Bordeaux Sciences Agro, INRAE, 33140 Villenave d'Ornon, France.
- 9 ^b EA 4592, G&E, Bordeaux INP, 33600 Pessac, France.
- [°] EA 4592, G&E, Université Bordeaux Montaigne, 33600 Pessac, France.
- ^d ONF, 9, rue Raymond Manaud, 33524 Bruges, France.
- 12 * Correspondence: arthur.guignabert@gmail.com
- 13

14 Abstract:

15 Stand regeneration is a crucial step in the management of many forests and its failure can 16 jeopardize future forest growth and production. Thus, adapting forestry practices to improve 17 seedling establishment is of prime importance to maintain sustainable forest management. In the 18 coastal dune forests of maritime pine in SW France, regeneration failure after clear-cutting has 19 greatly increased in the last decades. The aim of this study was to quantify the different stages 20 involved in the regeneration process (seed rain, germination, survival), and to assess the impact of 21 harvesting methods (partial cutting vs. clear-cutting) and of the use of direct seeding (seeding vs. no 22 seeding) on these stages. We established five trials located in areas with contrasting regeneration statuses (of which two sites were in an area characterized by chronic regeneration failure), and we 23 24 investigated the effect of the harvesting method and the use of direct seeding in a factorial design. 25 We monitored the seed rain, germination and first-year survival for three years. Due to the transient 26 nature of the seed bank, we found that the seed rain of the previous year was the only possible seed source for tree regeneration, and it increased with the proximity and size of surrounding 27 28 mature trees. Nonetheless, seed rain did not limit regeneration in these stands. In fact, germination 29 was the bottleneck stage of the regeneration process in all sites because it was short-lived and 30 consequently exposed to failure in case of unfavourable conditions. Once established, seedling death was mainly due to summer drought. Clear-cutting without seeding resulted in severe 31 32 regeneration failures in 4 out of 5 sites, with the most severe in the failure area. Direct seeding 33 increased seedling emergence only in the first year, while partial cutting had a longer-lasting effect 34 by providing a regular seed supply over several years. Partial cutting also decreased biotic and 35 abiotic stresses through microhabitat modification under the tree canopy, resulting in a higher 36 number of germinations and greater seedling survival. This positive effect was more pronounced in 37 the two sites within the failure area, suggesting that conditions were more stressful in this part of the forests. Consequently, we recommend avoiding clear-cutting in favour of partial cutting in all parts of 38 39 these forests, and ensuring the maintenance of sufficient forest cover to promote regeneration. 40 Direct seeding could be used in addition to partial cutting to maximize the chances of success, but 41 only in areas where natural regeneration is low.

42

Key words: Direct seeding, Drought, Forest management, Germination, Partial cutting, *Pinus pinaster*, Seed availability, Seedling survival, Tree regeneration.

- 45
- 46

47 **1. Introduction**

48 The renewal of forest stands is a key step in forest dynamics and for their sustainable 49 management. Natural regeneration is one way to do this and involves the stages of seed production, seed dispersal, germination and seedling survival, each being influenced by many 50 51 interacting biotic and abiotic factors (Price et al., 2001; Kozlowski, 2002; Calama et al., 2017). The establishment of a species results from the interactions between the quantity of available seeds and 52 53 the number of suitable microsites for regeneration (Harper, 1977; Schupp, 1995). The density of seeds reaching the ground depends on the seed production, on the spatial arrangement and the 54 55 characteristics of the seed trees, and on the seed dispersal capacity (Greene et al., 1999; Viglas et 56 al., 2013; Montoro Girona et al., 2018). Then, seedling emergence and survival depends on favourable local conditions such as soil moisture, soil temperature, or light (Harper, 1977; 57 58 Kozlowski, 2002). These ecological conditions vary widely, both spatially and over time (Beckage

and Clark, 2003; Gómez-Aparicio et al., 2005), and can be influenced by biotic interactions with
overstorey trees, herbaceous species or herbivore populations (Harmer, 2001; Wagner et al., 2011;
Lavoie et al., 2019). Consequently, regeneration failure can also depend on the local environmental
conditions.

63 Natural regeneration is frequently unsuccessful in temperate and Mediterranean managed forests (Calama et al., 2017; Dey et al., 2019), and could be increasingly threatened in the current 64 context of global change in areas where droughts are expected to increase in frequency and 65 66 intensity (IPCC, 2014). Thus, understanding the regeneration processes is very important in the adaptation of forest management practices to ensure the natural regeneration of stands. Forest 67 harvesting, which is the starting point of the regeneration cycle, can be performed in two different 68 ways in even-aged forests: i) clear-cutting, when all the trees are removed in a single harvesting 69 70 operation and the new seedlings emerge from the soil seed bank. In the most difficult cases, natural 71 regeneration can be assisted with a direct seeding using local seeds to overcome establishment 72 limitations (Grossnickle and Ivetić, 2017); ii) partial cutting, when some of the mature trees are left 73 unlogged in the stand to promote regeneration, and are harvested few years later, once 74 regeneration is sufficient (Nyland, 2016). The degree of canopy closure induced by partial cuts can 75 potentially influence several ecological processes and is therefore a key factor that can affect 76 regeneration. First, the remaining trees can supply the viable soil seed bank for a few years to offset 77 low seed availability for regeneration (Nyland, 2016). Second, forest cover can create more suitable 78 conditions for seedling establishment by buffering microclimatic stresses (e.g. vapour pressure 79 deficit, soil moisture or solar radiation) (Aussenac, 2000; Heithecker and Halpern, 2006). Partial 80 cuts also modify light availability at the ground level and can influence the development of herbaceous and understorey species that may limit seedling establishment by competing for 81 82 resources (Löf, 2000; Wagner et al., 2011). Finally, this habitat modification may also have an 83 indirect impact on the behaviour of herbivores and therefore on the potential damage caused to seedlings (Reimoser and Gossow, 1996; Côté et al., 2004). In addition, partial cutting, which is 84 85 increasingly used to ensure the sustainable management of many forests, can have other effects 86 than promoting natural regeneration. Partial cutting also promote both economic and ecological 87 features of mature stands, by increasing the radial growth of residual trees following the decrease in

stand density (Montoro Girona et al., 2016), and by supporting higher richness and greater abundance of flora and fauna than clear-cuts (Fedrowitz et al., 2014). However, these positive effects could be reduced in case of mortality of residual trees, as partially cut stands tend to be more sensitive to disturbance (particularly windthrow) than unharvested stands (Montoro Girona et al., 2019).

93 Maritime pine (Pinus pinaster Aiton) is a widely distributed conifer in the Mediterranean 94 basin and along the Atlantic coast of Western Europe, and is of high ecological, economic and 95 social importance in its native area (Alía and Martín, 2003) and in countries where it has been 96 introduced (mainly Australia, New-Zealand, and South Africa). The coastal forests of maritime pine 97 in southwestern France cover about 100 000 ha (IFN, 2017). Natural regeneration following clear-98 cutting has been used traditionally for many decades for the renewal of these forests to fulfil their 99 multifunctional role (i.e. wood production, soil erosion protection, preservation of biodiversity, 100 recreation usage and aesthetic value) and to improve the conservation of genetic diversity and the 101 capacity to adapt to global change. However, regeneration failures (i.e. less than 3000 seedlings ha-102 ¹ – or 0.3 m⁻² – three years after harvest (Sardin, 2009)) have steadily increased in recent years, 103 and persisted in spite of an additional supply of local seeds in some stands at the harvesting period. 104 These failures were mainly observed in the central part of the coastal fringe occupied by dune 105 forests (up to half of the regenerating stands may fail some years (Ouallet, 2012)), but this is 106 beginning to spread to other areas, causing great concern among forest managers. In this type of 107 ecosystem with freely draining sandy soils and regular severe summer drought, partial cutting is a 108 method often used in other regions to renew pine forests, such as Spain (Calama et al., 2017). This 109 method regularly results in sufficient seedling density in maritime pine forests (e.g. Rodríguez-110 García et al., 2010), but failures are also observed (e.g. González-Alday et al., 2009). This is 111 because natural regeneration is highly dependent on site characteristics and on local climatic 112 conditions, and is also variable between pine populations (Rodríguez-García et al., 2011b; 113 Vergarechea et al., 2019).

The aim of this study was to identify the processes involved in failures of natural regeneration in the coastal forests of maritime pine, in interaction with silvicultural practices. For this purpose, we established five stands located in areas with contrasting regeneration successes, and

117 investigated the effect of the harvesting method (partial cutting compared to the usual clear-cutting) 118 and of the use of direct seeding (compared to the absence of seeding). We hypothesized that: i) 119 partial cutting could compensate for poor-regeneration years by supplying extra seeds each year, 120 and by increasing germination and seedling survival rates due to improved microclimatic conditions, 121 especially in the summer; ii) this positive effect would be more pronounced on sites within areas of 122 high regeneration failures, so that the threshold of 3,000 seedlings ha⁻¹ necessary for a successful 123 regeneration would be exceeded; and iii) direct seeding would improve the number of seedlings 124 regardless of the harvesting method, but it might be less useful than partial cutting that provides a 125 combination of seed supply and microhabitat amelioration. By improving our understanding of the 126 ecology of maritime pine regeneration, our findings should enable us to propose adequate 127 management strategies to ensure natural regeneration in these forests.

128 2. Materials and methods

129 2.1. Study sites

130 Our study combined experimentation (harvesting method and seeding as treatments, see 131 2.2.) and exploration on a regional scale, integrating a major regeneration failure area. Thus, we 132 established five trials along the coast in the maritime pine (Pinus pinaster Aït) sand dune forests of 133 SW France in areas with contrasting regeneration statuses (Fig. 1A): two of the sites were within 134 the area of chronic regeneration failure (hereafter referenced to as sites F1 and F2), while the three 135 other sites were in areas with frequent regeneration success (S1, S2 and S3). The lower number of 136 site in the failure area is explained by its geographically smaller area compared to the whole dune 137 forests (Fig. 1A). Stand area ranged from 7.4 to 11.5 ha and tree age ranged from 58 to 77 years 138 old (Table S1). The climate in the region is temperate oceanic. Average, minimum and maximum 139 annual temperatures were 13.6 °C, 8.8 °C and 18.5 °C for S1 and about 14.1 °C, 9.3 °C and 19.2 140 °C for the four other sites (Table S1). Average annual precipitation ranged from 840 mm to 1007 141 mm (Table S1), with the wettest period in winter and the driest in July-August. All sites were chosen 142 on westerly facing slopes at about 2.5 km from the ocean, with an average slope of 10°. Soils are young sandy soils (WRB classification: arenosols; USDA classification: entisols) mainly composed 143 144 of coarse sands (96–97%), are slightly acidic (topsoil values of pH = 4.5-5.0), and are extremely poor in nutrients (Augusto et al., 2010). These soils have a low water holding capacity which,
combined with low summer rainfall, results in an extremely low soil moisture content and a high
water stress for seedlings in summer (Guignabert et al., 2020).



148

Fig. 1: A) Location of the five study sites. B) Aerial photography of S3 showing the two harvesting treatments,
and the distribution of the monitoring areas. C) Schematic of one monitoring area and of the seed
arrangement pattern.

152 **2.2. Experimental design**

Two silvicultural practices were tested factorially at each site in early 2015: i) the harvesting 153 154 method (usual clear-cut vs newly tested partial cut using the uniform shelterwood system), and ii) 155 the use of direct seeding (vs no seeding). To that end, each of the two harvesting methods was 156 carried out on half of all stands (Fig. 1B). Tree harvesting was performed mechanically with a feller-157 buncher between January and March 2015. For the partial cut treatment, trees were chosen on the basis of their vigour and appearance, and tree density after harvest was ~70 stems/ha in all sites. 158 159 The basal area after partial cutting was between 7.5 and 13.2 m² ha⁻¹ (Table S1). Other 160 management practices were carried out before logging in all experimental treatments: understorey 161 vegetation was removed mechanically to limit post-logging competition and was combined with light 162 mechanical tillage to increase soil aeration and nutrient availability. Then, at each site, we 163 positioned ten monitoring areas in the centre of each harvesting treatment (Fig. 1B), each area 164 being split into four 4 m² guadrats (Fig. 1C). Three guadrats were sown with five seeds m⁻², which 165 corresponds to the quantity currently used by foresters in regeneration failure areas. Seeds were 166 sown following an identical pattern (Fig. 1C; ca. 0.5 cm deep) and came from cones collected from 167 stands of maritime pine located in the same forests. The fourth quadrat was unsown, and its 168 position was determined randomly. Finally, we installed a 1.5 m² seed trap one meter away of each 169 monitoring area in the partial cut treatment (Fig. 1C) to quantify the seed rain (10 seed traps per 170 site). Traps were wooden frames filled with high-density polyethylene mesh and with 25 cm high 171 edges in soft PVC, and were set up at about 25 cm above the ground (Fig. S1).

172 2.3. Regeneration monitoring

173 Seed rain was collected monthly in seed traps and counted from May to October in 2015, 174 2016 and 2017, then air-dried and stored at 4 °C. Seed weight and seed length were measured on 175 a random sample of 80 seeds per site of the 2015 seed rain, using a precision balance and a digital 176 calliper respectively. Since seed rain is known to be dependent on tree basal area and tree density 177 (Ruano et al., 2015b; Viglas et al., 2013), we measured the circumference of all trees within a 20 m 178 radius of each seed trap and their distance from the trap. Then, from these two variables we 179 calculated a seed source abundance index (SSAI) for each seed trap as follows: SSAI = $\sum_{i=1}^{n} (TBA_i/d_i)$ with $TBA_i(m^2) = (c_i/100)^2/4\pi$ and where c (cm) is the circumference at 1.30 m 180 181 of tree *i*, *TBA* is the basal area of tree *i* (m^2) and *d* the distance (m) of tree *i* from the seed trap. The 182 closer and bigger the tree, the more it contributes to the index value. The choice of a 20 m radius 183 circle around each seed trap was based on previous studies about the dispersal distance 184 corresponding to the majority of dispersed pine seeds (Juez et al., 2014; Ruano et al., 2015b).

Seedling emergence and first-year survival were monitored each year from 2015 to 2017. Censuses were carried out monthly from May to October in 2015 and 2016, and only three censuses were made in 2017: in late March, before the germination peak; in mid-June, after the germination peak and before the drought period; and in late September, after the drought period. Cause of seedling death was recorded: drought, when the seedling turned from green to red/brown; or due to herbivores, when the seedling died after browsing by ungulates or rodents (see Guignabert et al. (2020) for more details on the distinction). Non-lethal ungulate browsing events were also noted at each survey. For each monitoring year, the survival of seedlings that emerged in the spring was only surveyed until the end of that specific year, and not throughout the whole study. Thus, a final count of all living seedlings was performed in March 2018 to obtain an assessment of the regeneration density comparable to the density value defined by foresters as the threshold value necessary for a successful regeneration (i.e. 0.3 seedlings m⁻²; Sardin, 2009).

197 **2.4. Complementary measurements**

198 2.4.1. Soil seed bank

Soil samples were collected from all sites to assess the soil seed bank, both in the clear-cut and in the partial cut. This sampling was performed at the beginning of the experiment in April 2015, after tree harvesting and before the beginning of the seed rain. A soil sample of $1/32 \text{ m}^2$ and 5 cm deep was collected in the proximity of the four corners of each monitoring area, giving a total of 400 samples (5 sites x 2 harvesting methods x 10 monitoring areas x 4 samples) from a total surface area of 12.5 m². Samples were then sieved using a 2 mm mesh and undamaged seeds were counted and stored at 4 °C.

206 *2.4.2. Seed germinative capacity*

207 We carried out a glasshouse germination test in March/April 2017 to evaluate the 208 germinative capacity of the seeds in the five sites. This test was performed using the three pools of 209 potential seed sources: i) seed rain, using seeds collected from seed traps in summer 2016; ii) soil 210 seed bank, using seeds collected from soil sampled in April 2016 following the same procedure 211 described above; and iii) seeds used for direct seeding by the French National Forestry Office 212 (seeds ref: "PPA 303, dunes littorales de Gascogne", collected in winter 2015-16). All seeds were 213 stored at 4 °C in the same place between their collection and the test. We sowed 11 batches of 50 214 seeds in independent trays (50 x 30 cm) on March 15: five batches of soil seed bank and of seed 215 rain (one per site), and one batch of seeds used for seeding. The soil used in all trays was collected 216 from the F2 site and sieved to remove seeds potentially present in the collected soil. Trays were 217 installed in a glasshouse at ambient temperature and watered every two days. New emergence was 218 surveyed once a week for 8 weeks.

219 2.4.3. Dendrometric features of partial cuts

220 The dendrometric features of a mature stand (e.g. basal area, canopy cover) are partly 221 related to the silviculture carried out during the forest rotation (e.g. thinning). We carried out this 222 additional study in winter 2018-2019 in other forest stands to find out whether the stand structure of 223 partial cuts might be a factor that should be integrated into the management considerations to 224 improve regeneration. It should be noted that this study was not carried out at our five experimental 225 sites, but at the scale of a large management canton. This area was located within the main failure 226 area (i.e. near F1 and F2), the only area where partial cuts had been implemented in the region so 227 far. We selected forest stands which had been harvested at least 3 years before (i.e. harvested in 228 winter 2013-14, 2014-15 and 2015-16), which represented 29 stands and 280 ha (stand average = 229 9.7 ha, min = 3.4 ha, max = 19.7 ha). For each stand, we measured: i) the seedling density, by 230 counting the number of seedlings in 80 plots of 6 m² positioned according to a systematic grid 231 pattern (all were summed and then reported as seedlings m^{-2}); ii) the tree density by counting all 232 trees in the stand; iii) the basal area, estimated from the tree diameter. The diameter of each tree 233 (d, cm) was measured using a forest compensated calliper (i.e. graduated in 5 cm classes), the 234 method which is regularly used in silviculture to carry out inventories and provides a good estimate 235 of the basal area (Cordonnier et al., 2007). Stand basal area (BA) was then calculated for each stand as: $BA(m^2/ha) = (\Sigma TBA)/stand area$, with $TBA(m^2) = \pi (d/100)^2/4$; iv) the forest cover, 236 237 which was estimated using the normalised difference vegetation index (NDVI) based on analysis of 238 multispectral images of the Pléiades satellite taken on 14 April 2018 and 03 September 2018, and 239 performed with ArcGIS. This variable, obtained in m², was then expressed as a percentage value by 240 dividing the value by the stand area; and v) the individual tree crown surface area (m² tree⁻¹), by 241 dividing the forest cover (m²) by the number of trees.

242 **2.5. Data treatment and statistics**

All statistical analyses were performed with R software version 3.5.2. (R Core Team, 2018). Data regarding seed and seedling density (i.e. seed rain, soil seed bank, germination and final seedling density) are expressed in number m⁻² and are continuous quantitative variables. Seedling survival and browsing occurrence are binomial variables, while the different types of mortality are expressed as a percentage. Assumption of normality and homoscedasticity were checked for all
models mentioned below, and the response variable was square root transformed when necessary.
Post-hoc Tukey pairwise multiple comparisons were carried out when we found that a multilevel
factor was significant.

251 *2.5.1. Seed rain and seed bank*

We tested the effects of site and year on seed rain, the effects of site and harvesting method on soil seed bank, and the effect of site on seed morphology using analysis of variance. To investigate the influence of tree size and distance on seed rain, we performed linear mixed modelling with our seed source abundance index as an explanatory variable of the mean annual seed rain and the site as random effect. Differences in germinative capacity of the seeds between the three pools of seed sources and between sites were compared using chi-square tests.

258 *2.5.2. Germination and seedling establishment*

259 Analyses regarding germination, survival, browsing occurrence and final seedling density 260 were performed following a two-step procedure: i) full analyses with the site, harvesting method and 261 seeding as explanatory factors, and all two-and three-way interactions; and ii) intra-site analyses of 262 harvesting method and seeding separately. A random intercept for each monitoring area was 263 included in all models to take into account the spatial dependence of the quadrats of the same 264 monitoring area. Germination, tested for each year separately, and final seedling density were 265 analysed using linear mixed models. Longitudinal analyses of seedling survival were performed on 266 the first cohort of emerged seedlings (i.e. cohort of May, which represents 75.4% of the seedlings in 267 2015) using Cox proportional hazard mixed models. We examined seedling survival only in 2015 268 because the number of seedlings in clear-cuts was too low to compare harvesting treatments and to 269 perform statistical tests for three out of five sites in 2016, and for all sites in 2017. Browsing 270 occurrence was tested with generalized linear mixed models for binary data with a logit link function. 271 The effects of the harvesting method on the distribution of causes of mortality (i.e. drought, rodents 272 or ungulates) were tested using chi-square tests between sites and within sites.

273 In addition, one-sample *t*-tests were used to determine whether the final seeding densities 274 were significantly different from the regeneration success threshold for the four possible 275 management scenarios (i.e. clear-cut, clear-cut + seeding, partial cut, partial cut + seeding).

276 2.5.3. Overall probability of recruitment

In order to identify the limiting stages of regeneration capacity, we calculated the transition probabilities of passing the successive stages of the regeneration process successfully, for each harvest treatment of each site. We estimated the following stages:

i) Seed germinative capacity, estimated via our glasshouse test. .

ii) Germination, as the germination number divided by the viable seed rain (estimated by
combining total seed rain and seed germinative capacity) plus the viable seeds from direct
seeding. Since the germinations of a year resulted from the seed rain of the previous summer,
we used the seed rain data from 2014-2016 and germination data from 2015-2017. Seed rain
before harvest (i.e. 2014) was not measured in our study and was thus calculated from the
2015-2017 data by estimating the contribution of one tree to the seed rain (number of
seeds m⁻² year⁻¹), and relating it to stand density before harvest.

- iii) 1st year survival, as the number of living seedlings one year after emergence divided by the
 germination number.
- iv) 2nd/3rd year survival, as the number of seedlings counted three years after harvest divided by
 the number of living seedlings at the end of each year. This represented the seedlings from
 2015 that died in 2016 and 2017, and those from 2016 that died in 2017.

We finally calculated the overall probability of recruitment as the product of the successive probabilities.

295 *2.5.4. Stand dendrometric features*

Relationships between dendrometric features of partial cuts (tree density, basal area, forest cover, and tree crown surface – see Table S2 for descriptive statistics of these variables) and seedling density were prospected with linear regression. As the year of harvesting had no effect on seedling density (p = 0.33), these regressions were carried out using data from the 29 stands together.

301 3. Results

309

302 3.1. Seed availability

The number of seeds reaching the ground varied depending on the year and the site (both *p* < 0.001; Fig. 2), with a different pattern of annual variation between sites (interaction *p* < 0.001). All sites combined, seed rain averaged 11.1 ± 0.7 seeds m⁻² year⁻¹, with a minimum of 6.1 ± 0.8 seeds m⁻² year⁻¹ in S3 and a maximum of 24 ± 2.3 seeds m⁻² year⁻¹ in S2. Seed rain was influenced by tree size and density of mature trees in the vicinity of the seed traps, as seed rain was positively related to the seed source abundance index (F_(1,44) = 9.59, *p* = 0.004).



Fig. 2: Amount of seed rain for each site per year (n = 10 for each boxplot). Different uppercase letters
 indicate significant differences between sites, whereas different lowercase letters indicate significant
 differences between years within each site (post-hoc Tukey test).

Seed weight and seed length were significantly different between sites (Fig. S2). With an average weight ranging from $38.2 \pm 1.6 \text{ mg}$ (F2) to $47.3 \pm 1.8 \text{ mg}$ (S2), and an average length ranging from $7.28 \pm 0.1 \text{ mm}$ (F2) to $7.71 \pm 0.1 \text{ mm}$ (S2), the seeds from all sites were within the range of known values for maritime pine seeds in Atlantic and Mediterranean forests (Fig. S2).

The amount of seeds in the soil seed bank also varied according to the site (p < 0.001; Fig. S3), but not between clear-cut and partial cut areas (p = 0.215). The average number of seeds was

319 119 ± 6.5 seeds m⁻², with the largest amount in S1 with 196.6 \pm 1.6 seeds m⁻² and the lowest in F1 320 with 79.6 \pm 1.1 seeds m⁻².

The germinative capacity of seeds showed very significant differences between the various seed pools ($\chi^{2}_{(2)} = 202.7$, p < 0.001; Fig. S4). The seeds used for the direct seeding had the highest germinative capacity (86%), followed by the seeds from the seed rain (26 to 32% depending on the site) and then the seed bank, of which no seeds from any stand germinated (Fig. S4). No difference in germinative capacity was observed between the seed rain of the five sites ($\chi^{2}_{(4)} = 0.57$, p = 0.97).

326 3.2. Germination

327 There was a significant site effect with, overall, a higher number of germinations in S2 and 328 S3, and a fewer in S1 (Table 1; Fig. 3). The interactions "site x harvesting method" and "site x 329 seeding" were also significant (Table 1), suggesting a major influence of site-specific conditions on 330 germination. The harvesting method had a highly significant effect on germination each 331 year (Table 1), with a higher number of germinations in partial cuts than in clear-cuts (Fig. 3). The 332 average number of germinations in partial cuts was 0.93 seedlings m⁻² in 2015, and this decreased 333 in the following two years (0.55 \pm 0.06 seedlings m⁻² in 2016; 0.47 \pm 0.09 seedlings m⁻² in 2017). In 334 clear-cuts, the number of germinations decreased sharply between the first and the second year 335 $(0.65 \pm 0.07 \text{ seedlings m}^2 \text{ in } 2015; 0.14 \pm 0.02 \text{ seedlings m}^2 \text{ in } 2016)$, to reach nearly zero in the 336 third year (0.04 \pm 0.02 seedlings m⁻² in 2017). Interestingly, when we examined the results site by 337 site, the positive effect of partial cuts was only observed in sites within the failure area in the first 338 year (Fig. 2A) while it was observed in all sites in the second and third year (Fig. 2B-C). The use of 339 direct seeding had a positive impact on germination only during the first year (Table 1; Fig. 2), 340 where the average number of germinations was 0.92 ± 0.08 seedlings m⁻² in the seeded quadrats 341 and 0.39 \pm 0.07 seedlings m⁻² in the unseeded ones. This positive effect of seeding in the first year 342 was observed in all sites (Fig. 2D).

Table 1: Results of the linear mixed models for the effects of site, harvesting method, seeding and their interactions on number of germinations for the three years. Significant effects (p < 0.05) are indicated in bold.

		2015		2016		2017		
df	F	p	F	p	F	p		

Site	4,90	13.09	<0.001	23.41	<0.001	8.89	<0.001
Harvesting Method	1,90	23.93	<0.001	106.58	<0.001	195.77	<0.001
Seeding	1,290	95.72	<0.001	0.02	0.884	2.75	0.099
Site x Harvesting	4,90	6.35	<0.001	7.95	<0.001	4.90	0.001
Site x Seeding	4,290	4.52	0.002	0.67	0.615	0.98	0.422
Harvesting x Seeding	1,290	0.11	0.741	0.31	0.577	0.38	0.540
Site x Harvesting x Seeding	4,290	0.05	0.994	1.08	0.367	0.80	0.523

and panels C & F to 2017. Asterisks indicate a significant difference between the two modalities within a site (linear mixed models: *** p < 0.001; ** p < 0.01; * p < 0.05; " p < 0.1).

350 **3.3. Survival**

Seedling survival was influenced by the harvesting method ($\chi^2_{(1)} = 8.44$, p = 0.004), but not by the site ($\chi^2_{(4)} = 8.48$, p = 0.075) or their interaction ($\chi^2_{(4)} = 9.16$, p = 0.057). Partial cutting had a positive impact on survival by decreasing mortality by 10.8% compared to clearcutting. On a site-bysite basis, partial cutting had a positive effect on three out of five sites, including the two in the failure area (mortality reduction of 17.5%, 21.5% and 25.6% for S2, F1 and F2 respectively; Fig. 4B-D-E). In addition, F1 and F2 were the only two sites where clear-cut survival rate was lower than 50% (Fig. 4D-E).

358

Fig. 4: Seedling survival probability as a function of harvesting method in 2015, for each site separately. Statistical significance between harvesting method is shown in the bottom right corner of each panel (Cox mixed models: *** p < 0.001; * p < 0.05; ns = non-significant).

Drought was the main cause of mortality (Fig. 5A). All sites combined, we observed a higher mortality due to drought in clear-cuts than in partial cuts ($\chi^{2}_{(4)} = 11.18$, p < 0.001). Within sites, partial cutting significantly reduced the percentage of drought-killed seedlings in S2 ($\chi^{2}_{(1)} = 7.39$, p =0.007), F1 ($\chi^{2}_{(1)} = 6.28$, p = 0.012) and F2 ($\chi^{2}_{(1)} = 8.28$, p = 0.004; Fig. 5A). Seedling death by herbivores was mainly caused by ungulates, while damage caused by rodents was almost nil, except in F1 (Fig. 5A), and harvesting method had no influence on mortality caused by either typeof herbivore.

Conversely to herbivory-killed seedlings, there was a strong effect of site ($\chi^{2}_{(4)} = 72.67$, p < 0.001) and harvesting method ($\chi^{2}_{(1)} = 17.99$, p < 0.001) on the percentage of non-lethal browsed seedlings. Browsing was more frequent in clear-cuts and in the two sites within the failure area (Fig. 5B). It was also on these two sites that the positive effect of partial cutting was the most pronounced, reducing the browsing occurrence by 12.6% in F1 and 22.5% in F2 (Fig. 5B).

374

Fig. 5: A) Percentage mortality of each cause of death in 2015 according the site and harvesting method. Yellow asterisks indicate a significant difference in the distribution of drought-killed seedlings between the two harvesting method within a site (chi-square test: ** p < 0.01; * p < 0.5). B) Seedlings non-lethally browsed by ungulates in 2015 according to the site and the harvesting method. Different lowercase letters indicate significant differences between sites (post-hoc Tukey test). Asterisks indicate a significant difference in browsing occurrence between the two harvesting methods within a site (generalized linear mixed models: *** p < 0.001; * p < 0.5).

382 3.4. Overall regeneration success

Three years after starting regeneration, seedling density was significantly higher in S2 and S3 compared to the three other sites ($F_{(4,90)} = 35.25$, p < 0.001). The harvesting method had a very significant effect on all the sites (Table S3) with a greater density of seedlings in partial cuts (Fig. 6), whereas the use of seeding had a significant positive effect only in three sites (S2, S3 and F1; Table S3). Interaction between harvesting method and seeding was not significant in all sites (Table S3), suggesting that these two practices had independent effects on seedling densities. Clearcutting without seeding did not result in a sufficient seedling density to secure regeneration in any site except for S3 (Fig. 6). Using direct seeding in addition to clear-cutting attained or exceeded the threshold of 0.3 seedlings m⁻² for S1, S2 and S3, but was still largely insufficient for both sites in the failure area (Fig. 6). Partial cutting without seeding had a very positive effect, as it allowed four out of five sites to have a satisfactory regeneration, with F1 being around the predefined threshold (Fig. 6). Adding seeds to partial cuts increased seedling density in all sites, allowing all of them to regenerate easily (Fig. 6).

396

In addition, results from the regional survey of the 29 stands harvested with partial cuts
showed that seedling density was not related to stand density (Fig. 7A) or basal area (Fig. 7B).
Conversely, stands with a higher forest cover (Fig. 7C) or with trees with well-developed crowns
(Fig. 7D) were associated with a more abundant regeneration.

In terms of recruitment probability, we observed strong differences between sites with a minimal overall probability of 0.7% in F1 and a maximum probability of 3.4% in S3. The probability of finding a seedling three years after seed dispersal was higher in partial cuts than in clear-cuts in all sites because of higher germination and greater survival (Table 2). Between all stages, the transition from viable seed rain to germination showed the lowest probability in all sites (Table 2).
Sites in failure areas showed the lowest recruitment probability in clear-cuts because of the lowest
level of survival, while lowest probabilities of germination were observed in S1 (Table 2). Overall,
the probability of final recruitment was between 0.4% and 3.1% in clear-cuts and between 0.9% and
3.6% in partial cuts.

414

Fig. 7: Influence of dendrometric characteristics of partial cuts on seedling density. A) tree density; B) basal area; C) forest cover; and D) tree crown surface area. *P*-values and r^2 of the regressions are shown in the top left corner. Blue lines represent significant regressions, with the 95% confidence intervals in light blue. The dashed line represents the threshold of 0.3 seedlings m⁻² indicating sufficient regeneration.

Table 2: Stage-specific transition probabilities leading to the recruitment of *Pinus pinaster* seedlings,
 according to the site and harvesting method.

	ç	S1		S2		S3		F1		F2	
Stage	Clear- cut	Partial cut	Clear- cut	Partial cut							
Seed germinative capacity	0.300	0.300	0.320	0.320	0.300	0.300	0.260	0.260	0.320	0.320	
Germination	0.036	0.053	0.042	0.073	0.154	0.182	0.055	0.079	0.037	0.114	
1 st year survival	0.707	0.678	0.537	0.690	0.681	0.688	0.465	0.631	0.438	0.632	
2 nd /3 rd year survival	0.891	0.965	0.771	0.885	0.981	0.961	0.693	0.709	0.822	0.834	
Overall probabilty of recruitment for a given seed	0.007	0.010	0.006	0.014	0.031	0.036	0.005	0.009	0.004	0.019	

421

422 4. Discussion

Partial cutting and direct seeding, the two practices evaluated as alternatives to clearcutting, had a positive effect on regeneration in these dune forests. However, their effects on each stage showed site-specific differences, suggesting the importance of local factors on the success of the regeneration process (Rodríguez-García et al., 2010, 2011b). These two practices have two main axes of influence on the regeneration process: the increase of seed availability, improving seed rain and germination; and the modification of microhabitat beneath the tree canopy, affecting both germination and survival.

430 **4.1. Influence of forestry practices on seed availability**

431 Seed rain showed large spatial and temporal variations but did not appear to be a limiting 432 factor in our context, as the two failure sites did not differ from the others and as S3, with the lowest 433 seed rain, was one of the sites with the highest number of germinations and with the greatest 434 regeneration three years after harvest. Two recent studies on maritime pine stands suggesting that 435 neither seed production nor seed dispersal are limiting factors for regeneration showed similar 436 results, with 7-24 seeds m⁻² (Ruano et al., 2015b) and 7-15 seeds m⁻² (Juez et al., 2014). 437 However, it will be important to pay attention to these issues in the future as extreme climatic 438 conditions are expected to increase and could lead to a decrease in seed production (Mutke et al.,

439 2005; Ruano et al., 2015b). In our context, the main problem seems to be seed storage and 440 conservation in the belowground seed bank until germination, which is a highly variable stage 441 between provenances and populations of this species (Wahid and Bounoua, 2013; Calvo et al., 442 2016). In particular, the Atlantic provenance, which includes our coastal forests (de la Mata et al., 443 2012), showed the lowest germination rates in response to drought (Nuñez Paniagua et al., 2013). 444 Here, we observed a probability of germination of only 4–18%, a fairly low percentage compared to 445 other studies on maritime pine in situ (13–62%; Ruano et al., 2015a) or under controlled conditions 446 (14-27%; Cruz et al., 2019). As seed weight was similar in our study and previous ones, this low 447 germination rate is more likely to be explained by: high post-dispersal predation by many different 448 organisms (Ruano et al., 2015a), the presence of a litter layer and harvest debris acting as a 449 physical barrier preventing seeds from reaching the soil (Facelli and Pickett, 1991), or a very poor 450 conservation of seed germinative capacity in the soil (Kozlowski, 2002). Indeed, the number of 451 germinations in 2015 was slightly over one seedling m⁻² at the most, while the soil seed bank was 452 high, with an average of 119 seeds m⁻². Although pine species form only a short-lived soil seed 453 bank with an expected minor role for regeneration (Izhaki et al., 2000), such quantities of seeds in 454 the soil is not surprising with regard to other pine forests, where 150 and 187 seeds m² were 455 reported for *P. pinaster* and *P. halepensis* stands respectively (Daskalakou and Thanos, 1996; Luis-456 Calabuig et al., 2002). However, most seeds in the belowground seed bank are not viable, as 457 shown by two studies on Aleppo pine: Izhaki et al. (2000) found that the density of germinable pine seeds was 0.83 seeds m⁻², while Daskalakou and Thanos (1996) showed that only 8 to 18% of the 458 459 seeds stored in the soil were sound. Our glasshouse test of germination capacity correspond to 460 those results, showing that only seeds from the seed rain of the previous summer could contribute 461 to germination in the following spring, while those stored in the seed bank were not viable for 462 germination. Moreover, the significant decrease in number of germinations in clear-cuts between 463 the first and second year, and then near-zero in the third year confirmed that seeds do not remain 464 viable for more than a year in the soil in most cases. Finally, although seeds from direct seeding 465 had a germination capacity of 86%, their effect on germination was only observed in the first year. 466 This confirms the short life span of the soil seed bank, and that the regeneration of maritime pine 467 depends almost exclusively upon the seed rain falling during the year before harvest.

468 Partial cutting and direct seeding were therefore two effective practices to increase seed 469 stocks and improve germination. Direct seeding had an instantaneous impact by significantly 470 increasing the number of germinations on all sites, up to 4-fold greater than in the unseeded plots, 471 but only in the first year. This strong impact can be explained by their higher germination capacity 472 as a result of a prior selection during which only the heaviest -and hence those most likely to be 473 viable- seeds were selected. It can also be explained by the shorter time spent in the soil by these 474 seeds, as seed rain occurred in summer while direct seeding was done in winter, and was therefore 475 less vulnerable to predation or detrimental environmental conditions. However, the effect of seeding 476 in our study was probably overestimated because seeds were sown manually one-by-one and were 477 thus incorporated into the soil under better conditions, in contrast to the broadcast seeding 478 practiced by foresters. Conversely to seeding, partial cutting had a longer-lasting effect with a 479 regular yearly seed supply, leading to higher germination in all sites in the second and third years.

480 **4.2. Influence of partial cutting on microhabitat**

481 Our results showed a positive effect of partial cuts, both for germination and survival stages. 482 However, the outcome of partial cutting is affected by local site conditions (Rodríguez-García et al., 483 2010), and this was also observed in our study, with large variations between the five sites. Effect of 484 partial cutting was significantly positive in two sites for germination (F1 and F2) and three sites for 485 survival (S2, F1 and F2). The similar differences in germination found in S1, S2 and S3, which also 486 had very different survival rates, and the fact that S2 was similar to F1 and F2 for survival but not 487 germination, suggests that the mechanisms of canopy-induced effects are different depending on 488 the stage of regeneration considered. In general, the positive effects of partial cutting increased 489 gradually with the increase of forest cover, as demonstrated in our regional survey (Fig. 7).

However, the influence of partial cutting on the microhabitat can only be discussed for the first year of the study for two reasons: i) germination in 2015 was linked to the seed rain in 2014 (before exploitation and so were similar in the two harvest modalities), and differences observed that year were therefore mainly due to the influence of harvesting method and not related to seed availability. In contrast, the role of microhabitat on germination became indistinguishable in the second and third years because differences in germination were essentially the consequence of an additional seed supply by seed-trees; and ii) we could not perform statistical analyses regarding
seedling survival as the number of seedlings was too low in clear-cuts in the second and third
years.

499 4.2.1. Influence on germination

500 Soil moisture and soil temperature, which are closely related to the light reaching the forest 501 floor, are the main drivers of germination and could be affected by forest canopy structure 502 (Aussenac, 2000; Castro et al., 2005). In our study, the effect of canopy cover had a strong impact 503 in the two sites within the failure area, as well as a slightly positive effect in the other three sites, as 504 confirmed by the greater probability of germination in partial cuts than in clear-cuts for all sites. This 505 suggests that partial cutting induced slightly more suitable conditions for germination in all sites, and 506 that another factor was involved at this stage within the failure area which was reduced by the 507 presence of a canopy. Canopy mitigation of light intensity in partial cuts was similar in the five 508 stands (Guignabert et al., 2018). This could explain the slight positive effect on all sites because it 509 was the only factor similarly impacted by the partial cutting in all sites. This positive effect was more 510 likely to be due to a decrease in soil temperature than an increase in soil moisture, since the effects 511 of harvesting method were identical in the driest site and the wettest site (see S2 and S3 in Table 512 S1). However, soil moisture was probably a primordial factor for germination in our forests because 513 the site with the highest amount of precipitation (S3) was by far the one with the highest probability 514 of germination. Thus, we may not have preserved enough forest cover in the experimental partial 515 cuts to have a real impact on soil moisture, because the regional sandy soils do not hold enough 516 water, or to maintain a low light intensity as pine forests usually have low LAI values (Gonzalez et 517 al., 2013). Further studies are therefore needed to investigate the role of light and water availability 518 on the germination process to better understand the canopy-induced effects on seedling 519 emergence. The large difference in the number of germinations between the two harvesting 520 methods in the failure area is intriguing and could be the result of a biotic component rather than 521 improved microclimate, such as rodent seed predation. Indeed, rodent seed predation is one of the 522 main bottlenecks in some Spanish pine stands (Ruano et al., 2015a), and a former study has 523 already shown higher rodent damage in this area of our coastal forests (GEREA, 1990). As rodent

foraging activity increases with the volume of coarse woody debris on the stand (Puig-Gironès et al., 2020), this may explain the lower number of germinations in clear-cuts where woody debris was more abundant.

527 *4.2.2. Influence on seedling survival*

528 Summer drought was the main cause of mortality in all sites, and was reduced by partial 529 cuts in three sites (S2, F1 and F2). This is in agreement with previous studies on maritime pine that 530 demonstrated enhancement of seedling recruitment beneath conspecific mature trees where 531 canopy cover buffered abiotic stresses (Ruano et al., 2009; Rodríguez-García et al., 2011a, 2011b). 532 The positive impact of canopy cover on survival would mainly be characterized by a decrease in 533 vapour pressure deficit, as demonstrated on oak seedlings in the same coastal forests (Muhamed et al., 2013), where higher vapour pressure deficit increased seedling transpiration demand and 534 535 seedling sensitivity to drought (Will et al., 2013). Even if we did not have accurate microclimate 536 measurements in each harvesting method x site locations, we observed that S1 had lower 537 temperatures and S3 had higher precipitation. Consequently, they should both have a lower aridity level than the other three sites especially in summer (Ouallet, 2012). This may explain why the 538 539 effect of partial cutting was positive only in S2, F1 and F2. It is also important to mention that the 540 summer of 2015 was an average summer, while the summer of 2016 was the driest of the last 541 decade, particularly the July-August period (precipitation: 94-110 mm in 2015 vs. 15-26 mm in 542 2016). We were unable to compare survival during that year because we lacked seedlings in clear-543 cuts. However, survival in partial cuts in 2016 was between 42% and 76% (Fig. S5), whereas in 544 comparison with another study carried out in the same year in a clear-cut within the failure area, 545 only 10% of seedlings survived (Fig. S5; Guignabert et al., 2020). This suggests that the benefit of 546 partial cutting is even more positive during an extremely dry summer. This drier summer in 2016 547 also showed that mortality after the first year could be high as well (up to 30% in F1) although it 548 remained lower than the first summer mortality. In this previous study, we also found that 549 understorey vegetation could facilitate seedling establishment at intermediate water stress and that 550 damage by deer and rodents, which were quite important, could be modulated by shrub presence 551 (Guignabert et al., 2020). Therefore, further studies investigated the effect of partial cutting on

seedling establishment should consider these biotic interactions, as overstorey canopy may modify
the outcome of plant-plant interactions (Rodríguez-García et al., 2011a).

554 At a larger scale than the seed-seedling microsite, harvesting method also affects habitat 555 and therefore the behaviour of ungulate and rodent species (Reimoser and Gossow, 1996; Fisher 556 and Wilkinson, 2005). Browsing by herbivores can kill seedlings or reduce height growth leading to 557 recruitment failure of tree species or to poorly shaped trees, thus reducing their commercial value 558 (Côté et al., 2004). In our study, ungulate browsing on pine seedlings was higher at both sites in the 559 failure area, and higher in clear-cuts than in partial cuts. Post-harvest development of herbaceous 560 vegetation is greater in clear-cuts than partial cuts because of differences in light intensity, 561 increasing food supplies for ungulates. This results in higher risk of browsing damage in clear-cuts 562 than partial cuts (Reimoser and Gossow, 1996), a risk that is more severe in areas with high 563 ungulate abundance (Tremblay et al., 2007). This was potentially the case for the sites within the 564 failure area, because they were located within a military area where hunting is controlled and lower 565 in comparison to other parts of the coastal forests (Guignabert, 2018). In addition, it has been 566 shown that rodents could have a negative impact on seedling survival in a recently clear-cut stand 567 in the failure area (Guignabert et al., 2020), but this seems not systematic as it was rarely observed 568 in the five sites of the present study.

569 **4.3. Conclusion and management implications**

570 Overall, seedling density in all sites three years after harvest was higher in partial cuts than 571 clear-cuts, and was higher in seeded than unseeded plots (but only significant in three sites). These 572 practices influenced the whole regeneration process by various mechanisms discussed in the 573 foregoing paragraphs, validating our three initial hypotheses. Indeed, partial cutting had a positive 574 effect in the long term by providing seeds each year, and by reducing biotic and abiotic stresses, 575 which improved germination and survival (hyp. 1). This positive effect was actually more 576 pronounced in the failure area, as the only two sites where partial cutting had an impact on both 577 germination and survival were within this area (hyp. 2). Lastly, direct seeding had a positive impact 578 on germination, but seedling density was always higher in partial cuts than in clear-cuts + seeding, 579 which makes its use less beneficial (hyp. 3). Thus, sufficient regeneration can be achieved in all 580 stands by using these two practices appropriately. Their effects are independent, thus enabling us 581 to propose a revision of silvicultural practices currently applied in these forests, while additional 582 research with longer-term monitoring and more replications, particularly in the area of chronic 583 regeneration, should be pursued to optimize these management recommendations.

584 Clear-cutting without seeding led to severe regeneration failure in 4 out of 5 sites, the last 585 one being around the threshold limit. Applying direct seeding enabled only two sites to significantly 586 exceed the required seedling density. Consequently, the practice of clear-cutting currently used in 587 these forests should be avoided, even with the addition of direct seeding. Indeed, seeding had an 588 immediate effect and if the harvesting followed a poor year of seed production or if the spring and/or 589 summer following harvest was very dry, regeneration would not be sufficient resulting in a 590 silvicultural stalemate in the stand, making forest renewal impossible. In contrast, partial cutting had 591 a strong positive impact on all sites and should be the method preferred here, as already performed 592 in the Iberian peninsula in the same type of pine forests with sandy soils and severe summer 593 drought (Calama et al., 2017). The forest cover and to a lesser extent the basal area should be 594 considered when carrying out partial cutting, which should not be based on a specific tree density. 595 The forest cover retained for partial cutting should be at least 18%, preferably with trees with well-596 developed crowns (>33 m²). In areas of regeneration failures where the canopy effects are 597 important for both germination and survival, leaving a denser forest cover (around 30%) would be 598 better. Even if the basal area had no effects on the final seedling density, it was positively related to 599 a higher seed rain. Thus, keeping trees with high diameter and great vigour is also recommended as they are a good indicator of higher cone production (Bravo et al., 2017), and should be 600 601 distributed as homogeneously as possible throughout the stand. In addition, partial cutting could be 602 combined with a direct seeding to maximize the chances of success, but only in failure areas.

603

604 Acknowledgements

The ECODUNE project would have not been possible without the commitment of Didier Canteloup (Office National des Forêts – ONF) who was the first to identify the problem of forest regeneration in the study area, and who built up a consortium of scientists to investigate this major threat to the regional forests. All the authors are grateful to him. We are also very grateful to the ONF staff

- 609 working on the management of the study sites for the valuable help provided for field measurement.
- 610 The ONF staff were in charge of the implementation of the harvesting methods. We also thank
- 611 Céline Birolleau for assistance with field and laboratory work, and Aldyth Nys for revising the
- 612 English. A. Guignabert was funded by a PhD grant by the Région Nouvelle-Aquitaine and Bordeaux
- 613 Sciences Agro.
- 614

615 References

- Alía, R., Martín, S., 2003. EUFORGEN: Technical guidelines for genetic conservation and use for
 Maritime pine (*Pinus pinaster*). Int. Plant Genet. Resour. Inst.
 https://doi.org/10.1016/j.jhazmat.2010.10.029
- Augusto, L., Bakker, M.R., Morel, C., Meredieu, C., Trichet, P., Badeau, V., Arrouays, D., Plassard,
 C., Achat, D.L., Gallet-Budynek, A., Merzeau, D., Canteloup, D., Najar, M., Ranger, J., 2010. Is
 'grey literature' a reliable source of data to characterize soils at the scale of a region? A case
 study in a maritime pine forest in southwestern France. Eur. J. Soil Sci. 61, 807–822.
 https://doi.org/10.1111/j.1365-2389.2010.01286.x
- Aussenac, G., 2000. Interactions between forest stands and microclimate: Ecophysiological aspects
 and consequences for silviculture. Ann. For. Sci. 57, 287–301.
 https://doi.org/10.1051/forest:2000119
- 627 Beckage, B., Clark, J.S., 2003. Seedling survival and growth of three forest tree species: the role of 628 spatial heterogeneity. Ecology 84, 1849–1861.
- Bravo, F., Maguire, D.A., González-Martínez, S.C., 2017. Factors affecting cone production in *Pinus pinaster* Ait.: lack of growth-reproduction trade-offs but significant effects of climate and tree and stand characteristics. For. Syst. 26, e07S. https://doi.org/10.5424/fs/2017262-11200
- Calama, R., Manso, R., Lucas-Borja, M.E., Espelta, J.M., Piqué, M., Bravo, F., Del Peso, C.,
 Pardos, M., 2017. Natural regeneration in Iberian pines: A review of dynamic processes and
 proposals for management. For. Syst. 26. https://doi.org/10.5424/fs/2017262-11255
- Calvo, L., Hernández, V., Valbuena, L., Taboada, A., 2016. Provenance and seed mass determine
 seed tolerance to high temperatures associated to forest fires in *Pinus pinaster*. Ann. For. Sci.
 73, 381–391. https://doi.org/10.1007/s13595-015-0527-0
- Castro, J., Zamora, R., Hodar, J.A., Gómez, J.M., 2005. Ecology of seed germination of *Pinus sylvestris* L. at its southern, Mediterranean distribution range. Investig. Agrar. Sist. y Recur.
 For. 14, 143. https://doi.org/10.5424/srf/2005142-00879
- 641 Cordonnier, T., Tran-ha, M., Piat, J., François, D., 2007. La surface terrière: méthodes de mesure et
 642 intérêts. Rendez-vous Tech. ONF n°18, 9–16.
- Côté, S.D., Rooney, T.P., Tremblay, J.-P., Dussault, C., Waller, D.M., 2004. Ecological Impacts of
 Deer Overabundance. Annu. Rev. Ecol. Evol. Syst. 35, 113–147.
 https://doi.org/10.1146/annurev.ecolsys.35.021103.105725
- 646 Cruz, O., García-Duro, J., Casal, M., Reyes, O., 2019. Role of serotiny on *Pinus pinaster* Aiton
 647 germination and its relation to mother plant age and fire severity. iForest 12, 491–497.
 648 https://doi.org/10.3832/ifor2968-012
- 649 Daskalakou, E.N., Thanos, C.A., 1996. Aleppo pine (*Pinus halepensis*) postfire regeneration: The 59-66. 650 seed banks. Int. J. Wildl. Fire role of canopy and soil 6. 651 https://doi.org/10.1071/WF9960059

- de la Mata, R., Voltas, J., Zas, R., 2012. Phenotypic plasticity and climatic adaptation in an Atlantic
 maritime pine breeding population. Ann. For. Sci. 69, 477–487. https://doi.org/10.1007/s13595011-0173-0
- Dey, D.C., Knapp, B.O., Battaglia, M.A., Deal, R.L., Hart, J.L., O'Hara, K.L., Schweitzer, C.J.,
 Schuler, T.M., 2019. Barriers to natural regeneration in temperate forests across the USA.
 New For. 50, 11–40. https://doi.org/10.1007/s11056-018-09694-6
- Facelli, J.M., Pickett, S.T.A., 1991. Plant litter: Its dynamics and effects on plant community
 structure. Bot. Rev. 57, 1–32. https://doi.org/10.1007/BF02858763
- Fedrowitz, K., Koricheva, J., Baker, S.C., Lindenmayer, D.B., Palik, B., Rosenvald, R., Beese, W.,
 Franklin, J.F., Kouki, J., Macdonald, E., Messier, C., Sverdrup-Thygeson, A., Gustafsson, L.,
 2014. REVIEW: Can retention forestry help conserve biodiversity? A meta-analysis. J. Appl.
 Ecol. 51, 1669–1679. https://doi.org/10.1111/1365-2664.12289
- Fisher, J.T., Wilkinson, L., 2005. The response of mammals to forest fire and timber harvest in the
 North American boreal forest. Mamm. Rev. 35, 51–81. https://doi.org/10.1111/j.13652907.2005.00053.x
- 667 GEREA, 1990. Etude de la prédation exercée par les vertébrés sur les graines de pin maritime 668 dans les forêts dunaires. ONF, Université de Bordeaux I.
- Gómez-Aparicio, L., Valladares, F., Zamora, R., Luis Quero, J., 2005. Response of tree seedlings to
 the abiotic heterogeneity generated by nurse shrubs: an experimental approach at different
 scales. Ecography (Cop.). 28, 757–768. https://doi.org/10.1111/j.2005.0906-7590.04337.x
- González-Alday, J., Martínez-Ruiz, C., Bravo, F., 2009. Evaluating different harvest intensities over
 understory plant diversity and pine seedlings, in a Pinus pinaster Ait. natural stand of Spain.
 Plant Ecol. 201, 211–220. https://doi.org/10.1007/s11258-008-9490-2
- 675 Gonzalez, M., Augusto, L., Gallet-Budynek, A., Xue, J., Yauschew-Raguenes, N., Guyon, D., 676 Trichet, P., Delerue, F., Niollet, S., Andreasson, F., Achat, D.L., Bakker, M.R., 2013. 677 Contribution of understory species to total ecosystem aboveground and belowground biomass Manage. 678 temperate *Pinus pinaster* Ait. forests. For. Ecol. 38–47. in 289. 679 https://doi.org/10.1016/j.foreco.2012.10.026
- Greene, D.F., Zasada, J.C., Sirois, L., Kneeshaw, D., Morin, H., Charron, I., Simard, M.J., 1999. A
 review of the regeneration dynamics of North American boreal forest tree species. Can. J. For.
 Res. 29, 824–839. https://doi.org/10.1139/x98-112
- 683 Grossnickle, S., Ivetić, V., 2017. Direct Seeding in Reforestation A Field Performance Review. 684 REFORESTA 94–142. https://doi.org/10.21750/REFOR.4.07.46
- Guignabert, A., 2018. Etude des processus de régénération naturelle du pin maritime en contexte
 de dune forestière gérée. Influence de la sylviculture, du climat et des interactions biotiques.
 PhD Thesis Université de Bordeaux.
- Guignabert, A., Augusto, L., Gonzalez, M., Chipeaux, C., Delerue, F., 2020. Complex biotic
 interactions mediated by shrubs: Revisiting the stress-gradient hypothesis and consequences
 for tree seedling survival. J. Appl. Ecol. 57, 1341–1350. https://doi.org/10.1111/13652664.13641
- Guignabert, A., Delerue, F., Gonzalez, M., Augusto, L., Bakker, M., 2018. Effects of Management
 Practices and Topography on Ectomycorrhizal Fungi of Maritime Pine during Seedling
 Recruitment. Forests 9, 245. https://doi.org/10.3390/f9050245
- Harmer, R., 2001. The effect of plant competition and simulated summer browsing by deer on tree
 regeneration. J. Appl. Ecol. 38, 1094–1103. https://doi.org/10.1046/j.1365-2664.2001.00664.x
- Harper, J.L., 1977. Population biology of plants. Academic Press, London.
- Heithecker, T.D., Halpern, C.B., 2006. Variation in microclimate associated with dispersed-retention

- harvests in coniferous forests of western Washington. For. Ecol. Manage. 226, 60–71.
 https://doi.org/10.1016/j.foreco.2006.01.024
- IFN, 2017. Inventaire Forestier National [WWW Document]. URL https://inventaire-forestier.ign.fr/
 (accessed 12.18.17).
- 703 IPCC, 2014. Climate Change 2014: Synthesis Report. Cambridge University Press, Cambridge.
- Izhaki, I., Henig-Sever, N., Ne'Eman, G., 2000. Soil seed banks in Mediterranean Aleppo pine
 forests: the effect of heat, cover and ash on seedling emergence. J. Ecol. 88, 667–675.
 https://doi.org/10.1046/j.1365-2745.2000.00486.x
- 707 Juez, L., González-Martínez, S.C., Nanos, N., De-Lucas, A.I., Ordóñez, C., del Peso, C., Bravo, F., 708 2014. Can seed production and restricted dispersal limit recruitment in Pinus pinaster Aiton 709 Northern Plateau? Manage. from the Spanish For. Ecol. 313. 329-339. 710 https://doi.org/10.1016/j.foreco.2013.10.033
- Kozlowski, T.T., 2002. Physiological ecology of natural regeneration of harvested and disturbed
 forest stands: implications for forest management. For. Ecol. Manage. 158, 195–221.
 https://doi.org/10.1016/S0378-1127(00)00712-X
- Lavoie, J., Montoro Girona, M., Morin, H., 2019. Vulnerability of Conifer Regeneration to Spruce
 Budworm Outbreaks in the Eastern Canadian Boreal Forest. Forests 10, 850.
 https://doi.org/10.3390/f10100850
- Löf, M., 2000. Establishment and growth in seedlings of *Fagus sylvatica* and *Quercus robur*:
 influence of interference from herbaceous vegetation. Can. J. For. Res. 30, 855–864.
 https://doi.org/10.1139/x99-257
- Luis-Calabuig, E., Torres, O., Valbuena, L., Calvo, L., Marcos, E., 2002. Impact of large fires on a community of *Pinus pinaster*. Fire Biol. Process. 1–12.
- Montoro Girona, M., Lussier, J.-M., Morin, H., Thiffault, N., 2018. Conifer Regeneration After
 Experimental Shelterwood and Seed-Tree Treatments in Boreal Forests: Finding Silvicultural
 Alternatives. Front. Plant Sci. 9, 1–14. https://doi.org/10.3389/fpls.2018.01145
- Montoro Girona, M., Morin, H., Lussier, J.-M., Ruel, J.-C., 2019. Post-cutting Mortality Following
 Experimental Silvicultural Treatments in Unmanaged Boreal Forest Stands. Front. For. Glob.
 Chang. 2, 1–16. https://doi.org/10.3389/ffgc.2019.00004
- Montoro Girona, M., Morin, H., Lussier, J.-M., Walsh, D., 2016. Radial Growth Response of Black
 Spruce Stands Ten Years after Experimental Shelterwoods and Seed-Tree Cuttings in Boreal
 Forest. Forests 7, 240. https://doi.org/10.3390/f7100240
- Muhamed, H., Touzard, B., Le Bagousse-Pinguet, Y., Michalet, R., 2013. The role of biotic
 interactions for the early establishment of oak seedlings in coastal dune forest communities.
 For. Ecol. Manage. 297, 67–74. https://doi.org/10.1016/j.foreco.2013.02.023
- Mutke, S., Gordo, J., Gil, L., 2005. Variability of Mediterranean Stone pine cone production: Yield
 loss as response to climate change. Agric. For. Meteorol. 132, 263–272.
 https://doi.org/10.1016/j.agrformet.2005.08.002
- Nuñez Paniagua, M., Sierra de Grado, R., Alía, R., Bravo, F., 2013. Efecto del estrés hídrico y la oscilación de las temperaturas sobre la germinación de semillas de diversas procedencias de Pinus pinaster Ait., in: 6º Congreso Forestal Español. Vitoria, junio, pp. 1–12.
- Nyland, R.D., 2016. Silviculture: Concepts and Applications. 3rd edn. Waveland Press, Long Grove,
 Illinois.
- Ouallet, P., 2012. Quels peuvent-être les facteurs écologiques responsables des échecs de régénération naturelle du pin maritime sur les dunes littorales des forêts domaniales de Biscarrosse et de Sainte-Eulalie? MsC Thesis - BSA, ONF.
- 745 Price, D.T., Zimmermann, N.E., van der Meer, P.J., Lexer, M.J., Leadley, P., Jorritsma, I.T.M.,

- Schahber, J., Clark, D.F., Lasch, P., McNulty, S., Wu, J., Smith, B., 2001. Regeneration in gap
 models: priority issues for studying forest responses to climate change. Clim. Change 51, 475–
 508. https://doi.org/10.1023/A:1012579107129
- Puig-Gironès, R., Imbeau, L., Clavero, M., Rost, J., Pons, P., 2020. Does post-fire salvage logging
 affect foraging activity by rodents? Eur. J. For. Res. https://doi.org/10.1007/s10342-020-012855
- R Core Team, 2018. R: A language and environment for statistical computing. R Foundation for
 Statistical Computing, Vienna, Austria. https://www.r-project.org/.
- Reimoser, F., Gossow, H., 1996. Impact of ungulates on the silvicultural system. For. Ecol. Manage.
 88, 107–119.
- Rodríguez-García, E., Bravo, F., Spies, T.A., 2011a. Effects of overstorey canopy, plant–plant
 interactions and soil properties on Mediterranean maritime pine seedling dynamics. For. Ecol.
 Manage. 262, 244–251. https://doi.org/10.1016/j.foreco.2011.03.029
- Rodríguez-García, E., Gratzer, G., Bravo, F., 2011b. Climatic variability and other site factor
 influences on natural regeneration of *Pinus pinaster* Ait. in Mediterranean forests. Ann. For.
 Sci. 68, 811–823. https://doi.org/10.1007/s13595-011-0078-y
- Rodríguez-García, E., Juez, L., Bravo, F., 2010. Environmental influences on post-harvest natural
 regeneration of *Pinus pinaster* Ait. in Mediterranean forest stands submitted to the seed-tree
 selection method. Eur. J. For. Res. 129, 1119–1128. https://doi.org/10.1007/s10342-010-03997
- Ruano, I., del Peso, C., Bravo, F., 2015a. Post-dispersal predation of *Pinus pinaster* Aiton seeds:
 key factors and effects on belowground seed bank. Eur. J. For. Res. 134, 309–318.
 https://doi.org/10.1007/s10342-014-0853-z
- Ruano, I., Manso, R., Fortin, M., Bravo, F., 2015b. Extreme climate conditions limit seed availability
 to successfully attain natural regeneration of Pinus pinaster in sandy areas of central Spain.
 Can. J. For. Res. 45, 1795–1802. https://doi.org/10.1139/cjfr-2015-0257
- Ruano, I., Pando, V., Bravo, F., 2009. How do light and water influence *Pinus pinaster* Ait.
 germination and early seedling development? For. Ecol. Manage. 258, 2647–2653.
 https://doi.org/10.1016/j.foreco.2009.09.027
- Sardin, T., 2009. Guide des sylvicultures Forêts littorales atlantiques dunaires. Office National des
 Forêts, Paris.
- Schupp, E.W., 1995. Seed-seedling conflicts, habitat choice, and patterns of plant recruitment. Am.
 J. Bot. 82, 399–409. https://doi.org/10.1002/j.1537-2197.1995.tb12645.x
- Tremblay, J.P., Huot, J., Potvin, F., 2007. Density-related effects of deer browsing on the
 regeneration dynamics of boreal forests. J. Appl. Ecol. 44, 552–562.
 https://doi.org/10.1111/j.1365-2664.2007.01290.x
- Vergarechea, M., del Río, M., Gordo, J., Martín, R., Cubero, D., Calama, R., 2019. Spatio-temporal
 variation of natural regeneration in *Pinus pinea* and *Pinus pinaster* Mediterranean forests in
 Spain. Eur. J. For. Res. https://doi.org/10.1007/s10342-019-01172-8
- Viglas, J.N., Brown, C.D., Johnstone, J.F., 2013. Age and size effects on seed productivity of
 northern black spruce. Can. J. For. Res. 43, 534–543. https://doi.org/10.1139/cjfr-2013-0022
- Wagner, S., Fischer, H., Huth, F., 2011. Canopy effects on vegetation caused by harvesting and
 regeneration treatments. Eur. J. For. Res. 130, 17–40. https://doi.org/10.1007/s10342-010 0378-z
- Wahid, N., Bounoua, L., 2013. The relationship between seed weight, germination and biochemical
 reserves of maritime pine (*Pinus pinaster* Ait.) in Morocco. New For. 44, 385–397.
 https://doi.org/10.1007/s11056-012-9348-2

Will, R.E., Wilson, S.M., Zou, C.B., Hennessey, T.C., 2013. Increased vapor pressure deficit due to
 higher temperature leads to greater transpiration and faster mortality during drought for tree
 seedlings common to the forest-grassland ecotone. New Phytol. 200, 366–374.
 https://doi.org/10.1111/nph.12321