



HAL
open science

Debunking the Myth of the Endogenous Antiangiogenic Vegfxxx_b Transcripts

Hugues Dardente, William English, Manoj Valluru, Chryso Kanthou, David Simpson

► **To cite this version:**

Hugues Dardente, William English, Manoj Valluru, Chryso Kanthou, David Simpson. Debunking the Myth of the Endogenous Antiangiogenic Vegfxxx_b Transcripts. Trends in Endocrinology and Metabolism = Trends in Endocrinology & Metabolism , 2020, 31 (6), pp.398-409. 10.1016/j.tem.2020.01.014 . hal-02927071

HAL Id: hal-02927071

<https://hal.inrae.fr/hal-02927071v1>

Submitted on 28 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



This is a repository copy of *Debunking the myth of the endogenous antiangiogenic Vegfxxx b transcripts.*

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/157656/>

Version: Accepted Version

Article:

Dardente, H., English, W.R. orcid.org/0000-0003-3024-2441, Valluru, M.K. et al. (2 more authors) (2020) Debunking the myth of the endogenous antiangiogenic Vegfxxx b transcripts. *Trends in Endocrinology & Metabolism*, 31 (6). pp. 398-409. ISSN 1043-2760

<https://doi.org/10.1016/j.tem.2020.01.014>

Article available under the terms of the CC-BY-NC-ND licence
(<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

1 **Debunking the myth of the endogenous anti-angiogenic *Vegfaxxb* transcripts.**

2

3 Hugues Dardente^{1*}, William R English², Manoj K Valluru², Chryso Kanthou² & David Simpson³

4

5 ¹PRC, INRA, CNRS, IFCE, Université de Tours, 37380 Nouzilly, France

6 ²Department of Oncology and Metabolism, Tumour Microcirculation Group, University of

7 Sheffield, School of Medicine, Beech Hill Road, Sheffield, S10 2RX, United Kingdom

8 ³Wellcome-Wolfson Institute for Experimental Medicine, Queen's University Belfast, Belfast,

9 BT7 1NN, Northern Ireland

10

11 *Corresponding author: hugues.dardente@inra.fr

12

13 **Keywords:** VEGFA, transcription, RNA-seq, alternative splicing, angiogenesis, cancer, retina.

14 **Conflicts of interest:** the authors declare that no conflict of interest exists.

15 **Author contributions:** HD wrote the first draft and prepared the figures; CK, WRE, MJK and

16 DS edited the MS and contributed additional data analysis; all authors approved the final draft.

17

18 **Abstract**

19 In this article we critically assess evidence for the existence of a family of anti-angiogenic

20 *Vegfaxxb* transcripts, arising from the use of a phylogenetically conserved alternative distal

21 splice site within exon 8 of the *VEGFA* gene. We explain that prior evidence for *Vegfaxxb*

22 transcripts in tissues rests heavily upon flawed RT-PCR methodologies, with the extensive use of

23 5'-tailing in primer design being the main issue. Furthermore, our analysis of large RNA-seq

24 datasets (human and ovine) fails to identify a single *Vegfaxxb* transcript. Therefore, we

25 challenge the very existence of *Vegfaxxb* transcripts, which further questions the physiological

26 relevance of studies based on the use of “anti-VEGFA_{xxx}b” antibodies. Our analysis has
27 implications for the proposed therapeutic use of isoform-specific anti-VEGFA strategies for
28 treating cancer and retinopathies.

29

30 **Classical angiogenic VEGFA isoforms**

31 VEGFA is a key regulator of vascular homeostasis and therapeutic target in pathological
32 angiogenesis. The *VEGFA* gene is subject to alternative splicing, which occurs primarily in
33 exons 6 and 7 (**Figure 1**) [1-4]. In humans, the most abundant mature translated isoform is
34 VEGFA165 (i.e. this isoform comprises 165 amino acids), which lacks exon 6. Three other
35 isoforms are present in moderate/high amounts: VEGFA121, which lacks both exons 6 and 7,
36 VEGFA189, which includes exon 7 but uses an alternative 5'-donor site in exon 6 and
37 VEGFA206, which includes both exons 6 and 7 (**Figure 1B & Figure 1E**). These VEGFA
38 isoforms exist in other species, including mouse, sheep and cattle, but they are one amino acid
39 shorter than their human counterparts. Other less abundant isoforms have been described, some
40 of which are also generated through alternative splicing within exons 6/7 (VEGFA145,
41 VEGFA148, VEGFA183). The amino acid stretches encoded by exons 6 and 7 are enriched in
42 clusters of basic amino acids, which confer VEGFA with the ability to bind the extracellular
43 matrix and thus mitigate its diffusion. Consequently, VEGFA121 is highly diffusible, while
44 VEGFA189 and VEGFA206 remain bound to the extracellular matrix and VEGFA165 has
45 intermediate matrix binding characteristics [1-4]. VEGFA isoforms bind with varying affinities
46 to the tyrosine kinase receptors VEGFR1 and VEGFR2 and recruit the co-receptor Neuropilin 1
47 to activate intracellular signaling pathways as illustrated on the left side of **Figure 1G** [5,6].
48 Critically, all VEGFA isoforms promote angiogenesis, which is key to the progression of
49 tumorigenesis [7].

50

51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71

Non-classical anti-angiogenic VEGFAxxx_b and VEGF-Ax isoforms

In 2002, Bates and colleagues [8] reported a novel human *Vegfa* splice variant, generated through use of an alternative distal splice site within the last exon of the *VEGFA* gene, exon 8 (see **Figure 1C** and **Figure 1D**). The transcript was dubbed *Vegfa165b*. Follow-up studies suggested that the encoded VEGFA165b protein product belongs to an entire VEGFAxxx_b family of proteins, which are derived from transcripts that use the distal splice site in exon 8 (**Figure 1E**) and bear anti-angiogenic properties [3,9-14]. From a functional standpoint, VEGFA165b was shown to bind to VEGFR2 with the same affinity as VEGFA165 [15-18] but failed to bind to Neuropilin 1 as illustrated on the right part of **Figure 1G** [16,17,19]. VEGFA165b induces VEGFR2 tyrosine phosphorylation in a dose-dependent manner, albeit less efficiently and more transiently than VEGFA165 [15-17,19]. However, this impaired ability of VEGFA165b to activate intracellular signaling pathways may not apply to all cell types [20,21]. Another isoform, dubbed VEGF-Ax, has recently been described [22]. VEGF-Ax arise from programmed translational read-through, such that the stop-codon of classical VEGFA isoforms leads to insertion of a Serine residue, and the in-frame stop-codon of putative VEGFAxxx_b isoforms is used instead. In other words, VEGF-Ax and VEGFAxxx_b isoforms would share the same C-term sequence. However, the *in vivo* existence of VEGF-Ax is disputed and *in vitro* data demonstrate pro-angiogenic properties [23].

Existence of VEGFAxxx_b isoforms: clinical implications

Overall, VEGFA165b appears to behave as a partial agonist [16], which competes with VEGFA165 for VEGFR2/Neuropilin 1 binding. This may explain the reduced angiogenic potential of VEGFA165b [20,23,24]. Based on this, Harper & Bates proposed a model in which

76 the balance between endogenous levels of angiogenic VEGFA_{xxx} and anti-angiogenic
77 VEGFA_{xxxb} isoforms sets the angiogenic potential of a tissue [3]. They further proposed that
78 VEGFA_{xxxb} isoforms predominate in many normal cells and tissues where they can amount to
79 more than 50% of total VEGFA protein [10]. In normal colonic tissues in particular,
80 VEGFA_{xxxb} variants have been reported to comprise as much as 95% of all VEGFA [25]. The
81 proposed use of the alternative distal splice site within exon 8 is therefore not a rare event, and
82 its prevalence implies major physiological relevance. However, in pathological conditions such
83 as cancer and retinopathies, VEGFA_{xxxb} levels appear to be substantially downregulated.
84 Considering the crucial role of VEGFA in angiogenesis of various cancers and retinopathies
85 [1,26-28] such a finding may have broad clinical implications.

86
87 Indeed, current strategies to target VEGFA, such as the anti-VEGFA antibody bevacizumab,
88 Avastin®) or aflibercept (VEGFA-trap), would indiscriminately inhibit both angiogenic
89 VEGFA_{xxx} and anti-angiogenic VEGFA_{xxxb} isoforms. Bates *et al* showed that targeting
90 VEGFA in colorectal cancer was more efficient if endogenous levels of VEGFA_{xxxb} were low
91 and proposed that assessment of the ratio of VEGFA_{xxxb} to VEGFA_{xxx} could potentially
92 predict response to bevacizumab and other therapies directed against VEGFA [3]. Therefore,
93 modification of VEGFA splicing to alter the VEGFA_{xxxb}/VEGFA_{xxx} balance, or development
94 of compounds to target only VEGFA_{xxx} isoforms, may have therapeutic value for treating
95 cancers and retinopathies [29]. In 2009, D.O. Bates and S.J. Harper filed a patent for the “Novel
96 use of VEGFA_{xxxb}”, which covers the use of these isoforms (primarily VEGFA165b) for
97 several potential therapeutic applications.

98

99 **The Controversy: Do VEGFA_{xxxb} isoforms really exist?**

100 Despite a substantial amount of literature reporting evidence for the existence of VEGFA_{xxx}b
101 isoforms, the VEGFA scientific community is divided on this issue. We and others have yet to
102 detect these isoforms in normal and pathological tissues and have thus concluded that
103 VEGFA_{xxx}b isoforms, if they do exist, are likely not of physiological relevance. In recent
104 reviews on VEGFA signaling and disease, VEGFA_{xxx}b biology is largely disregarded [7] or
105 notably overlooked [30]. Until now, evidence for the existence of VEGFA_{xxx}b isoforms has
106 been provided through PCR-based approaches and studies using an antibody generated against
107 the sequence TCRSLTRKD encoded by putative exon 8b of the human *Vegfa* gene. Here, we
108 critically review and reassess the evidence which led to the assumption that "there is an
109 important role for VEGFA_{xxx}b isoforms in normal physiology" [10]. Our own findings [31-33]
110 and a thorough analysis of the literature has led us to question and re-evaluate the existence of
111 *Vegfa_{xxx}b* transcripts

112

113 **Absence of Evidence: Detection of *Vegfa_{xxx}b* transcripts with flanking primers.**

114 Since alternative splicing affects exons 6, 7 and 8, the best way to identify all *Vegfa* splice
115 variants is to use primers flanking these regions. Such a primer design is expected to amplify all
116 *Vegfa* isoforms. To do so, the forward primer has to be located within exons 1-5 (common to all
117 splice variants) and the reverse primer located downstream of the putative exon 8b splice site. If
118 *Vegfa_{xxx}* and *Vegfa_{xxx}b* splice variants are sister families and *Vegfa_{xxx}b* mRNAs represent a
119 large fraction of total *Vegfa* mRNA, at least in normal tissues [10], a standard RT-PCR with the
120 aforementioned primers would be expected to yield at least 6 main products/bands corresponding
121 to mRNA for the major isoforms VEGFA189, VEGFA165 and VEGFA121 and their "sister
122 bands" for VEGFA189b, VEGFA165b and VEGFA121b, albeit at different abundances. The
123 bands encoding *Vegfa_{xxx}b* isoforms would be 66bp shorter than their respective counterparts
124 (see **Figure 1D**) thereby yielding 3 "doublets" easily discernible on an agarose gel. However, in

125 their seminal paper reporting on the discovery of *Vegfa165b* [8], Bates *et al* stated that "PCR of
126 the full-length product using primers V165K (complementary to the translation initiation site of
127 the other isoforms of VEGF) and V165X (a primer downstream of the original 3'-UTR) resulted
128 in one strong band at ~670bp". Considering the strategy used, it is difficult to explain why other
129 *Vegfaxxx* and *Vegfaxxxb* transcripts were not identified by this initial study. Indeed, as mentioned
130 above, at least 6 bands, corresponding to the 3 "doublets" (VEGFA189/189b, VEGFA165/165b
131 and VEGFA121/121b), should have been observed by agarose electrophoresis of PCR products.
132 ~~Furthermore, nested PCR with a 3'-UTR and exon 7a primers resulted in a strong band at~~
133 ~~~130bp confirming that the full-length was VEGFA165b".~~

134
135 We and others have adopted this strategy in order to identify *Vegfaxxxb* transcripts in human,
136 mouse, sheep and cattle but failed to identify *Vegfa165b* mRNA - or other *Vegfaxxxb* products
137 [22,31,33,34]. Instead, only three bands were observed (see **Figure 1B** and **Figure 1F**) and
138 sequencing of the PCR products showed unequivocally that they encoded VEGFA189,
139 VEGFA165 and VEGFA121 [22,24,32-34]. Therefore, an unbiased and straightforward RT-PCR
140 strategy to simultaneously amplify all *Vegfa* isoforms failed to support the existence of
141 *Vegfaxxxb* mRNA. Since these flanking primers could not discriminate *Vegfaxxx* from putative
142 *Vegfaxxxb* transcripts, isoform-specific primers were developed.

143
144 **Detection of *Vegfaxxxb*-like transcripts with "isoform-specific primers": a tale of a tail.**
145 Over time, there has been a considerable drift in the design of primers aimed at the specific
146 identification of *Vegfa165b* that span the exon 7 – exon 8b splice site. However, in most studies,
147 the exon 7-specific stretch has been greatly lengthened at the expense of the exon 8b-specific
148 region (see **Figure 2A** and **Table S1**). Since exon 7 would be shared by both *Vegfa165* and
149 *Vegfa165b*, the isoform-specificity of these reverse primers is questionable. Technically, the

150 necessary and sufficient primer design to allow specific detection of *Vegfa165b* mRNA would
151 include the shortest possible 3'-anchor within exon 7; based on standard PCR principles (see
152 **Figure 3**) one base specific to exon 7 on the 3'-end of the primer would suffice. However, in the
153 case of the human sequence, since the AT motif would be shared by both *Vegfa165* and
154 *Vegfa165b* (see **Figure 1**) this means that the last 3bp at the 3'-end of such a primer would
155 anneal to exon 7 with the last nucleotide discriminating between *Vegfa165* and *Vegfa165b*
156 isoforms (see **Figure 2A**, Primer P1). Surprisingly, this most simple strategy has never been used
157 in the literature.

158
159 Rather than using such a “minimal and sufficient” primer design for detection of *Vegfaxxb*
160 isoforms, the design of the primer has been "adapted". The initial design, published in the
161 seminal *Vegfa165b* paper by Bates *et al* [8], comprised of 7bp complementary to either *Vegfa165*
162 or *Vegfa165b* (**Figure 2A**, Primer P2). This primer has been used in a number of subsequent
163 studies by the same group [15,35-39] and other research teams that used RT-PCR to demonstrate
164 the existence of *Vegfaxxb* transcripts [40,41]. In 2007, Ribeiro *et al* used a modified version of
165 this primer for their studies in pig [42], with 9bp complementary to either *Vegfa165* or
166 *Vegfa165b* (**Figure 2A**, Primer P3), while Baba *et al* used 8bp complementary to either
167 *Vegfa165* or *Vegfa165b* [43]. Overall, the number of bases complementary to either *Vegfa165* or
168 *Vegfa165b* has been on the rise since 2007: 13 bp in 2008 ([44]; **Figure 2A**, Primer P4), 14bp in
169 2009 (study in rat; [45]) and 15bp in 2010-2011 ([46,47]; **Figure 2A**, Primer P5). Finally, a
170 design including no less than 17bp common to both *Vegfa165* or *Vegfa165b* was proposed to
171 specifically amplify *Vegfa165b* mRNA ([48]; **Figure 2A**, Primer P6). We also note that the same
172 methodology has been used for the design of a *Vegfa165b*-specific forward primer, which
173 includes 14bp that would also anneal to *Vegfa165* ([34]; **Figure 2A**, Primer P7). This approach
174 has also been used by others to "specifically" detect *Vegfa121b* isoforms ([29,44]; see **Table S1**).

175
176 We demonstrated that this primer design is responsible for PCR artefacts, due to 5'-tailing, that
177 could be mis-interpreted as *Vegfaxxb* transcripts ([31,33]; see **Figure 3**). The conclusion of
178 these studies was that PCR products correspond to putative *Vegfaxxb* transcripts only because
179 primers were designed to include on their 5'-end a few bases of the *Vegfaxxb* sequence itself.
180 This method has been used for decades, most often to add restriction sites to aid cloning. Good
181 examples are provided by Catena *et al* [20] and Ganta *et al* [49], who flanked the 5'-end of their
182 reverse primers with an *HindIII* (or *BamHI*, respectively) restriction site followed by the entire
183 predicted sequence for the putative exon 8b and 20-23 bp complementary to either exon 5 or
184 exon 7 in order to generate synthetic expression vectors for *Vegfa121b* or *Vegfa165b* (see **Table**
185 **S1**). We demonstrated that *Vegfa* amplification by RT-PCR could be obtained when bases
186 "specific for *Vegfaxxb*" on the primer 5'-end were changed to a GGGGG or an AAAAA stretch
187 [33]. The minimal primer design we defined to be sufficient to obtain artefactual *Vegfaxxb*-like
188 PCR products perfectly matches the P4 primer described above ([33]; see **Figure 2B**). Therefore,
189 studies which claimed identification of *Vegfaxxb* transcripts using primers with ≥ 13 bp common
190 to *Vegfaxx* and putative *Vegfaxxb* mRNA do not reach required standards of evidence, and
191 their conclusions should be re-evaluated.

192
193 By 2013, the use of the questionable P4 primer design had become widespread in the literature
194 [44,50-52]. This design constitutes a significant departure from initial studies that used the P2
195 primer with 7bp complementarity to either *Vegfa165* or *Vegfa165b*. The reasons for this drift in
196 primer design are not clear. However, as early as 2006, Bates and colleagues [37] acknowledged
197 that "...RT-PCR is not quantitative and it has not been possible so far to develop isoform-
198 specific qPCR, due to the lack of exon-specific sequences". It seems surprising that primers

199 deemed suitable for specific detection of *Vegfa165b* in standard RT-PCR would not be suitable
200 for qPCR.

201 In conclusion, the detection of “*Vegfaxxb*-like” transcripts could be explained by the use of
202 inadequate RT-PCR methodologies involving 5'-tailing of the primers (also see **BOX1**).

203

204

205

206 **Bioinformatic analysis of alternative splicing: absence of evidence for *Vegfaxxb* isoforms.**

207 Using AVISPA, a tool for prediction and analysis of alternative splicing, Barash *et al* reported
208 that "prediction of other splice variations of *Vegfa*, such as the 3' splice site variation in exon 8,
209 are currently not supported by the tool [53] ". Another splice site prediction software ([54];
210 available at http://www.fruitfly.org/seq_tools/splice.html) identified the canonical acceptor
211 splice site of exon 8 (score of 0.97) but failed to identify the putative exon 8b splice site.
212 Therefore, efficient *in silico* tools do not currently support the existence of the proposed distal
213 splice site in exon 8 of the *Vegfa* gene.

214

215 **Detection of the *Vegfaxxb* transcripts: evidence of absence by unbiased RNA-seq analysis.**

216 As detailed above, the most parsimonious explanation for the lack of *Vegfaxxb* PCR products is
217 that such transcripts do not exist. However, a theoretical possibility remains that failure to detect
218 *Vegfaxxb* transcripts results from the targeted nature of the approach (also see **BOX2**). If so, the
219 analysis of RNA datasets obtained through the use of an alternative, unbiased methodology, such
220 as RNA-seq, should clarify this matter. Indeed, “RNA-seq represents the method of choice for
221 the discovery of alternative splicing events across tissues [55]”.

222

223 Therefore, we investigated existence of *Vegfaxxb* transcripts in publicly available RNA-seq
224 datasets generated from multiple human tissues [29]. Our extensive analysis confirmed all
225 known *Vegfa* splice variants and also identified novel transcripts. In particular >40000 *Vegfa*
226 transcripts were uncovered that spanned the junction between exon 7 and exon 8 in >10 different
227 human tissues. None of these reads supported the existence of an exon 8b splice site [29].

228
229 We then analysed two independent RNA-seq datasets of hypothalamus / pituitary from castrated
230 rams [57] and ovariectomized, estradiol-implanted ewes [56]. This analysis confirmed that *Vegfa*
231 splice variants are phylogenetically conserved, as transcripts identified in sheep correspond to
232 those present in human [33]. Indeed, we also identified several novel ovine *Vegfa* transcripts that
233 correspond to those identified by Bridgett *et al* in human [29]. Crucially, we identified 2693
234 *Vegfa* transcripts that cover the junction between exon 5 and exon 8 or between exon 7 and exon
235 8. None of the reads supported the existence of an exon 8b splice site, hence the existence of
236 ovine *Vegfaxxb* transcripts [33]. These findings [33] rule out the proposed role for VEGFA_{xxx}b
237 isoforms in the control of ovine seasonal breeding proposed by Castle-Miller *et al* [58]. We
238 further note that the 43bp-long reverse primer (see **Table S1**), as provided in the erratum of the
239 aforementioned study [58] to “specifically” amplify *Vegfa165b* by qPCR, would indeed lead to
240 co-amplification of the classical *Vegfa165* isoform, as its sequence is complementary to a stretch
241 of exon 8 located downstream of the putative distal splice site. This implies that qPCR would
242 yield two amplicons, which makes the primer pair inadequate for this use and leads to questions
243 on the validity of the data published by Castler-Miller *et al* [58]. **In summary, the failure to**
244 **detect *Vegfaxxb* mRNA in human and ovine RNA-seq data strongly suggests that these**
245 **transcripts do not exist** (also see **BOX3**).

246

247 **No *Vegfaxxb* transcripts: detection of VEGFA_{xxx}b proteins is an artefact.**

248 Absence of endogenous *Vegfxxx* mRNA implies absence of endogenous VEGFAxxx
249 proteins. Consequently, studies which investigated VEGFAxxx levels in tissues or plasma,
250 using anti-VEGFAxxx antibodies and ELISA kits, or those which relied exclusively on over-
251 expression or injection of recombinant VEGFAxxx proteins have questionable physiological
252 relevance (see **Table S1**; e.g. [18,59-74]).

253
254 The first antibody against VEGFAxxx isoforms (MVRL56/1) was developed by Woolard *et al*
255 2004 [15] and was raised against the synthetic peptide TCRSLTRKD which corresponds to the
256 nine amino acid C-terminal sequence of human VEGFA165b (see **Figure 1C**). This peptide is
257 only six amino acids different from the peptide that corresponds to the C-terminal sequence of
258 VEGFA165 (TCRC DKPRR). This antibody was made commercially available and distributed
259 by R&D (MAB3045) and Abcam (Ab149940; [41]). A sandwich ELISA kit that utilizes the
260 MAB3045 antibody is also available via R&D (#DY3045) who also supply a recombinant
261 human VEGFA165b protein (#3045-VE-025). Interestingly, Abcam removed Ab149940 from
262 their catalog in 2015 as it "did not meet the quality criteria". Specifically, the antibody yielded
263 "low signal-to-noise ratio in immunofluorescence and detected multiple non-specific bands in
264 western-blot" (communication from Abcam). The use of MVRL56/1 for analysis in the same
265 tissue has sometimes led to opposite findings. For instance, using normal breast tissue, Catena *et*
266 *al* [20] found no VEGFAxxx staining while Qiu *et al* [75] observed strong staining. When used
267 in western-blot studies, MVRL56/1 yields multiple bands (i.e. smear) that span a broad range of
268 molecular weights. These bands have been considered to represent monomers, dimers or large
269 complexes of VEGFAxxx isoforms [36,37]. However, definitive evidence that these bands
270 correspond to various forms of VEGFAxxx is still lacking.

271

272 As summarized in **Table S1**, virtually all studies aimed at detection of VEGFA_{xxx}b proteins
273 have relied on the use of the MVRL56/1 antibody. Origene Europe also sells a different antibody
274 against hVEGFA165b C-Terminal peptide (the exact sequence of the epitope is not provided;
275 #DM3615P). Finally, an "Anti-VEGFA111b antibody" has been raised [76] using "synthetic
276 peptide fragments of the 8 amino acids CRSLTRKD". Contrary to what the authors claim, this
277 antibody would not be specific for the VEGFA111b isoform, [since the epitope corresponds to the](#)
278 [8 amino acid C-terminal sequence shared by all predicted human VEGFA_{xxx}b isoforms.](#)

279
280 Surprisingly, the original anti-human VEGFA_{xxx}b antibody (MVRL56/1) has been used to
281 demonstrate the presence of VEGFA_{xxx}b in multiple species that do not share the same C-
282 terminal sequence (see **Figure S1**). For instance, it has been used in sheep [58] for which the
283 predicted sequence for the last 9 amino acids of the C-Terminal would be TCRCLTRKD,
284 therefore slightly divergent from that in human (underlined, Cys instead of Ser). This would
285 reduce the size of the VEGFA_{xxx}b-specific epitope recognized by the anti-VEGFA_{xxx}b
286 antibody to the LTRKD sequence. Having shown that *Vegfaxxxb* transcripts do not exist, we
287 surmised that VEGFA_{xxx}b-like immunostaining is accounted for by cross-reactivity of the
288 antibody with one or several protein(s) unrelated to VEGFA but bearing an epitope of similar
289 sequence to that of the putative LTRKD sequence of VEGFA_{xxx}b. We searched human protein
290 databases using BLASTP, which led to the unambiguous identification of 10 proteins harboring
291 an LTRKD motif [33]. The MVRL56/1 anti-human VEGFA_{xxx}b antibody was also used to
292 detect endogenous VEGFA_{xxx}b isoforms in tissues from mouse [65,75,77] and rat [45,60],
293 species which, as already noted by us and others ([13,28]; see **Figure S1**), would present a
294 distinct CRPLTGKTD motif at the C-term of VEGFA_{xxx}b, divergent from the human sequence.
295 Specifically, we found that in mouse, this antibody detects unidentified proteins that can be
296 mistaken for VEGFA isoforms and have raised this as a serious concern [28]. These non-specific

297 proteins were present in mouse cell extracts and their conditioned media as well as mouse
298 tissues. Although the suggestion by Bates *et al* [50] that these findings might be due to
299 artefactual detection of mouse IgG could potentially be correct when testing mouse tissues, this
300 possibility is excluded in cell culture and serum-free conditioned media. Taken together, we
301 conclude that the anti-VEGFA_{xxx}b antibody might cross-react with a large number of
302 endogenous proteins (>10) across different species but none of these belong to the VEGFA_{xxx}b
303 family.

304

305

306 **Concluding remarks**

307 We have reviewed current evidence for the proposed existence of a *Vegfaxxxb* family of
308 transcripts which would yield VEGFA_{xxx}b proteins bearing anti-angiogenic properties.
309 Inadequate PCR methodology which likely led to the erroneous identification of such transcripts
310 has been identified and characterized. Multiple published studies, along with our additional
311 analyses of RNA-seq data have failed to identify any *Vegfaxxxb*-specific splicing events. Thus
312 we conclude that *Vegfaxxxb* transcripts do not exist *in vivo* and therefore challenge the view that
313 endogenous VEGFA_{xxx}b proteins have any physiological relevance. Consequently, efforts to
314 develop therapeutics to modulate VEGFA activity should not be based upon modification of a
315 splicing event that is not supported by evidence. [The story of *Vegfaxxxb* splicing reinforces the](#)
316 [importance of supporting new findings using orthogonal techniques prior to basing subsequent](#)
317 [studies upon them \(see Outstanding Questions\).](#)

318

319 **Acknowledgements**

320 We thank Stephen Bridgett for assistance with analysis of RNA-seq data.

321

322 **References**

- 323 1 Ferrara, N. (2004) Vascular endothelial growth factor: basic science and clinical progress. *Endocr Rev* 25, 581-611
- 324 2 Takahashi, H. and Shibuya, M. (2005) The vascular endothelial growth factor (VEGF)/VEGF receptor system and
325 its role under physiological and pathological conditions. *Clin Sci (Lond)* 109, 227-241
- 326 3 Harper, S.J. and Bates, D.O. (2008) VEGF-A splicing: the key to anti-angiogenic therapeutics? *Nat Rev Cancer* 8,
327 880-887
- 328 4 Giacca, M. (2010) Non-redundant functions of the protein isoforms arising from alternative splicing of the VEGF-
329 A pre-mRNA. *Transcription* 1, 149-153
- 330 5 Gitay-Goren, H. *et al.* (1996) Selective binding of VEGF121 to one of the three vascular endothelial growth factor
331 receptors of vascular endothelial cells. *J Biol Chem* 271, 5519-5523
- 332 6 Peach, C.J. *et al.* (2018) Real-Time Ligand Binding of Fluorescent VEGF-A Isoforms that Discriminate between
333 VEGFR2 and NRP1 in Living Cells. *Cell Chem Biol* 25, 1208-1218.e5
- 334 7 Apte, R.S. *et al.* (2019) VEGF in Signaling and Disease: Beyond Discovery and Development. *Cell* 176, 1248-
335 1264
- 336 8 Bates, D.O. *et al.* (2002) VEGF165b, an inhibitory splice variant of vascular endothelial growth factor, is down-
337 regulated in renal cell carcinoma. *Cancer Res* 62, 4123-4131
- 338 9 Qiu, Y. *et al.* (2009) The anti-angiogenic isoforms of VEGF in health and disease. *Biochem Soc Trans* 37, 1207-
339 1213
- 340 10 Woolard, J. *et al.* (2009) Molecular diversity of VEGF-A as a regulator of its biological activity.
341 *Microcirculation* 16, 572-592
- 342 11 Peiris-Pages, M. (2012) The role of VEGF 165b in pathophysiology. *Cell Adh Migr* 6, 561-568
- 343 12 Biselli-Chicote, P.M. *et al.* (2012) VEGF gene alternative splicing: pro- and anti-angiogenic isoforms in cancer. *J*
344 *Cancer Res Clin Oncol* 138, 363-370
- 345 13 Guyot, M. and Pagès, G. (2015) VEGF splicing and the role of VEGF splice variants: from physiological-
346 pathological conditions to specific pre-mRNA splicing. (Springer Science Business Media New York 2015) *VEGF*
347 *Signaling: Methods and Protocols, Methods in Molecular Biology, Vol 1332* 3-23
- 348 14 Bates, D.O. *et al.* (2018) Physiological Role of Vascular Endothelial Growth Factors as Homeostatic Regulators.
349 *Compr Physiol* 8, 955-979
- 350 15 Woolard, J. *et al.* (2004) VEGF165b, an inhibitory vascular endothelial growth factor splice variant: mechanism
351 of action, in vivo effect on angiogenesis and endogenous protein expression. *Cancer Res* 64, 7822-7835
- 352 16 Cebe Suarez, S. *et al.* (2006) A VEGF-A splice variant defective for heparan sulfate and neuropilin-1 binding
353 shows attenuated signaling through VEGFR-2. *Cell Mol Life Sci* 63, 2067-2077
- 354 17 Kawamura, H. *et al.* (2008) Vascular endothelial growth factor (VEGF)-A165b is a weak in vitro agonist for
355 VEGF receptor-2 due to lack of coreceptor binding and deficient regulation of kinase activity. *Cancer Res* 68, 4683-
356 4692
- 357 18 Hua, J. *et al.* (2010) Recombinant human VEGF165b inhibits experimental choroidal neovascularization. *Invest*
358 *Ophthalmol Vis Sci* 51, 4282-4288

- 359 19 Delcombel, R. *et al.* (2013) New prospects in the roles of the C-terminal domains of VEGF-A and their
360 cooperation for ligand binding, cellular signaling and vessels formation. *Angiogenesis* 16, 353-371
- 361 20 Catena, R. *et al.* (2010) VEGF(1)(2)(1)b and VEGF(1)(6)(5)b are weakly angiogenic isoforms of VEGF-A. *Mol*
362 *Cancer* 9, 320-4598-9-320
- 363 21 Boudria, A. *et al.* (2018) VEGF165b, a splice variant of VEGF-A, promotes lung tumor progression and escape
364 from anti-angiogenic therapies through a beta1 integrin/VEGFR autocrine loop. *Oncogene* 38-1050-1066
- 365 22 Eswarappa, S.M. *et al.* (2014) Programmed translational readthrough generates antiangiogenic VEGF-Ax. *Cell*
366 157, 1605-1618
- 367 23 Xin, H. *et al.* (2016) Evidence for pro-angiogenic functions of VEGF-ax. *Cell* 167, 275-284.
- 368 24 Guyot, M. *et al.* (2017) Targeting the pro-angiogenic forms of VEGF or inhibiting their expression as anti-cancer
369 strategies. *Oncotarget* 8, 9174-9188
- 370 25 Varey, A.H. *et al.* (2008) VEGF 165 b, an antiangiogenic VEGF-A isoform, binds and inhibits bevacizumab
371 treatment in experimental colorectal carcinoma: balance of pro- and antiangiogenic VEGF-A isoforms has
372 implications for therapy. *Br J Cancer* 98, 1366-1379
- 373 26 Ferrara, N. (2002) VEGF and the quest for tumour angiogenesis factors. *Nat Rev Cancer* 2, 795-803
- 374 27 Ferrara, N. *et al.* (2003) The biology of VEGF and its receptors. *Nat Med* 9, 669-676
- 375 28 Diabetic Retinopathy Clinical Research Network *et al.* (2015) Aflibercept, bevacizumab, or ranibizumab for
376 diabetic macular edema. *N Engl J Med* 372, 1193-1203
- 377 29 Rennel, E.S. *et al.* (2009) VEGF(121)b, a new member of the VEGF(xxx)b family of VEGF-A splice isoforms,
378 inhibits neovascularisation and tumour growth in vivo. *Br J Cancer* 101, 1183-1193
- 379 30 Mackenzie, F. and Ruhrberg, C. (2012) Diverse roles for VEGF-A in the nervous system. *Development* 139,
380 1371-1380
- 381 31 Harris, S. *et al.* (2012) Do anti-angiogenic VEGF (VEGFxxx) isoforms exist? A cautionary tale. *PLoS One* 7,
382 e35231
- 383 32 Bridgett, S. *et al.* (2017) RNA-Sequencing data supports the existence of novel VEGFA splicing events but not
384 of VEGFAxxx isoforms. *Sci Rep* 7, 58-017-00100-3
- 385 33 Lomet, D. *et al.* (2018) Anti-angiogenic VEGFAxxx transcripts are not expressed in the medio-basal
386 hypothalamus of the seasonal sheep. *PLoS One* 13, e0197123
- 387 34 Chesnokov, M.S. *et al.* (2018) Shift in VEGFA isoform balance towards more angiogenic variants is associated
388 with tumor stage and differentiation of human hepatocellular carcinoma. *Peerj* 6, e4915
- 389 35 Cui, T.G. *et al.* (2004) Differentiated human podocytes endogenously express an inhibitory isoform of vascular
390 endothelial growth factor (VEGF165b) mRNA and protein. *Am J Physiol Renal Physiol* 286, F767-73
- 391 36 Perrin, R.M. *et al.* (2005) Diabetic retinopathy is associated with a switch in splicing from anti- to pro-angiogenic
392 isoforms of vascular endothelial growth factor. *Diabetologia* 48, 2422-2427
- 393 37 Bates, D.O. *et al.* (2006) The endogenous anti-angiogenic family of splice variants of VEGF, VEGFxxx, are
394 down-regulated in pre-eclamptic placentae at term. *Clin Sci (Lond)* 110, 575-585

- 395 38 Pritchard-Jones, R.O. *et al.* (2007) Expression of VEGF(xxx)b, the inhibitory isoforms of VEGF, in malignant
396 melanoma. *Br J Cancer* 97, 223-230
- 397 39 Rennel, E. *et al.* (2008) The endogenous anti-angiogenic VEGF isoform, VEGF165b inhibits human tumour
398 growth in mice. *Br J Cancer* 98, 1250-1257
- 399 40 Diaz, R. *et al.* (2008) p73 Isoforms affect VEGF, VEGF165b and PEDF expression in human colorectal tumors:
400 VEGF165b downregulation as a marker of poor prognosis. *Int J Cancer* 123, 1060-1067
- 401 41 Manetti, M. *et al.* (2011) Overexpression of VEGF165b, an inhibitory splice variant of vascular endothelial
402 growth factor, leads to insufficient angiogenesis in patients with systemic sclerosis. *Circ Res* 109, e14-26
- 403 42 Ribeiro, L.A. *et al.* (2007) Characterization and differential expression of vascular endothelial growth factor
404 isoforms and receptors in swine corpus luteum throughout estrous cycle. *Mol Reprod Dev* 74, 163-171
- 405 43 Baba, T. *et al.* (2012) VEGF 165 b in the developing vasculatures of the fetal human eye. *Dev Dyn* 241, 595-607
- 406 44 Miller-Kasprzak, E. and Jagodzinski, P.P. (2008) 5-Aza-2'-deoxycytidine increases the expression of anti-
407 angiogenic vascular endothelial growth factor 189b variant in human lung microvascular endothelial cells. *Biomed*
408 *Pharmacother* 62, 158-163
- 409 45 Artac, R.A. *et al.* (2009) Neutralization of vascular endothelial growth factor antiangiogenic isoforms is more
410 effective than treatment with proangiogenic isoforms in stimulating vascular development and follicle progression in
411 the perinatal rat ovary. *Biol Reprod* 81, 978-988
- 412 46 Merdzhanova, G. *et al.* (2010) The transcription factor E2F1 and the SR protein SC35 control the ratio of pro-
413 angiogenic versus antiangiogenic isoforms of vascular endothelial growth factor-A to inhibit neovascularization in
414 vivo. *Oncogene* 29, 5392-5403
- 415 47 Xu, J. *et al.* (2011) The evolution of alternative splicing exons in vascular endothelial growth factor A. *Gene* 487,
416 143-150
- 417 48 Clifford, R.L. *et al.* (2012) Abnormal histone methylation is responsible for increased vascular endothelial
418 growth factor 165a secretion from airway smooth muscle cells in asthma. *J Immunol* 189, 819-831
- 419 49. Ganta, V.C. *et al.* (2019) Antiangiogenic VEGF165b Regulates Macrophage Polarization via S100A8/S100A9
420 in Peripheral Artery Disease. *Circulation* 139, 226-242
- 421 50 Bates, D.O. *et al.* (2013) Detection of VEGF-A(xxx)b isoforms in human tissues. *PLoS One* 8, e68399
- 422 51 Kikuchi, R. *et al.* (2014) An antiangiogenic isoform of VEGF-A contributes to impaired vascularization in
423 peripheral artery disease. *Nat Med* 20, 1464-1471
- 424 52 Oltean, S. *et al.* (2015) Vascular Endothelial Growth Factor-A165b Is Protective and Restores Endothelial
425 Glycocalyx in Diabetic Nephropathy. *J Am Soc Nephrol* 26, 1889-1904
- 426 53 Barash, Y. *et al.* (2013) AVISPA: a web tool for the prediction and analysis of alternative splicing. *Genome Biol*
427 14, R114-2013-14-10-r114
- 428 54 Reese, M.G. *et al.* (1997) Improved splice site detection in Genie. *J Comput Biol* 4, 311-323
- 429 55 Raj, B. and Blencowe, B.J. (2015) Alternative Splicing in the Mammalian Nervous System: Recent Insights into
430 Mechanisms and Functional Roles. *Neuron* 87, 14-27
- 431 56 Lomet, D. *et al.* (2018) The impact of thyroid hormone in seasonal breeding has a restricted transcriptional
432 signature. *Cell Mol Life Sci* 75, 905-919

- 433 57 Wood, S.H. *et al.* (2015) Binary Switching of Calendar Cells in the Pituitary Defines the Phase of the Circannual
434 Cycle in Mammals. *Curr Biol* 25, 2651-2662
- 435 58 Castle-Miller, J. *et al.* (2017) Mechanisms regulating angiogenesis underlie seasonal control of pituitary function.
436 *Proc Natl Acad Sci U S A* 114, E2514-E2523
- 437 59 Konopatskaya, O. *et al.* (2006) VEGF165b, an endogenous C-terminal splice variant of VEGF, inhibits retinal
438 neovascularization in mice. *Mol Vis* 12, 626-632
- 439 60 Ergorul, C. *et al.* (2008) Levels of vascular endothelial growth factor-A165b (VEGF-A165b) are elevated in
440 experimental glaucoma. *Mol Vis* 14, 1517-1524
- 441 61 Bevan, H.S. *et al.* (2008) The alternatively spliced anti-angiogenic family of VEGF isoforms VEGF_{xxx}b in
442 human kidney development. *Nephron Physiol* 110, 57-67
- 443 62 Rennel, E.S. *et al.* (2008) Recombinant human VEGF165b protein is an effective anti-cancer agent in mice. *Eur*
444 *J Cancer* 44, 1883-1894
- 445 63 Bills, V.L. *et al.* (2009) Failure to up-regulate VEGF165b in maternal plasma is a first trimester predictive
446 marker for pre-eclampsia. *Clin Sci (Lond)* 116, 265-272
- 447 64 Magnussen, A.L. *et al.* (2010) VEGF-A165b is cytoprotective and antiangiogenic in the retina. *Invest*
448 *Ophthalmol Vis Sci* 51, 4273-4281
- 449 65 Zhao, M. *et al.* (2011) Expression of pro- and anti-angiogenic isoforms of VEGF in the mouse model of oxygen-
450 induced retinopathy. *Exp Eye Res* 93, 921-926
- 451 66 Oltean, S. *et al.* (2012) VEGF165b overexpression restores normal glomerular water permeability in VEGF164-
452 overexpressing adult mice. *Am J Physiol Renal Physiol* 303, F1026-36
- 453 67 Qiu, Y. *et al.* (2012) Ovarian VEGF(165)b expression regulates follicular development, corpus luteum function
454 and fertility. *Reproduction* 143, 501-511
- 455 68 Beazley-Long, N. *et al.* (2013) VEGF-A165b is an endogenous neuroprotective splice isoform of vascular
456 endothelial growth factor A in vivo and in vitro. *Am J Pathol* 183, 918-929
- 457 69 Hulse, R.P. *et al.* (2015) Vascular endothelial growth factor-A165b prevents diabetic neuropathic pain and
458 sensory neuronal degeneration. *Clin Sci (Lond)* 129, 741-756
- 459 70 Vencappa, S. *et al.* (2015) Cisplatin induced sensory neuropathy is prevented by vascular endothelial growth
460 factor-A. *Am J Transl Res* 7, 1032-1044
- 461 71 Hulse, R.P. *et al.* (2016) The control of alternative splicing by SRSF1 in myelinated afferents contributes to the
462 development of neuropathic pain. *Neurobiol Dis* 96, 186-200
- 463 72 Ved, N. *et al.* (2017) Vascular endothelial growth factor-A165b ameliorates outer-retinal barrier and vascular
464 dysfunction in the diabetic retina. *Clin Sci (Lond)* 131, 1225-1243
- 465 73 Pruszko, M. *et al.* (2017) The mutant p53-ID4 complex controls VEGFA isoforms by recruiting lncRNA
466 MALAT1. *EMBO Rep* 18, 1331-1351
- 467 74 Blochowiak, K.J. *et al.* (2018) Expression of VEGF(1)(6)(5)b, VEGFR1, VEGFR2 and CD34 in benign and
468 malignant tumors of parotid glands. *Adv Clin Exp Med* 27, 83-90
- 469 75 Qiu, Y. *et al.* (2008) Mammary alveolar development during lactation is inhibited by the endogenous
470 antiangiogenic growth factor isoform, VEGF165b. *Faseb j* 22, 1104-1112

- 471 76 Gu, F. *et al.* (2013) VEGF111b, a new member of VEGFxxx isoforms and induced by mitomycin C, inhibits
472 angiogenesis. *Biochem Biophys Res Commun* 441, 18-24
- 473 77 Shi, X. *et al.* (2013) Inhibition of neovascularization and expression shift of pro-/anti-angiogenic vascular
474 endothelial growth factor isoforms after intravitreal bevacizumab injection in oxygen-induced-retinopathy mouse
475 model. *Chin Med J (Engl)* 126, 345-352
- 476 78 Bustin, S.A. *et al.* (2009) The MIQE guidelines: minimum information for publication of quantitative real-time
477 PCR experiments. *Clin Chem* 55, 611-622
- 478 79 Hulse, R.P. *et al.* (2014) Regulation of alternative VEGF-A mRNA splicing is a therapeutic target for analgesia.
479 *Neurobiol Dis* 71, 245-259
- 480 80 Gustafsson, T. *et al.* (2005) VEGF-A splice variants and related receptor expression in human skeletal muscle
481 following submaximal exercise. *J Appl Physiol (1985)* 98, 2137-2146
- 482 81 Dokun, A.O. and Annex, B.H. (2011) The VEGF165b "ICE-o-form" puts a chill on the VEGF story. *Circ Res*
483 109, 246-247
- 484 82 Irimia, M. and Blencowe, B.J. (2012) Alternative splicing: decoding an expansive regulatory layer. *Curr Opin*
485 *Cell Biol* 24, 323-332
- 486 83 Braunschweig, U. *et al.* (2013) Dynamic integration of splicing within gene regulatory pathways. *Cell* 152, 1252-
487 1269
- 488 84 Wang, B. *et al.* (2019) Reviving the Transcriptome Studies: An Insight Into the Emergence of Single-Molecule
489 Transcriptome Sequencing. *Front Genet* 10, 384
- 490 85 Workman, R.E. *et al.* (2018) Nanopore native RNA sequencing of a human poly(A) transcriptome. *BioRxiv Cold*
491 *Spring Harbor Laboratory*. <https://doi.org/10.1101/459529>.
- 492 86 Buckley, P.T. *et al.* (2014) Cytoplasmic intron retention, function, splicing, and the sentinel RNA hypothesis.
493 *Wiley Interdiscip Rev RNA* 5, 223-230
- 494 87 Jacob, A.G. and Smith, C.W.J. (2017) Intron retention as a component of regulated gene expression programs.
495 *Hum Genet* 136, 1043-1057
- 496 88 English, W.R. *et al.* (2017) Differential expression of VEGFA isoforms regulates metastasis and response to anti-
497 VEGFA therapy in sarcoma. *Cancer Res.* 77, 2633-46.

498

499 **Glossary**

500 **Alternative splicing:** process by which splice sites in precursor (pre)-mRNA are differentially
501 selected to produce multiple mRNA and protein isoforms. This process diversifies the functional
502 characteristics of genes and drastically expands the potential repertoire of protein variants.
503 Nearly all multi-exon genes have at least one splice variant.

504 **VEGFA** (Vascular Endothelial Growth Factor A): a key member of the family of growth factors
505 which plays a prominent role in angiogenesis - the growth of blood vessels - both in health and
506 disease (tumorigenesis and retinopathies).

507 **5'-tailing**: addition of nucleotides on the 5'-end of a PCR primer. These nucleotides are not
508 complementary to the target mRNA but do not impair PCR processivity and aid further cloning
509 of the PCR amplicon.

510

511

512

513 **Text BOXES**

514 **BOX1**

515 **Further technical issues with the use of qRT-PCR**

516 Since primers for standard PCR were not suitable for qPCR, Bates and co-workers developed an
517 alternative method, which we might call "indirect subtractive strategy" [25]. This requires the
518 use of two distinct primer pairs. The first pair consists of "an exon 7b forward primer and a 3'-
519 UTR Primer (both pan-VEGF)", the sequences of which were not provided. The second pair of
520 primers consists of "an exon 7a forward primer and a reverse primer specific for exon 8a that do
521 not detect *Vegfaxxb* isoforms". The authors then assume that subtracting what is found with the
522 second primer pair (*Vegfaxx* only) to what is found with the first primer pair (total *Vegfa*:
523 *Vegfaxx* + *Vegfaxxb*) yields the amount of *Vegfaxxb*. This is an unusual strategy, which is
524 likely to be flawed for a number of reasons. The primary concern is that these are two different
525 pairs of primers, with different sequences, hence divergent optimal T_m 's, which amplify
526 fragments of different lengths and nucleotide composition and therefore have differing
527 efficiencies. The use of qPCR is based on the principle that the number of copies doubles at
528 each cycle. This rule is used as a proxy to determine the initial number of copies. Using the

529 values of the slopes for the calibration curves (see Figure 1 in [25]) for both primer pairs allows
530 calculation of an amplification factor (which should ideally be ~2), which informs on the
531 efficiency of the qPCR assay. The efficiency should be ~100%, but values between 85-115% are
532 usually judged acceptable, providing the dissociation curves are good [78]. The calculated
533 efficiency values for the primer pairs used by Varey *et al* [25] are 83.7% and 71.8%. The
534 meaning of this is twofold. First, neither primer pair is appropriate for use in qPCR. Second,
535 different efficiencies are enough to generate a difference when data obtained with the two primer
536 pairs are compared. A broadly similar technique, but with different primers, was used by the
537 same team in a later study [79]. Other issues with RT-PCR and primer design were also found,
538 which are detailed in **Supplemental text**.

539 **BOX2**

540 **Ruling out potential issues with mRNA secondary structure**

541 The idea that PCR amplification of *Vegfa165b* might be difficult due to secondary structure of
542 the mRNA in the 3'-UTR has been invoked [9] to explain why we [31,33] and other authors
543 [13,79,80] failed to detect these transcripts. The existence of an unusually stable secondary
544 structure seems unlikely because most regulatory splicing mechanisms are based on the
545 recognition of short degenerate RNA motifs at the exon/intron boundaries rather than secondary
546 structures [82,83]. Indeed, bioinformatic analysis of the ribonucleotidic sequence does not
547 predict formation of potential hairpins that might inhibit reverse transcription (data not shown).
548 Furthermore, classical *Vegfaxxx* isoforms are readily amplified using primers spanning the
549 putative alternative splice site (see main text). There is therefore no evidence to support the
550 notion that *Vegfaxxxb* isoforms, which would lack a 66bp stretch present in *Vegfaxxx* isoforms,
551 would resist PCR amplification. Furthermore, *Vegfa165b* transcripts expressed from a
552 recombinant construct were readily detected by RNA-seq [32] and since the library preparation
553 involved both RT and PCR steps, neither are inherently blocked by the transcript secondary

554 structure. Therefore, the inability to detect an abundant mRNA such as *Vegfa165b* at least in
555 normal tissues by classical RT-PCR can not be explained by the secondary structure of the
556 mRNA.

557 **BOX3**

558 **Further RNA-seq evidence for absence of *Vegfaxxb* mRNAs**

559 Recently developed long read RNA sequencing technologies (e.g. PacBio Iso-Seq and Oxford
560 Nanopore Technology) can measure full length mRNA transcripts [85]. Both individual
561 alternative splicing events and the co-occurrence of multiple events in transcripts such as *Vegfa*
562 are detected. To determine whether data generated with this newly available approach could
563 detect *Vegfaxxb* mRNAs we analysed a publicly available dataset [85]. Full length transcripts
564 encoding all the commonly reported exon 5 and exon 6 isoforms (VEGFA121, VEGFA165,
565 VEGFA189) were present but no evidence of splicing to generate *Vegfaxxb* was detected
566 (**Supplemental Figure S2**). A notable feature apparent from this full length transcript data is the
567 retention of the intron between exons 3 and 4 in almost 20% of transcripts (which is supported
568 by previous EST data). The regulatory potential of such intron retention events has been widely
569 discussed [86,87]. Our recent analysis of RNA-seq data from tumours grown from fibrosarcomas
570 expressing VEGFA188 [88] has also failed to identify reads corresponding to the exon 7-8b
571 splice site, even though endogenous expression of multiple *Vegfaxxx* isoforms could be detected
572 from the stromal cells (**Supplemental Figure S3**). To better understand the role of alternative
573 splicing in the regulation of *Vegfa* expression, research should be focused upon such events that
574 are supported by strong experimental evidence.

575

576 **Figure Legends**

577 **Figure 1:** Structure and sequence of the human *Vegfa* gene and VEGFA_{xxx}/VEGFA_{xxxb}
578 isoforms. A/ Schematic of the human *Vegfa* gene locus. The gene is comprised of 8 exons. B/

579 Schematic of the longest human VEGFA protein (VEGF206, see GenBank accession number
580 NM_001171623). The stop-codon (TGA, in red) is located in exon 8 and the sequence of the last
581 9 amino acids at the C-terminus of the protein is provided below. The location of flanking PCR
582 primers O14I/O16I is also indicated (see panel F). C/ Schematic of the putative human
583 VEGFA206b protein (GenBank accession number NM_001033756). The stop-codon (TGA, in
584 red) is located in exon 8 – renamed exon 8b due to the usage of a distal splice site – and the
585 sequence of the last 9 amino acids at the C-terminus of the protein is provided below. Note that
586 only the sequence of the last 6 amino acids at the C-terminus differ between VEGFA and
587 VEGFAb isoforms. D/ Nucleotide sequence (from nt 1716 to nt 1808) of the 5'-end of human
588 *Vegfa* exon 8 and deduced amino acid sequences corresponding to the C-term of VEGFAxxx and
589 VEGFAxxx isoforms ; corresponding nucleotide triplets are underlined. The reference sequence
590 used for nucleotide annotation is Genbank NM_001171623. The usual splice site (mauve and
591 underlined) leading to VEGFAxxx isoforms (as shown in B) and the putative alternative splice
592 site (green and underlined) leading to VEGFAxxx isoforms (as shown in C) are shown. The
593 respective stop codons are also shown (red and bold). Also note that exon 8 and exon 8b would
594 share an AT dinucleotide sequence at their 5'-end (grey boxes and bold). E/ Schematic of the
595 four most abundant classical hVEGFA splicing isoforms and of their corresponding sister
596 isoforms of the proposed hVEGFAxxx family : both families differ only by the alternative use
597 of either exon 8a or 8b, as described in panels B and C using VEGFA206 as an example.
598 Adapted from Bridgett *et al* [32]. F/ Agarose gel electrophoresis of PCR products obtained using
599 flanking primers O14I/O16I and cDNA from ovine medio-basal hypothalamus (see panel B for
600 location). Gel extraction, cloning and sequencing of the three bands revealed products encoding
601 ovine homologs of human VEGFA206 and VEGFA189 (562 and 544bp ; sizes too close to be
602 separated on the gel), VEGFA165 (490bp) and VEGFA121 (358bp). No single VEGFAxxx
603 PCR product was obtained. For further information, see Lomet *et al.* [33]. G/ Schematic of the

604 signaling pathway elicited by VEGFA isoforms. Classical VEGFA proteins bind both VEGFR2
605 homodimer and the Neuropilin1 (NRP1) co-receptor while VEGFA_{xxx}b isoforms fail to bind
606 NRP1, impairing intracellular transduction pathways.

607
608 **Figure 2:** Issues with RT-PCR primer design: how 5'-tailing led to erroneous identification of
609 *Vegfaxxxb* transcripts. A/ Schematic of the putative junction between exon 7 (blue) and the
610 putative exon 8b (green and black) of *Vegfaxxxb* transcripts and location of the PCR primers
611 used to identify *Vegfaxxxb* transcripts. Note that exon 7 and the AT dinucleotide (green) at the
612 end of exon 8 sequences would be common to both classical *Vegfaxxx* and *Vegfaxxxb* isoforms.
613 The number of non-isoform specific nucleotides is provided in blue; note that this number has
614 substantially increased (from 7bp to 17bp) throughout the years (primers P1 through to P7).
615 Sequences in orange represent 5'-tailing. Primers P1-P6 are reverse primers while primer P7 is a
616 forward primer, as indicated by the arrows. B/ Minimal sequence requirement to obtain PCR
617 amplification of ovine *Vegfaxxxb*-like transcripts through 5'-tailing as defined in Lomet *et al*
618 [33]. The error-prone primer consists of 11 nucleotides in exon 7 and 2-3 nucleotides of exon 8
619 (in green, shared by isoforms), then any nucleotide can be added on the 5'-end of the primer
620 providing the number is sufficient (6 or more) to reach a T_m compatible with the PCR
621 methodology (see **Figure 3**). Note that this primer design corresponds to primer P4.

622
623 **Figure 3:** Basic principles of PCR and how 5'-tailing is achieved. A/ Principle of standard PCR.
624 After reverse transcription, cDNAs are submitted to 30-40 cycles of denaturation/annealing and
625 extension during which a pair of primers (forward and reverse) allow efficient amplification of
626 the target sequence they flank. B/ To anneal to the cDNA, the melting temperature (T_m) of the
627 primer must be higher than the annealing temperature. C/ The T_m of the primer depends on its
628 nucleotide composition and its length, with long GC-rich primers having higher T_ms than short

629 AT-rich primers. A series of 3 efficient primers is shown, the one on the top is the shortest. D/
630 The shortest efficient reverse primer as defined in C can be flanked on its 5'-end with a large
631 stretch of additional non-specific extra-bases (for instance restriction sites for further cloning
632 purposes), which do not anneal to the target and do not impair PCR efficiency and processivity:
633 this is 5'-tailing. E/ The additional non-specific 5'-tail is added to the amplicon as the PCR
634 progresses. At the end of the 30-40 cycles, all amplicons are flanked with the additional, non-
635 specific stretch of the reverse primer.

636

637 **Supplementary Material**

638 Includes Supplemental Text, Supplemental table S1, Supplemental Figures S1-S3

Authors	Journal	Issue:Page	Year	Species/Tissue	PCR	IHC WB ELISA	Recomb. prot Over-exp
Bates <i>et al</i>	<i>Cancer Res</i>	62:4123-4131	2002	Human / Kidney	VEGF165b exon 8b/ 7 - Reverse primer : 5'-TCAGTCTTTCCTGGTGAGAGATCTGCA-3' ----- Amplification of full-length VEGF165b with a reverse primer located downstream in the 3'UTR.		Home made
Cui <i>et al</i>	<i>Am J Physiol Renal Physiol</i>	286:F767-773	2004	Human / Kidney	Same primers as Bates <i>et al</i> 2002.		
Woolard <i>et al</i>	<i>Cancer Res</i>	64:7822-7835	2004	Various	Same primers as Bates <i>et al</i> 2002.	Production monoclonal antibody hVEGF165b	Home made
Perrin <i>et al</i>	<i>Diabetologia</i>	48:2422-2427	2005	Human / Eye	Same primers as Bates <i>et al</i> 2002.	Home made ELISA	
Konopatskaya <i>et al</i>	<i>Mol Vis</i>	12:626-632	2006	Mouse / Eye	No		3045-VE-025
Bates <i>et al</i>	<i>Clin Sci (Lond)</i>	110:575-585	2006	Human / Placenta	Same primers as Bates <i>et al</i> 2002.	MAB3045 Home made ELISA	3045-VE-025
Cebe Suarez <i>et al</i>	<i>Cell Mol Life Sci</i>	63:2067-2077	2006	Human cell line (HUVEC)	No	MAB3045	Home made
Ribeiro <i>et al</i>	<i>Mol Reprod Dev</i>	74:163-171	2007	Swine / Ovary	VEGF164b exon 8b/ 7 - Reverse primer : 5'-TCCTGGTGAGAGATCTGCAAG-3' Note : level << VEGF164 (2 ¹² fold)		
Pritchard-Jones <i>et al</i>	<i>Br J Cancer</i>	97:223-230	2007	Human / Melanoma tissue	VEGF165b HindIII RS - exon 8b/ 7 - Reverse primer : 5'-TTAAGCTTTCAGTCTTTCCTGGTGAGACTGCA-3' Note : GAT missing from oligo ; see Bates <i>et al</i> 2002 Invalid primer.	MAB3045	
Schumacher <i>et al</i>	<i>J Am Soc Nephrol</i>	18:719-729	2007	Human / Kidney	VEGF both isoforms BamHI RS – 3'UTR Reverse primer : 5'-ATGGATCCGTATCAGTCTTTCCT-3' Note : short primer, theoretical fusion T° = 41°C		
Ergorul <i>et al</i>	<i>Mol Vision</i>	14:1517-1524	2008	Rat / Retina	No	Ab14994	
Qiu <i>et al</i>	<i>Faseb J</i>	22:1104-1112	2008	Mouse / Various tissues	Primers use to « detect specifically the transgene » Forward : 5'-TCAGCGCAGCTACTGCCATC-3' Reverse : 5'-GTGCTGGCCTTGGTGAGGTT-3' Note : Forward is within exon3 and Reverse is within exon4 → actually detect all Vegfa.	MAB3045	Yes – TG mice

					Invalid primers → Invalid screening of TG mice.		
Bevan <i>et al</i>	<i>Nephron Physiol</i>	110:57-67	2008	Human / Kidney	No	MAB3045	
Kawamura <i>et al</i>	<i>Cancer Res</i>	68:4683-4692	2008	Cell lines	No		Home made
Varey <i>et al</i>	<i>Br J Cancer</i>	98:1366-1379	2008	Human / Colon	Original indirect subtractive strategy : Use of « An exon 7b forward primer and a 3'UTR Primer (both pan-VEGF) » - Sequences not provided Then use of « an exon 7a forward primer and a reverse primer specific for exon 8a that did not detect Vegfxxx isoforms ». Then subtraction would yield Vegfxxx. Invalid method.	MAB3045	Yes
Diaz <i>et al</i>	<i>Int J Cancer</i>	123 :1060-1067	2008	Human / Colon	Same primers as Bates <i>et al</i> 2002.	MRVL56/1	
Nowak <i>et al</i>	<i>J Cell Sci</i>	121:3487-3495	2008	Cell lines	VEGF both isoforms BamHI RS – 3'UTR 5'-ATGGATCCGTATCAGTCTTCTCTGG-3'	MAB3045 & clone 264610/1	Home made
Rennel <i>et al</i>	<i>Br J Cancer</i>	98:1250-1257	2008	Human / Prostate Mouse	VEGF165b HindIII RS - exon 8b/ 7 5'-TTAAGCTTTCAGTCTTCTCTGGTGAGAGATCTGCA-3'	Home made ELISA	Home made
Rennel <i>et al</i>	<i>Eur J Cancer</i>	44:1883-1894	2008	Cell lines Mouse	No	MAB3045	Home made
Miller-Kasprzak and Jagodinski	<i>Biomed Pharmacother</i>	62:158-163	2008	Human / Lung cell line	VEGF121b exon 5/8b - Forward primer : 5'-GAAAATCTCTCACCAGGAAA-3' Note : one A missing from oligo. VEGF165b exon 8b/ 7 - Reverse primer : 5'-GTGAGAGATCTGCAAGTACG-3'	MAB3045	
Artac <i>et al</i>	<i>Biol Reprod</i>	81:978-988	2009	Rat / Ovary	VEGF165b exon 8b/ 7 - Reverse primer : 5'-GGTGAGAGGCTGCAAGTACGTT-3'	MAB3045	
Bills <i>et al</i>	<i>Clin Sci (Lond)</i>	116:265-272	2009	Human / Plasma	No	MAB3045 Home made ELISA	
Rennel <i>et al</i>	<i>Br J Cancer</i>	101:1183-1193	2009	Human / Colon	VEGF121b exon 5/8b - Forward primer : 5'-GAAAATCTCTCACCAGGAAA-3'	MAB3045	DY3045
Baltes-Breitwisch <i>et al</i>	<i>Reproduction</i>	140:319-329	2010	Rat / Testes	Same primers as Artac <i>et al</i> . 2009.	Ab14994	
Catena <i>et al</i>	<i>Molecular Cancer</i>	9 :320	2010	Human cell lines	VEGF121b – HindIII RS – exon 8b/ exon5 Reverse primer : 5'-TTAAGCTTTCAGTCTTCTCTGGTGAGAGATTTTCTTGTCTTGCTCTATC - 3' VEGF165b – HindIII RS – exon 8b/ exon7 Reverse primer : 5'-TTAAGCTTTCAGTCTTCTCTGGTGAGAGATCTGCAAGTACGTTCTGTTAACTC - 3'	MAB3045	Yes

Hua <i>et al</i>	<i>Invest Ophthalmol Vis Sci</i>	51:4282-4288	2010	Human / Eye	No	MAB3045	
Magnussen <i>et al</i>	<i>Invest Ophthalmol Vis Sci</i>	51:4273-4281	2010	Mouse / Eye	No	MAB3045	3045-VE-025
Nowak <i>et al</i>	<i>J Biol Chem</i>	285:5532-5540	2010	Cell lines	Same as Nowak <i>et al</i> 2008	MAB3045	
Peiris-Pagès <i>et al</i>	<i>J.Pathol</i>	222:138-147	2010	Cell lines	Same as Nowak <i>et al</i> 2008	MAB3045	DY3045 3045-VE-025
Qiu <i>et al</i>	<i>J Am Soc Nephrol</i>	21:1498-1509	2010	Cell lines Mouse	Same invalid primers as Qiu <i>et al</i> 2008 to « detect specifically the transgene » → Invalid screening of TG mice.	Home made ELISA & clone 264610/1	TG-mice DY3045
Merdzhanova <i>et al</i>	<i>Oncogene</i>	29: 5392-5403	2010	Mouse & Human Cell lines	VEGF165b exon 8b/ 7 - Reverse primer : 5'- TGGTGAGAGATCTGCAAGTACGTT -3'		
Zhao <i>et al</i>	<i>Exp Eye Res</i>	93:921-926	2011	Mouse / Retina	No	Ab14994	
Amin <i>et al</i>	<i>Cancer Cell</i>	20:768-780	2011	Cell lines	Same primer as Nowak <i>et al</i> 2008.		
Xu <i>et al</i>	<i>Gene</i>	487:143-150	2011	Human-Cat-Rabbit- Rat/ Various tissues	Human : VEGF165b exon 8b/ 7 Reverse primer : 5'- TGGTGAGAGATCTGCAAGTACGTT -3' Cat : VEGF165b exon 8b/ 7 Reverse primer : 5'- TGGTGAGAGGCTGCAAGTACGTT -3' Rabbit : VEGF165b exon 8b/ 7 Reverse primer : 5'- CGGTGAGAGGCTGCAAGTACGTT -3' Rat : VEGF165b exon 8b/ 7 Reverse primer : 5'- GGTGAGAGGCTGCAAGTACGTT -3'		
Manetti <i>et al</i>	<i>Circ Res</i>	109:e14-26	2011	Human / Skin	Same primer as Rennel <i>et al.</i> 2008	Ab14994	
Baba <i>et al</i>	<i>Dev Dyn</i>	241:595-607	2012	Human / Eye	VEGF165b exon 8b/ 7 - Reverse primer : 5'- TCCTGGTGAGAGATCTGCAA -3'	Ab14994	
Oltean <i>et al</i>	<i>Am J Physiol Renal Physiol</i>	303:F1026-1036	2012	Mouse / Kidney	Same invalid primers as Qiu <i>et al</i> 2008 to « detect specifically the transgene » → Invalid screening of TG mice.		TG-mice
Qiu <i>et al</i>	<i>Reproduction</i>	143:501-511	2012	Marmoset / Ovary Mouse	No	Home made ELISA & clone 264610/1	
Clifford <i>et al</i>	<i>J Immunol</i>	189:819-831	2012	Human / Cell lines	VEGF165b exon 8b/ 7 - Reverse primer : 5'- AGAGAGATCTGCAAGTACGTTCCG -3' VEGF189b exon 8b/ 7 - Reverse primer : 5'- GTGAGAGATCTGCAAGTACG -3'		
Bates <i>et al</i>	<i>PLoS One</i>	8:e68399	2013	Human / Cell lines	VEGF165b exon 8b/7 - Reverse primer :	MAB3045	

					5'- GTGAGAGATCTGCAAGTACG -3'		
Beazley-Long <i>et al</i>	<i>Am J Pathol</i>	183:918-929	2013	Human / Brain Rat / Brain	No	MAB3045	3045-VE-025
Gu <i>et al</i>	<i>Biochem Biophys Res Comm</i>	441:18-24	2013	Human / Cell lines	VEGF111b – « Reverse primer » : 5' – AATGCAGATGTGACAAGCCGAG – 3' Actually a forward primer at junction exon4/8a . Invalid → Identification of Vegf111b invalid. VEGF165b HindIII RS - exon 8b/ 7 - Reverse primer : 5'- TTAAGCTTTCAGTCTTTCCTGGTGAGAGATCTGCA -3'	MRVL56/1 & Home Made αVEGF111b	
Delcombel <i>et al</i>	<i>Angiogenesis</i>	16:353-371	2013	HEK293 cells	VEGF111b exon8b/4 – Reverse primer : 5'- TCCTGGTGAGAGATCTGCATTCAC -3' VEGF121b exon8b/5 – Reverse primer : 5'- GTCTTTCCTGGTGAGAGAGTTTCTT -3' Note : primer issue → AT missing between exon8b and exon5. Invalid. VEGF165b exon8b – Reverse primer : 5'- CGATCGTTCTGTATCAGTCTTTCCT -3' Note : internal to exon 8 → PCR product = classical mVEG Invalid.		
Hulse <i>et al</i>	<i>Neurobiol Dis</i>	71:245-259	2014		Strategy similar to Varey <i>et al</i> 2008 but different Primers ; « primers specific for VEGF-A165a » are: Forward primer exon 7: 5'-GTTCAAGCGGAGAAAGCAT-3' Reverse primer exon 8a : 5'- TCACA TCTGCAAGTACGTTCG -3' → This primer covers exon8a/7	Ab14994	DY3045
Kikuchi <i>et al</i>	<i>Nat Med</i>	20:1464-1471	2014	Human	hVEGF165b exon 8b/7 - Reverse primer : 5'- GTGAGAGATCTGCAAGTACG -3' ----- mVEGF165b exon 8b/7 - Reverse primer : 5'- CTTCCGGTGAGAGGTCTGC -3'		3045-VE-025
Hulse <i>et al</i>	<i>Clin Sci</i>	129:741-756	2015	Cell culture Rat	No		3045-VE-025
Oltean <i>et al</i>	<i>J Am Soc Nephrol</i>	26:1889-1904	2015	Human / Kidney	VEGF165b exon 8b/7 - Reverse primer : 5'- GTGAGAGATCTGCAAGTACG -3'	DY3045	
Vencappa <i>et al</i>	<i>Am J Transl Res</i>	7:1032-1044	2015	Cell culture Mouse	No	Ab14994	3045-VE-025
Li <i>et al</i>	<i>J Translat Med</i>	13 :164	2015	Human cell lines	Same primers as Gu <i>et al</i> 2013	Same as	Yes

						Gu <i>et al</i> 2013	
Hulse <i>et al</i>	<i>Neurobiol Dis</i>	96:186-200	2016	Rat	No	Ab14994	
Castle-Miller <i>et al</i>	<i>Proc Natl Acad Sci USA</i>	114:E2517-2523	2017	Sheep / Pituitary	Multiple issues with primers leading to an erratum. VEGF _{Axxx} -specific reverse primer : 5'-CGGCGGCTATGGGTCGTTCTGTGTCAGTCTTTCCT GGTGAGAC-3' 43bp. Unusual. Targets a sequence downstream of putative distal splice located in exon 8. Non-isoform specific. ➔ qRT-PCR data invalid.	Home made Clone56/8 & DY3045	
Ved <i>et al</i>	<i>Clin Sci (Lond)</i>	131:1225-1243	2017	Rat / Retina	No	Home made Clone56/8	Yes
Pruszko <i>et al</i>	<i>EMBO reports</i>	18:1331-1351	2017	Human cell lines	VEGF121b exon 8b/5 - Reverse primer : 5'- CTTTCCTGGTGAGAGATTTTCTTGTC -3' VEGF165b exon 8b/7 - Reverse primer : 5'- CCTGGTGAGAGATCTGCAAGTAC -3'	Ab14994	
Hueso <i>et al</i>	<i>Scientific Reports</i>	7 :9962	2017	Human Serum / Heart	No	MAB3045	MBS109074
Chesnokov <i>et al</i>	<i>PeerJ</i>	6:e4915	2018	Human / Tissues	VEGF _{xxx} exon 7/8b - Forward primer : 5'- ACGTACTTGCAGATCTCACCA -3' Underlined bases correspond to either end of exon7 OR exon 8.		
Blochowiak <i>et al</i>	<i>Adv Clin Exp Med</i>	27:83-90	2018	Human / Parotid	No	DM3615P	
Boudria <i>et al</i>	<i>Oncogene</i>	AOP		Mice / Cell lines	No	Yes	Yes
Stevens <i>et al</i>	<i>Nephron</i>	139:51-62	2018	Mice	No	No	Yes / TG mice










Figure 1





Figure 2




Figure 3

Supplemental Text

Debunking the myth of the endogenous anti-angiogenic *Vegfaxxb* transcripts.

Hugues Dardente^{1*}, William R English², Manoj K Valluru², Chryso Kanthou² & David Simpson³

¹PRC, INRA, CNRS, IFCE, Université de Tours, 37380 Nouzilly, France

²Department of Oncology and Metabolism, Tumour Microcirculation Group, University of Sheffield, School of Medicine, Beech Hill Road, Sheffield, S10 2RX, United Kingdom

³Centre for Experimental Medicine, Queen's University Belfast, Belfast, BT7 1NN, Northern Ireland

*Corresponding author: hugues.dardente@inra.fr

Other methodological issues with RT-PCR

We noted further inconsistencies in the literature and not-readily-explainable RT-PCR findings. First, to identify *Vegfa165b*, Bates *et al* [1] used primers designed to detect *Vegfa148*, a transcript identified from kidney by the same team [2] and recently shown to be enriched in some tissues, including kidney [3]. The pair of primers used by Bates *et al* [1] consisted of a forward primer located within exon 7a and a reverse primer located in the 3'-UTR, downstream of the putative alternative 8b splice site (i.e. flanking primers; **Table S1**). This should yield at least two PCR products of 185bp and 151bp, corresponding to *Vegfa165* and *Vegfa148*. Should the *Vegfa165b* transcript exist, one would expect an extra PCR product at 120bp. Hence, 3 bands should be discernible on the gel, with the one corresponding to *Vegfa148* located in between those for *Vegfa165* and *Vegfa165b*. This was not the case. This is puzzling considering these primers were in the first instance "designed to detect *Vegfa148*". The same issue applies to the study by Schumacher *et al* [4] who used the same primers. Furthermore, *Vegfaxxb* transcripts were not identified by

Whittle *et al* [2], who used a pair of *ad hoc* primers for this (forward primer in exon 7a and reverse primer downstream of the putative 8b splice site).

In follow-up studies from the team led by Bates and Harper, DNA rulers were often missing and sequencing of the PCR products was not reported [e.g.; 5-8], and therefore the identity of PCR products could not be confirmed. Rather, to validate the existence of an entire family, the authors used assertions such as "size compatible with *Vegfa165b*" [9], "bands corresponding to VEGFA_{xxx}b isoforms (VEGFA165b, VEGFA189b, etc.) were seen ..." [6] but unfortunately not shown, or "multiple isoforms were detected, with bands seen consistent with VEGFA121b (108 bp), VEGFA145b (219 bp) and VEGFA165b (240 bp)". In particular, Cui *et al* [9] used isoform-specific reverse primers to amplify either *Vegfa165* or *Vegfa165b*. These primers were used in combination with the same forward primer, located in exon 4. Therefore RT-PCR for *Vegfa165* is expected to yield at least 3 major bands. Considering the "sister family of *Vegfa_{xxx}b*", RT-PCR for *Vegfa165b* should yield all transcripts that include exon 7b (at least *Vegfa189b* and *Vegfa165b*). However, only products of "sizes compatible with" the presence of *Vegfa189* and *Vegfa165b* were evident on the gel (see [9]; their Fig.3).

Finally, primers used to "detect specifically the transgene" [10,11] in a transgenic mouse model, which over-expresses *Vegfa165b* in the mammary gland, are located within exon3 (forward primer) and exon4 (reverse primer). Therefore, this primer pair will detect all *Vegfa* isoforms (see **Table S1**), which casts doubts on this mouse model.

1 Bates, D.O. *et al.* (2002) VEGF165b, an inhibitory splice variant of vascular endothelial growth factor, is down-regulated in renal cell carcinoma. *Cancer Res* 62, 4123-4131

2 Whittle, C. *et al.* (1999) Heterogeneous vascular endothelial growth factor (VEGF) isoform mRNA and receptor mRNA expression in human glomeruli, and the identification of VEGF148 mRNA, a novel truncated splice variant. *Clin Sci (Lond)* 97, 303-312


- 3 Bridgett, S. *et al.* (2017) RNA-Sequencing data supports the existence of novel VEGFA splicing events but not of VEGFAxxx isoforms. *Sci Rep* 7, 58-017-00100-3
- 4 Schumacher, V.A. *et al.* (2007) Impaired glomerular maturation and lack of VEGF165b in Denys-Drash syndrome. *J Am Soc Nephrol* 18, 719-729
- 5 Woolard, J. *et al.* (2004) VEGF165b, an inhibitory vascular endothelial growth factor splice variant: mechanism of action, in vivo effect on angiogenesis and endogenous protein expression. *Cancer Res* 64, 7822-7835
- 6 Perrin, R.M. *et al.* (2005) Diabetic retinopathy is associated with a switch in splicing from anti- to pro-angiogenic isoforms of vascular endothelial growth factor. *Diabetologia* 48, 2422-2427
- 7 Bates, D.O. *et al.* (2006) The endogenous anti-angiogenic family of splice variants of VEGF, VEGFxxx, are down-regulated in pre-eclamptic placentae at term. *Clin Sci (Lond)* 110, 575-585
- 8 Peiris-Pages, M. *et al.* (2010) Balance of pro- versus anti-angiogenic splice isoforms of vascular endothelial growth factor as a regulator of neuroblastoma growth. *J Pathol* 222, 138-147
- 9 Cui, T.G. *et al.* (2004) Differentiated human podocytes endogenously express an inhibitory isoform of vascular endothelial growth factor (VEGF165b) mRNA and protein. *Am J Physiol Renal Physiol* 286, F767-73
- 10 Qiu, Y. *et al.* (2008) Mammary alveolar development during lactation is inhibited by the endogenous antiangiogenic growth factor isoform, VEGF165b. *Faseb j* 22, 1104-1112
- 11 Qiu, Y. *et al.* (2010) Overexpression of VEGF165b in podocytes reduces glomerular permeability. *J Am Soc Nephrol* 21, 1498-1509

Antibodies and ELISA kits (see text and Supplemental Table1):

- MAB3045 from R&D : Monoclonal Mouse IgG₁ Clone # 56-1 ; directed against human VEGF_{165b} synthetic peptideTCRSLTRKD.
https://www.rndsystems.com/products/human-vegf-165b-antibody-56-1_mab3045
Note : another Ab (raised against the same 9 amino acids) from clone #264610/1 was used in Nowak et al 2008 – no further information could be found.
Note : for Castle-Miller et al 2017 and Ved et al 2017 another clone (56/8) was used. No validation provided.
- DM3615P from Origene : Monoclonal Mouse IgG1 Clone #7F17 ; directed against human recombinant Human VEGF-165b C-terminal peptide
<https://www.acris-anticorps.fr/antibodies/primary-antibodies/vegf165b-dm3615p.htm>
- DY3045 from R&D : human VEGF_{165b} Solid Phase Sandwich ELISA
https://www.rndsystems.com/products/human-vegf-165b-duoset-elisa_dy3045#product-details
- 3045-VE-025 : Recombinant human VEGFA165b protein
https://www.rndsystems.com/products/recombinant-human-vegf-165b-protein_3045-ve
- Ab14994 (clone MRVL56/1) from Abcam : mouse monoclonal antibody – discontinued since 2015.
<https://www.abcam.com/vegf165b-antibody-mrvl561-ab14994.html>
- MBS109074 form MyBioSource : hVEGFA165b ELISA kit
https://www.mybiosource.com/prods/ELISA-Kit/Human/Vascular-Endothelial-Growth-Factor-165B-VEGF-165B/VEGF-165B/datasheet.php?products_id=109074

List of additional references only cited in Supplemental Table1 :

- 1 Amin, E.M. *et al.* (2011) WT1 mutants reveal SRPK1 to be a downstream angiogenesis target by altering VEGF splicing. *Cancer Cell* 20, 768-780
- 2 Baltes-Breitwisch, M.M. *et al.* (2010) Neutralization of vascular endothelial growth factor antiangiogenic isoforms or administration of proangiogenic isoforms stimulates vascular development in the rat testis. *Reproduction* 140, 319-329
- 3 Hueso, L. *et al.* (2017) Dynamics and implications of circulating anti-angiogenic VEGF-A165b isoform in patients with ST-elevation myocardial infarction. *Sci Rep* 7, 9962-017-10505-9
- 4 Li, X. *et al.* (2015) VEGF111b, a C-terminal splice variant of VEGF-A and induced by mitomycin C, inhibits ovarian cancer growth. *J Transl Med* 13, 164-015-0522-0
- 5 Nowak, D.G. *et al.* (2008) Expression of pro- and anti-angiogenic isoforms of VEGF is differentially regulated by splicing and growth factors. *J Cell Sci* 121, 3487-3495
- 6 Nowak, D.G. *et al.* (2010) Regulation of vascular endothelial growth factor (VEGF) splicing from pro-angiogenic to anti-angiogenic isoforms: a novel therapeutic strategy for angiogenesis. *J Biol Chem* 285, 5532-5540
- 7 Qiu, Y. *et al.* (2010) Overexpression of VEGF165b in podocytes reduces glomerular permeability. *J Am Soc Nephrol* 21, 1498-1509
- 8 Stevens, M. *et al.* (2018) Vascular Endothelial Growth Factor-A165b Restores Normal Glomerular Water Permeability in a Diphtheria-Toxin Mouse Model of Glomerular Injury. *Nephron* 139, 51-62



Canonical splice site

Alternative splice site

CCA/CCG/CCC/CCT = Proline.
22 seq, all identical = CDKPRR

	Putative VEGFAxxx sequences
Chinese hamster	PVTRKTD
Deer mice/Mole rat	PLTRKTD
Mouse/Rat	PLTGKTD
Star-nosed mole	PLTGKD
Sheep/Alpaca	CLTRKD
Goat/Cow	RLTRKD
Human/Baboon/Macaque	SLTRKD
Beaver/Cat	PLTRKD
Vole	PLTMSPPHRHRHRQNSP
Bat	PLTRKDGyRMTDTETTPLPpHHH
Bat2	PLTRRDGyRMTDTETTPLPpHHHHHRQNNP
Mouse	PLTRKDRHRTDTEATPLPpHRHRHRQNR
Donkey	PLKERLIGNDRNRNHAATTSPTSEQSLIQKPEMKEETAQSTLGPGETPAEAFPPG
Coquerel's sifaka	PLTRKDRHRSHAAATTPSPSTEQSLDQKPEMKEEETLRRALWVRRARLRQKHSRAGDQARSLLLELDSPFrfSCC
	:

Figure S1

