



## A reassessment of explanations for discordant introgressions of mitochondrial and nuclear genomes

Timothée Bonnet, Raphaël Leblois, Francois Rousset, Pierre-André Crochet

### ► To cite this version:

Timothée Bonnet, Raphaël Leblois, Francois Rousset, Pierre-André Crochet. A reassessment of explanations for discordant introgressions of mitochondrial and nuclear genomes. Conférence Jacques Monod : Coalescence des approches théoriques et expérimentales en génomique évolutive et biologie des systèmes, Oct 2016, Roscoff, France. hal-02932269

**HAL Id: hal-02932269**

**<https://hal.inrae.fr/hal-02932269>**

Submitted on 7 Sep 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# A reassessment of explanations for discordant introgressions of mitochondrial and nuclear genomes

Timothée Bonnet   Raphaël Leblois   François Rousset   Pierre-André Crochet

UMRs **CBGP** & **ISEM** & **CEFE**, (INRA - CNRS, Montpellier, France) [raphael.leblois@supagro.inra.fr](mailto:raphael.leblois@supagro.inra.fr)




## Motivation

Examples of massive mitochondrial introgressions combined with no or very limited nuclear introgressions (i.e. **mito-nuclear discordance**) are accumulating, stimulating the development of various hypotheses to explain this pattern. On the one hand, **selective hypotheses** propose a selective advantage of the introgressing mitochondrial variants, or counter-selection of the nuclear genome; On the other hand, **neutral hypotheses** invoke sex-biased processes and/or drift and/or spatial invasions. The neutral hypotheses however are mostly verbal and have not been quantitatively evaluated. Here we reassess all these hypotheses using simulations under a wide range of demographic and genetic scenarios.

### General simulation settings

► **Spatially explicit simulations of a secondary contact:**

- 2 habitats, each with 15 x 10 demes of  $N=10$  diploid adult pairs ( $N_1=N_2=3,000$ )
- Forward individuals-based simulations of 6,000 generations of secondary contact considering **spatially limited dispersal (IBD)**, **sex-biased processes**, **spatial invasion**, and **selection**
- Standard coalescent to simulate 80,000 generations of **divergence** between the ancestral populations



► **Multi-locus nuclear genome:**

- One chromosome with 20 pairs of loci (1 neutral & 1 selected)
- Recombination within and between pairs of loci

► **Mitochondrial genome:**

- One chromosome with one pair of linked loci (1 neutral + 1 selected)

► **Local adaptation** is modeled with one nuclear "speciation" locus (proxy for many co-adapted loci in LD):

- Lower fitness in the non-original habitat
- Lower fitness for hybrids in both habitats

### Additional demographic and genetic processes explored

► **Sex-biased processes:**

- asymmetric crosses: one sex is preferred by both taxa e.g. the female cow in the Bison/Cow example
- sex-specific dispersal: one sex disperses more than the other
- sex-specific survival: one sex has higher mortality than the other

► **Spatial invasion combined with sex-biased dispersal:**

- One habitat expands over the other after secondary contact, gradually but completely.
- Direction and strength of sex-biased dispersal varies among different scenarios

► **Strong drift** (single hybridization event in small populations):

- Only 3 x 10 demes in each habitat ( $N_1=N_2=600$ )
- Single inter-taxa mating event allowed between two pure individuals
- Then no restrictions on matings between hybrids
- No sex-biased processes

For all scenarios, we considered **both high** (0.1) and **low migration rates** (0.001)

### Outputs of the simulation study: introgression measures

$M$ : proportion of introgressed Mt copies in the most introgressed taxon


$\mathbf{a}$ : vector of proportions of introgressed nuclear gene copies

We are mostly interested in the following output:

►  $M - \bar{\mathbf{a}}$  the **mito-nuclear introgression discordance**

and we define a Massively Discordant Mitochondrial Introgression (**MDMI**) when  $M - \bar{\mathbf{a}} > 0.8$

## Results



## Conclusions

Gene flow	Sex-biases	Invasion	Nu Selection	Mt Selection	MDMI
High	-	-	-	-	No
	Yes	-	-	-	No <sup>1</sup>
	-	Yes	-	-	No
	-	-	Yes	-	Yes
Low	-	-	-	Yes	No
	Yes	-	-	-	No
	-	Yes	-	-	No
	-	-	-	Yes	Yes
Single hybridization	-	-	-	-	No <sup>2</sup>

- Neutral processes (sex-biases, strong drift and spatial invasion) fail to explain **MDMI**
- Only positive selection on Mt with low migration rates or counter selection on the whole nuclear genome can create **MDMI**
- but no detection of positive selection using Tajima's D or Fu' Fs on Mt polymorphism observed in the introgressed taxon

### Take home message

- **Neutral processes can't dissociate nuclear and mitochondrial introgression...** .... because females too have nuclear genes!
- Our results are in line with recent and increasing evidence that **selection on Mt DNA** may be common
- Beware of verbal models, do not trust but **test them using models**

**Acknowledgements:** This work was partly funded by the Institut de Biologie Computationnelle (IBC), and recurrent funds from INRA and CNRS.

