



Does soil organic matter stoichiometry varied with agricultural practices on the long-term?

Fabien Ferchaud, Alain Mollier, Isabelle Bertrand

► To cite this version:

Fabien Ferchaud, Alain Mollier, Isabelle Bertrand. Does soil organic matter stoichiometry varied with agricultural practices on the long-term?. 7. International Symposium on Soil Organic Matter, Oct 2019, Adelaide, Australia. 10.13140/RG.2.2.25072.20486 . hal-02940593

HAL Id: hal-02940593

<https://hal.inrae.fr/hal-02940593>

Submitted on 23 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

FERCHAUD, F.¹, MOLLIER, A.², BERTRAND, I.³


¹ UR AgroImpact, Laon, France; ²UMR ISPA, Bordeaux, France; ³UMR Eco&Sols, Univ Montpellier, CIRAD, INRA, IRD, Montpellier, SupAgro, Montpellier, France

Aim


Soil stoichiometry constraints may limit the impacts of agroecological practices to increase nutrient recycling and foster C storage in cropped soils (Bertrand et al. 2019). However, little is known about the long term impact of agricultural practices on soil stoichiometry. Our aim is to analyse long-term (8-49 yr) field experiments in France including several treatments with contrasted N and/or P budgets.

Methods

We compiled and completed a dataset of long-term (8-49 yr) field experiments in France.



Site	Name and duration	Tested agricultural practices
1	La Cage 1998-2014 (16 years)	4 arable cropping systems: conventional, low input, conservation agriculture, organic farming
2	SOERE QualiAgro 1998-2013 (15 years)	Organic waste products (4 types) * N fertilisation (2 rates) Wheat-maize rotation
3	SOERE ACBB 2009-2015 (6 years)	4 treatments: conventional, reduced tillage, crop residue removal, reduced N fertilisation Arable cropping system
4	Biomass & Environment 2006-2016 (10 years)	Crop type (perennials vs annuals) * N fertilisation (2 rates)
5	Auzeville 1969-2017 (48 years)	P fertilization (superphosphate): 0, 11, 22 and 33 kg P ha ⁻¹ yr ⁻¹ Arable cropping system
6	Mant 1975-1992 (17 years)	P fertilization (superphosphate): 0, 27, 79 kg P ha ⁻¹ yr ⁻¹ Continuous irrigated maize
7	Tartas 1972-2004 (32 years)	P fertilization (superphosphate): 0, 44, 96 kg P ha ⁻¹ yr ⁻¹ Continuous irrigated maize
8	Pierroton 1995-2015 (20 years)	P fertilization (superphosphate): 10, 15, 20, 40, 80 kg P ha ⁻¹ yr ⁻¹ Continuous irrigated maize





Calcul of the balance of C, N and P in (kg X ha⁻¹) in all sites

$$X_{\text{cumulated budget}} = \sum_{i=1}^n (X_{\text{input}} - X_{\text{export}})i \quad X_{\text{input}} = X_{\text{fert}} + \text{BNF} + X_{\text{deposition}}$$


With X_{export} = exportations at harvest; X_{input} = amount of fertilizers (mineral or organic).
For N, atmospheric deposition ($X_{\text{deposition}}$) (http://www.emep.int/mscm_ydata.html) and N fixation by leguminous (BNF) (Anglade et al. 2015) were taken into account.

Results




Site 4, 10 years, N fertilisation * type of crops






Changes in soil C, N and P contents between initial and final dates



Site 2, 15 years, N fertilisation * organic amendments



Site 5, 48 years, P fertilisation



Take home messages

The site has a stronger effect on soil C, N and P contents than practices. Soil C:N ratios were very constrained and not influence by the different agricultural practices, even after 48 years. The N:P and C:P ratios (data not shown) were more flexible. However, the forms of P considered modify the relationships with soil organic C. Organic P is more strongly correlated with organic C underlining the role of soil heterotrophic microorganisms. Such high level of stoichiometry constraint implies that N and P will be necessary to store C in soils.

References: - Bertrand et al. (2019) Stoichiometry constraints challenge the potential of agroecological practices for the soil C storage. A review. Agronomy for Sustainable Development. In Press
- Anglade et al. (2015). Relationships for estimating N₂ fixation in legumes: incidence for N balance of legume-based cropping systems in Europe. *Ecosphere*, 6(3), 1-24.