



HAL
open science

Evidence for autophagy attenuation during post-mortem maturation of hypertrophied muscle in myostatin deficient mice

Rim Nassar, Barbara Vernus, Gilles Fouret, Benedicte Goustard, François Casas, Lionel Tintignac, Isabelle Cassar-Malek, Brigitte Picard, Iban Seiliez, Arnaud Chatonnet, et al.

► To cite this version:

Rim Nassar, Barbara Vernus, Gilles Fouret, Benedicte Goustard, François Casas, et al.. Evidence for autophagy attenuation during post-mortem maturation of hypertrophied muscle in myostatin deficient mice. 7ème congrès du club français Autophagie CFATG, Nov 2017, Paris, France. , 2017. hal-02940597

HAL Id: hal-02940597

<https://hal.inrae.fr/hal-02940597>

Submitted on 16 Sep 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Rim Nassar^{1,2}, Barbara Vernus¹, Gilles Fouret¹, Bénédicte Goustard¹, François Casas¹, Lionel Tintignac³, Isabelle Cassar-Malek⁴, Brigitte Picard⁴, Iban Seliez⁵, Arnaud Chatonnet¹, Aline Hamade², Fadia Najjar², Anne Bonniou¹ et Béatrice Chabi¹

1- INRA, UMR 866 Dynamique Musculaire et Métabolisme, Montpellier, France ; 2 - Laboratoire d'Innovation thérapeutique, Université Libanaise, Beyrouth, Liban; 3 - Département de Biomédecine, Université de Bâle, Bâle, Suisse; 4 - INRA, UMR1213 Herbivores, Saint-Genès-Champagnelle, France; 5 - INRA, UMR1419 Nutrition, Métabolisme, Aquaculture, Saint Pée sur Nivelle, France.


Introduction

The conversion of skeletal muscle into meat, i.e. maturation, is a complex process where muscle undergoes different biochemical and physiological changes (Ouali et al, 2006). In agronomic field, the study of these events is of particular interest, in order to improve the quality of the final product put on the market. If the characterization of the proteolytic mechanisms involved in skeletal muscle maturation is still ongoing, the participation of autophagy in this process is still under debate. The aim of the study was therefore to assess the involvement of autophagy during skeletal muscle maturation and to study the interaction between autophagy and myostatin, a negative regulator of skeletal muscle mass (Mc Pherron et al, 1997), within a 72h postmortem (PM) time frame in mice.

Methods and model validation

Animal model

Healthy 6 month-old Mstn^{+/+} (WT) and Mstn^{-/-} (Mstn KO; Grobet et al, 2003) male mice (n=72 for each genotype) were used in this study.




Muscle sampling

Mice were sacrificed using cervical dislocation. After decapitation and bleeding, mice were stored at room temperature (22±2°C) during 1h and then at 4°C until the dissection time.



Longissimus dorsi and gastrocnemius muscles were harvested and used for pH and western-blotting analyses respectively.


1) pH evolution in the Longissimus muscle



n=8 for each time point x genotype, two-way ANOVA, main effect * p<0,05 WT vs. KO

Postmortem muscle pH decreased faster in Mstn KO mice compared to WT mice.

2) Myofibrillar protein degradation




n=8 for each time point x genotype, two-way ANOVA, *p<0,05; **p<0,01; ***p<0,001 vs. respective T=0

Myofibrillar proteins (e.g. Troponin T) showed a significant and progressive degradation over the 48h PM period and is more pronounced in gastrocnemius muscle of Mstn KO mice compared to WT mice.

=> 3) Following PM, autophagy was assessed in skeletal muscle from the two genotypes using signaling pathways and autophagic flux analyses.


Results

1) Phosphorylation state of proteins involved in autophagic signaling pathway (AMPK, FOXO and ULK)




n=8 for each time point x genotype, two-way ANOVA, *p<0,05; ****p<0,0001

Activation of AMPK is greater in WT mice compared to Mstn KO mice
Energetic stress or/and Autophagy ?



n=6 for each time point x genotype, two-way ANOVA, *p<0,05 vs. respective T=0

Inhibition of Foxo1 is relieved in the late PM interval (8h PM) in both genotypes => Activation of autophagy and Ub proteasome system




n=6 for each time point x genotype, two-way ANOVA, ****p<0,0001 vs. KO T=0


The decrease in ULK1 (Ser757) phosphorylation in Mstn KO mice suggests an early activation (1h PM) of autophagy

2) Autophagy flux measurement


Autophagic flux was monitored using autophagy blockade with four colchicine injections (i.p., 0.1 mg/ml in NaCl 0.9%, every 12h) before sacrifice.



Autophagic flux is calculated as LC3BII_{colchicine} - LC3BII_{NaCl 0.9%} (Jeong-Sun Jun et al, 2010)



Colchicine administration resulted in LC3BII accumulation (NaCl 0.9% vs. colchicine)



n=4 for each time point x genotype, two-way ANOVA, main effect ***p<0,001 WT vs. KO

Basal and postmortem autophagic fluxes are greater in WT versus Mstn KO mice.

Conclusion

In conclusion, our data showed that autophagy is preserved during skeletal muscle PM maturation, but to a lower extent in Mstn-deficient mice, suggesting a relationship between myostatin and autophagy. This work will be completed with microscopic analyses of skeletal muscle tissue sections to visualize *in situ* autophagic events.