



# Quantifying ecosystem capacity, modulation by agricultural practices and actual use of ecosystem services by farmers

Manon Dardonville, Christian Bockstaller, Baptiste Legrand, Olivier Therond

## ► To cite this version:

Manon Dardonville, Christian Bockstaller, Baptiste Legrand, Olivier Therond. Quantifying ecosystem capacity, modulation by agricultural practices and actual use of ecosystem services by farmers. EASY Conference, Groupe “ Ecologie & Agriculture ” de la Société Française d’Ecologie (Sfe), Ecology & Agriculture Summit for Young scientists, 2019, Nice, France. hal-02947656

**HAL Id: hal-02947656**

**<https://hal.inrae.fr/hal-02947656>**

Submitted on 24 Sep 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L’archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d’enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Quantifying ecosystem capacity, modulation by agricultural practices and actual use of ecosystem services by farmers

Manon DARDONVILLE\*<sup>1,2</sup>, Christian BOCKSTALLER<sup>1</sup>, Baptiste LEGRAND<sup>1</sup>, Olivier THEROND<sup>1</sup>

<sup>1</sup> LAE, Université de Lorraine, INRA, Colmar, France

<sup>2</sup>Agrosolutions, Paris, France

\* [manondardonville@inra.fr](mailto:manondardonville@inra.fr) ; +33630862209

Ecosystem services (ES) provided to farmers are now well-considered as natural-capital-based production factors. Developing them is a way to reduce use of anthropic inputs and so to develop a sustainable agriculture. However, it is still a challenge to evaluate quantitatively part of ES relatively of anthropic inputs in production.

Based on literature review, we develop a generic assessment method based on operational indicators to describe ES bundles provided to farmer according to field, farm and landscape characteristics. Application of this method in 30 contrasted farms in terms of agricultural practices enable to assess their relationships with 4-ES dimensions, their role and relative weight in the production of agricultural goods.

According to the spatiotemporal configuration of cropping systems (rotations, cover crops, semi-natural habitats etc.), we assessed the level of (i) ES capacity (potential) available to farmer. Further, we evaluate to which extent ES capacity is (ii) modulated due to beneficial or detrimental agricultural practices and forward the impact on sustainability of (iii) the natural capital. Then, we determine (iv) how much farmer use ES capacity instead of relying on anthropic inputs to produce.

We identified (i) the main ESs limiting agricultural production, and (ii) the ways to enhance their furniture and assessed (iii) the weight of ES in production.

This ES assessment method enable user to categorize farms according to the level of ES and their role in agricultural production process. It allows to finely determine the biotechnical functioning of farming systems and overcome coarse classification like conventional vs. organic farming.