



Deciphering molecular origin and functional impact of structural variation in maize through genome sequences comparison and integrative analysis of genetic variation, transcriptome and phenotype data

Johann J. Joets, Sylvie Coursol, Olivier Turc, Llorenç Cabrera-Bosquet, Marie-Laure Magniette Martin-Magniette, Silvio Salvi, Anthony Venon, Sandrine Chaignon, Jérôme Laplaige, Ghislaine Gendrot, et al.

► To cite this version:

Johann J. Joets, Sylvie Coursol, Olivier Turc, Llorenç Cabrera-Bosquet, Marie-Laure Magniette Martin-Magniette, et al.. Deciphering molecular origin and functional impact of structural variation in maize through genome sequences comparison and integrative analysis of genetic variation, transcriptome and phenotype data. First International Plant Systems Biology meeting, Sep 2018, Roscoff, France. hal-02955192

HAL Id: hal-02955192

<https://hal.inrae.fr/hal-02955192>

Submitted on 1 Oct 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Deciphering molecular origin and functional impact of structural variation in maize through genome sequences comparison and integrative analysis of genetic variation, transcriptome and phenotype data

Johann Joets¹, Sylvie Coursol², Olivier Turc³, Llorenç Cabrera-Bosquet³, Marie-Laure Martin-Magniette^{4,5,6}, Silvio Salvi⁷, Anthony Venon¹, Sandrine Chaignon², Jérôme Laplaige⁸, Ghislaine Gendrot⁸, Carine Palaffre⁹, Stéphanie Castel⁹, Stéphanie Pateyron^{4,5}, Véronique Brunaud^{4,5}, Ludivine Soubigou-Taconnat^{4,5}, Fabrice Dumas¹, William Marande¹⁰, Agnès Rousselet¹, Harry Belcrum¹, Alain Charcosset¹, François Tardieu³, Claude Welcker³, Peter Rogowsky⁸, Clémentine Vitte¹

¹GQE-Le Moulon, INRA, Université Paris-Sud, CNRS, AgroParisTech, Université Paris-Saclay, 91190 Gif-sur-Yvette, France; ²Institut Jean-Pierre Bourgin, INRA, AgroParisTech, CNRS, Université Paris-Saclay, RD10, 78026 Versailles Cedex, France; ³LEPSE, INRA, Université Montpellier, 34060 Montpellier Cedex 1, France; ⁴Institute of Plant Sciences Paris-Saclay IPS2, CNRS, INRA, Université Paris-Sud, Université Evry, Université Paris-Saclay, Bâtiment 630, 91405 Orsay, France; ⁵Institute of Plant Sciences Paris-Saclay IPS2, Paris-Diderot, Sorbonne Paris-Cité, Bâtiment 630, 91405 Orsay, France; ⁶UMR MIA-Paris, AgroParisTech, INRA, Université Paris-Saclay, 75005 Paris, France; ⁷University of Bologna, Department of Agricultural and Food Sciences, DISTAL, Bologna, Italie; ⁸Laboratoire Reproduction et Développement des Plantes, Université de Lyon, ENS de Lyon, UCB Lyon 1 CNRS, INRA, 69000 Lyon, France; ⁹UE 0394 SMH Maïs, INRA, France; ¹⁰Centre National des Ressources Génomiques Végétales, INRA UPR 1258, Castanet-Tolosan, France.

sylvie.coursol@inra.fr

Structural variation (SV) is a major driver of plant adaptation and genome evolution. It originates from transposable element insertion, as well as gene Copy Number (CNV) and Presence/Absence Variation (PAV). Maize is a crop species with a complex genome, and exhibits extensive structural variation among lines, as well as strong phenotypic differences. It is therefore a good model to explore the diverse molecular mechanisms leading to SV, and to investigate to what extent SV impacts phenotypic variation. Finally, the origin of the different maize inbred lines is well described, allowing for linking structural variation to adaptation. Here, we present whole genome assemblies from seven European and American maize lines of various geographical origins and phenotypes, and with contrasted genome size. This dataset allows unprecedented genome-wide comparisons and characterization of maize SV with high sequence accuracy, thus offering opportunity to evaluate the prevalence of the molecular mechanisms underlying these variations and to characterize the features responsible for genome size variation. These seven maize lines together with B73 were cultivated under contrasted water conditions in the PhenoArch phenotyping platform allowing precise characterization of growth and development together with precise measurements of environmental conditions. Thirteen different organs harvested at various developmental stages have been used for RNA-seq-based transcriptome analysis. This massive dataset will be used to evidence the possible role of SVs in quantitative responses to water deficit as well as the impact of SVs in gene regulation networks. Overall, this work will provide insights on the molecular origins and functional consequences of SV.

Funding acknowledgment: French National Research Agency (Amaizing, ANR-10-BTBR-03), France Agrimer and LabEx Saclay Plant Sciences-SPS (ANR-10-LABX-0040-SPS).