

## Water deficit responsive QTLs for cell wall degradability and composition in maize at silage stage

Laetitia Virlovet, Fadi Hage, Yves Griveau, Marie-Pierre Jacquemot, Emilie Gineau, Aurélie Baldy, Sylvain Legay, Christine Horlow, Valérie Combes, Cyril Bauland, et al.

► **To cite this version:**

Laetitia Virlovet, Fadi Hage, Yves Griveau, Marie-Pierre Jacquemot, Emilie Gineau, et al.. Water deficit responsive QTLs for cell wall degradability and composition in maize at silage stage. 3rd European Maize Meeting, 2019, Montpellier, France. hal-02955451

**HAL Id: hal-02955451**

**<https://hal.inrae.fr/hal-02955451>**

Submitted on 1 Oct 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Sylvie COURSOL

### P5: Water deficit responsive QTLs for cell wall degradability and composition in maize at silage stage

Laëtitia Virlovet<sup>1</sup>, Fadi El Hage<sup>1</sup>, Yves Griveau<sup>1</sup>, Marie-Pierre Jacquemot<sup>1</sup>, Emilie Gineau<sup>1</sup>, Aurélie Baldy<sup>1</sup>, Sylvain Legay<sup>1</sup>, Christine Horlow<sup>1</sup>, Valérie Combes<sup>2</sup>, Cyril Bauland<sup>2</sup>, Carine Palafre<sup>3</sup>, Matthieu Falque<sup>2</sup>, Laurence Moreau<sup>2</sup>, Valérie Méchin<sup>1</sup>, Matthieu Reymond<sup>1</sup>

<sup>1</sup>Institut Jean-Pierre Bourgin, INRA, AgroParisTech, CNRS, Université Paris-Saclay, 78000, Versailles, France

<sup>2</sup>Génétique Quantitative et Evolution - Le Moulon, INRA, Université Paris-Sud, CNRS, AgroParisTech, Université Paris-Saclay, 91190, Gif-Sur-Yvette, France

<sup>3</sup>Unité Expérimentale du Maïs, INRA, 40390, Saint Martin de Hinx, France

The use of lignocellulosic biomass for animal feed or biorefinery requires the optimization of its degradability. Moreover, biomass crops need to be better adapted to the changing climate and in particular to periods of drought. Although the negative impact of water deficit on biomass yield has often been mentioned, its impact on biomass quality has only been recently reported in a few species. In the present study, we combined the mapping power of a maize recombinant inbred line population with robust near infrared spectroscopy predictive equations to track the response to water deficit of traits associated with biomass quality. The population was cultivated under two contrasted water regimes over three consecutive years in the south of France and harvested at silage stage. We showed that cell wall degradability and  $\beta$ -O-4-linked H lignin subunits were increased in response to water deficit, while lignin and p-coumaric acid contents were reduced. A mixed linear model was fitted to map quantitative trait loci (QTLs) for agronomical and cell wall-related traits. These QTLs were categorized as 'constitutive' (QTL with an effect whatever the irrigation condition) or 'responsive' (QTL involved in the response to water deficit) QTLs. Fifteen clusters of QTLs encompassed more than two third of the 213 constitutive QTLs and 13 clusters encompassed more than 60% of the 149 responsive QTLs. Interestingly, we showed that only half of the responsive QTLs colocalized with constitutive and yield QTLs, suggesting that specific genetic factors support biomass quality response to water deficit. Overall, our results demonstrate that water deficit favors cell wall degradability and that breeding of varieties that reconcile improved drought-tolerance and biomass degradability is possible.

This study was supported by LabEx Saclay Plant Sciences-SPS, Grants ANR-10-LABX- 0040-SPS and ANR-11-BTBR-0006 BIOMASS FOR THE FUTURE. The authors benefited from the IJPB "Plant Observatory" facilities.