



HAL
open science

Analysis of surface water reveals land pesticide contamination: an application for the determination of chlordecone-polluted areas in Guadeloupe, French West Indies

Romain Rochette, Vincent Bonnal, Patrick Andrieux, Philippe Cattan

► **To cite this version:**

Romain Rochette, Vincent Bonnal, Patrick Andrieux, Philippe Cattan. Analysis of surface water reveals land pesticide contamination: an application for the determination of chlordecone-polluted areas in Guadeloupe, French West Indies. *Environmental Science and Pollution Research*, 2020, 10.1007/s11356-020-10718-y . hal-02958848

HAL Id: hal-02958848

<https://hal.inrae.fr/hal-02958848v1>

Submitted on 8 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

1 Analysis of surface water reveals land pesticide contamination
2 An application for the determination of chlordecone-polluted areas in
3 Guadeloupe, French West Indies

4
5
6 Romain ROCHETTE^{1,*}, Vincent BONNAL², Patrick ANDRIEUX¹, Philippe CATTAN³

7
8 1 ASTRO Agrosystèmes Tropicaux, INRA, 97170, Petit-Bourg (Guadeloupe), France

9 2 CIRAD, UMR TETIS, F-97130 Capesterre-Belle-Eau (Guadeloupe), France

10 2 TETIS, Univ Montpellier, AgroParisTech, CIRAD, CNRS, IRSTEA, Montpellier, France

11 3 CIRAD, UPR GECO, F-34398, Montpellier, France

12 3 GECO, Univ Montpellier, CIRAD, Montpellier, France

13 * Corresponding author

14 *Email: romain.rochette@inra.fr*

15
16 **Summary**

17 In Guadeloupe, the use between 1972 and 1993 of chlordecone, an organochlorine insecticide, has permanently
18 contaminated the island's soil, thus contaminating the food chain at its very beginning. There is today a strong societal
19 requirement for an improved mapping of the contaminated zones. Given the extent of the areas to be covered, carrying
20 out soil tests on each plot of the territory would be a long and expensive process. In this article, we explore a method of
21 demarcating polluted areas. The approach adopted consists in carrying out, using surface water analyses, a hydrological
22 delimitation that makes it possible to distinguish contaminated watersheds from uncontaminated ones. The selection of
23 sampling points was based on the spatial analysis of the actual and potential contamination data existing at the
24 beginning of the study. The approach was validated by soil analyses, after having compared the contamination data of
25 the watersheds with the soil contamination data of the plots within them. The study thus made it possible to highlight
26 new contaminated areas and also those at risk of contamination, and to identify the plots to be targeted as a priority
27 during future analysis campaigns by State services.

1 **Keywords**

2 Organochlorine pesticide, Watersheds, Contamination, Water, Soil

3

4 **1. Introduction**

5 The health safety of food produced by agriculture is a major societal concern. Agricultural products can become
6 contaminated by pesticides, either by their direct application to crops or indirectly by their transfer from the soil, to
7 which they had become fixed after application. The latter is especially the case with persistent molecules which were
8 used in the past and still persist in the environment (Barron et al. 2017; Wu and Zhu 2019). When originating from old
9 practices, this contamination is hard to predict, making it difficult to control the risk of exposure of human populations.
10 An obvious prerequisite is the identification and mapping of contaminated areas.

11 This problem is particularly relevant to the French West Indies where an organochlorine insecticide, chlordecone, was
12 used between 1972 and 1993 on banana crops to control the banana weevil (*Cosmopolites sordidus*). This persistent
13 pesticide is still present in soils (Cattan et al. 2016) and continues to contaminate crops grown there, rendering them
14 unfit for consumption (Cabidoche and Lesueur-Jannoyer 2012; Clostre et al. 2015). Following the discovery of this
15 health problem, maps of soil contamination were drawn up. In Guadeloupe, a first mapping of plots likely to be
16 contaminated was carried out in 2006 (Tillieut 2006) and was based on a classification of areas according to their
17 banana cultivation history. Since chlordecone was used mainly in banana plantations, and in a quasi-systematic manner
18 (Levillain et al. 2012), this hypothesis was based on the assumption that the historical use of the land for banana
19 cultivation should account for the level of soil contamination.

20 However, historical information on land use was not comprehensive or accurate (unregistered or unmapped banana
21 plots; diverted use of chlordecone on other crops, etc.). There was therefore uncertainty about the state of contamination
22 of the plots: uncontaminated plots could be located in areas identified as being at risk (based on land use history), and,
23 more importantly, contaminated plots could be located in areas identified as being at zero to negligible risk. To reduce
24 these uncertainties, many soil analyses have been undertaken in the territory by different entities (Department of Food,
25 Agriculture and Forestry; Chamber of Agriculture; Regional Health Agency; National Institute for Agronomic
26 Research). The results were collected in 2009 in a spatialized database within a geographical information system (GIS)
27 for an effective mapping of contamination of plots. However, these analyses were not based on systematic campaigns to
28 examine the environment, which were too expensive to undertake. The plots that were analysed were limited mainly to
29 within areas defined as being at risk and thus did not cover the entire territory in a systematic manner. In fact, there is

1 today a strong need to undertake a new mapping to better determine the areas that are contaminated by chlordecone, and
2 also those that are not, so that their use for agricultural purposes can be allowed.

3 To draw up contamination maps, it is customary to conduct soil analyses (e.g. (Zhao et al. 2013)), and then, to keep
4 costs down, use spatial interpolation between them to represent potentially polluted areas (e.g. (Frangi and Richard
5 1997)). However, in the case of chlordecone, soil concentrations are dependent on farmer categories and are not
6 spatially structured (highly contaminated plots may be adjacent to uncontaminated plots), making spatial interpolation
7 unsuitable for this molecule (Levillain et al. 2012). However, other methods can be envisaged based on chlordecone's
8 intrinsic characteristics: it is particularly stable and strongly sequestered in soil organic matter (Cabidoche et al. 2009;
9 Woignier et al. 2013). Although it is not very mobile, some of it is leachable by water drainage, in more or less
10 significant quantities depending on soil type. The knowledge acquired on the transfer of this molecule (Cabidoche et al.
11 2009; Crabit et al. 2016; Della Rossa et al. 2017) shows that permanent contamination of watercourses can be linked to
12 the state of soil contamination at the watershed scale. On this basis, we can hypothesize that if river water is
13 contaminated, some of the land in its watershed is contaminated. Conversely, the absence of water contamination at the
14 point of sampling should imply that no or very low soil contamination of the associated watershed. So it can be
15 surmised that water analyses can account for the state of contamination of the lands of the watershed associated with the
16 sampling point.

17 The purpose of this paper is to explore a method to improve chlordecone pollution maps based on this hypothesis. The
18 case of Guadeloupe was selected for the study. The approach adopted consisted of analysing, for different watersheds,
19 both the contamination of water and of the soil. To this end, we carried out a hydrological division of the Guadeloupean
20 territory into watersheds and sub-watersheds, making it possible to distinguish, by means of water analyses at the outlet
21 of each of them, those contaminated by chlordecone from those that are not. This method makes it possible to better
22 manage the health risk associated with the use of the soil for cultivation.


23 **2. Materials and methods**

24 **2.1 Study site**

25 Guadeloupe (16°15'N, 61°32'W) is an archipelago located on the arc of the Lesser Antilles, which marks the boundary
26 between the Atlantic Ocean and the Caribbean Sea (Figure 1). The study focuses on 'continental' Guadeloupe, which is
27 composed of two main islands: Basse-Terre, a mountainous volcanic island with an area of 848 km², which rises to an
28 altitude of 1467 m; and Grande-Terre, limestone plateau of 587 km², whose altitude does not exceed 136 m.

29 The study site is subject to an oceanic tropical climate, regulated by a steady flow of hot and humid trade winds. The
30 year is divided into two clearly distinct seasons: a so-called dry season (*carême*) from December to May and a wet

1 season (*hivernage*) from June to November, which is marked by the occurrence of tropical waves. Rainfall shows great
2 variation over the territory, with annual cumulative amounts ranging from 950 mm on the east of Grande-Terre and on
3 the west coast of Basse-Terre, to more than 10,000 mm at the highest altitudes of Basse-Terre. The hydrographic
4 networks of the two islands are also very different. In Basse-Terre, where rainfall is high, the rivers are perennial,
5 sustained during periods of low precipitation by groundwater (Charlier et al. 2011; Crabit et al. 2016). In Grande-Terre,
6 the absence of marked relief and the much lower rainfall make surface flows much more limited and very irregular,
7 giving rise to numerous phenomena of endorheism (when the surface flows do not reach the sea and get lost in closed
8 depressions) (Morell and Jérémie 1994).



9
10 **Figure 1** Geographical location of Guadeloupe and water and soil sampling points

11 **2.2 Approach**

12 The approach we adopted was to link the water contamination at the outlet of a watershed with that of its soils. On the
13 basis of the spatial analysis of the actual and potential contamination data existing at the beginning of the study, we
14 carried out a hydrological delimitation of the territory into watersheds and sub-watersheds to distinguish, by means of
15 water analyses at the outlet of each of them, those that are contaminated by chlordecone from those that are not. If the
16 water is contaminated, then the watershed is considered to be at high risk and will require soil tests before plots can be
17 used for cultivation and increased monitoring of the agricultural production originating from the watershed. If the water
18 is not contaminated, the watershed is considered low-risk and will not require such increased monitoring. In this
19 approach, we did not take into account the soil type. In fact, although the soil type is obviously a determinant of the

1 level of river contamination, we considered that the fact that the soils were contaminated, regardless of their type, was
2 sufficient to explain the binary state of the river: to be or not to be contaminated.

3 The method was validated by comparing the results of the water analyses with the initial data of soil contamination of
4 watersheds: risk map drawn up in 2006 on the basis of historical land use (Tillieut 2006) and GIS database of
5 chlordecone soil analyses. For the watersheds for which there were inconsistencies, new soil analyses were carried out
6 on some plots to:

- 7 - verify the contamination of plots previously analysed for chlordecone;
- 8 - verify the contamination – or not – of plots identified as at risk;
- 9 - if no information was available, find a terrestrial origin for the detected water contamination.

10 **2.3 Delimitation of watersheds**

11 Watersheds and their drainage systems have been previously defined, delimited and generated through spatial analysis
12 tools under Esri ArcMap®. We used the RGE ALTI® digital elevation model (DEM) from the French National Institute
13 of Geographic and Forest Information (IGN), which covers the whole of continental Guadeloupe at a resolution of 1 m.
14 However, this DEM had to be modified because the elevated elements of the road network were not open, i.e., they
15 formed barriers to water flow. It was therefore decided to resample it at a resolution of 25 m to minimize this problem.
16 We then used the AGREE method (Hellweger 1997) to force the DEM by IGN's BD TOPO® hydrographic network,
17 considered as a reference, by verifying 1) the correspondence between the generated hydrographic network and that of
18 the reference network, 2) the coherence between the generated watersheds and the reference hydrographic network, and
19 3) the non-modification/conservation of ridges, by comparing the watersheds generated by the forcing with those that
20 did not undergo forcing.

21 **2.4 Water sampling**

22 **2.4.1 Water sampling plan**

23 The water sampling points were located at the outlets of watersheds with a minimum area of 50 ha, which represented a
24 good compromise between the desire to cover the territory as widely as possible and the budgetary constraints which
25 allowed only a limited number of water analyses. In cases in which the outlets were not accessible (mangroves, private
26 access, steep relief, etc.), the sampling points were moved upstream.

27 Additional sampling points were then distributed over certain watersheds to obtain information on the state of soil
28 contamination at the sub-watershed scale and therefore to be able to distinguish contaminated sub-watersheds from
29 uncontaminated ones, especially in areas that have not been extensively studied so far. These sampling points were

1 selected on the basis of the spatial analysis of actual and potential contamination data existing at the beginning of the
2 study (chlordecone soil contamination risk map and spatialized water and soil contamination data for Guadeloupe).
3 In Basse-Terre, in watersheds with an available agricultural area (AAA – (SIRS 2015)) greater than 25 ha, a size we
4 considered sufficiently big to warrant further analysis, it was possible to identify sub-watersheds bigger than 50 ha that
5 did not have known chlordecone contamination. Sampling points were identified along their watercourses at the
6 frequency of one point every 1.2 km or so. This allowed us to obtain a grid sufficiently dense to identify possible
7 contamination as well as to follow its evolution along the watershed's hydrological slope, i.e., upstream to downstream.
8 For watersheds with an AAA of less than 25 ha, it was considered that the analysis conducted at their outlet was
9 sufficient to assess their chlordecone contamination.

10 In Grande-Terre, flows are not perennial and chlordecone contamination data were sparse. As a result, we mapped a
11 grid across the territory, but this time with the objective of obtaining information on the contamination of areas that are
12 hydrologically disconnected from one another for most of the year. Samples were thus planned on the flow axes of all
13 the sub-watersheds with an area greater than 50 ha, with an average sampling frequency of 1.2 km. Since flows are rare,
14 the majority of the sampling points were in ponds/reservoirs along the flow axes.

15 On the basis of this sampling plan, 219 different points were sampled in Basse-Terre, spread over 97 watersheds
16 emptying into the sea and representing 83% of the island's surface area. In Grande-Terre, samples were collected from
17 198 different points, spread over 33 watersheds emptying into the sea and representing 71% of the island's surface area.
18 All the water sampling points are shown in Figure 1.

19 **2.4.2 Periods of water sampling**

20 Water samples were collected during different hydrological periods depending on the sampling location.

21 In Basse-Terre, previous measurements, made on different soil types, have shown that runoff is far less significant than
22 infiltration. Indeed, runoff rates are only 5 to 10% for low-intensity rainfall, and rise to a maximum of 30% for intense
23 events (Cattan et al. 2006). Surface runoff is therefore not very effective in transporting chlordecone under the
24 conditions prevailing in the French West Indies (Cabidoche et al. 2009). Contamination of rivers is mainly via
25 groundwater, which provides flows during low-water periods (Charlier et al. 2009; Crabit et al. 2016). This period also
26 represents the most stable state of the hydrological system. For Basse-Terre, water sampling was therefore scheduled
27 during low-water periods.

28 In Grande-Terre, too, infiltration is much more significant than surface runoff, with the latter (estimated at 1% of annual
29 rainfall) being almost negligible (Cottez 1972). Moreover, since rainfall is much lower than in Basse-Terre, the flows in
30 the ravines are very limited and very irregular, giving rise to a number of endorheism phenomena. Surface water is

1 therefore not present throughout the year, and depends heavily on rainfall during the rainy season. For Grande-Terre,
2 the samples were therefore planned during the rainy season.
3 As far as possible, samples from each watershed were made on the same day and/or in similar hydrological conditions.
4 If a stream was dry on the day planned for the sampling, the sample was rescheduled, if possible, after a rain event
5 sufficient to cause runoff.

6 **2.4.3 Water sampling method**

7 The water samples were collected in 1 litre brown-coloured glass bottles. On return from the field, these bottles were
8 stored in a refrigerator at 4 °C before being shipped in refrigerated coolers the next day to metropolitan France for
9 analysis. The shipments were made through an express delivery service (Chronopost) and took 2 to 3 days to reach the
10 laboratory.

11 **2.5 Soil sampling**

12 **2.5.1 Soil sampling plan**

13 A typology of all the watersheds was carried out according to the contamination data available for each of them,
14 namely: (1) the state of contamination of the river waters at the watershed's outlet, according to the analyses performed
15 (WC); (2) the presence of contaminated soils in the watersheds, according to the GIS database of already existing soil
16 tests (SC); (3) the presence of plots at risk, according to the risk map drawn up in 2006 (PC). One of two values was
17 assigned to each of these three variables (WC, SC and PC): uncontaminated = 0; contaminated = 1. Four watersheds
18 with plots known to be contaminated (SC 1) presented a particular context, with contaminated water but with only a few
19 number of contaminated plots, of small areas or which were far from the main flow axes, i.e. contributing not much to
20 water pollution. These watersheds could therefore contain other contaminated plots not referenced in the GIS database.
21 We then created a new type denoted SC1' for these 4 watersheds. Soil analyses were then conducted on watersheds with
22 inconsistencies: (case 1) presence of contaminated soils or plots at risk of contamination while the rivers were not
23 contaminated, (case 2) absence of contaminated soils or plots at risk of contamination while the rivers are contaminated,
24 and (case 3) presence of contaminated rivers while the contribution of actual contaminated plots to water pollution is
25 low.

26 The decision rules concerning the plots to be targeted for soil analyses were thus formalized as follows:

- 27 - for watersheds of type [WC 0, SC 1, PC 0] and [WC 0, SC 1, PC 1] (case 1): verify the contamination of plots
28 indicated as contaminated. For Grande-Terre, only plots located in the immediate vicinity of a water sampling
29 point were considered, as it was difficult to define the catchment area of a given water sampling point with
30 accuracy (for reasons already mentioned above);

- 1 - for watersheds of type [WC 1, SC 1, PC 1] (case 3), [WC 1, SC 0, PC 1] (case 2) and [WC 0, SC 0, PC 1]
2 (case 1): verify the contamination status of plots said to be at risk;
- 3 - for watersheds of type [WC 1, SC 1, PC 0] (case 3) and [WC 1, SC 0, PC 0] (case 2): locate an as yet
4 unknown terrestrial contribution to the water contamination. In this case, the plots to be analysed were selected
5 from the analysis of old aerial photos and historical land use maps (analysis of the documents used to produce
6 the 2006 risk map, as well as two additional maps), and were then validated by field investigations.

7 In Basse-Terre, of the 219 watersheds analysed through water analyses, 60 presented inconsistencies. For Grande-Terre,
8 only 3 watersheds out of the 199 analysed presented inconsistencies. Because of cost considerations, additional analyses
9 were conducted on only 36 plots: 32 on Basse-Terre and 4 on Grande-Terre (Figure 1). The targeted watersheds were
10 primarily those that each had an unknown source of contamination and those that had only one doubtful plot to verify.
11 Watersheds with several doubtful plots were also analysed. In these, the suspect plot that was closest to the river and
12 was easiest to access was selected for analysis.

13 **2.5.2 Soil sampling method**

14 Chlordecone exhibits strong spatial heterogeneity at the plot scale (Clostre et al. 2014). The scientific literature
15 recommends 20 sampling points per plot for soil analyses [21]. However, this high number of samples is difficult to
16 implement on a large number of plots. For our study, we decided on a density of 15 samples per hectare, with a
17 minimum of 8 samples for plots with areas less than 1 ha. These sampling points were located on transects on each plot,
18 spaced 10 to 30 m apart and at least 7 m distant from the plot's boundaries to limit edge effects. The samples were
19 collected only from the 0-30 cm horizon, to be in line with the existing reference base.

20 Samples from the same plot were collected in a thick plastic bag, which was closed once filled. The auger was cleaned
21 with brush and water after sampling each plot. On return from the field, each sample was manually homogenised after
22 removal of extraneous material and reduction of aggregates. Then, a sub-sample of 500 g was formed by quartering
23 method. It was placed in a polypropylene sampling pot. The samples were stored in the refrigerator at 4 °C before
24 shipment to metropolitan France at the beginning of each week for analysis. The shipments were made through an
25 express delivery service (Chronopost) and took 2 to 3 days to reach the laboratory.

26 **2.6 Analyses of samples**

27 The water and soil samples were analysed by the La Drôme Departmental Laboratory (LDA 26).

28 For the analysis of chlordecone in water, 1 L of sample was extracted in liquid/liquid partition with a
29 dichloromethane/ethyl acetate mixture (80/20) at several pH values. Extraction tracers HBB/TPP/DIA-DS were added
30 prior to extraction. The organic phases were collected and frozen at -18 °C to remove traces of water. The extract was

1 then concentrated under nitrogen flow in a 35 °C bath (Turbo-Vap) and a volume tracer, propazine D6, was added. A
2 drop of pentanol was brought to the extract and the solvent was evaporated. 0.5 mL of the extract were introduced in a
3 water/acetonitrile mixture (50/50) and internal standards, 24-D D3 and atrazine D5, were then added. The analysis was
4 performed by HPLC-MS/MS, with an analytical uncertainty of 30% and detection and quantification thresholds of 0.01
5 µg/L.

6 For the analysis of chlordecone in soils, 10 g of sample were introduced into an extraction cell and tracers were added
7 (HBB/TPP). Accelerated solvent extraction (ASE) was carried out with a dichloromethane/acetone mixture at 100°C
8 and under pressure (120 bar). The resulting extract was concentrated with a vacuum centrifuge (GENEVAC EZ2)
9 which greatly reduces the loss of volatile compounds. The extract was concentrated to 10 mL and an aliquot of 1 mL
10 was taken. A drop of pentanol was added to the extract and the solvent was evaporated in a GENEVAC MiVac system
11 to preserve the volatile compounds. The extract was then taken up by a mixture of acetonitrile and water with the
12 Chlordecone C13 internal standard. The analysis was performed by HPLC-MS/MS, with an analytical uncertainty of
13 40%, a detection threshold of 2 µg/kg of dry soil and a quantification threshold of 5 µg/kg of dry soil.


14 **3. Results**

15 **3.1 Contamination of watersheds**

16 The level of water contamination by chlordecone was assessed against thresholds defined in French law by the decree
17 of January 11, 2007, concerning the limits and references of quality of raw water and water intended for human
18 consumption, in application of European Directives 75/440/EEC and 98/83/EC. The first threshold, of 0.1 µg/L,
19 corresponds to the limit of the concentration of chlordecone that is allowed in tap water intended for human
20 consumption. The second, of 2 µg/L, is the chlordecone concentration threshold above which raw water can no longer
21 be made safe for drinking even after treatment.

22 **3.1.1 Highly contaminated water in Basse-Terre**

23 In Basse-Terre, chlordecone was found in 36% of 277 analyses, with concentrations ranging from 0.01 to 42.9 µg L⁻¹
24 (Figure 2). 24.1% of rivers have levels above the French consumption threshold for this pesticide (0.1 µg L⁻¹), and 7.9%
25 of them have water contamination above the higher threshold of 2 µg L⁻¹, i.e., water that is considered unfit for
26 consumption even after treatment. Furthermore, of the 110 watersheds analysed at their outlets, 43 (39%) were
27 identified as discharging contaminated water into the sea. In Grande-Terre, 5 points of contamination out of 198 were
28 detected, each of them at low levels, lower than the consumption norm.



1


2 **Figure 2** Distribution in different classes of contamination of the results of analyses of chlordecone in the waters of Basse-Terre and
 3 Grande-Terre

4 **3.1.2 A contamination structured by banana cultivation**

5 A map of watershed contamination by chlordecone was drawn up (Figure 3).

6 In Basse-Terre, no chlordecone contamination was detected in the watercourses in the northern half of the west coast or
 7 the northern part of the east coast, except in an upstream tributary of a small stream, where this contamination was
 8 measured at $0.011 \mu\text{g L}^{-1}$ (slightly higher than the quantification threshold of $0.01 \mu\text{g L}^{-1}$). In the centre of the east coast,
 9 contamination was recorded in various rivers and some of their tributaries, with concentrations ranging from $0.016 \mu\text{g L}^{-1}$
 10 L^{-1} to $0.41 \mu\text{g L}^{-1}$. Finally, in the island's southern part, where chlordecone had seen massive use, it was not possible to
 11 identify uncontaminated sub-watersheds belonging to contaminated watersheds.

12 In Grande-Terre, 5 points of contamination were recorded, with relatively low levels for each (close to the
 13 quantification threshold). Field investigations that were undertaken allowed 3 of these points to be eliminated since the
 14 source was not land contamination but a contaminated water supply (water of the agricultural irrigation network coming
 15 from Basse-Terre).



1
2 **Figure 3** Mapping of chlordecone contamination of watersheds in continental Guadeloupe

3 **3.2 Comparisons between chlordecone databases**

4 These comparisons were made only for Basse-Terre, for which the three variables WC, SC and PC were available.
5 Indeed, the risk of soil contamination by the chlordecone on Grande-Terre appears to range from nil to negligible on the
6 entirety of its territory. In addition, as already noted, it is impossible to define the supply watershed of a given water
7 sampling point with any accuracy, which makes it difficult to establish a correspondence between soil data and water
8 data.

9 **3.2.1 Soil analyses consistent with the plot contamination risk map**

10 Table 1 shows that for 81% of the watersheds analysed in Basse-Terre there exists a correlation between the presence of
11 contaminated soils and plots at risk of contamination (i.e. [PC 0, SC 0] and [PC 1, SC 1]). The absence of contaminated
12 soils, while plots at risk are present in the watersheds ([PC 1, SC 0], 9% of cases), can be explained by the fact that no
13 plot has yet been analysed in these watersheds. On the other hand, the presence of contaminated soils in watersheds
14 without plots at risk of contamination [PC 0, SC 1] reflects either the lack of exhaustiveness of the historical analysis of
15 the plots on which banana was cultivated, or shows the existence of historical uses of the pollutant diverted to plots
16 other than those of banana, or is due to the uncertainty associated with the contamination data. In the end, the risk of
17 having contaminated soils (SC 1) in watersheds without a plot at risk (PC 0) appears to be significant and amounts to
18 $23/(120 + 23) = 16\%$ (percentage calculated on the number of watersheds without a plot risk only).

Table 1 Number of watersheds with/without the presence of plots at risk of contamination and contaminated soils in Basse-Terre.

The percentages are calculated on the total number of watersheds

	PC 0	PC 1
SC 0	120	19
	55%	9%
SC 1	23	57
	11%	26%

3.2.2 Correspondence between water contamination and soil contamination

Table 2 shows that for 80% of the watersheds in Basse-Terre, there is a correspondence between the existence or the absence of soil contamination and the existence or the absence of water contamination ([SC 0, WC 0] and [SC 1, WC 1]).

Table 2 Number of watersheds according to soil and water contamination status in Basse-Terre. The percentages are calculated on the total number of watersheds

	WC 0	WC 1
SC 0	124	15
	57%	7%
SC 1	29	51
	13%	23%

For the remaining 20%, 7% is due to a lack of information on soil contamination. In contrast, for the other 13%, contamination of plots is not detectable in watercourses. This problem reflects either the uncertainty related to the measurement of soil contamination, or a very large dilution effect of the soil contamination in the water feeding the river.

3.3 Verification of contamination of plots

3.3.1 Identification of watersheds with inconsistencies between WC, SC and PC

In Basse-Terre, of the 153 watersheds whose water was found to be uncontaminated, 39 (25%) contained plots that had to be verified by new soil analyses (Table 3a):

- 29 watersheds with plots known to be contaminated, to confirm that they were not 'false positives';
- 10 watersheds with plots at risk, to verify that these plots are uncontaminated.

1 Conversely, of the 66 watersheds whose water was found to be contaminated, 21 (32%) contained plots that had to be
 2 verified by soil analyses in order to locate a terrestrial origin to the water contamination (Table 3b):

- 3 - 15 watersheds with plots at risk, to check whether these plots contributed to the contamination found in the
 4 water;
- 5 - 6 watersheds without any indication of contamination and without risk, to try to locate a terrestrial origin for
 6 the contamination.

7 **Table 3** Number of watersheds with inconsistencies when: a. no water contamination by chlordecone was detected; b. water
 8 contamination by chlordecone was detected

a.	Num. of watersheds	[SC 1, PC 0]	[SC 0, PC 1]
		[SC 1, PC 1]	[SC 0, PC 1]
WC 0	153	29	10
	100%	19%	7%

b.	Num. of watersheds	[SC 0, PC 1]	[SC 0, PC 0]
		[SC 1, PC 1]	[SC 1, PC 0]
WC 1	66	15	6
	100%	23%	9%

9
 10 In Grande-Terre, of the 197 watersheds whose water was found to be uncontaminated, only one contained a plot known
 11 to be contaminated immediately upstream from the water sampling point.

12 Conversely, in 2 watersheds whose waters were found to be contaminated, we carried out soil analyses in order to try to
 13 locate a terrestrial origin for the contamination found in the water.

14 **3.3.2 Results of the soil analysis**

15 In Basse-Terre, in the watersheds that did not show water contamination (WC 0), 24 soil analyses were carried out
 16 (Table 4a). Of the 14 plots known to be contaminated that were verified, 11 were found to be chlordecone free (79%).
 17 Similarly, of the 10 plots at risk of contamination that were analysed, 8 were found to be uncontaminated (80%).

18 In the watersheds whose waters were found to be contaminated (WC 1), 11 soil analyses were carried out (Table 4b). In
 19 9 of the 11 watersheds analysed (82%), we managed to find terrestrial origins for the chlordecone contamination. Of the
 20 9 plots at risk of contamination that were analysed, 7 were found to be contaminated (78%).

21 **Table 4** Summary of the results of soil analyses carried out in Basse-Terre watersheds where: a. no water contamination by
 22 chlordecone was detected; b. water contamination by chlordecone was detected

a.	Type of plots analysed	Num. of plots analysed	Num. of negative analyses	% negatives
	Contaminated plots	14	11	79%
	Plots at risk	10	8	80%

1

b.	Type of plots analysed	Num. of plots analysed	Num. of positive analyses	% positives
	Plots at risk	9	7	78%
	Plots with unknown status	2	2	100%

2 In Grande-Terre, the only plot known to be contaminated that was re-analysed did not contain chlordecone. For
3 watersheds in which the terrestrial origin of water contamination was sought to be located, we could not identify, from
4 the documents of historical land use, any real leads allowing us to identify a possible source of pollution. 3 plots were
5 nevertheless analysed but no contamination was found. This reflects the difficulty, in some watersheds, of finding a
6 source of contamination without having any information beforehand as to where to start looking.

7 **3.3.3 Assessment of the approach**

8 Our approach consisted of characterizing the contamination of the soil of a watershed on the basis of the quality of its
9 watercourse. We assessed this approach on the basis of the correlation between water contamination data (WC) and soil
10 contamination data (SC) or soil contamination risk (PC).

11 In Basse-Terre, the approach proved to be robust in the sense that a watershed identified as uncontaminated by our
12 water analyses (WC 0) had a very good chance of not being contaminated. Indeed, 75% of these watersheds had no
13 contamination or known risk of contamination of their soils ([WC0, PC0, SC0]). In the remaining 25%, 80% of the
14 plots that were verified by soil analyses were actually found to be uncontaminated. Either these were plots at risk that
15 happened to be uncontaminated, or they were plots that were falsely identified as contaminated in the GIS database
16 (because of a methodological or technical error during the sampling, or an error when entering data). Given the time
17 elapsed between the old and new analyses (6 to 11 years), this cannot be due to the degradation process, unless a high
18 degradation rate is retained, which is not consistent with the current state of soil contamination in Guadeloupe. For the
19 other 20% of these plots, the limitation of the approach seems to be the cause and raises the question of the sources of
20 errors in the different methods of assessing contamination (see 4. Discussion). In watersheds identified as contaminated,
21 68% of water contaminations can be explained by pre-existing soil contamination data. For the remaining 32%, for
22 which there is therefore a lack of information, a terrestrial origin of the contamination could be identified in 82% of the
23 watersheds analysed. In Grande-Terre, our approach has shown certain limitations in view of the limited period during

1 which surface water is present and the low hydrological connectivity between the different points of a watershed (due to
2 the geomorphology of Grande-Terre). However, contaminations could still be detected.

3 **4. Discussion**

4 Our goal was to define a method to determine the state of chlordecone contamination in areas where few analyses had
5 been carried out. This method is based on the assumption that water quality at the outlet of a watershed reflects the state
6 of contamination of its soils. On the whole, our results showed good agreement between the contamination of rivers and
7 the contamination of soils in their watersheds. However, there are some limitations of this method.

8 The first pertains to the physical environment. Indeed, the hypothesis adopted assumes that the watershed corresponds
9 to the hydrogeological basin supplying the river. However, this hypothesis can be challenged (Crabit et al. 2016),
10 especially in the volcanic context of Basse-Terre where water flow paths are affected by successive eruptive episodes
11 and where the transfer routes are mostly underground (Charlier et al. 2008). Thus a river may receive flows from
12 adjacent watersheds or not receive all the flows from the plots within its watershed. These situations may explain the
13 fact that contaminated plots may be present in watersheds that do not present any water contamination at their outlets.

14 The risk map was also found to be incomplete. This map is based on only 3 years of land use for banana cultivation
15 (1969, 1985 and 1997) and only concerns plots that were still declared in 2003 as being used for agriculture (Tillieut
16 2006). It thus incorporates a large amount of uncertainty as to land use during the period of use of chlordecone in
17 Guadeloupe. Furthermore, it cannot account for the diverted use of the chemical on other crops. Although these cases
18 were, no doubt, limited (Levillain et al. 2012), they are nevertheless likely to cause water contamination. These
19 observations lead to the conclusion that the risk map under-represents the state of contamination and explain why
20 certain watersheds can present water contamination without having plots at risk within them (case [WC1, PC0] of table
21 3b). On the other hand, the results of the soil analysis show that an inverse bias exists: plots are classified at risk but are
22 uncontaminated. This phenomenon can arise from the fact that the risks of contamination of plots used to cultivate
23 banana in 1969 or 1997 were taken into account in the map, but they may not have been actually used to cultivate
24 banana during the period of chlordecone use (1972-1993). It can also arise from the extreme variability of agricultural
25 practices (Levillain et al. 2012; Clostre et al. 2014) from one plot to the next, and the difficulty of interpolating levels of
26 practice to a set of plots. Another explanation is that although the molecule is stable, it is nevertheless carried away by
27 the water and thus the contamination of the plots decreases over time (Cabidoche et al. 2009). Depending on rainfall
28 conditions and soil types, some plots at risk may no longer be sources of contamination. Notice that today, the
29 degradation process cannot be invoked to explain why some plots are no longer contaminated. Recent work (Chevallier
30 et al. 2019) shows the value of a better understanding of chlordecone degradation conditions in order to understand the
31 evolution of soil contamination in watersheds.

1 Consequently, the map of the contaminated soil raises questions regarding its reliability. The abnormally high rate of
2 false positives found by comparing soil analyses guided by the map and our campaign of analysis highlights the
3 problem of undertaking the analyses. There are many factors that can skew the results of an analysis: minimum number
4 of samples per plot, improper cleaning of tools before collecting each sample, changes in analytical precision, etc.
5 Finally, the analytical bias can also concern water. To begin with and considering the consistency of river
6 contamination by chlordecone (Crabit et al. 2016; Cattani et al. 2019), collecting a single river sample does not lead to
7 any significant bias in the assessment of pollution: the permanence of contamination suggests that irrespective of the
8 sampling period, the analysis will be positive. A negative sample, on the other hand, may be related to the absence of
9 contamination or correspond to a very low level of contamination, around the detection threshold. Water contamination
10 by the container (glass bottle) seems unlikely given the process of cleaning the flasks by the laboratory and the
11 sampling protocol put in place by our team. Finally, the time taken to send samples to the laboratory, less than 5 days, is
12 sufficiently short to avoid a significant degradation of the intrinsically very stable molecule transported in a refrigerated
13 condition.

14 All these biases cannot be controlled, irrespective of the environment. Therefore, the delimitation of the areas at risk
15 and those not at risk of contamination must combine several approaches. In our case, these approaches are: 1) a
16 hydrological approach based on the dispersal process of the chemical, 2) a historical approach based on practices, and
17 3) a pedological approach based on contamination data that leads to the zoning of the environment for risk management
18 (restrictions on cultivation depending on the danger of contamination, for example). It should also be noted that the
19 method has limited applicability in the case of low water-flow conditions (case of Grande-Terre) and of low levels of
20 contamination since it will require a large number of samples to be taken from across the territory. Finally, the difficulty
21 in drawing up contamination maps highlights the need for plot-level information on agricultural practices. The
22 monitoring of practices remains the principal way to effectively manage the risks of contamination.

23 **5. Conclusions**

24 Our study focused on the delimitation of areas contaminated by a persistent organochlorine used more than 30 years ago
25 in banana plantations in the French West Indies. The originality of our approach was to determine terrestrial
26 contamination of a watershed from the quality of its watercourses, by dividing the territory on a hydrological basis. To
27 our knowledge, this is a new approach in the field of pesticide contamination of soil and water that attempts to take
28 advantage of the functional relationship between soil and river contamination. This approach has made it possible to
29 identify areas at high risk of contamination that were as yet unknown, as well as areas where the probability of
30 contamination was low.

1 The different biases that exist in the acquisition of water data as also in soil analyses and knowledge of practices do not
2 allow for an unequivocal confirmation of the absence of contamination of a geographical area. The saving in costs
3 compared to a systematic soil analysis is however significant (in this case about 219 sampling points led to information
4 on 83% of the surface area of Basse-Terre). This method of analysis of the water at the watershed's outlet thus acquires
5 relevance in the context of exploratory surveys. Furthermore, as it reflects soil pollution trends, this approach may
6 complement monitoring of soil pollution over time. So, when used in combination with all the other information
7 produced (risk map, soil contamination map), this method can become part of an approach for long-term risk
8 management, including the identification of agricultural plots to be targeted as a priority during future analysis
9 campaigns.

10 **Acknowledgements**

11 We would like to thank the French Government as well as the Guadeloupe Water Office which, as part of the National
12 Chlordecone Action Plan III, co-funded this study, as well as the drafting of this article with the help of the European
13 Union and the Guadeloupe Region (ERDF RIVAGE project). We also thank the Department of Food, Agriculture and
14 Forestry (Daaf) of Guadeloupe for providing us with access to their data and for their logistical support; the La Drôme
15 Departmental Laboratory (LDA 26) for having carried out all the analyses; and the technicians of the National Institute
16 of Agronomic Research (Inra) and the Centre for International Cooperation in Agronomic Research for Development
17 (Cirad) for having carried out the sampling campaigns.

18 **References**

- 19 Barron MG, Ashurova ZJ, Kukaniev MA, Avloev HK, Khaidarov KK, Jamshedov JN, Rahmatullova OS, Atolikshoeva
20 SS, Mamadshova SS, Manzenyuk O (2017) Residues of organochlorine pesticides in surface soil and raw
21 foods from rural areas of the Republic of Tajikistan. *Environ Pollut* 224:494–502 .
22 <https://doi.org/10.1016/j.envpol.2017.02.031>
- 23 Cabidoche Y-M, Achard R, Cattan P, Clermont-Dauphin C, Massat F, Sansoulet J (2009) Long-term pollution by
24 chlordecone of tropical volcanic soils in the French West Indies: A simple leaching model accounts for current
25 residue. *Environ Pollut Spec Issue Sect Ozone Mediterr Ecol Plants People Probl* 157:1697–1705
- 26 Cabidoche Y-M, Lesueur-Jannoyer M (2012) Contamination of Harvested Organs in Root Crops Grown on
27 Chlordecone-Polluted Soils. *Pedosphere* 22:562–571 . [https://doi.org/10.1016/S1002-0160\(12\)60041-1](https://doi.org/10.1016/S1002-0160(12)60041-1)
- 28 Cattan P, Cabidoche Y-M, Lacas J-G, Voltz M (2006) Effects of Tillage and mulching on runoff under banana (*Musa*
29 *spp.*) on a tropical Andosol. *Soil Tillage Res* 86:38–51
- 30 Cattan P, Charlier J-B, Clostre F, Letourmy P, Arnaud L, Gresser J, Jannoyer M (2019) A conceptual model of
31 organochlorine fate from a combined analysis of spatial and mid- to long-term trends of surface and ground
32 water contamination in tropical areas (FWI). *Hydrol Earth Syst Sci* 23:691–709 . [https://doi.org/10.5194/hess-](https://doi.org/10.5194/hess-23-691-2019)
33 [23-691-2019](https://doi.org/10.5194/hess-23-691-2019)

- 1 Cattan P, Woignier T, Clostre F, Lesueur Jannoyer M (2016) Heterogeneity of soil pollution. In: Lesueur Jannoyer M,
2 Cattan P, Woignier T, Clostre F (eds) Crisis management of chronic pollution: contaminated soil and human
3 health. CRC Press, Boca Raton, États-Unis, pp 31–44
- 4 Charlier J-B, Cattan P, Moussa R, Voltz M (2008) Hydrological behaviour and modelling of a volcanic tropical
5 cultivated catchment. *Hydrol Process* 22:4355–4370
- 6 Charlier J-B, Cattan P, Voltz M, Moussa R (2009) Transport of a Nematicide in Surface and Groundwaters in a
7 Tropical Volcanic Catchment 10.2134/jeq2008.0355. *J Env Qual* 38:1031–1041
- 8 Charlier J-B, Lachassagne P, Ladouche B, Cattan P, Moussa R, Voltz M (2011) Structure and hydrogeological
9 functioning of an insular tropical humid andesitic volcanic watershed: A multi-disciplinary experimental
10 approach. *J Hydrol* 398:155–170 . <https://doi.org/doi: 10.1016/j.jhydrol.2010.10.006>
- 11 Chevallier ML, Della-Negra O, Chaussonnerie S, Barbance A, Muselet D, Lagarde F, Darii E, Ugarte E, Lescop E,
12 Fonknechten N, Weissenbach J, Woignier T, Gallard J-F, Vuilleumier S, Imfeld G, Le Paslier D, Saaidi P-L
13 (2019) Natural Chlordecone Degradation Revealed by Numerous Transformation Products Characterized in
14 Key French West Indies Environmental Compartments. *Environ Sci Technol* 53:6133–6143 .
15 <https://doi.org/10.1021/acs.est.8b06305>
- 16 Clostre F, Cattan P, Gaude J-M, Carles C, Letourmy P, Lesueur-Jannoyer M (2015) Comparative fate of an
17 organochlorine, chlordecone, and a related compound, chlordecone-5b-hydro, in soils and plants. *Sci Total*
18 *Environ* 532:292–300 . <https://doi.org/10.1016/j.scitotenv.2015.06.026>
- 19 Clostre F, Lesueur-Jannoyer M, Achard R, Letourmy P, Cabidoche Y-M, Cattan P (2014) Decision support tool for soil
20 sampling of heterogeneous pesticide (chlordecone) pollution. *Environ Sci Pollut Res* 21:1980–1992 .
21 <https://doi.org/10.1007/s11356-013-2095-x>
- 22 Cottez S (1972) S.Cottez, Etude des ressources en eau souterraine de la Grande-Terre (Guadeloupe), Rapport de
23 synthèse des résultats acquis entre 1969 et 1972 par le B.R.G.M. et atlas des eaux souterraines de la Grande-
24 Terre, 1972. Fort de France, Martinique
- 25 Crabit A, Cattan P, Colin F, Voltz M (2016) Soil and river contamination patterns of chlordecone in a tropical volcanic
26 catchment in the French West Indies (Guadeloupe). *Environ Pollut* 212:615–626 .
27 <https://doi.org/10.1016/j.envpol.2016.02.055>
- 28 Della Rossa P, Jannoyer M, Mottes C, Plet J, Bazizi A, Arnaud L, Jestin A, Woignier T, Gaude J-M, Cattan P (2017)
29 Linking current river pollution to historical pesticide use: Insights for territorial management? *Sci Total*
30 *Environ* 574:1232–1242 . <https://doi.org/10.1016/j.scitotenv.2016.07.065>
- 31 Frangi J-P, Richard D (1997) Heavy metal soil pollution cartography in northern France. *Sci Total Environ* 205:71–79 .
32 [https://doi.org/10.1016/S0048-9697\(97\)00184-8](https://doi.org/10.1016/S0048-9697(97)00184-8)
- 33 Hellweger FL (1997) AGREE DEM Surface Reconditioning System. University of Texas, Austin, USA
- 34 Levillain J, Cattan P, Colin F, Voltz M, Cabidoche Y-M (2012) Analysis of environmental and farming factors of soil
35 contamination by a persistent organic pollutant, chlordecone, in a banana production area of French West
36 Indies. *Agric Ecosyst Environ* 159:123–132 . <https://doi.org/10.1016/j.agee.2012.07.005>
- 37 Morell M, Jérémie J-J (1994) La Guadeloupe au fil de l'eau. 1994
- 38 SIRS (2015) Délimitation et caractérisation des espaces agricoles en Guadeloupe et à Saint-Martin
- 39 Tillieut O (2006) Cartographie de la pollution des sols de Guadeloupe par la Chlordecone. DAAF & INRA, Basse-Terre
40 (Guadeloupe)
- 41 Woignier T, Fernandes P, Soler A, Clostre F, Carles C, Rangon L, Lesueur-Jannoyer M (2013) Soil microstructure and
42 organic matter: Keys for chlordecone sequestration. *J Hazard Mater* 262:357–364 .
43 <https://doi.org/10.1016/j.jhazmat.2013.08.070>
- 44 Wu X, Zhu L (2019) Prediction of organic contaminant uptake by plants: Modified partition-limited model based on a
45 sequential ultrasonic extraction procedure. *Environ Pollut* 246:124–130 .
46 <https://doi.org/10.1016/j.envpol.2018.11.066>

1 Zhao C, Xie H, Zhang J, Xu J, Liang S (2013) Spatial distribution of organochlorine pesticides (OCPs) and effect of
2 soil characters: A case study of a pesticide producing factory. *Chemosphere* 90:2381–2387 .
3 <https://doi.org/10.1016/j.chemosphere.2012.10.029>

4