



HAL
open science

Immobilized enzymes at work: when surface density matters Context and Objective

Cédric Montanier, Mathieu Fanuel, Hélène Rogniaux, David Ropartz,
Anne-Marie Di Guilmi, Antoine Bouchoux

► **To cite this version:**

Cédric Montanier, Mathieu Fanuel, Hélène Rogniaux, David Ropartz, Anne-Marie Di Guilmi, et al.. Immobilized enzymes at work: when surface density matters Context and Objective. 13th Carbohydrate Bioengineering Meeting, May 2019, Toulouse, France. hal-02966459

HAL Id: hal-02966459

<https://hal.inrae.fr/hal-02966459>

Submitted on 14 Oct 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Immobilized enzymes at work: when surface density matters

Cédric Montanier¹, Mathieu Fanuel², Hélène Rogniaux², David Ropartz²,
Anne-Marie Di Guilmi³, Antoine Bouchoux¹


1- LISBP, Université de Toulouse, CNRS, INRA, INSA, Toulouse, France,
2- INRA, UR1268 Biopolymers Interactions Assemblies, F-44316 Nantes, France,
3- CEA, 18 route du panorama, 92265 Fontenay-aux-roses, France

Context and Objective

In nature, the plant-based organic carbon contained within plant cell walls is mainly recycled by the action of cellulolytic microorganisms, such as bacteria and fungi, which produce complex arrays of cell wall-degrading enzymes. Many of the enzymes that degrade plant cell wall polysaccharides are modular proteins, which contain single or multiple copies of both catalytic domain(s) and CBM(s). Thus, substrate is attacked by different enzymes acting together at different regions, and much more than sum of individual different enzymatic activities, synergism drives the efficiency of such system. Furthermore, in the case of the cellulosome, a large multi-component cell-bound structures, enzymes are localized at close distance to each-other. The benefits of this spatial proximity on the efficiency of the enzymatic reaction are still poorly understood. We investigate this question by using an in-house developed system, Jo-In, where enzymes are immobilized with controlled densities - therefore distances - that can be controlled precisely. The enzyme used is a xylanase that participates to the hydrolysis of plant cell wall polymers, the Xyn11A from *Neocallimastix patriciarum*. **Our approach preserved the intrinsic activity of the enzyme, making the density of grafting the only parameters that is tuned¹.**

Results

1 Precise control of the distance between immobilized enzymes using Jo and In recombinant proteins^{2,3}.



2 Immobilization does not affect enzyme activity; only distance does.

Specific activity using small chromogenic substrate pNP-X₃


Beads	Specific activity SA (μmol.min ⁻¹ .mg ⁻¹)		Free enzymes
	(A) at constant beads volume fraction = 0.3%	(B) at constant total concentration of InNpXyn11A in solution = 4.47 mg/L	
1	2.12 (4.47) ^a	2.14 (0.3%) ^b	2.29 (4.47) ^c
2	2.88 (1.15) ^a	2.14 (0.7%) ^b	2.48 (1.15) ^c
3	2.52 (0.67) ^a	2.10 (1.3%) ^b	2.45 (0.67) ^c
4	2.56 (0.35) ^a	2.12 (2.7%) ^b	2.38 (0.35) ^c
5	2.19 (0.15) ^a	2.10 (5.9%) ^b	2.14 (0.15) ^c

^a the equivalent concentration of enzyme that corresponds to 0.3% of beads volume fraction, as expressed in mg/L, ^b the volume fraction of beads that corresponds to 4.47 mg/L of InNpXyn11A, ^c the concentration of free InNpXyn11A in solution, as expressed in mg/L

Kinetic parameters using beechwood xylan as long chain polymer

	Free InNpXyn11A	Beads 0	Beads 2	Beads 4	Beads 5
Surface density (nmol.m ⁻²)	—	21.2	3.5	1.1	0.5
K _m (mg.mL ⁻¹)	1.8 ± 0.7	3.6 ± 0.5	3.6 ± 0.5	2.6	2.1
k _{cat} (10 ³ min ⁻¹)	46.1 ± 8.8	8.9 ± 0.9	22.4 ± 2.7	43.5	55.1
k _{cat} /K _m (10 ³ min ⁻¹ .mg ⁻¹ .mL)	25.6 ± 6.5	2.5 ± 0.07	6.2 ± 9.5	16.5	26.2

3 Enzyme proximity and hydrolytic profile.



Conclusions

Overall, results show that xylanase molecules can be distanced from 9.5 to 64.4 nm center-to-center. Using small 4-nitrophenyl-β-D-xylotriose as substrate, no modification of the kinetic parameters is observed compare to the enzyme in solution. However, when long polymer beechwood xylan is used as substrate, kinetic parameters are affected with higher density of grafting. The product profile was analyzed by HPAEC-PAD and MALDI-TOF. Data indicate that immobilized enzyme product profiles are different from those produced by the enzymes dispersed in solution; the immobilized enzymes release more short oligo-saccharides and oligomers with average DP more homogenous. Our results question the relationship between spatial proximity and synergistic effect as encountered in the cellulosome.