



**HAL**  
open science

# Seedbed structure of major field crops as affected by cropping systems and climate: Results of a 15-year field trial

Jay-Ram Lamichhane, Jean Boiffin, Hubert Boizard, Carolyne Dürr, Guy Richard

## ► To cite this version:

Jay-Ram Lamichhane, Jean Boiffin, Hubert Boizard, Carolyne Dürr, Guy Richard. Seedbed structure of major field crops as affected by cropping systems and climate: Results of a 15-year field trial. *Soil and Tillage Research*, 2021, 206, 10p. 10.1016/j.still.2020.104845 . hal-03012868

**HAL Id: hal-03012868**

**<https://hal.inrae.fr/hal-03012868v1>**

Submitted on 7 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Seedbed structure of major field crops as affected by cropping systems and**  
2 **climate: Results of a 15-year field trial**

3

4 Jay Ram Lamichhane<sup>1\*</sup>, Jean Boiffin<sup>2</sup>, Hubert Boizard<sup>2</sup>, Carolyne Dürr<sup>2</sup>, Guy Richard<sup>3</sup>

5

6 <sup>1</sup>INRAE, Université Fédérale de Toulouse, UMR 1248 AGIR, F-31326 Castanet-Tolosan,

7 France

8 <sup>2</sup>Formerly INRA, UPR1158 Agro-Impact, site d'Estrées-Mons, F-80203 Péronne, France,

9 and now retired

10 <sup>3</sup>INRAE, UAR1241 DEPE, 147 rue de l'Université, 75338 PARIS Cedex 07, France;

11

12 \*Corresponding author: [jay-ram.lamichhane@inrae.fr](mailto:jay-ram.lamichhane@inrae.fr)

13 Tel: +33 (0)5 61 28 52 50; Fax: +33 (0)5 61 28 55 37

14

15 **Abstract**

16 Seedbed structure directly or indirectly affects crop establishment by modifying seed-

17 soil contact, acting as mechanical obstacles or modifying temperature, moisture and

18 oxygen contents of seedbed as well as the dynamic of pests, pathogens and weeds.

19 However, very few detailed descriptions of seedbed's structure of major field crops exist

20 to date, in terms of precise aggregate size distributions in relation to different factors

21 including the cropping system (CS), soil and climatic conditions and their interactions.

22 Here, we characterized seedbeds of major European field crops across three CSs over a

23 15-year period (1991-2005) of a long-term field experiment. CS I was the succession of

24 spring pea/winter wheat/oilseed rape/winter wheat. Likewise, CSs II and III were the

25 succession of sugar beet/winter wheat/maize/winter wheat with different sowing dates  
26 based on two different decision rules aimed at: minimizing the risk of soil compaction in  
27 the CS II or maximizing the duration of the crop in the CS III. We classified three types of  
28 seedbed structure, namely fine (with >20mm soil aggregates <15%), intermediate (with  
29 >20mm soil aggregates >15<25%), and coarse (with >20mm soil aggregates >25%). We  
30 found a statistically significant effect of CSs ( $P < 0.05$ ), crops ( $P < 0.001$ ), previous crops  
31 ( $P < 0.01$ ) and year ( $P < 0.001$ ) while no significant effect of wheel traffic ( $P > 0.05$ ) and  
32 tillage ( $P > 0.05$ ) was observed on the soil aggregates > 20 mm. No irreversible effect of  
33 the CSs was observed over the study period on seedbed structure and the consequent  
34 risks of generating coarser seedbeds, which is unfavorable for crop establishment. This  
35 dataset offers a unique description of the seedbed structure variations for major  
36 European field crops. This information can be used for future simulation studies of crop  
37 emergence, using, for example, the SIMPLE model that has seedbed structure as one of  
38 the input variables.

39

40

41 **Key words:** seedbed preparation, seedbed structure, soil aggregates, cropping systems

42

43

## 44 **1. Introduction**

45 Seedbed structure can significantly affect early growth and development of the crop, by  
46 altering abiotic components of the seedbed including soil-plant contact, mechanical  
47 forces exerted on plant or plant parts, soil aeration, thermal regime and water balance  
48 (Dexter 1988, 2004a). In addition, seedbed structure can also affect the impact of early  
49 biotic stresses and the effectiveness of weed control (Glen et al. 1989; Bale et al. 1992;  
50 Otten and Gilligan 2006; Finney and Creamer 2008; Melander and Kristensen 2011). A  
51 seedbed containing a high proportion of large soil aggregates not only leads to a poor  
52 seed-soil contact but it also cools down more rapidly thereby slowing down the seed  
53 imbibition and germination process (Brown et al. 1996; Håkansson et al. 2002).  
54 Consequently, the time interval between the seed germination and seedling emergence  
55 phase in a seedbed comprising bigger soil aggregates is longer, due to the increased  
56 tortuosity of the seedling path before reaching the soil surface (Boiffin et al. 1992).  
57 Likewise, the risk of seedling death before emergence is higher in coarser seedbeds as  
58 seedlings can be trapped under the soil aggregates encountered during its elongation  
59 from the sowing depth (Dürr and Aubertot 2000). Delayed emergence also increases the  
60 risk of seedling death in an indirect way; for example, by increasing the probability of  
61 attacks due to soil-borne pests and pathogens owing to longer heterotrophic phase or  
62 topsoil crust formation impeding the emergence (Gallardo-Carrera et al. 2007). Lastly,  
63 delayed emergence may lead to a reduced growth rate (Tamet et al. 1995; Dürr and  
64 Boiffin 1995). On the other hand, a high proportion of very fine aggregates in the topsoil  
65 layer is not always beneficial as it induces a higher sensitivity to crust formation under  
66 rainfall (Boiffin 1986). The soil structural state of a given field plot evolves over time  
67 because of a dynamic combination of compaction and fragmentation process (Richard et

68 al. 1999; Obour et al. 2017;). This evolution results from the interaction between climate  
69 and crop management history including crop rotation, management techniques applied  
70 to these crops such as sowing and harvesting operations, equipment used for these  
71 operations, and the soil water content at the time of field operations (Guérif et al. 2001;  
72 Boizard et al. 2002; Dexter 2004b; Tagar et al. 2020). Following the harvest of a crop, the  
73 soil profile contains a more or less important volume of compacted zones created by  
74 wheel tracks, especially when harvest or antecedent field operations occur under wet  
75 conditions and with heavy machineries. Beginning from this “initial” structural state,  
76 seedbed structure of the following crop results from a more or less simplified sequence  
77 of tillage operations, including (or not) mouldboard ploughing, and, finally, the action of  
78 the driller itself. Depending on the soil moisture profile and on the related soil  
79 consistency, these tillage operations not only induce fragmentation of previously  
80 compacted soil volumes and big clods but also compaction or coalescence of fine  
81 aggregates, especially across deep layers located under wheel tracks. In addition to  
82 tillage, other types of field operations with a passage of wheel tracks may also lead to  
83 soil compaction (e.g. spraying fertilizers). Soil structure can also be greatly altered by  
84 climatic alternations (e. g. freezing–thawing or wetting-drying), which exert a more or  
85 less intense and deep effect of fragmentation, depending on soil texture, and on the  
86 frequency and amplitude of these alternations (Stengel et al. 1984; Kværnø and  
87 Øygarden 2006). Taken together, the seedbed structure obtained with a given set of  
88 tillage equipment reflects the combined prints of the last sequence of tillage and/or  
89 drilling, the recent intercultural phase, and the anterior history of the field. Based on  
90 these factors and their interactions, cropping systems (CSs) could result in different and  
91 more or less favorable trends affecting seedbed structure. These trends may or may not  
92 be reversible. For example, a progressive increase of the compacted volumes in the soil

93 profile results in an increased occurrence of seedbeds with a coarse or heterogeneous  
94 soil structure (Boizard et al. 2002) and this could affect the quality of crop  
95 establishment. Therefore, it is important to describe and analyze the inter-annual  
96 variability of seedbed structure, in relation to contrasted and well-characterized CSs,  
97 especially in order to detect potential patterns of progressive evolution.

98 A previous study (Boizard et al. 2002), based on a field experiment, investigated  
99 cumulative effects of CSs on the structural state of the tilled layer, in particular the  
100 proportion of compacted zones in a loamy soil that is characteristic of northern  
101 European soils. The authors showed that the compaction level of a soil was dependent  
102 on the soil moisture at the time of field operations as well as the characteristics of the  
103 machinery used and that there was no indication of irreversible cumulative degradation.

104 Another study (Boizard et al. 2013), compared the impact of conventional *versus*  
105 reduced tillage on the soil structure evolution and showed that the soil structure in the  
106 untilled layer mainly depends on the soil compaction intensity and that regeneration of  
107 this compacted layer over time was slower compared with that of the tilled layer.

108 Although seedbeds in this long-term experiment were characterized, no study yet  
109 reported whether and to what extent the seedbed structure of major field crops can be  
110 affected by crop management history of a plot in interaction with climate. Seedbed  
111 characterization is not only time consuming and resource intensive but also difficult to  
112 perform due to limited field access, especially under rainy seasons and high moisture  
113 conditions. Only little knowledge is available to date concerning precise numerical data  
114 characterizing seedbed structure and its variations (Braunack and Dexter 1989;  
115 Braunack and McPhee 1991; Aubertot et al. 1999; Gallardo-Carrera et al. 2007).

116 Therefore, the key objective of this work was to characterize seedbed structures of  
117 major field crops grown in Europe, and analyze how they are affected by CSs, climate

118 and their interactions. The generated information will be useful for future simulation  
119 studies concerning crop establishment in a context of climatic and agricultural change.

120

## 121 **2. Material and methods**

### 122 2.1. Study sites

123 Most of the data presented in this study comes from a long-term field experiment (1991-  
124 2005) carried out at Mons-en-Chaussée (49°52'44"N 3°00'27"E), Northern France. A  
125 detailed description of the study site and soil texture and chemical characteristics have  
126 been published (Boizard et al. 2002). The soil is a Haplic Luvisol (FAO classification).  
127 Briefly, the soil at the 0-30 cm soil horizon had the following characteristics: 0.197 g.g<sup>-1</sup>  
128 clay, 0.747 g.g<sup>-1</sup> silt and 0.056 g.g<sup>-1</sup> sand; 0 g.g<sup>-1</sup> CaCO<sub>3</sub>, 0.095 g.g<sup>-1</sup> C, 0.001 g.g<sup>-1</sup>N, C/N  
129 ratio 9.3, and pH 7.7. Soil water contents at -10, -50, -100 and -1500 kPa were 0.252,  
130 0.213, 0.164 and 0.083 g.g<sup>-1</sup>, respectively. Water content at field capacity, measured  
131 during winter in field 2–3 days after excess water had drained away (Hillel 1971), was  
132 0.24 g g<sup>-1</sup>. The average annual air temperature and cumulative rainfall were 9.6 °C and  
133 690 mm, respectively. The cumulated rainfall one week before seedbed preparation for  
134 each crop is presented in **Table 1**.

### 135 2.2. Cropping systems

136 The long-term field experiment was set up to investigate the potential impact of  
137 contrasted CSs on soil structure changes, in the tilled or non-tilled layers (Guérif et al.  
138 2001; Boizard et al. 2002). Three CSs were designed taking into account key agricultural  
139 features of the region. These factors included possible crops and rotations, and time  
140 schedule for field operations. The rotation in the cropping system I was spring pea (SP;  
141 *Pisum sativum* L.)/winter wheat (WW; *Triticum aestivum* L.)/winter oilseed rape (OR;  
142 *Brassica napus* L.)/ WW. Sowing and harvesting were always carried out either in

143 summer or early autumn, *i.e.* during the dry period of the year, except for pea sowing in  
144 early spring. SP was sown between 7 March and 7 April, OR was sown between 10 and  
145 31 August, while WW was sown between 8 and 20 October. The rotation in the CSs II  
146 and III was sugar beet (SB; *Beta vulgaris* L.)/WW/maize (M; *Zea mays* L.)/WW. The CS II  
147 was managed to avoid possible soil compaction during sowing and harvesting of the  
148 three crops. Consequently, SB was sown between 20 March and 30 April, M was sown  
149 between 3 April and 4 May while WW was sown between 8 October and 19 November.  
150 The soil water content was measured for each field operation by taking samples at 0-5,  
151 5-10, 10-15 et 15-20 cm soil horizon (four replicates at each depth) per plot, weighed  
152 and dried 24 h at 105 °C as described previously (Boizard et al. 2002). When the  
153 decision had to be made quickly during the day, drying is carried out with a microwave  
154 on a reduced sample (60 g). The decision-making rules applied for seedbed preparation  
155 and sowing of SB and M in the CS I and CS II were with  $<0.20 \text{ g g}^{-1}$  and  $<0.22 \text{ g g}^{-1}$  water  
156 content in the 0–10 cm and 10–20 cm soil layer, respectively. In contrast, the CS III was  
157 managed to maximize light interception and yield by SB and M canopies without taking  
158 into account any possible compaction during field operations. The decision rules applied  
159 for seedbed preparation and sowing was with  $<0.25 \text{ g g}^{-1}$  water content in the 0–10 cm  
160 soil layer. Therefore, SB and M were sown in early spring (between 4 and 17 March and  
161 between 1 and 23 April, respectively) and harvested in late autumn, during wet periods  
162 of the year. Consequently, sowing of WW in the CSs I and II occurred earlier and they  
163 were very similar in terms of dates while WW in the CS III occurred later (*i.e.* between 4  
164 and 21 November) compared with the previous two CSs. The harvesting dates of WW  
165 were similar independent of the CSs.

166 Each crop from each CS was grown every year leading to four plots per CS (12 plots in  
167 total) that were replicated in two blocks (24 plots in total). As explained above, the rules



168 for decision making in each CS, depended on a combination of crop physiology  
169 requirements and field access for seedbed preparation (Boizard et al. 2002). The key  
170 agronomic rule consisted in avoiding soil compaction by considering pre-determined  
171 thresholds of soil water content at the 0-10 and 10-20 cm soil horizons. Major field  
172 operations used following the harvest of a preceding crop and before tillage were:  
173 chopping (only for M) and stubble disking (only for SP and OR). Seedbed preparation  
174 was followed by sowing within a maximum of 24 hours for all crops.

175 A detailed description of the equipment used for seedbed preparation and sowing is  
176 presented in **Table 2**. Conventional tillage was applied to each crop every year from the  
177 beginning of the experiment up to 1999. A new treatment, which consisted of reduced  
178 tillage at a 4 to 8 cm depth, was introduced in 1999 into the experiment in order to  
179 compare the effects of conventional and reduced tillage on soil structure evolution. For  
180 conventional tillage treatment, the field plots were subjected to a 30 cm depth  
181 mouldboard ploughing (for all crops) followed either by seedbed preparation with a  
182 combination harrow (for SB, M and OR), or combined seedbed preparation (for WW and  
183 SP). Mouldboard ploughing before spring crops was carried out between November and  
184 January to take advantage of the effect of climatic conditions (freezing thawing and  
185 wetting drying) on the soil surface horizon. Mouldboard ploughing for WW and OR was  
186 carried out just before sowing. The depth of the seedbed layer ranged from 3 to 9 cm  
187 (average 5.5 cm). Stubble tillage was performed every year after WW, M, SP and OR at a  
188 depth ranging from 6.5 to 10 cm (average 8 cm). Seedbed preparation was performed  
189 with a combination harrow (with several rows of small tines and two rows of rollers) at  
190 6-8 cm depth, for SB, M and OR. WW and SP were sown using a combined rotary harrow  
191 and disc drill carrying out tillage and sowing in one pass. The equipment used for  
192 seedbed preparation had similar characteristics in terms of weight (6.5–8 Mg), tire

193 width (0.70 m), working widths (3 m), and inflation pressure (70 kPa). In contrast, the  
194 harvesting equipment was much heavier (about 15 Mg) with a high inflation pressure  
195 (200-300 kPa), and a wide variation in the percentage of the experimental plot covered  
196 by wheel tracks (29–77%). In reduced tillage treatment, the only difference was that a  
197 compact disc cultivator at 6 cm average depth (ranging from 4 to 8 cm; **Table 2**)  
198 replaced mouldboard ploughing.

### 199 2.3. Cumulated rainfall before seedbed preparation and sowing

200 Rainfalls were daily recorded at the meteorological station located at the study site.  
201 Rainfalls one week before seedbed preparation were cumulated to get an indicator of  
202 the soil moisture conditions for soil tillage and sowing.

203 Cumulated rainfall values 7 days before sowing for all three CSs are presented in **Table**  
204 **1**. These values could markedly vary between years, especially for WW sowing (0 to 52  
205 mm). As expected, the highest and the most variable cumulated rainfall values were  
206 observed in the CS III while the lowest and least variable values were registered in the  
207 CS I, because of the sown crops and the decision rules. For all CSs, the highest cumulated  
208 rainfall values were for WW while the lowest values were in the CS I for OR

209

### 210 2.4. Characterization of soil compaction level

211 The soil compaction level caused by soil traffic, in interaction with soil moisture, can  
212 affect the seedbed structure. A higher level of soil compaction may lead to the formation  
213 of bigger soil aggregates that may affect the quality of crop establishment with a higher  
214 rate of seedling mortality under clods. The soil compaction level was assessed following  
215 each crop establishment as previously described (Boizard et al. 2002, 2013). A  
216 morphological description of the soil structure of the 0–30 cm layer was carried after  
217 each sowing, from a randomly located 3 m wide soil profile perpendicular to the tillage

218 and wheeling directions. We used the “profil cultural” method proposed by Manichon  
219 (1987) and presented in detail in Boizard et al. (2002, 2013) and Roger-Estrade et al.  
220 (2004), to map soil structure in this tilled layer. The highly compacted zones containing  
221 specific morphological features (no visible macro-pores, a massive structure and a  
222 smooth breaking surface) were identified in slight relief on the observation face. The  
223 proportion of highly compacted zones in the tilled layer under the seedbed was  
224 calculated. The working depth was measured by digging a pit perpendicular to the  
225 direction of tillage following each tillage operation (3 m wide at sowing and 1 m wide at  
226 stubble tillage). The tilled layer was visually delimited and its thickness was measured  
227 every 10 cm laterally.

228 In addition to assessing the soil compaction level at the plot scale, we investigated the  
229 potential effect of wheel-induced compaction along the row due to wheel traffic on  
230 seedbed structure, compared with the control treatment (*i.e.* the zone outside the  
231 passage of wheel) as described previously (Boizard et al. 2002, 2013). Working widths  
232 markedly differed from one operation to the others (2.7–5.4 m) that led to an important  
233 variation in the percentage of the experimental plot covered by wheel tracks (11–44%).  
234 The location of the wheel tracks was recorded after each field operation on a transect  
235 over the plot width.

## 236 2.5. Seedbed characterization

237 The distribution of seedbed aggregate size was characterized as described previously  
238 (Aubertot et al. 1999). Seedbed samples were taken just after sowing. A surface was  
239 delimited along the row of the seedbed with combs (20 length x 10 width x 10 cm depth)  
240 to determine either the numbers, mass or percentage of aggregates or all of them in a  
241 precise soil volume. The number of replicate/year/crop ranged from 10 to 48 for SP, 16  
242 to 24 for OR, 10 to 48 for WW, 12 to 48 for SB, and 16 to 48 for M. Soil samples from all

243 the treatments (i.e. the zones subjected and non-subjected to wheel traffic from  
244 conventionally tilled plots as well as from plots with reduced tillage) were carefully  
245 extracted with a spoon, brought to the laboratory, and air-dried. The samples were  
246 sieved with a gently shaking machine (30 s, 50-mm amplitude) and grades <5, 5-10, 10-  
247 20, 20-30, 30-40, 40-50, and >50 mm diameter were obtained. The number, the mass  
248 and the volume for each of these grades were determined. Finally, the observed  
249 variations, in terms of percentages of soil aggregates, measured for each seedbed class of  
250 crops belonging to the three cropping system, were used to define three types of  
251 seedbed structures: a fine seedbed (with >20mm soil aggregates <15%), an  
252 intermediate seedbed (with >20mm soil aggregates >15<25%), and a coarse seedbed  
253 (with >20mm soil aggregates >25%). A visual aspect of these three seedbed types is  
254 reported in **Figure 1**.

## 255 2.6. Statistical analyses

256 The variability of seedbed structure can be related to several fixed factors, mainly the CS,  
257 crop and year. These fixed factors influence seedbed structure by determining sowing  
258 equipment and the soil conditions at sowing, including soil moisture and compactness at  
259 the time of seedbed preparation. A 3-step statistical analysis was performed: i) the effect  
260 of the main fixed factors on soil moisture and compactness at sowing was tested through  
261 a one-way ANOVA, ii) possible relationships between seedbed structure and soil  
262 moisture or compactness were investigated through linear correlation tests, and iii) an  
263 ANCOVA, with the dependent variable (the seedbed structure represented by % soil  
264 aggregates >20 mm), the independent variables considered as fixed factors (the CS, crop,  
265 previous crop and year, and additionally, the type of tillage and the location of wheel  
266 tracks in the seedbed), and covariates (the % of soil moisture at 0-15 cm and that of the

267 topsoil compaction level). All statistical analyses were computed using JASP version  
268 0.13.0 (Love et al. 2019).

269

### 270 **3. Results**

271 3.1. Types of seedbed structure in relation to year, crops and cropping systems

272 The types of seedbed structure found over the 15 years of the experiment are reported  
273 in **Figure 2**. When analyzed crop-wise for each individual year, the fine seedbed  
274 structure was the most frequently found for SP, SB and M in all three CSs tested,  
275 followed by the intermediate and coarse seedbed structure. In contrast, the coarse  
276 seedbed structure followed by the intermediate one prevailed for WW, especially in the  
277 CS III and, to a lower extent, in the CS II. When analyzed year-wise for each CS, the  
278 intermediate seedbed structure was the most frequent followed by fine and coarse  
279 seedbed for all but in the CSs III for which coarse and intermediate seedbed structures  
280 were the most frequent and in equal number.

281 3.2. Topsoil moisture and compaction at sowing

282 Results of the seedbed moisture level at 0-15 cm soil horizon at the time of seedbed  
283 preparation are reported in **Figure 3**. The moisture level ranged from 19 to 21% and it  
284 was significantly higher ( $P < 0.001$ ) in the CS III followed by the CSs I and II. When  
285 analyzed by crop, the moisture level at 0-15 cm was significantly higher ( $P < 0.001$ ) for SP  
286 (22%), followed by SB (21%), WW (20%) and M (19%). No data were recorded for OR.  
287 The topsoil compaction level, observed after sowing, as analyzed by ANOVA ranged from  
288 0.17 to 0.41% for all crops and the CSs combined (**Figure 4**). When individually  
289 analyzed by the CS, the compaction level was significantly higher ( $P < 0.001$ ) in the CS III  
290 (0.41%) followed by the CSs II (0.28%) and I (0.17%). Likewise, when analyzed by crop,

291 the topsoil compaction level of SB (0.36%) and M (0.30%) was significantly higher  
292 ( $P < 0.001$ ) compared with that of WW (0.27%) and SP (0.20%).

### 293 3.3. Correlations between seedbed structure and soil moisture or topsoil 294 compaction level

295 When analyzed by CSs, there was a positive correlation between the seedbed moisture  
296 at 0-15 cm soil horizon at the time of seedbed preparation and the % of soil aggregates >  
297 20 mm (**Table 3**). Likewise, a positive correlation was observed between the topsoil  
298 compaction level and the soil aggregates > 20 mm (Table 3). When analyzed by crop, the  
299 seedbed moisture level at 0-15 cm soil horizon was positively correlated with the soil  
300 aggregates > 20 mm (Table 4). Similarly, there was a positive correlation between the  
301 topsoil compaction level and the soil aggregates > 20 mm (**Table 4**).

### 302 3.4. Combined effects of fixed factors and covariables on seedbed structure

303 ANCOVA showed a statistically significant effect ( $P < 0.001$ ) of both the covariates (the  
304 % of soil moisture at 0-15 cm and that of the topsoil compaction level) on the soil  
305 aggregates > 20 mm (**Table 5**). There was also a significant effect of CSs ( $P < 0.05$ ), crops  
306 ( $P < 0.001$ ), previous crops ( $P < 0.01$ ) and year ( $P < 0.001$ ) while no significant effect of  
307 wheel traffic ( $P > 0.05$ ) and tillage ( $P > 0.05$ ) was observed on the soil aggregates > 20 mm.  
308 The seedbed structure, expressed as the % of soil aggregates > 20 mm, was the coarsest  
309 in the CS III ( $26 \pm 15$  %) followed by the CS II ( $22 \pm 14$  %) and the CSI ( $18 \pm 10$  %).  
310 Among crops, WW seedbeds were the coarsest ( $26 \pm 14$  %) followed by those of SB ( $19$   
311  $\pm 13$  %), M ( $16 \pm 10$  %) and SP ( $15 \pm 9$  %). As for previous crops, WW seedbeds  
312 prepared after the harvest of SB were the coarsest ( $31 \pm 19$  %), followed by M ( $30 \pm 18$   
313 %), SP ( $22 \pm 13$  %) and OR ( $20 \pm 13$  %). Finally, a significant ( $P < 0.001$ ) inter-annual  
314 fluctuation of the seedbed structure was observed over the 15-year period with the least

315 and most coarsest seedbed structure observed in 2003 ( $14 \pm 14$  %) and 2001 ( $30 \pm 18$   
316 %), respectively.

#### 317 **4. Discussion**

318 In this study we analyzed a unique dataset on seedbed structure, resulting from various  
319 crop types and CSs put in place and observed during 15 consecutive years. This  
320 combination of crops, CSs and years generated a wide range of soil structure dynamics  
321 and seedbed structures. Differences between CS and crop types, large inter-annual  
322 fluctuations, and a variable differentiation of the CSs according to year and crop types  
323 were the dominant features of the variability in seedbed structure.

324 Independent of the fixed factors, aggregate size in the seedbed was correlated to topsoil  
325 compactness and moisture at the time of seedbed preparation and sowing, with a  
326 tendency to coarser seedbeds associated with profiles that are more compact and wet.  
327 This trend is consistent with the initial hypotheses that (i) seedbed structure partly  
328 reflects the importance of continuous and compact soil volumes at the time of seedbed  
329 preparation, and that (ii) this influence is modulated by the loosening efficiency of tillage  
330 tools, which itself depends on soil consistency resulting from soil moisture. Starting  
331 from a dry soil with a maximum level of cohesion, the increase of soil water content  
332 decreases soil cohesion and facilitates fragmentation up to an optimal threshold value  
333 above which pressures exerted by the working utensils result in plastic deformations  
334 rather than ruptures (Guérif 1982; Guérif et al. 1988). Once this threshold is reached,  
335 any increase in soil water content should then reduce fragmentation, and induce the  
336 formation of new clods by coalescence of small aggregates.

337 The ranking of CS (III II and I, in decreasing order of coarseness) is consistent with the  
338 effect of these systems on soil compactness. Harvests and seedbed preparations  
339 generally occurred in wetter soil conditions in the CS III than in the CS I and II. They

340 induced higher pressures on topsoil, as well as larger wheel tracks areas, for the CS II  
341 and III than for the CS I. Consequently, the CS III corresponded to higher levels of soil  
342 compaction, followed by the CS II and I, as previously mentioned by Boizard et al (2013)  
343 and confirmed here with supplementary data (**Figs. S1-S3**).

344 The observed contrast between seedbeds of different crop types (coarser seedbeds for  
345 WW compared with those of SB and M) is commonly observed in the agricultural  
346 context of Northern Europe, in relation with the fact that emergence rate and seedbed  
347 structure requirements for precision sowing crops are higher than those for WW or OR.  
348 Consequently, tillage operations for preparing SB or M seedbeds are generally higher  
349 than for WW or OR (2 vs 1 in our case). The contrast also might result from two  
350 additional effects. First, a higher proportion of WW plots resulted from the late harvest  
351 of the preceding crop (SB and M) under wet conditions leading to a high level of soil  
352 compaction. Second, WW sowings followed shorter intercultural periods with less  
353 wetting-drying and no freezing-thawing alternations.

354 The seedbed types distribution does not reflect simple effects of CS or crop type, but a  
355 stronger differentiation of CS for WW than for SB and M seedbeds. Crop-specific  
356 modalities of seedbed preparation and sowing (autumn vs spring, 1 vs 2 tillage  
357 operations, specific sowing equipment) combined with different durations of the  
358 intercrop sequences (short vs long) can modulate the influence of antecedent  
359 compaction on seedbed structure. Short intervals between harvests and sowing dates of  
360 the preceding crop (case of WW) tend to maintain the antecedent contrasts in soil  
361 structure, while long intervals including freezing-thawing and wetting-drying  
362 alternations (case of SB and M) tend to reduce them.

363 The influence of CSs and crop types on soil moisture at sowing is less contrasted than  
364 their effect on topsoil compaction level. Higher soil moisture in the CS III than in the II



365 on average is again consistent with the differences in sowing times induced by the  
366 decision rules, with later sowing dates for autumn-sown crops and earlier sowing dates  
367 for spring-sown crops in the CS III, and consequently higher rainfall before sowing on  
368 average. Independent of the fixed factors and except for OR, most of the variation of  
369 topsoil moisture at sowing occur in a relatively narrow range of water potential (-100 to  
370 -50 kPa). For this type of soil texture, such range corresponds to well drained but still  
371 plastic soils. The trend to positive correlations between soil moisture at sowing and the  
372 proportion of coarse aggregates, which was observed on average, confirms that soil  
373 consistency at seedbed preparation varied within the plastic domain. Then higher soil  
374 moistures obtained some years in the CS III in relation to decision rules, were an  
375 additional cause of coarser seedbeds.

376 There was no difference in seedbed structure due to the presence or absence of wheel  
377 tracks during seedbed preparation. This confirms that the soil conditions for seedbed  
378 preparation and seeding were appropriately chosen to avoid significant compaction  
379 under the wheel tracks that, otherwise, may have led to reduced loosening and a coarser  
380 structure. This means that variations in seedbed structure were generally determined  
381 by the topsoil compaction generated before rather than during seedbed preparation.  
382 The type of tillage (conventional vs. reduced) did not have any effect on seedbed  
383 structure although, depending on the tillage system. The under representation of the  
384 reduced tillage treatment, introduced only in 2000, could explain the lack of significant  
385 effect that warrants an in-depth further investigation.

386 The importance of inter-annual fluctuations of seedbed structure, in interaction with  
387 the crop types and CSs, reflects the influence of climatic conditions on the different  
388 processes influencing soil structural dynamics, from the harvest of the preceding crop to  
389 the sowing of the following crop. Years with high or low levels of large aggregates in

390 seedbed alternate in an apparently erratic way. Indeed, no irreversible trend could be  
391 obviously detected within or among the CSs and that the seedbed structural differences  
392 between the CSs observed at the end of the period (**Fig. 5**) were not larger from those  
393 detected at the beginning. The year sequences 1991-1995-1999-2003, 1992-1996-2000-  
394 2004, 1993-1997-2001-2005 (only for WW), exactly corresponded to the same groups  
395 of experimental fields in each case, and did not exhibit a progressive trend. However, for  
396 the year sequences 1994-1998-2002 and 1995-1999-2003, which again corresponded  
397 to the same groups of experimental fields, the largest differences between the CSs  
398 occurred at the end of the sequence. For these sequences, it is impossible to neither  
399 exclude nor emphasize a cumulative effect of the CSs, due to the lack of the monitoring  
400 and observations in 2006. This is again consistent with the fact that seedbed structure  
401 variability was mainly inherited from a relatively short-term soil dynamics. The impact  
402 of the CSs on seedbed structure was important, but essentially summed up to an effect of  
403 the antecedent crop on the subsequent seedbed structure. This antecedent effect not  
404 only depends on the crop species, but also on the decision-making rules applied to the  
405 different CSs. The different set of rules applied in the experiment have led to various  
406 conditions of harvest and seedbed preparation, with contrasted inter-annual rainfall  
407 accumulations the week before seedbed preparations, and contrasted soil moisture  
408 levels at 0-15 cm soil horizon. Our findings corroborate with a previous study that  
409 showed no long-term irreversible effect of CSs on the tilled soil layer (Boizard et al.  
410 2002). The apparent reversibility of cumulative degradation of seedbed structure should  
411 be related to soil texture, and more precisely to the nature and amount of the clay  
412 content. This latter determines the soil aptitude to cracking (Stengel and Guérif 1985).  
413 However, this reversibility should not be extrapolated to soils with clay content <16%

414 neither to CS more aggressive than the CS III, as for example CS with sequences of  
415 successive maize with late harvests.

## 416 **5. Conclusions**

417 The results presented in this study represent an important dataset on seedbed  
418 structures obtained across a wide diversity of sowing contexts, in terms of sowing  
419 season (spring, summer and autumn), impact of cropping systems with regard to topsoil  
420 compaction risks, tillage intensity and climatic conditions during the year affecting  
421 seedbed preparation. Consequently, the observed variability of seedbed structure was  
422 large, with a higher proportion of coarser seedbeds when all three factors were  
423 combined. First, a high level of topsoil compaction at the time of seedbed preparation,  
424 related to harvest conditions of the preceding crop. Second, a short interval between  
425 sowing and harvest of the preceding crop, reducing the occurrence of climatic episodes  
426 able to induce fragmentation of compacted zones in the topsoil. Third, a high soil  
427 moisture at the time of seedbed preparation, resulting in less loosening and even  
428 clumping of previous aggregates.

429 Our results confirm the initial assumption of an influence of cropping systems on  
430 seedbed structure. Cropping systems with high risks of topsoil compaction correspond  
431 to coarser seedbeds, especially for crops sown in autumn following late harvests.  
432 However, no cumulative and irreversible differentiation between CSs could be detected  
433 in our experiment. This means that the impact of repeated soil degradation on seedbed  
434 structure does not prevail on the reversible and annually variable effects of the  
435 intercultural periods. Even so, the influence of CSs on seedbed structure is important,  
436 and involves not only the nature of crops included in the crop rotation, but also the  
437 strategies which determine the techniques of crop management, and especially the time  
438 period for field operations. The soil aptitude to cracking increases the reversibility of

439 compaction that results from more or less aggressive CSs and therefore it should be  
440 considered as a key criterion for extending or restricting the trends observed in this  
441 experiment.

442 The dataset presented in this work represents an important input variable for  
443 simulation studies using the SIMPLE crop emergence models (Dürr et al. 2001), which  
444 takes into account the impact of seedbed structure on seedling emergence. The seedbed  
445 classification proposed in this study may also facilitate the identification of any seedbed  
446 structure, for which no data are available to date.

447

#### 448 **Acknowledgements**

449 We thank the technical team of the INRA's domain in Mons-en-Chaussée for supporting  
450 the long-term experiment, and in particular Paul Régnier, Charles Leforestier, Daniel  
451 Boitez, Caroline Dominiarczyk, Bertrand Chauchard, Frédéric Mahut who efficiently  
452 contributed to the management of the long-term experiment and characterization of  
453 seedbed structure.

454

#### 455 **References**

456

457

458 Aubertot, J. N., Dürr, C., Kieu, K., and Richard, G. 1999. Characterization of Sugar Beet  
459 Seedbed Structure. *Soil Sci. Soc. Am. J.* 63:1377–1384.

460

461 Bale, J. S., Ekebuisi, M., and Wright, C. 1992. Effect of seed bed preparation, soil structure  
462 and release time on the toxicity of a range of grassland pesticides to the carabid beetle  
463 *Pterostichus melanarius* (Ill.) (Col., Carabidae) using a microplot technique. *J. Appl.*  
464 *Entomol.* 113:175–182.

465

466

467 Boiffin, J. 1986. Stages and time-dependency of soil crusting in situ. In *Proceedings of the*  
468 *International Symposium on the Assessment of Soil Surface Sealing and Crusting*, eds. F  
469 Callebaut, D Gabriels, and M de Boodt. Ghent, Belgium, p. 91–98.

470

471 Boiffin, J., Dürr, C., Fleury, A., Marinlafleche, A., and Maillet, I. 1992. Analysis of the  
472 variability of sugar-beet (*Beta vulgaris* L.) growth during the early stages. *Agronomie.*  
473 12:515–525.

474

475 Boizard, H., Richard, G., Roger-Estrade, J., Dürr, C., and Boiffin, J. 2002. Cumulative effects  
476 of cropping systems on the structure of the tilled layer in northern France. *Soil Tillage*  
477 *Res.* 64:149–164.

478  
479 Boizard, H., Yoon, S. W., Leonard, J., Lheureux, S., Cousin, I., Roger-Estrade, J., et al. 2013.  
480 Using a morphological approach to evaluate the effect of traffic and weather conditions  
481 on the structure of a loamy soil in reduced tillage. *Soil Tillage Res.* 127:34–44.  
482  
483 Braunack, M. V, and Dexter, A. R. 1989. Soil aggregation in the seedbed: A review. I.  
484 Properties of aggregates and beds of aggregates. *Soil Tillage Res.* 14:259–279.  
485  
486 Braunack, M. V, and McPhee, J. E. 1991. The effect of initial soil water content and tillage  
487 implement on seedbed formation. *Soil Tillage Res.* 20: 5–17.  
488  
489 Brown, A. D., Dexter, A. R., Chamen, W. C. T., and Spoor, G. 1996. Effect of soil  
490 macroporosity and aggregate size on seed-soil contact. *Soil Tillage Res.* 38: 203-2016.  
491  
492 Dexter, A. R. 1988. Advances in characterization of soil structure. *Soil Tillage Res.*  
493 11:199–238.  
494  
495 Dexter, A. R. 2004a. Soil physical quality: Part I. Theory, effects of soil texture, density,  
496 and organic matter, and effects on root growth. *Geoderma.* 120:201–214.  
497  
498 Dexter, A. R. 2004b. Soil physical quality: Part II. Friability, tillage, tilth and hard-setting.  
499 *Geoderma.* 120:215–225.  
500  
501 Dürr, C., and Aubertot, J.-N. 2000. Emergence of seedlings of sugar beet (*Beta vulgaris* L.)  
502 as affected by the size, roughness and position of aggregates in the seedbed. *Plant Soil.*  
503 219:211–220.  
504  
505 Dürr, C., Aubertot, J. N., Richard, G., Dubrulle, P., Duval, Y., and Boiffin, J. 2001. SIMPLE: a  
506 model for SIMulation of PLant Emergence predicting the effects of soil tillage and sowing  
507 operations. *Soil Sci. Soc. Am. J.* 65:414–442.  
508  
509 Dürr, C., and Boiffin, J. 1995. Sugarbeet seedling growth from germination to first leaf  
510 stage. *J. Agric. Sci.* 124:427–435.  
511  
512 Finney, D. M., and Creamer, N. G. 2008. Weed management on organic farms. The  
513 Organic Production Publication Series, CEFS, p 1–34.  
514  
515 Gallardo-Carrera, A., Léonard, J., Duval, Y., and Dürr, C. 2007. Effects of seedbed structure  
516 and water content at sowing on the development of soil surface crusting under rainfall.  
517 *Soil Tillage Res.* 95:207–217.  
518  
519 Gardarin, A., Coste, F., Wagner, M.-H., and Dürr, C. 2016. How do seed and seedling traits  
520 influence germination and emergence parameters in crop species? A comparative  
521 analysis. *Seed Sci. Res.* 26:317–331.  
522  
523 Glen, D. M., Milsom, N. F., and Wiltshire, C. W. 1989. Effects of seed-bed conditions on  
524 slug numbers and damage to winter wheat in a clay soil. *Ann. Appl. Biol.* 115:177–190.  
525  
526 Guérif, J. 1982. Compactage d'un massif d'agrégats : effet de la teneur en eau et de la

527 pression appliquée. *Agronomie*. 2:287–294.  
528  
529 Guérif, J., Richard, G., Dürr, C., Machet, J. ., Recous, S., and Roger-Estrade, J. 2001. A review  
530 of tillage effects on crop residue management, seedbed conditions and seedling  
531 establishment. *Soil Tillage Res.* 61:13–32.  
532  
533 Guérif, J., Royère, J., and Grison, D. 1988. Résistance en traction des agrégats terreux :  
534 influence de la texture, de la matière organique et de la teneur en eau. *Agronomie*.  
535 8:379–386.  
536  
537 Håkansson, I., Myrbeck, Å., and Etana, A. 2002. A review of research on seedbed  
538 preparation for small grains in Sweden. *Soil Tillage Res.* 64: 23-40.  
539  
540 Hillel, D. 1971. *Soil and water: physical principles and processes. Physiological Ecology:*  
541 *A series of Monographs, Texts and Treatises.* Academic Press, New York, pp. 162–165.  
542  
543 Kværnø, S. H., and Øygarden, L. 2006. The influence of freeze–thaw cycles and soil  
544 moisture on aggregate stability of three soils in Norway. *CATENA*. 67:175–182.  
545  
546  
547 Love, J., Selker, R., Marsman, M., Jamil, T., Dropmann, D., Verhagen J., et al. 2019. JASP:  
548 Graphical statistical software for common statistical designs. *J. Stat. Softw.* 88:  
549 10.18637/jss.v088.i02.  
550  
551 Manichon, H. 1982. Influence des systé`mes de culture sur le profil cultural: e´laboration  
552 dune me´thode de diagnostic base´e sur l’observation morphologique. The`se. Institut  
553 National Agronomique Paris-Grignon, Paris.  
554  
555 Melander, B., and Kristensen, J. K. 2011. Soil steaming effects on weed seedling  
556 emergence under the influence of soil type, soil moisture, soil structure and heat  
557 duration. *Ann. Appl. Biol.* 158:194–203  
558  
559 Obour, P. B., Lamandé, M., Edwards, G., Sørensen, C. G., and Munkholm, L. J. 2017.  
560 Predicting soil workability and fragmentation in tillage: a review. *Soil Use Manag.*  
561 33:288–298.  
562  
563 Otten, W., and Gilligan, C. A. 2006. Soil structure and soil-borne diseases: using  
564 epidemiological concepts to scale from fungal spread to plant epidemics. *Eur. J. Soil Sci.*  
565 57:26–37 Available at: <https://doi.org/10.1111/j.1365-2389.2006.00766.x>.  
566  
567 Richard, G., Boizard, H., Roger-Estrade, J., Boiffin, J., and Guérif, J. 1999. Field study of soil  
568 compaction due to traffic in northern France: pore space and morphological analysis of  
569 the compacted zones. *Soil Tillage Res.* 51:151–160  
570  
571 Roger-Estrade, J., Richard, G., Caneill, J., Boizard, H., Coquet, Y., Defossez, P., et al. 2004.  
572 Morphological characterisation of soil structure in tilled fields: from a diagnosis method  
573 to the modelling of structural changes over time. *Soil Tillage Res.* 79:33–49  
574

575 Stengel, P., Douglas, J. T., Guérif, J., Goss, M. J., Monnier, G., and Cannell, R. Q. 1984.  
576 Factors influencing the variation of some properties of soils in relation to their  
577 suitability for direct drilling. *Soil Tillage Res.* 4:35–53

578  
579 Stengel, P., and Guérif, J. 1985. Effect of tillage techniques on cracking of a heavy clay  
580 soil. In the *10<sup>th</sup> Conference of ISTRO, Guelph, Canada*, p. 29.

581  
582 Tagar, A. A., Adamowski, J., Memon, M. S., Do, M. C., Mashori, A. S., Soomro, A. S., et al.  
583 2020. Soil fragmentation and aggregate stability as affected by conventional tillage  
584 implements and relations with fractal dimensions. *Soil Tillage Res.* 197:104494

585  
586 Tamet, V., Souty, N., and Rode, C. 1995. Emergence des plantules de carotte (*Daucus*  
587 *carotta* L) sous des obstacles mecaniques super ciels. *Agronomie.* 15:109–121.

588  
589  
590

### 591 **Figure legends:**

592 **Figure 1.** Characteristic seedbed structure of major field crops sown in Europe. Three  
593 types of seedbed structure are identified based on the percentage of soil aggregates >20  
594 mm. Coarse seedbed (>25% of soil aggregates >20 mm; A), intermediate seedbed  
595 (>15<25% volume of soil aggregates >20 mm; B), and fine seedbed (<15% of soil  
596 aggregates >20 mm; C).

597  
598 **Figure 2.** Percentage of mean annual soil aggregates >20 mm diameter in the seedbed of  
599 each crop for the three CSs. Results are based on the combined data measured from  
600 plots subjected to both conventional and reduced tillage. CS: cropping system; NT: not  
601 tested; SP: spring pea; WW: winter wheat; OR: oilseed rape, SB: sugar beet; M: maize.  
602 The values in each cell represents the percentage of soil aggregates >20mm diameter  
603 measured in the seedbed. Red, yellow and green colours indicate coarse, intermediate  
604 and fine seedbed, respectively.

605  
606 **Figure 3.** Moisture level (%) at 0-15 cm soil horizon at the time of seedbed preparation  
607 as determined by cropping system (a) and crop (b). CS I: cropping system I  
608 (pea/wheat/oilseed rape/wheat), CS II: second cropping system (sugar  
609 beet/wheat/maize/wheat), and CS III: third cropping system (sugar  
610 beet/wheat/maize/wheat). Error bars on the graph represent standard deviation.  
611 Means followed by the same letter are not significantly different at  $P < 0.05$

612  
613 **Figure 4.** Topsoil compaction level (%) as affected by cropping system (a) and crop (b).  
614 CS I: cropping system I (pea/wheat/oilseed rape/wheat), CS II: second cropping system  
615 (sugar beet/wheat/maize/wheat), and CS III: third cropping system (sugar  
616 beet/wheat/maize/wheat). Error bars on the graph represent standard deviation. No  
617 data were available for oilseed rape. Means followed by the same letter are not  
618 significantly different at  $P < 0.05$ .

619  
620  
621

622 **Figure 3.** Correlation between the topsoil moisture (%) and the percentage of soil  
623 aggregates >20 mm (a), and the topsoil compaction level (%) and the percentage of soil  
624 aggregates >20 mm (b) as affected by cropping systems. Results are based on the  
625 combined data measured from plots subjected to both conventional and reduced tillage.  
626 CS I: cropping system I (pea/wheat/oilseed rape/wheat), CS II: second cropping system  
627 (sugar beet/wheat/maize/wheat), and CS III: third cropping system (sugar  
628 beet/wheat/maize/wheat). No data were available for oilseed rape. Specific values of  
629 Pearson's correlation coefficients are detailed in Table 3.

630

631 **Figure 4.** Correlation between the topsoil moisture (%) and the percentage of soil  
632 aggregates >20 mm (a), and the topsoil compaction level (%) and the percentage of soil  
633 aggregates >20 mm (b) as affected by tillage practice. Results are based on the combined  
634 data measured from all cropping systems. Specific values of Pearson's correlation  
635 coefficients are detailed in Table 3.

636

637 **Figure 5.** Seedbed of spring (a) and winter (b) crops containing >20 mm aggregates  
638 over the period 1991-2005. Results are based on the combined data measured from  
639 plots subjected to both conventional and reduced tillage. Spring crops were represented  
640 by sugar beet and maize while winter crops were represented by wheat. The soil  
641 aggregate similarity observed between the beginning and the end of the period for all  
642 cropping systems indicates no irreversible effect of cropping system over time. Values  
643 related to every four year period correspond to the same groups of experimental fields.  
644 Error bars on the graph represent standard deviation.



645 **Table 1.** Cumulated rainfall (mm  $\pm$  standard deviation) one week before seedbed preparation for each crop belonging to the three  
646 cropping systems tested from 1991 to 2005. The time range of the cumulated rainfall was 1<sup>st</sup> February for pea, 1<sup>st</sup> to 20<sup>th</sup> March for sugar  
647 beet, 1<sup>st</sup> to 20<sup>th</sup> April for maize, 25<sup>th</sup> August for oilseed rape, and 10<sup>th</sup> October for winter wheat. Seedbed preparation was followed by  
648 sowing within a maximum of 24 hours for all crops.  
649

Year	Cropping system I			Cropping system II				Cropping system III				
	Pea	Wheat	Oilseed rape	Wheat	Sugar beet	Wheat	Maize	Wheat	Sugar beet	Wheat	Maize	Wheat
1990	ND	2 $\pm$ 0	ND	2 $\pm$ 0	ND	2 $\pm$ 0	ND	2 $\pm$ 0	ND	2 $\pm$ 0	ND	2 $\pm$ 0
1991	11 $\pm$ 2	5 $\pm$ 1	ND	5 $\pm$ 1	8 $\pm$ 1	5 $\pm$ 1	6 $\pm$ 1	5 $\pm$ 1	28 $\pm$ 7	9 $\pm$ 2	0 $\pm$ 0	9 $\pm$ 2
1992	2 $\pm$ 1	5 $\pm$ 2	ND	5 $\pm$ 2	3 $\pm$ 1	5 $\pm$ 2	6 $\pm$ 1	5 $\pm$ 2	22 $\pm$ 6	14 $\pm$ 3	16 $\pm$ 3	14 $\pm$ 3
1993	1 $\pm$ 0	28 $\pm$ 7	ND	28 $\pm$ 7	1 $\pm$ 0	28 $\pm$ 7	6 $\pm$ 1	28 $\pm$ 7	1 $\pm$ 0	6 $\pm$ 2	17 $\pm$ 2	6 $\pm$ 2
1994	10 $\pm$ 2	0 $\pm$ 0	ND	0 $\pm$ 0	7 $\pm$ 1	0 $\pm$ 0	1 $\pm$ 0	0 $\pm$ 0	8 $\pm$ 2	20 $\pm$ 2	19 $\pm$ 3	20 $\pm$ 2
1995	16 $\pm$ 2	1 $\pm$ 0	ND	1 $\pm$ 0	0 $\pm$ 0	1 $\pm$ 0	7 $\pm$ 2	1 $\pm$ 0	10 $\pm$ 3	1 $\pm$ 0	15 $\pm$ 4	1 $\pm$ 0
1996	0 $\pm$ 0	3 $\pm$ 1	ND	3 $\pm$ 1	1 $\pm$ 0	3 $\pm$ 1	0 $\pm$ 0	3 $\pm$ 1	1 $\pm$ 0	19 $\pm$ 4	1 $\pm$ 0	19 $\pm$ 4
1997	1 $\pm$ 0	24 $\pm$ 8	ND	24 $\pm$ 8	1 $\pm$ 0	7 $\pm$ 2	13 $\pm$ 3	7 $\pm$ 2	4 $\pm$ 1	17 $\pm$ 4	0 $\pm$ 0	17 $\pm$ 4
1998	0 $\pm$ 0	7 $\pm$ 2	1 $\pm$ 0	7 $\pm$ 2	1 $\pm$ 0	7 $\pm$ 2	2 $\pm$ 1	7 $\pm$ 2	4 $\pm$ 1	46 $\pm$ 8	1 $\pm$ 0	46 $\pm$ 8
1999	0 $\pm$ 0	4 $\pm$ 1	0 $\pm$ 0	4 $\pm$ 1	3 $\pm$ 1	4 $\pm$ 1	1 $\pm$ 0	4 $\pm$ 1	8 $\pm$ 2	1 $\pm$ 0	3 $\pm$ 1	1 $\pm$ 0
2000	3 $\pm$ 1	36 $\pm$ 6	ND	36 $\pm$ 6	4 $\pm$ 1	36 $\pm$ 6	ND	36 $\pm$ 6	ND	ND	11 $\pm$ 2	ND
2001	12 $\pm$ 2	33 $\pm$ 8	ND	33 $\pm$ 8	0 $\pm$ 0	1 $\pm$ 0	20 $\pm$ 3	1 $\pm$ 0	ND	32 $\pm$ 9	12 $\pm$ 2	32 $\pm$ 9
2002	0 $\pm$ 0	6 $\pm$ 1	ND	6 $\pm$ 1	0 $\pm$ 0	8 $\pm$ 2	0 $\pm$ 0	8 $\pm$ 2	10 $\pm$ 2	52 $\pm$ 10	0 $\pm$ 0	52 $\pm$ 10
2003	0 $\pm$ 0	11 $\pm$ 2	ND	11 $\pm$ 2	1 $\pm$ 0	11 $\pm$ 2	0 $\pm$ 0	11 $\pm$ 2	1 $\pm$ 0	11 $\pm$ 2	0 $\pm$ 0	11 $\pm$ 2
2004	2 $\pm$ 0	9 $\pm$ 3	ND	9 $\pm$ 3	ND	9 $\pm$ 3	ND	9 $\pm$ 3	ND	1 $\pm$ 0	ND	1 $\pm$ 0
2005	ND	ND	ND	ND	6 $\pm$ 1	ND	26 $\pm$ 5	ND	0 $\pm$ 0	ND	5 $\pm$ 2	ND

ND: not determined

650 **Table 2.** Main characteristics of the equipment used for tillage, seedbed preparation and  
 651 sowing for different crops belonging to the three cropping systems  
 652

Cropping system	Previous crop	Crop	Conventional tillage <sup>a</sup>	Reduced tillage <sup>b</sup>	Seedbed preparation
I	Pea	Winter wheat	Mouldboard ploughing	NP	CSP & S (unique passage)
	Winter wheat	Oilseed rape	Mouldboard ploughing	NP	CSP & S (unique passage)
	Oilseed rape	Winter wheat	Mouldboard ploughing	NP	CH
	Winter wheat	Pea	Mouldboard ploughing	NP	CSP & S (unique passage)
	Pea	Winter wheat	NP	Compact disc cultivator	CSP & S (unique passage)
	Winter wheat	Oilseed rape	NP	Compact disc cultivator	CSP & S (unique passage)
	Oilseed rape	Winter wheat	NP	Compact disc cultivator	CH (1 <sup>st</sup> passage)
	Winter wheat	Pea	NP	Compact disc cultivator	CSP & S (unique passage)
II and III	Sugar beet	Winter wheat	Mouldboard ploughing	NP	CH (1 <sup>st</sup> passage)
	Winter wheat	Maize	Mouldboard ploughing	NP	CSP & S (unique passage)
	Maize	Winter wheat	Mouldboard ploughing	NP	CH (1 <sup>st</sup> passage)
	Winter wheat	Sugar beet	Mouldboard ploughing	NP	CSP & S (unique passage)
	Sugar beet	Winter wheat	NP	Compact disc cultivator	CH (1 <sup>st</sup> passage)
	Winter wheat	Maize	NP	Compact disc cultivator	CSP & S (unique passage)
	Maize	Winter wheat	NP	Compact disc cultivator	CH (1 <sup>st</sup> passage)
	Winter wheat	Sugar beet	NP	Compact disc cultivator	CSP & S (unique passage)

<sup>a</sup>Conventional tillage was performed at 30 cm depth, using a 4-bottom mouldboard plough (41 cm wide) pulled by a tractor (Case IH 956).  
<sup>b</sup>Reduced tillage was performed at 10 cm depth, using Rubbin (4m width) combining two rows of discs and two rows of rotors (MX 150).  
 CSP & S: Combined seedbed preparation and sowing was performed with an unique passage of Rotary harrow and seeder (Case IH 956) for winter wheat and pea; CH & S: Combination harrow, which was performed combining several rows of small tines pulled by a tractor (Case IH 956) for oilseed rape, sugar beet and maize, followed by S: sowing (2<sup>nd</sup> passage) using a 12-row seeder (Renault 851.4 for oilseed rape and MF 575 for sugar beet); NP: Not performed.

653  
 654

655 **Table 3.** Pearson's correlation between soil moisture (%) at 0-15 cm and soil aggregates  
 656 >20 mm (%), and between the topsoil compaction level (%) and soil aggregates >20 mm  
 657 (%) as affected by cropping systems.  
 658

<b>Treatment</b>	<b>Correlation between soil aggregates &gt;2 cm (%) and</b>	<b>All CSs combined</b>	<b>CS I</b>	<b>CS II</b>	<b>CS III</b>
Conventional tillage	Soil moisture at 0-15 cm (%)	0.25	0.0 8	0.2 1	0.31
	Soil compaction (%)	0.41	0.3 5	0.3 2	0.47
Reduced tillage	Soil moisture at 0-15 cm (%)	0.50	0.6 0	0.5 2	0.37
	Soil compaction (%)	0.20	- 0.0 2	0.0 1	- 0.12
Combined (Conventional & reduced tillage)	Soil moisture at 0-15 cm (%)	0.31	0.1 8	0.3 5	0.32
	Soil compaction (%)	0.34	0.2 1	0.2 4	0.33

CSs: cropping ststems; CS I: cropping system I (pea/wheat/oilseed rape/wheat), CS II: second cropping system (sugar beet/wheat/maize/wheat), and CS III: third cropping system (sugar beet/wheat/maize/wheat).

659  
660

661 **Table 4.** Pearson's correlation between soil moisture (%) at 0-15 cm and soil aggregates  
 662 >20 mm (%), and between the topsoil compaction level (%) and soil aggregates >20 mm  
 663 (%) as affected by crops.  
 664  
 665

Treatment	Correlation between soil aggregates >2 cm (%) and	Pea	Whe at	Sugar beet	Maiz e
Conventional tillage	Soil moisture at 0-15 cm (%)	0.59	0.30	0.32	0.25
	Soil compaction (%)	0.42	0.65	0.49	0.12
Reduced tillage	Soil moisture at 0-15 cm (%)	0.67	0.73	0.31	0.20
	Soil compaction (%)	-	0.23	-0.18	-0.26
Combined (Conventional & reduced tillage)	Soil moisture at 0-15 cm (%)	0.57	0.40	0.33	0.32
	Soil compaction (%)	0.28	0.48	0.35	0.07

666  
 667


668 **Table 5.** Mean ( $\pm$ SD) values of seedbed structure (% aggregates > 2 cm) computed by an  
669 analysis of co-variance (ANCOVA). The analysis was performed taking into account the  
670 soil moisture at 0-15 cm and the soil compaction level as a co-variables, and cropping  
671 system, crop, previous crop, wheel traffic, tillage, and year as fixed treatments.  
672

Treatment	Fixed factor	n	Aggregates > 2cm $\pm$ SD (%)	Effect of treatment	Effect of co-variables	
					Soil moisture at 0-15 cm (%)	Soil compaction level at 0-15 cm (%)
Cropping system	CS I	230	18 $\pm$ 10			
	CS II	258	22 $\pm$ 14	*	***	***
	CS III	252	26 $\pm$ 15			
Crop	Maize	128	16 $\pm$ 10			
	Pea	62	15 $\pm$ 9	***	***	***
	Sugar Beet	130	19 $\pm$ 13			
	Wheat	390	26 $\pm$ 14			
Previous crop	Maize	680	30 $\pm$ 18			
	Oilseed rape	342	20 $\pm$ 13	**	***	***
	Pea	296	22 $\pm$ 13			
	Sugar beet	481	31 $\pm$ 19			
Wheel traffic	Along the wheel traffic	369	22 $\pm$ 14			
	Outside the wheel traffic	371	22 $\pm$ 14	ns	***	***
Tillage	Conventional tillage	590	21 $\pm$ 12			
	Reduced tillage	150	24 $\pm$ 19	ns	***	***
Year	1991	199	16 $\pm$ 11			
	1992	211	22 $\pm$ 17			
	1993	218	26 $\pm$ 18			
	1994	285	27 $\pm$ 16			
	1995	192	24 $\pm$ 14			
	1996	409	27 $\pm$ 13			
	1997	128	17 $\pm$ 10			
	1998	220	17 $\pm$ 11	***	***	***
	1999	294	21 $\pm$ 14			
	2000	178	24 $\pm$ 14			
	2001	176	30 $\pm$ 18			
	2002	196	25 $\pm$ 23			
	2003	177	14 $\pm$ 14			
2004	191	15 $\pm$ 15				
2005	154	21 $\pm$ 16				

\*\*\*P < 0.001; \*\*P < 0.01; \*P < 0.05; ns: not significant; n: sample size; SD: standard deviation

674  
675  
676

**Figure 1.**



677  
678

679 **Figure 2.**

680

Year	Cropping system I				Cropping system II				Cropping system III			
	SP	WW	OR	WW	SB	WW	M	WW	SB	WW	M	WW
1991	12	NT	NT	NT	12	NT	13	NT	16	NT	12	NT
1992	20	15	NT	9	10	13	13	17	6	33	11	17
1993	13	23	NT	22	17	36	26	27	10	54	10	35
1994	37	33	NT	31	20	34	18	29	21	50	17	38
1995	23	32	NT	23	22	37	22	33	32	39	29	34
1996	17	25	NT	21	6	27	17	24	16	31	7	37
1997	10	17	NT	21	12	26	11	25	20	NT	16	18
1998	10	17	NT	13	6	16	8	10	14	40	5	19
1999	14	19	26	21	12	17	13	26	27	40	16	39
2000	19	27	16	20	12	23	12	43	27	41	14	34
2001	10	28	NT	29	12	35	19	26	13	62	23	24
2002	12	31	NT	NT	17	36	11	42	19	55	11	56
2003	9	12	NT	NT	20	56	15	15	15	17	7	23
2004	NT	13	NT	NT	17	25	26	29	18	23	9	15
2005	NT	13	NT	NT	NT	8	NT	NT	NT	NT	NT	NT

Year	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
CS I	12	15	19	34	26	21	16	13	20	21	22	22	11	13	13
CS II	13	13	27	25	29	19	19	10	17	23	23	27	27	24	8
CS III	14	17	27	32	34	23	18	20	31	29	31	35	16	16	NT

CS I	SP	WW	OR	WW	CS II	SB	WW	M	WW	CS III	SB	WW	M	WW
		17	22	21		23		18	29		17	26		21

Previous crop	SP	22	OR	23	SB	29	M	26	SB	29	M	30

	Coarse seedbed (>20 mm soil aggregates >25%)
	Intermediate seedbed (>20 mm soil aggregates >15<25%)
	Fine seedbed (>20 mm soil aggregates <15%)

681

682



683

684

685

686


687 **Figure 3.**



688  
689  
690  
691




692 **Figure 4.**  
693  
694  
695



696  
697  
698  
699  
700  
701

702 **Figure 5.**  
703



704  
705  
706