Transcriptional integration of the plant responses to iron availability

Fei Gao, Christian Dubos

► To cite this version:

Fei Gao, Christian Dubos. Transcriptional integration of the plant responses to iron availability. *Journal of Experimental Botany*, 2020, 2056-2070 (6), pp.2056-2070. 10.1093/jxb/eraa556 . hal-03031725

HAL Id: hal-03031725

<https://hal.inrae.fr/hal-03031725>

Submitted on 30 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives| 4.0 International License

Transcriptional integration of the plant responses to iron availability

Fei Gao¹ & Christian Dubos^{1,*}

¹ BPMP, Univ Montpellier, CNRS, INRAE, Institut Agro, Montpellier, France

* Corresponding author (Tel: 0033 (0)499 61 28 18)

E-mails: Fei Gao, fei.gao@supagro.fr; Christian Dubos, christian.dubos@inrae.fr

Highlights: Within this review, current knowledge on the control of iron homeostasis in plants is presented and future prospects to improve our understanding of this complex mechanism are provided.

Abstract

Iron is one of the most important micronutrient for plant growth and development. It functions as the enzyme cofactor or component of electron transport chains in various vital metabolic processes, including photosynthesis, respiration and amino acid biosynthesis. For maintaining iron homeostasis, and therefore preventing any deficiency or excess that could be detrimental, plants have evolved complex transcriptional regulatory networks to tightly control iron uptake, translocation, assimilation and storage. Such regulatory networks are composed of various transcription factors among them members from the basic helix-loop-helix (bHLH) family play an essential role. Here, we first review recent advances in understanding the roles of bHLH transcription factors involved in the regulatory cascade controlling iron homeostasis in the model plant *Arabidopsis*, extending it to rice and other plant species. The importance of other classes of transcription factors will also be discussed. Secondly, we elaborate on the posttranslational mechanisms involved in the regulation of these regulatory networks. Finally, we give some perspectives on future research that should be conducted in order to precisely depict how plants control the homeostasis of this micronutrient.

Key words: *Arabidopsis thaliana*, basic helix-loop-helix, bHLH, iron homeostasis, rice, transcription factor.

Introduction

Iron (Fe) is an essential micronutrient for almost all living organisms. In humans, iron deficiency anemia is a major global health issue affecting about one billion people worldwide (Camaschella, 2015). Therefore, increasing the iron content in plants, especially in crops (biofortification), would have enormous benefits for human health. To achieve this goal, a critical question is to uncover the mechanisms controlling iron homeostasis in plants. Like in humans, iron is an essential microelement for plant growth and development. It functions in various vital metabolic processes (e.g. photosynthesis, respiration, amino acid biosynthesis) by acting as cofactor for several metalloproteins or component of electron transport chains (Hänsch and Mendel, 2009; Touraine *et al.*, 2019). However, the excess of iron is deleterious to plants because of its capacity to interact with oxygen, generating reactive oxygen species (ROS) via the Fenton Reaction. Thus, the levels of iron in plant cells must be tightly regulated to avoid iron deficiency or iron excess, both of which severely affecting crops yield and the quality of their derived products (Briat *et al.*, 2015).

Although iron is the fourth most-abundant element on earth, much of it is not readily available for plant use due to the poor solubility of its main (hydro)oxides, especially in neutral-to-alkaline soils (Guerinot and Yi, 1994; Colombo *et al.*, 2018). To adapt to low iron conditions and acquire iron from soil, higher plants have evolved two different strategies (Marschner and Römheld, 1994) (Figure 1). Dicots and non-graminaceous monocots (non-grass species) employ the reduction strategy (named Strategy I). The graminaceous utilize the chelation strategy (named Strategy II). In strategy I, non-grass plants are able to acidify the rhizosphere to promote iron solubility through protons release and mobilize Fe^{3+} through coumarins (*i.e.* fraxetin and sideretin) or riboflavins secretion (Santi and Schmidt, 2009; Fourcroy *et al.*, 2016; Robe *et al.*, 2020). Following iron mobilization, Fe^{3+} is reduced into Fe^{2+} (a more soluble form of iron) that is subsequently transported into the root epidermal cells via high affinity iron transporters from the Zrt/Irt-like protein (ZIP) family.

In *Arabidopsis thaliana*, the release of protons into the rhizosphere is insured by the H^{+} -ATPASE 2 (AHA2; Santi and Schmidt, 2009) whereas the reduction of iron and its translocation into the roots are ensured by FERRIC REDUCTION OXIDASE 2 (FRO2) and IRON-REGULATED TRANSPORTER1 (IRT1), respectively (Brumbarova *et al.*, 2015; Connorton *et al.*, 2017). AHA2, FRO2 and IRT1 associate into a complex at the surface of the

root epidermal cells most likely allowing optimizing iron acquisition and therefore its uptake (Martín-Barranco *et al.*, 2020). Coumarins secretion is ensured by the PLEIOTROPIC DRUG RESISTANCE 9/ATP-BINDING CASSETTE G37 (PDR9/ABCG37) transporter (Fourcroy *et al.*, 2014; Fourcroy *et al.*, 2016). In strategy II, plants biosynthesize and secrete phytosiderophores of the mugineic acid family (MAs) into the rhizosphere to chelate Fe^{3+} (Kobayashi *et al.*, 2014). Fe^{3+} -MAs complexes are then transported into root cells by transporters of the YELLOW STRIPE 1 (YS1) and YELLOW STRIPE 1-like (YSL) family (Inoue *et al.*, 2009; Murata *et al.*, 2006). Interestingly, it was recently shown that some plant species might use both strategies. For instance, when grown in waterlogged soil condition, rice (*Oryza sativa*), a graminaceous species, acquires Fe^{2+} from the soil through the activity of two Fe^{2+} transporters, OsIRT1 and OsIRT2 (Ishimaru *et al.*, 2006). The secretion of caffeic and protocatechuic acids via the PHENOLICS EFFLUX ZERO 2 (PEZ2) transporter participates to Fe^{3+} mobilization and reduction into Fe^{2+} (Bashir *et al.*, 2011). If iron acquisition is the first step that participates to the maintenance of iron homeostasis in plants, it should be noted that it is not the sole mechanism involved in this process. Iron translocation, compartmentalization, assimilation and storage are also important processes required for the maintenance of iron homeostasis at the cellular and subcellular levels, throughout the whole plant body (Kobayashi and Nishizawa, 2012; Kobayashi *et al.*, 2019).

Gene regulation is a crucial step for coping with iron fluctuations. For instance, *FRO2* and *IRT1* expression is induced when iron availability is low whereas the expression of the genes encoding the iron storage ferritin proteins (i.e. *FER1*, 3 and 4) is induced in response to iron excess (Tissot *et al.*, 2019). How plants control iron homeostasis, by regulating the expression of genes involved in the various facets of this complex mechanism, was a critical question for the past three decades. To address this question, several studies mostly based on forward and reverse genetic approaches were conducted, leading to the identification and characterization of several transcription factors (TFs). These studies, mostly conducted in *Arabidopsis* and rice, allowed establishing regulatory networks controlling iron homeostasis in which basic helix-loop-helix (bHLH) TFs play a preponderant role (Gao *et al.*, 2019; Gao *et al.*, 2020a; Li *et al.*, 2019) (Figure 2 and Table 1). bHLH proteins form one of the largest families of TFs (Heim *et al.*, 2003) known to modulate several facets of plant growth and development, including cell differentiation, secondary metabolite biosynthesis, hormone signaling or the responses to environmental factors (Carretero-Paulet *et al.*, 2010). Indeed,

additional TFs and protein regulating TFs activity were also identified (Yan *et al.*, 2016; Palmer *et al.*, 2013; Rodríguez-Celma *et al.*, 2019a).

Within this review, current knowledge on the control of iron homeostasis by TFs, especially from the bHLH family, will be presented and discussed with a special emphasis on the latest findings. The transcriptional and posttranslational regulation of the iron homeostasis regulatory networks will also be documented. Last, some perspectives on future research to be conducted in order to improve our understanding of this complex mechanism will be provided.

The transcriptional regulation of iron homeostasis in strategy I plants, a preponderant role for bHLH TFs

The model plant *Arabidopsis* has allowed the identification of several factors involved in the regulation of iron homeostasis, notably by studying its response to iron deficiency. Such studies highlighted that the regulation of iron homeostasis was essentially occurring at the transcriptional level and was involving several TFs, in particular those of the bHLH family. This topic has also been covered in recent reviews by Gao *et al.*, 2019; Kobayashi, 2019; Kobayashi *et al.*, 2019; Wu and Ling, 2019; Schwarz and Bauer, 2020.

To date, at least six bHLH TF subfamilies (Heim *et al.*, 2003), encoding 17 different proteins, are known to participate to the maintenance of iron homeostasis in *Arabidopsis*. These bHLH TFs form an intricate regulatory network composed of two interconnected regulatory modules (Figure 3).

The first module relies on the activity of FIT/bHLH29 (FER-LIKE IRON DEFICIENCY INDUCED TRANSCRIPTION FACTOR), a clade IIIa bHLH TF (Colangelo and Gueriot, 2004; Jakoby *et al.*, 2004; Yuan *et al.*, 2005). FIT/bHLH29 is a direct regulator of *IRT1* and *FRO2* expression, highlighting its central role for the regulation of the iron uptake machinery (Wang *et al.*, 2013b). FIT/bHLH29 activity relies on its interaction with the four members of the Ib bHLH clade (*i.e.* bHLH38, bHLH39, bHLH100 and bHLH101), forming heterodimer complexes displaying partial redundant activities (Colangelo and Gueriot, 2004; Maurer *et al.*, 2014; Yuan *et al.*, 2008; Wang *et al.*, 2013b) (Figure 3). Recently, the members of the IVa bHLH clade (*i.e.* bHLH18, bHLH19, bHLH20 and bHLH25) were identified as FIT/bHLH29 interacting proteins (Cui *et al.*, 2018). These interactions were shown to

promote the degradation of FIT/bHLH29 via the 26S proteasome pathway, in a jasmonic acid-dependent manner (Cui *et al.*, 2018). It is noteworthy that the Ib bHLH and the IVa bHLH TFs antagonize the activity of each other in regulating FIT/bHLH29 protein accumulation to tightly regulate the iron uptake machinery in response to different environmental stimuli (Cui *et al.*, 2018) (Figure 3).

The second module acts upstream from FIT/bHLH29. It involves the four members of the IVc bHLH clade, namely ILR3/bHLH105 (IAA-LEUCINE RESISTANT 3), IDT1/bHLH34 (IRON DEFICIENCY TOLERANT 1), bHLH104 and bHLH115. These four TFs play additive roles in the iron deficiency responses and their activity is thought to rely, at least in part, on their ability to form homo- or heterodimers (Li *et al.*, 2016; Zhang *et al.*, 2015; Liang *et al.*, 2017) (Figure 3). In response to iron deficiency, these TFs directly activate the expression of clade Ib bHLH and indirectly the one of *FIT/bHLH29* (Li *et al.*, 2016; Zhang *et al.*, 2015; Liang *et al.*, 2017).

Clade IVb bHLH TFs (*i.e.* PYE/bHLH47, bHLH11 and URI/bHLH121), also participate to the regulation of iron homeostasis in Arabidopsis. PYE/bHLH47 is a negative regulator (Long *et al.*, 2010), which contains an EAR motif at its C-terminal region (*i.e.* DLNxxP; Kagale and Rozwadowski, 2011) that directly represses the expression of genes participating to the maintenance of iron homeostasis. Interaction studies highlighted that PYE/bHLH47 could heterodimerize with ILR3/bHLH105 and bHLH115 (Long *et al.*, 2010; Zhang *et al.*, 2015; Tissot *et al.*, 2019). However, the role of these interactions was still a matter of debate until recently. In a recent study, ILR3/bHLH105 was found to play a central role in the regulation of iron homeostasis where it acts as both transcriptional activator and repressor of the plant responses to iron deficiency and excess, respectively (Kroh and Pilon, 2019; Tissot *et al.*, 2019). In this study, the authors showed that the repressive activity of ILR3/bHLH105 was conferred by its dimerization with PYE/bHLH47 (Figure 3). The authors also highlighted that ILR3-PYE heterodimers might repress the expression of *PYE/bHLH47* when iron availability is not limiting *via* a negative feedback regulatory loop (Figure 3). bHLH11 is another transcriptional repressor also containing an EAR motif (*i.e.* LxLxL) in its C-terminal domain (Tanabe *et al.*, 2019). Overexpression studies suggest that bHLH11 inhibits the plant tolerance to iron deficiency and the expression of *IRT1* and *FRO2*, most probably by indirectly repressing the expression of *FIT/bHLH29* (Tanabe *et al.*, 2019) (Figure 3). In contrast to PYE/bHLH47 and bHLH11, URI/ bHLH121 (UPSTREAM REGULATOR OF

IRT1) has been recently identified and characterized by three different groups as a positive regulator of the plant responses to iron deficiency (Lei *et al.*, 2020; Lockhart, 2020; Gao *et al.*, 2020a; Kim *et al.*, 2019). URI/bHLH121 can form heterodimers with clade IVc bHLH TFs (Lei *et al.*, 2020; Gao *et al.*, 2020a; Kim *et al.*, 2019) (Figure 3). These interactions participate to the relocation of URI/bHLH121 from the cytosol into the nucleus (Lei *et al.*, 2020). It also suggests that the transcriptional activation of clade IVc bHLH target genes requires, at least in part, URI/bHLH121 (Lei *et al.*, 2020; Gao *et al.*, 2020a; Kim *et al.*, 2019). In support of this assertion, it was demonstrated that several genes directly targeted by URI/bHLH121 are identical to that of clade IVc bHLH TFs (Figure 3). The expression of *FIT/bHLH29* also relies on URI/bHLH121 activity, most probably *via* an indirect mechanism (Lei *et al.*, 2020; Gao *et al.*, 2020a; Kim *et al.*, 2019). It is noteworthy that under iron deficiency condition, URI/bHLH121 accumulates in its phosphorylated form (Figure 3) that increases its binding capacity to the promoter region of its target genes such as the clade Ib bHLH TFs (Kim *et al.*, 2019). Interestingly, both URI/bHLH121 transcript and protein accumulate constitutively regardless of iron status (Gao *et al.*, 2020a; Kim *et al.*, 2019). In contrast, URI/bHLH121 cellular localization in roots differs depending on iron availability (Gao *et al.*, 2020a). When iron is not limiting, URI/bHLH121 mainly localizes in the stele and the endodermis, whereas under iron deficiency condition, URI/bHLH121 is primarily observed in the cortex and the epidermis cells, where it promotes iron uptake (Gao *et al.*, 2020a). The thorough characterization of URI/bHLH121 indicates that it plays a key role in the control of plant iron homeostasis mainly because it directly or indirectly regulates the expression of most of the known genes involved in this regulatory network (Gao *et al.*, 2020a; Kim *et al.*, 2019) (Figure 3). Interestingly, it was recently reported that URI/bHLH121 directly activates *FER1*, *FER3* and *FER4* expression when iron availability is not in excess, indicating that URI/bHLH121 positively regulates the transient storage of iron in addition to the iron deficiency response (Gao *et al.*, 2020b).

Functional homologs of most of the above-described Arabidopsis bHLH TFs have been characterized in several dicots (Figure 2, Table 1) indicating that this regulatory mechanism is most likely conserved within strategy I plants.

bHLH TFs involved in the regulation of iron homeostasis in strategy II plants

As stated above, plants have developed two different strategies to take up iron from the soil, distinguishing the non-graminaceous (Strategy I, reduction strategy) and graminaceous (Strategy II, chelation strategy) species (Figure 1). However, the in depth study of the responses to iron deficiency in rice has highlighted that the regulation of iron homeostasis in strategy II plants involves the activity of several bHLH TFs belonging to the same clades than those identified in strategy I plant species (Figures 2 and 4, Table 1; Kobayashi, 2019; Kobayashi *et al.*, 2019).

OsFIT/OsbHLH156 was recently identified as a positive regulator of the iron deficiency responses (Liang *et al.*, 2020; Wang *et al.*, 2020). Loss-of-function of *OsFIT/OsbHLH156* resulted in strong iron deficiency symptoms under upland condition, whereas no iron deficiency symptoms were observed when plants were grown in waterlogged soil (Wang *et al.*, 2020). These results imply that the strategy II iron uptake system was impaired in the *Osfit* mutant, an hypothesis that is supported by the disruption of the expression of strategy II iron uptake related genes. It includes genes encoding enzyme involved in MAs biosynthesis (*e.g.* *NAS1* and *NAS2*, *NICOTIANAMINE SYNTHASE 1* and *2*) or Fe³⁺-MA transport (*e.g.* *OsYSL15*, *YELLOW STRIPE-LIKE 15*) (Figure 1) (Liang *et al.*, 2020; Wang *et al.*, 2020). Interestingly, it was demonstrated that OsFIT/OsbHLH156 regulates the expression of *OsIRT1* and therefore participates also to the regulation of strategy I iron uptake mechanism (Liang *et al.*, 2020) (Figure 1). OsFIT/OsbHLH156 interacts with OsIRO2/OsbHLH56, a clade Ib bHLH, and promotes its nuclear accumulation (Figure 4) (Liang *et al.*, 2020; Wang *et al.*, 2020). OsIRO2/OsbHLH56 was the first characterized bHLH TF involved in the control of iron homeostasis in rice, where it acts as a positive regulator of the iron deficiency response (Ogo *et al.*, 2006; Ogo *et al.*, 2007). Expression analysis indicated that the regulation of *OsIRT1* expression by OsFIT/OsbHLH156 might be different to that of *Arabidopsis* or that it might involve other Ib bHLH TFs than OsIRO2/OsbHLH56.

Three out of the four clade IVc bHLH TFs (*i.e.* OsPRI1/OsbHLH060, OsPRI2/bHLH058, OsPRI3/OsbHLH059; POSITIVE REGULATOR OF IRON HOMEOSTASIS 1, 2 and 3) have been identified in rice as playing a positive role in the iron deficiency responses (Zhang *et al.*, 2017; Kobayashi *et al.*, 2019; Zhang *et al.*, 2020b). The characterization of loss-of-function mutants suggested that these three TFs directly regulate the expression of

OsIRO2/OsbHLH56, and indirectly the expression of *OsNAS1*, *OsNAS2*, and *OsYSL15* via *OsIRO2/OsbHLH56* (Figure 4) (Zhang *et al.*, 2020b; Zhang *et al.*, 2017). Among the potential direct targets of *OsPRI1/OsbHLH060*, *OsPRI2/bHLH058* and *OsPRI3/OsbHLH059*, there is also *OsIRO3/OsbHLH63* (Figure 4) (Zhang *et al.*, 2020b; Zhang *et al.*, 2017). *OsIRO3/OsbHLH63* is a member of the IVb clade and the functional homolog of *PYE/bHLH47* (Zheng *et al.*, 2010). Like *PYE/bHLH47*, *OsIRO3/OsbHLH63* functions as a negative regulator of the iron deficiency responses (Zheng *et al.*, 2010). It is likely that *OsIRO3/OsbHLH63* activity might antagonize *OsIRO2/OsbHLH56* to tightly regulate iron uptake and avoid iron overload (Zhang *et al.*, 2012; Zhang *et al.*, 2020b; Zhang *et al.*, 2017; Zheng *et al.*, 2010). To date, there is no information on the role of *OsbHLH057*, the fourth member of the rice clade IVc bHLH. Indeed, based on the information gathered in Arabidopsis, it is likely that *OsbHLH057* participates to the control of iron homeostasis in rice. Whether *OsbHLH057* exerts a minor role in specific cell types or in specific environmental conditions remains to be determined.

OsbHLH133 functions as a negative regulator of iron translocation from roots to shoots (Wang *et al.*, 2013a). It should be noted that *OsbHLH133* is the only one member of the clade VIIIc reported to date as involved in the regulation of iron homeostasis in plants. The other members of this clade in Arabidopsis play a positive role in root hair development but have never been associated with the maintenance of iron homeostasis (Bruex *et al.*, 2012). Whether these bHLHs have a similar and conserved role, as that of *OsbHLH133* in the regulation of iron homeostasis in strategy II plants, is to be investigated.

The transcriptional regulation of iron homeostasis is not restricted to the activity of bHLH TFs

TFs from other families are involved in regulatory networks acting for iron homeostasis in both strategy I and strategy II plants (Figures 3 and 4, Table 2).

Several R2R3-MYB TFs from different plant species have been shown to be involved in the regulation of iron deficiency responses. In Arabidopsis, MYB10 and MYB72 are two iron deficiency inducible TFs required for proper iron uptake and whose expression is partially dependent on *FIT/bHLH29* and *URI/bHLH121* activities (Figure 3) (Gao *et al.*, 2020a; Palmer *et al.*, 2013; Zamioudis *et al.*, 2014). Furthermore, MYB72 has been identified as a

transcriptional activator of genes associated with the biosynthesis and secretion of iron-mobilizing coumarins, highlighting its role in strategy I iron uptake system (Stringlis *et al.*, 2018; Zamioudis *et al.*, 2014) (Figure 1). Recently, it was shown that the Arabidopsis MYB28 and MYB29 are at the interface of the plant sensitivity to ammonium stress and the modulation of iron homeostasis (Coletto *et al.*, 2020). Notably, the ammonium-dependent decrease of MYB28 and MYB29 expression (or the loss-of-function of both genes) leads to defects in iron translocation from roots to shoots and to the induction of the expression of *FIT/bHLH29*, clade Ib bHLHs, MYB72 as well as *IMA1/FEP3* and *IMA3/FEP1* in roots. MdMYB58, a close homolog of MYB72, was recently characterized as a positive regulator of iron uptake and translocation in apple (Wang *et al.*, 2018). Further investigation showed that MdMYB58 transcriptional activity is inhibited by its heterodimerization with MdSAT1/MdbHLH18, a IVa clade bHLH TF (Wang *et al.*, 2018). In contrast to the above-mentioned MYB TFs, MxMYB1 may function as a negative regulator of iron uptake and storage (Shen *et al.*, 2008).

In Arabidopsis, WRKY46 plays a role in iron translocation between root and shoot by directly regulating the expression of *VITL1* (*VACUOLAR IRON TRANSPORTER-LIKE 1*), a potential iron transporter involved in iron sequestration into the vacuoles (Yan *et al.*, 2016; Gollhofer *et al.*, 2014). HAP5A/NF-YC1 is also involved in iron translocation between root and shoot by regulating the expression of *NAS1* (Zhu *et al.*, 2020). ERF4 and ERF72 are two Arabidopsis TFs belonging to AP2/ERF family that have been reported as potential negative regulators of iron deficiency responses by repressing the expression of genes involved in iron uptake such as *IRT1* (Liu *et al.*, 2017a; Liu *et al.*, 2017b). Similarly, MbERF4 and MbERF72 from *Malus baccata* as well as MxERF4 from *Malus xiaojinensis* act as negative regulators of the iron deficiency responses in these two apple species (Liu *et al.*, 2018; Zhang *et al.*, 2020a). In contrast, ERF95 was recently proposed to promote iron storage in Arabidopsis seeds (Sun *et al.*, 2020). However, how iron distribution is controlled in seeds is not clearly established even if it was recently proposed that B3 TFs, which are involved in the regulation of embryo development and seed maturation, might be good candidates (Roschztardt *et al.*, 2020).

IDEF1 and IDEF2 (IRON DEFICIENCY-RESPONSIVE ELEMENT FACTOR 1 and 2) are two rice TFs belonging to two different families, the ABI3/VP1 and NAC, respectively (Figure 4) (Kobayashi *et al.*, 2007; Ogo *et al.*, 2008). These two TFs participate to the regulation of iron homeostasis in rice. IDEF1 is required for the coordinated activation of

genes related to iron uptake and translocation, including *OsIRT1*, *OsNAS1*, *OsNAS2* and *OsYSL15* (Kobayashi *et al.*, 2009; Kobayashi *et al.*, 2007). In addition, IDEF1 positively regulates the expression of *OsIRO2*, indicating that IDEF1 functions upstream of *OsIRO2* in the iron deficiency regulatory network (Figure 4) (Kobayashi *et al.*, 2009; Kobayashi *et al.*, 2007). Interestingly, it was shown that IDEF1 could bind to iron and zinc atoms and that this capacity was necessary for its activity (Kobayashi *et al.*, 2012). It was therefore proposed that IDEF1 could sense the cellular metal ion balance caused by changes in iron availability, suggesting that IDEF1 could be a cellular iron sensor allowing the tight regulation of the iron deficiency responses (Kobayashi *et al.*, 2012). Similarly to IDEF1, IDEF2 plays a positive role in the plant response to iron deficiency (Ogo *et al.*, 2008).

A family of peptides named IMA/FEP (IRONMAN/FE-UPTAKE-INDUCING PEPTIDE) has been reported to play a positive role in iron deficiency responses in Arabidopsis, by regulating a set of deficiency-inducible genes including Ib bHLH TFs, a function that seems to be conserved across plant species (Grillet *et al.*, 2018; Hirayama *et al.*, 2018). Two recent studies showed that URI/bHLH121 is a direct positive regulator of *IMA1/FEP3* and *IMA2/FEP2* expression (Gao *et al.*, 2020a; Kim *et al.*, 2019). These results indicate that IMAs/FEPs are implicated in the bHLH-dependent regulatory network regulating iron homeostasis. However, the precise regulatory mechanisms by which IMAs/FEPs act are still unknown.

Post-translational regulation of the iron homeostasis regulatory networks

Protein-protein interactions as well as post-translational modifications (*i.e.* ubiquitination, sumoylation, phosphorylation) can significantly affect the regulatory activities of TFs. Such mechanisms were shown to play an important role in the maintenance of iron homeostasis in plants. This topic has also been covered in recent reviews by Kobayashi, 2019; Rodríguez-Celma *et al.*, 2019a; Schwarz and Bauer, 2020; Spielmann and Vert, 2020; Wu and Ling, 2019.

As described earlier, the transcriptional activity of bHLH TFs involved in the control of iron homeostasis is extensively dependent on *in vivo* protein-protein interactions, in the form of homo- or heterodimers (Figures 3 and 4, Tables 1 and 2). For instance, FIT/bHLH29 heterodimerization with clade Ib bHLH TFs is required for its transcriptional activity and

stability, whereas its interaction with clade IVa members promotes its degradation *via* the 26S proteasome pathway (Cui *et al.*, 2018). bHLH39 nuclear localization also depends on its interaction with FIT/bHLH29 since in the cells lacking FIT/bHLH29, bHLH39 localizes predominantly in the cytoplasm (Trofimov *et al.*, 2019). Similarly, OsFIT/OsbHLH156 facilitates the nuclear localization of OsIRO2/OsbHLH156, the functional homolog of bHLH39 in rice, suggesting that this post-translational regulatory mechanism is conserved within the plant kingdom (Liang *et al.*, 2020; Wang *et al.*, 2020). FIT/bHLH29 activity and/or stability are also modulated by its interaction with several additional protein partners that do not belong to the bHLH family of TFs (Figures 3, Tables 1) (reviewed in Kobayashi, 2019; Schwarz and Bauer, 2020; Spielmann and Vert, 2020; Wu and Ling, 2019).

Several ubiquitin E3 ligases have been identified and characterized as negative regulators of iron uptake to avoid potential iron overload by targeting bHLH TFs for their degradation (for details, see Spielmann and Vert, 2020; Rodríguez-Celma *et al.*, 2019a) (Figures 3 and 4). BTS (BRUTUS), whose expression is induced by iron deficiency in roots, is proposed to be a critical iron sensing E3 ubiquitin ligase in Arabidopsis (Long *et al.*, 2010). BTS interacts with ILR3/bHLH105 and bHLH115 to facilitate their degradation via the 26S proteasome pathway, allowing the fine tuning of the expression of downstream iron deficiency response genes (Figure 3) (Selote *et al.*, 2015; Long *et al.*, 2010). Similarly, OsHRZ1 and OsHRZ2 (HAEMERYTHRIN MOTIF-CONTAINING REALLY INTERESTING NEW GENE (RING) AND ZINC-FINGER PROTEIN 1 and 2), two rice ubiquitin E3 ligases displaying strong sequence similarities with BTS, have been reported as potential iron sensors playing a negative role in iron acquisition under iron sufficient conditions (Kobayashi *et al.*, 2013). It was shown that OsHRZ1 could interact with OsPRI1/OsbHLH60, OsPRI2/OsbHLH58 and OsPRI3/OsbHLH59 and mediate their degradation via the 26S proteasome (Figure 4) (Zhang *et al.*, 2020b; Zhang *et al.*, 2017). However, different results were reported in another study only validating the sole interactions between OsHRZ1 and OsHRZ2 with OsPRI1/OsbHLH60 and OsPRI2/OsbHLH58 (Kobayashi *et al.*, 2019). Another *in vitro* ubiquitination study showed that neither OsPRI1/OsbHLH60, nor OsPRI2/OsbHLH58 or OsPRI3/OsbHLH59 were ubiquitinated by OsHRZ1 or by OsHRZ2 (Kobayashi *et al.*, 2019). These discrepancies may be due to the different methodologies and materials used in these studies. Whether these interactions participate to the ubiquitination and degradation of OsPRI1/OsbHLH60, OsPRI2/OsbHLH58 and OsPRI3/OsbHLH59 remains to be further demonstrated and confirmed. IDEF1, whose activity is necessary for *HRZs* expression, is also

degraded via the 26S proteasome pathway through a yet uncharacterized mechanism (Zhang *et al.*, 2014). However, IDEF1 degradation is inhibited by its interaction with the IBP1.1 (IDEF1-BINDING PROTEINS 1.1) Bowman–Birk trypsin inhibitor protein (Figure 4) (Zhang *et al.*, 2014). BTSL1 and BTSL2, two Arabidopsis homologs of BTS, function redundantly as negative regulators of the iron deficiency response (Hindt *et al.*, 2017). In addition, BTSL1 and BTSL2 could directly target FIT/bHLH29 and promote its ubiquitination and subsequent degradation via the 26S proteasome pathway, thus negatively regulating the expression of iron uptake related genes (Figure 3) (Rodríguez-Celma *et al.*, 2019b; Rodríguez-Celma *et al.*, 2019a). In apples, MdbT1 and MdbT2, two BTB-TAZ proteins, interact with MdbHLH104. MdbT proteins interact as well with MdCUL3 (CULLIN-RING UBIQUITIN LIGASE 3) to form MdbT^{MdCUL3} complexes required for MdbHLH104 ubiquitination and degradation *via* the 26S proteasome pathway (Zhao *et al.*, 2016a). In contrast, MdSIZ1, a SIZ/PIAS-type SUMO E3 ligase, directly sumoylates MdbHLH104, especially under iron deficiency conditions, to enhance its stability (Zhou *et al.*, 2019). Recently, it was shown that the mutation of an alanine into valine within IDT1/bHLH34 enhances both its stability and nuclear localization (Sharma and Yeh, 2020). Since this amino acid is conserved among clade IVc bHLH TFs in both monocots and dicots (Sharma and Yeh, 2020), one might speculate that this amino acid plays an important role in regulating the stability of clade IVc bHLH and thus their degradation via the 26S proteasome.

Phosphorylation plays also an important role in determining TF activity in the iron homeostasis networks (Figure 3). In Arabidopsis, FIT/bHLH29 phosphorylation by the calcium-dependent protein kinase CIPK11 (CBL-INTERACTING PROTEIN KINASE 11) positively regulates its activity by favoring its nuclear accumulation and dimerization with bHLH39 (Gratz *et al.*, 2019). *CIPK11* is induced and activated *via* a CBL1/9-mediated Ca²⁺ sensing pathway under iron deficiency conditions. CIPK21 could also interact with FIT/bHLH29, indicating that this protein kinase might have a potential role in the regulation of the FIT/bHLH29-dependent iron deficiency responses (Gratz *et al.*, 2019). Recently, the phosphorylation of URI/bHLH121 was also reported as a key mechanism that regulates its activity (Kim *et al.*, 2019). It was shown that the phosphorylated form of URI/bHLH121 only accumulates in response to iron deficiency, whereas the turnover of phosphorylated URI/bHLH121 is dependent on BTS activity (Kim *et al.*, 2019). Under iron deficiency, URI/bHLH121 showed enhanced heterodimerization capability with IVc bHLH TFs and increased binding ability to the promoter of its target genes, indicating that the

phosphorylation of URI/bHLH121 plays a positive role in the iron deficiency responses (Kim *et al.*, 2019). Nevertheless, since URI/bHLH121 can activate the expression of *FER1*, *FER3* and *FER4* in the stele when iron availability is not limiting (Gao *et al.*, 2020b), a growth condition where the phosphorylated form of bHLH121 is degraded (Kim *et al.*, 2019), it is likely that URI/bHLH121 is transcriptionally active independently of its phosphorylation state. Interestingly, as stated earlier, the pattern of accumulation of URI/bHLH121 within the root cells is controlled by an iron-dependent mechanism. URI/bHLH121 is preferentially accumulated in the epidermis and the cortex when plants are grown under iron deficiency and in the stele when grown under iron sufficiency. From these observations, one might hypothesize that URI/bHLH121 cellular distribution, rather than its transcriptional activity, is regulated by its phosphorylation state. However, the precise regulatory mechanism leading to the phosphorylation of URI/bHLH121 remains to be characterized.

Conclusions and future prospects

In the past two decades, remarkable progresses have been made in decrypting the molecular mechanisms that regulate iron homeostasis in both strategy I and strategy II plants (Figure 1). They highlighted that iron homeostasis in plants is regulated at the transcriptional level and involves several bHLH TFs that function in intricate regulatory networks (Figures 2 to 4).

The function of most of these bHLH TFs (*i.e.* clades Ib, IIIa, IVb and IVc) is conserved among grass and non-grass species (Figure 2), in contrast to the downstream target genes of iron acquisition machinery, which are distinctive in strategy I and strategy II plants (Figure 1). How these bHLH functional homologs have evolved to target different genes is an intriguing question. bHLH from clade IVa are involved in the regulation of iron homeostasis in *Arabidopsis* but whether or not this clade of bHLH TFs has a similar role in other plant species needs to be investigated. Additional investigations will also be necessary to determine if the non-grass homologs of OsbHLH133 (clade VIIIc) play a role in the control of iron homeostasis. Other regulatory proteins are also conserved between plants from both strategies. It is for instance the case for MYB and ERF TFs (Table 2) or for haemerythrin E3-ubiquitin ligases. Interestingly, BTB-TAZ-CUL3 ubiquitin ligase complexes participate to the degradation of clade IVc bHLH in apples. To date, such mechanism has only been observed in apple, raising the question of the conservation of this regulatory mechanism in other plants species, from non-grass to grass and from perennials to annuals. Ubiquitination,

as well as phosphorylation and sumoylation, are crucial posttranslational modifications for the regulation of key bHLH TFs activities involved in the iron homeostasis networks, notably clade IVc bHLH FIT/bHLH29 and URI/bHLH121. Whether such modifications participate to the regulation of other TF activities within this network is still to be assessed. Determining the degree of conservation of this regulatory network between plants from strategy I and strategy II, and between annual and perennial species, is still on its own a question that deserves to be addressed.

How plants sense iron status and switch on or repress the downstream regulatory network to regulate iron uptake, translocation, storage or assimilation has remained elusive. To date, haemerythrin E3-ubiquitin ligases are the main candidates (Rodríguez-Celma *et al.*, 2019a). These E3-ubiquitin ligases are induced in response to iron deficiency, participate to the degradation of key bHLH TFs (*i.e.* clade IVc and FIT/bHLH29) and are destabilized upon iron binding at their haemerythrin motifs. IDEF1, the rice iron binding TF mentioned earlier, was also proposed as a potential iron sensor (Kobayashi *et al.*, 2012; Kobayashi *et al.*, 2009; Kobayashi *et al.*, 2007). The characterization of IDEF1 functional homologs in non-grass species would be an additional element in support of this later hypothesis. Another important question concerns the regulation of the most upstream bHLH TFs. For instance, the main challenges would be to determine how the expression of clade IVc bHLH is controlled and to identify which protein kinase modulates URI/bHLH121 activity.

Recently, epigenetic regulation has emerged as playing important role in the control of iron homeostasis, by regulating DNA accessibility to promoters, and thus the expression of both TFs and iron uptake genes. This is for instance the case in *Arabidopsis* with the iron-dependent deposition of repressive marks on H3 histone (*i.e.* H3K27m3) at the promoter *loci* of *FIT/bHLH29*, *IRT1* and *FRO2* (Park *et al.*, 2019). This additional layer of regulation in the transcriptional networks controlling iron homeostasis raises questions on the importance and the significance of such regulatory mechanism in this process.

Within this review, it is possible to grasp the extent of the complexity of the regulatory networks that regulate iron homeostasis in plants. Nevertheless, another level of complexity still to be investigated concerns the apparent redundancies existing between several TFs, the localization of their target genes, and thus the physiological functions that are controlled. This will be achieved by the in depth study of the tissue and cellular localization of the

different TFs as well as the proteins involved in their regulation. Such knowledge is necessary to fully decrypt and understand the dynamics of this regulatory process. Unfortunately, such data are only available for a few of these proteins (Long *et al.*, 2010; Gao *et al.*, 2020a; Samira *et al.*, 2018).

In conclusion, much work still lies ahead to fully comprehend the transcriptional regulatory network that regulate iron homeostasis in plants, which might offer novel opportunities for improving plant growth and health and for generating iron-fortified crops.

Accepted Manuscript

Acknowledgements

This work was supported by grants from the “Agence Nationale de la Recherche (ANR)” to C.D. Support was provided by the China Scholarship Council to F.G. We thank Florence Vignols for help in preparing this article.

Author Contribution

FG and CD wrote the manuscript

Accepted Manuscript

References

- Bashir K, Ishimaru Y, Shimo H, Kakei Y, Senoura T, Takahashi R, Sato Y, Sato Y, Uozumi N, Nakanishi H, Nishizawa NK.** 2011. Rice phenolics efflux transporter 2 (PEZ2) plays an important role in solubilizing apoplasmic iron. *Soil Science and Plant Nutrition* **57**, 803-812.
- Briat J-F, Dubos C, Gaymard F.** 2015. Iron nutrition, biomass production, and plant product quality. *Trends in Plant Science* **20**, 33-40.
- Bruex A, Kainkaryam RM, Wieckowski Y, Kang YH, Bernhardt C, Xia Y, Zheng X, Wang JY, Lee MM, Benfey P.** 2012. A gene regulatory network for root epidermis cell differentiation in Arabidopsis. *PLoS genetics* **8**, e1002446.
- Brumbarova T, Bauer P, Ivanov R.** 2015. Molecular mechanisms governing Arabidopsis iron uptake. *Trends in Plant Science* **20**, 124-133.
- Camaschella C.** 2015. Iron-deficiency anemia. *New England journal of medicine* **372**, 1832-1843.
- Carretero-Paulet L, Galstyan A, Roig-Villanova I, Martínez-García JF, Bilbao-Castro JR, Robertson DL.** 2010. Genome-wide classification and evolutionary analysis of the bHLH family of transcription factors in Arabidopsis, poplar, rice, moss, and algae. *Plant Physiology* **153**, 1398-1412.
- Colangelo EP, Guerinot ML.** 2004. The essential basic helix-loop-helix protein FIT1 is required for the iron deficiency response. *The Plant Cell* **16**, 3400-3412.
- Coletto I, Bejarano I, Marín-Peña AJ, Medina J, Rioja C, Burow M, Marino D.** Arabidopsis thaliana transcription factors MYB28 AND MYB29 shape ammonium stress responses by regulating fe homeostasis. *New Phytol* doi: 10.1111/nph.16918.

- Colombo C, Iorio ED, Liu Q, Jiang Z, Barrón V.** 2018. Iron Oxide Nanoparticles in Soils: Environmental and Agronomic Importance. *Journal of Nanoscience and Nanotechnology*, **18**:761.
- Connorton JM, Balk J, Rodríguez-Celma J.** 2017. Iron homeostasis in plants—a brief overview. *Metallomics* **9**, 813-823.
- Cui Y, Chen C-L, Cui M, Zhou W-J, Wu H-L, Ling H-Q.** 2018. Four IVa bHLH transcription factors are novel interactors of FIT and mediate JA inhibition of iron uptake in Arabidopsis. *Molecular plant* **11**, 1166-1183.
- Du J, Huang Z, Wang B, Sun H, Chen C, Ling H-Q, Wu H.** 2015. SlbHLH068 interacts with FER to regulate the iron-deficiency response in tomato. *Annals of botany* **116**, 23-34.
- Fourcroy P, Sisó-Terraza P, Sudre D, Savirón M, Reyt G, Gaymard F, Abadía A, Abadía J, Álvarez-Fernández A, Briat JF.** 2014. Involvement of the ABCG 37 transporter in secretion of scopoletin and derivatives by Arabidopsis roots in response to iron deficiency. *New Phytologist* **201**, 155-167.
- Fourcroy P, Tissot N, Gaymard F, Briat J-F, Dubos C.** 2016. Facilitated Fe nutrition by phenolic compounds excreted by the Arabidopsis ABCG37/PDR9 transporter requires the IRT1/FRO2 high-affinity root Fe²⁺ transport system. *Molecular plant* **9**, 485-488.
- Gao F, Robe K, Bettembourg M, Navarro N, Rofidal V, Santoni V, Gaymard F, Vignols F, Roschztardt H, Izquierdo E.** 2020a. The transcription factor bHLH121 interacts with bHLH105 (ILR3) and its closest homologs to regulate iron homeostasis in Arabidopsis. *The Plant Cell* **32**, 508-524.
- Gao F, Robe K, Dubos C.** 2020b. Further insights into the role of bHLH121 in the regulation of iron homeostasis in Arabidopsis thaliana. *Plant Signaling and Behavior*, 1795582.
- Gao F, Robe K, Gaymard F, Izquierdo E, Dubos C.** 2019. The transcriptional control of iron homeostasis in plants: a tale of bHLH transcription factors? *Frontiers in plant science* **10**, 6.

- Gollhofer J, Timofeev R, Lan P, Schmidt W, Buckhout TJ.** 2014. Vacuolar-iron-transporter1-like proteins mediate iron homeostasis in Arabidopsis. *PLoS One* **9**, e110468.
- Gratz R, Manishankar P, Ivanov R, Köster P, Mohr I, Trofimov K, Steinhorst L, Meiser J, Mai H-J, Drerup M.** 2019. CIPK11-dependent phosphorylation modulates FIT activity to promote Arabidopsis iron acquisition in response to calcium signaling. *Developmental cell* **48**, 726-740. e710.
- Grillet L, Lan P, Li W, Mokkaapati G, Schmidt W.** 2018. IRON MAN is a ubiquitous family of peptides that control iron transport in plants. *Nature plants* **4**, 953-963.
- Guerinot ML, Yi Y.** 1994. Iron: nutritious, noxious, and not readily available. *Plant Physiology* **104**, 815-820.
- Hänsch R, Mendel RR.** 2009. Physiological functions of mineral micronutrients (Cu, Zn, Mn, Fe, Ni, Mo, B, Cl). *Current Opinion in Plant Biology* **12**, 259-266.
- Heim MA, Jakoby M, Werber M, Martin C, Weisshaar B, Bailey PC.** 2003. The basic helix-loop-helix transcription factor family in plants: a genome-wide study of protein structure and functional diversity. *Molecular biology and evolution* **20**, 735-747.
- Hindt MN, Akmakjian GZ, Pivarski KL, Punshon T, Baxter I, Salt DE, Guerinot ML.** 2017. BRUTUS and its paralogs, BTS LIKE1 and BTS LIKE2, encode important negative regulators of the iron deficiency response in Arabidopsis thaliana. *Metallomics* **9**, 876-890.
- Hirayama T, Lei GJ, Yamaji N, Nakagawa N, Ma JF.** 2018. The putative peptide gene FEP1 regulates iron deficiency response in Arabidopsis. *Plant and Cell Physiology* **59**, 1739-1752.
- Huang D, Dai W.** 2015. Molecular characterization of the basic helix-loop-helix (bHLH) genes that are differentially expressed and induced by iron deficiency in Populus. *Plant cell reports* **34**, 1211-1224.
- Inoue H, Kobayashi T, Nozoye T, Takahashi M, Kakei Y, Suzuki K, Nakazono M, Nakanishi H, Mori S, Nishizawa NK.** 2009. Rice OsYSL15 is an iron-regulated iron (III)-deoxymugineic acid transporter expressed in the roots and is essential for iron

uptake in early growth of the seedlings. *Journal of Biological Chemistry* **284**, 3470-3479.

Ishimaru Y, Suzuki M, Tsukamoto T, Suzuki K, Nakazono M, Kobayashi T, Wada Y, Watanabe S, Matsushashi S, Takahashi M. 2006. Rice plants take up iron as an Fe³⁺-phytosiderophore and as Fe²⁺. *The Plant Journal* **45**, 335-346.

Jakoby M, Wang H-Y, Reidt W, Weisshaar B, Bauer P. 2004. FRU (BHLH029) is required for induction of iron mobilization genes in *Arabidopsis thaliana*. *FEBS letters* **577**, 528-534.

Kagale S, Rozwadowski K. 2011. EAR motif-mediated transcriptional repression in plants: an underlying mechanism for epigenetic regulation of gene expression. *Epigenetics* **6**, 141-146.

Kim SA, LaCroix IS, Gerber SA, Guerinot ML. 2019. The iron deficiency response in *Arabidopsis thaliana* requires the phosphorylated transcription factor URI. *Proceedings of the National Academy of Sciences* **116**, 24933-24942.

Kobayashi T. 2019. Understanding the Complexity of Iron Sensing and Signaling Cascades in Plants. *Plant and Cell Physiology* **60**, 1440-1446.

Kobayashi T, Itai RN, Aung MS, Senoura T, Nakanishi H, Nishizawa NK. 2012. The rice transcription factor IDEF1 directly binds to iron and other divalent metals for sensing cellular iron status. *The Plant Journal* **69**, 81-91.

Kobayashi T, Itai RN, Ogo Y, Kakei Y, Nakanishi H, Takahashi M, Nishizawa NK. 2009. The rice transcription factor IDEF1 is essential for the early response to iron deficiency, and induces vegetative expression of late embryogenesis abundant genes. *The Plant Journal* **60**, 948-961.

Kobayashi T, Nagasaka S, Senoura T, Itai RN, Nakanishi H, Nishizawa NK. 2013. Iron-binding haemerythrin RING ubiquitin ligases regulate plant iron responses and accumulation. *Nature communications* **4**, 1-12.

Kobayashi T, Nakanishi Itai R, Nishizawa NK. 2014. Iron deficiency responses in rice roots. *Rice (N Y)* **7**, 27.

Kobayashi T, Nishizawa NK. 2012. Iron uptake, translocation, and regulation in higher plants. *Annual Review of Plant Biology* **63**, 131-152.

- Kobayashi T, Nozoye T, Nishizawa NK.** 2019. Iron transport and its regulation in plants. *Free Radical Biology and Medicine* **133**, 11-20.
- Kobayashi T, Ogo Y, Itai RN, Nakanishi H, Takahashi M, Mori S, Nishizawa NK.** 2007. The transcription factor IDEF1 regulates the response to and tolerance of iron deficiency in plants. *Proceedings of the National Academy of Sciences* **104**, 19150-19155.
- Kobayashi T, Ozu A, Kobayashi S, An G, Jeon J-S, Nishizawa NK.** 2019. OsbHLH058 and OsbHLH059 transcription factors positively regulate iron deficiency responses in rice. *Plant Molecular Biology* **101**, 471-486.
- Kroh GE, Pilon M.** 2019. Connecting the negatives and positives of plant iron homeostasis. *New Phytologist* **223**, 1052-1055.
- Le CTT, Brumbarova T, Ivanov R, Stoof C, Weber E, Mohrbacher J, Fink-Straube C, Bauer P.** 2016. Zinc finger of Arabidopsis thaliana12 (ZAT12) interacts with FER-like iron deficiency-induced transcription factor (FIT) linking iron deficiency and oxidative stress responses. *Plant Physiology* **170**, 540-557.
- Lei R, Li Y, Cai Y, Li C, Pu M, Lu C, Yang Y, Liang G.** 2020. bHLH121 Functions as a Direct Link that Facilitates the Activation of FIT by bHLH IVc Transcription Factors for Maintaining Fe Homeostasis in Arabidopsis. *Molecular plant* **13**, 634-649.
- Li L, Gao W, Peng Q, Zhou B, Kong Q, Ying Y, Shou H.** 2018. Two soybean bHLH factors regulate response to iron deficiency. *Journal of integrative plant biology* **60**, 608-622.
- Li Q, Chen L, Yang A.** 2019. The Molecular Mechanisms Underlying Iron Deficiency Responses in Rice. *International Journal of Molecular Sciences* **21**, 43.
- Li X, Zhang H, Ai Q, Liang G, Yu D.** 2016. Two bHLH transcription factors, bHLH34 and bHLH104, regulate iron homeostasis in Arabidopsis thaliana. *Plant Physiology* **170**, 2478-2493.
- Li Y-Y, Sui X-Y, Yang J-S, Xiang X-H, Li Z-Q, Wang Y-Y, Zhou Z-C, Hu R-S, Liu D.** 2020. A novel bHLH transcription factor, NtbHLH1, modulates iron homeostasis in tobacco (*Nicotiana tabacum* L.). *Biochemical and Biophysical Research Communications* **522**, 233-239.

- Liang G, Zhang H, Li X, Ai Q, Yu D.** 2017. bHLH transcription factor bHLH115 regulates iron homeostasis in *Arabidopsis thaliana*. *Journal of experimental botany* **68**, 1743-1755.
- Liang G, Zhang H, Li Y, Pu M, Yang Y, Li C, Lu C, Xu P, Yu D.** 2020. *Oryza sativa* Fer-like Fe deficiency-induced transcription factor (OsFIT/OsbHLH156) interacts with OsIRO2 to regulate iron homeostasis. *Journal of integrative plant biology* **62**, 668-689.
- Ling H-Q, Bauer P, Berczky Z, Keller B, Ganai M.** 2002. The tomato fer gene encoding a bHLH protein controls iron-uptake responses in roots. *Proceedings of the National Academy of Sciences* **99**, 13938-13943.
- Lingam S, Mohrbacher J, Brumbarova T, Potuschak T, Fink-Straube C, Blondet E, Genschik P, Bauer P.** 2011. Interaction between the bHLH transcription factor FIT and ETHYLENE INSENSITIVE3/ETHYLENE INSENSITIVE3-LIKE1 reveals molecular linkage between the regulation of iron acquisition and ethylene signaling in *Arabidopsis*. *The Plant Cell* **23**, 1815-1829.
- Liu W, Karemera NU, Wu T, Yang Y, Zhang X, Xu X, Wang Y, Han Z.** 2017a. The ethylene response factor AtERF4 negatively regulates the iron deficiency response in *Arabidopsis thaliana*. *PLoS One* **12**, e0186580.
- Liu W, Li Q, Wang Y, Wu T, Yang Y, Zhang X, Han Z, Xu X.** 2017b. Ethylene response factor AtERF72 negatively regulates *Arabidopsis thaliana* response to iron deficiency. *Biochemical and Biophysical Research Communications* **491**, 862-868.
- Liu W, Wu T, Li Q, Zhang X, Xu X, Li T, Han Z, Wang Y.** 2018. An ethylene response factor (MxERF4) functions as a repressor of Fe acquisition in *Malus xiaojinensis*. *Scientific reports* **8**, 1-13.
- Lockhart J.** 2020. Personal Trainer: bHLH121 Functions Upstream of a Transcriptional Network of Heavy Lifters Involved in Balancing Iron Levels. *The Plant Cell* **32**, 293-294.
- Long TA, Tsukagoshi H, Busch W, Lahner B, Salt DE, Benfey PN.** 2010. The bHLH transcription factor POPEYE regulates response to iron deficiency in *Arabidopsis* roots. *The Plant Cell* **22**, 2219-2236.

- Marschner H, Römheld V.** 1994. Strategies of plants for acquisition of iron. *Plant and soil* **165**, 261-274.
- Martín Barranco A, Spielmann J, Dubeaux G, Vert G, Zelazny E.** 2020. Dynamic control of the high-affinity iron uptake complex in root epidermal cells. *Plant Physiology*, doi:10.1104/pp.20.00234
- Maurer F, Arcos MAN, Bauer P.** 2014. Responses of a triple mutant defective in three iron deficiency-induced basic helix-loop-helix genes of the subgroup Ib to iron deficiency and salicylic acid. *PLoS One* **9**, e99234.
- Murata Y, Ma JF, Yamaji N, Ueno D, Nomoto K, Iwashita T.** 2006. A specific transporter for iron (III)-phytosiderophore in barley roots. *The Plant Journal* **46**, 563-572.
- Ogo Y, Itai RN, Nakanishi H, Inoue H, Kobayashi T, Suzuki M, Takahashi M, Mori S, Nishizawa NK.** 2006. Isolation and characterization of IRO2, a novel iron-regulated bHLH transcription factor in graminaceous plants. *Journal of experimental botany* **57**, 2867-2878.
- Ogo Y, Kobayashi T, Itai RN, Nakanishi H, Kakei Y, Takahashi M, Toki S, Mori S, Nishizawa NK.** 2008. A novel NAC transcription factor, IDEF2, that recognizes the iron deficiency-responsive element 2 regulates the genes involved in iron homeostasis in plants. *Journal of Biological Chemistry* **283**, 13407-13417.
- Ogo Y, Nakanishi Itai R, Nakanishi H, Kobayashi T, Takahashi M, Mori S, Nishizawa NK.** 2007. The rice bHLH protein OsIRO2 is an essential regulator of the genes involved in Fe uptake under Fe-deficient conditions. *The Plant Journal* **51**, 366-377.
- Palmer CM, Hindt MN, Schmidt H, Clemens S, Guerinot ML.** 2013. MYB10 and MYB72 are required for growth under iron-limiting conditions. *PLoS genetics* **9**, e1003953.
- Park EY, Tsuyuki KM, Hu F, Lee J, Jeong J.** 2019. PRC2-mediated H3K27me3 contributes to transcriptional regulation of FIT-dependent iron deficiency response. *Frontiers in plant science* **10**, 627.

- Qi Y, Wang S, Shen C, Zhang S, Chen Y, Xu Y, Liu Y, Wu Y, Jiang D.** 2012. OsARF12, a transcription activator on auxin response gene, regulates root elongation and affects iron accumulation in rice (*Oryza sativa*). *New Phytologist* **193**, 109-20.
- Ramamurthy RK, Waters BM.** 2017. Mapping and characterization of the fefe gene that controls iron uptake in melon (*Cucumis melo* L.). *Frontiers in plant science* **8**, 1003.
- Robe K, Conejero G, Gao F, Fourcroy P, Lefebvre-Legendre L, Sylvestre-Gonon E, Rofidal V, Hem S, Rouhier N, Barberon M, Hecker A, Gaymard F, Izquierdo E, Dubos C.** 2020. Coumarin accumulation and trafficking in *Arabidopsis thaliana*: a complex and dynamic process. *New Phytologist* In press.
- Rodríguez-Celma J, Chou H, Kobayashi T, Long TA, Balk J.** 2019a. Hemerythrin E3 ubiquitin ligases as negative regulators of iron homeostasis in plants. *Frontiers in plant science* **10**, 98.
- Rodríguez-Celma J, Connorton JM, Kruse I, Green RT, Franceschetti M, Chen Y-T, Cui Y, Ling H-Q, Yeh K-C, Balk J.** 2019b. Arabidopsis BRUTUS-LIKE E3 ligases negatively regulate iron uptake by targeting transcription factor FIT for recycling. *Proceedings of the National Academy of Sciences* **116**, 17584-17591.
- Roschttardt H, Gaymard F, Dubos C.** 2020. Transcriptional Regulation of Iron Distribution in Seeds: A Perspective. *Frontiers in Plant Science* **11**, 725.
- Samira R, Li B, Kliebenstein D, Li C, Davis E, Gillikin JW, Long TA.** 2018. The bHLH transcription factor ILR3 modulates multiple stress responses in *Arabidopsis*. *Plant Molecular Biology* **97**, 297-309.
- Santi S, Schmidt W.** 2009. Dissecting iron deficiency-induced proton extrusion in *Arabidopsis* roots. *New Phytologist* **183**, 1072-1084.
- Schwarz B, Bauer P.** 2020. FIT, a regulatory hub for iron deficiency and stress signaling in roots, and FIT-dependent and -independent gene signatures. *Journal of Experimental Botany* **71**, 1694-1705.
- Selote D, Samira R, Matthiadis A, Gillikin JW, Long TA.** 2015. Iron-binding E3 ligase mediates iron response in plants by targeting basic helix-loop-helix transcription factors. *Plant Physiology* **167**, 273-286.

- Sharma R, Yeh KC.** 2020. The dual benefit of a dominant mutation in *Arabidopsis* IRON DEFICIENCY TOLERANT1 for iron biofortification and heavy metal phytoremediation. *Plant biotechnology journal* **18**, 1200-1210.
- Shen J, Xu X, Li T, Cao D, Han Z.** 2008. An MYB transcription factor from *Malus xiaojinensis* has a potential role in iron nutrition. *Journal of integrative plant biology* **50**, 1300-1306.
- Spielmann J, Vert G.** 2020. The many facets of protein ubiquitination and degradation in plant root iron deficiency responses. *Journal of Experimental Botany*, doi: 10.1093/jxb.
- Shen C, Yue R, Sun T, Zhang L, Yang Y, Wang H.** 2015. OsARF16, a transcription factor regulating auxin redistribution, is required for iron deficiency response in rice (*Oryza sativa* L.). *Plant Science* **231**, 148-58.
- Sun Y, Li JQ, Yan JY, Yuan JJ, Li GX, Wu YR, Xu JM, Huang RF, Harberd NP, Ding ZJ, Zheng SJ.** 2020. Ethylene promotes seed iron storage during *Arabidopsis* seed maturation via ERF95 transcription factor. *Journal of Integrative Plant Biology* **62**, 1193-1212.
- Stringlis IA, Yu K, Feussner K, de Jonge R, Van Bentum S, Van Verk MC, Berendsen RL, Bakker PA, Feussner I, Pieterse CM.** 2018. MYB72-dependent coumarin exudation shapes root microbiome assembly to promote plant health. *Proceedings of the National Academy of Sciences* **115**, E5213-E5222.
- Tanabe N, Noshi M, Mori D, Nozawa K, Tamoi M, Shigeoka S.** 2019. The basic helix-loop-helix transcription factor, bHLH11 functions in the iron-uptake system in *Arabidopsis thaliana*. *Journal of plant research* **132**, 93-105.
- Tissot N, Robe K, Gao F, Grant-Grant S, Boucherez J, Bellegarde F, Maghiaoui A, Marcelin R, Izquierdo E, Benhamed M.** 2019. Transcriptional integration of the responses to iron availability in *Arabidopsis* by the bHLH factor ILR3. *New Phytologist* **223**, 1433-1446.
- Touraine B, Vignols F, Przybyla-Toscano J, Ischebeck T, Dhalleine T, Wu H-C, Magno C, Berger N, Couturier J, Dubos C.** 2019. Iron-sulfur protein NFU2 is required for branched-chain amino acid synthesis in *Arabidopsis* roots. *Journal of experimental botany* **70**, 1875-1889.

- Trofimov K, Ivanov R, Eutebach M, Acaroglu B, Mohr I, Bauer P, Brumbarova T.** 2019. Mobility and localization of the iron deficiency-induced transcription factor bHLH039 change in the presence of FIT. *Plant Direct* **3**, e00190.
- Wang F-P, Wang X-F, Zhang J, Ma F, Hao Y-J.** 2018. MdMYB58 modulates Fe homeostasis by directly binding to the MdMATE43 promoter in plants. *Plant and Cell Physiology* **59**, 2476-2489.
- Wang L, Ying Y, Narsai R, Ye L, Zheng L, Tian J, Whelan J, Shou H.** 2013a. Identification of OsbHLH133 as a regulator of iron distribution between roots and shoots in *Oryza sativa*. *Plant, cell & environment* **36**, 224-236.
- Wang N, Cui Y, Liu Y, Fan H, Du J, Huang Z, Yuan Y, Wu H, Ling H-Q.** 2013b. Requirement and functional redundancy of Ib subgroup bHLH proteins for iron deficiency responses and uptake in *Arabidopsis thaliana*. *Molecular plant* **6**, 503-513.
- Wang S, Li L, Ying Y, Wang J, Shao JF, Yamaji N, Whelan J, Ma JF, Shou H.** 2020. A transcription factor OsbHLH156 regulates Strategy II iron acquisition through localising IRO2 to the nucleus in rice. *New Phytologist* **225**, 1247-1260.
- Wu H, Ling H-Q.** 2019. FIT-Binding Proteins and Their Functions in the Regulation of Fe Homeostasis. *Frontiers in plant science* **10**, 844.
- Xu Z, Liu X, He X, Xu L, Huang Y, Shao H, Zhang D, Tang B, Ma H.** 2017. The soybean basic helix-loop-helix transcription factor ORG3-Like enhances cadmium tolerance via increased iron and reduced cadmium uptake and transport from roots to shoots. *Frontiers in plant science* **8**, 1098.
- Yan JY, Li CX, Sun L, Ren JY, Li GX, Ding ZJ, Zheng SJ.** 2016. A WRKY transcription factor regulates Fe translocation under Fe deficiency. *Plant Physiology* **171**, 2017-2027.
- Yin L, Wang Y, Yan M, Zhang X, Pan H, Xu X, Han Z.** 2013. Molecular cloning, polyclonal antibody preparation, and characterization of a functional iron-related transcription factor IRO2 from *Malus xiaojinensis*. *Plant physiology and biochemistry* **67**, 63-70.
- Yin L, Wang Y, Yuan M, Zhang X, Xu X, Han Z.** 2014. Characterization of MxFIT, an iron deficiency induced transcriptional factor in *Malus xiaojinensis*. *Plant physiology and biochemistry* **75**, 89-95.

- Yuan Y, Wu H, Wang N, Li J, Zhao W, Du J, Wang D, Ling H-Q.** 2008. FIT interacts with AtbHLH38 and AtbHLH39 in regulating iron uptake gene expression for iron homeostasis in Arabidopsis. *Cell research* **18**, 385-397.
- Yuan YX, Zhang J, Wang DW, Ling HQ.** 2005. AtbHLH29 of Arabidopsis thaliana is a functional ortholog of tomato FER involved in controlling iron acquisition in strategy I plants. *Cell research* **15**, 613-621.
- Zamioudis C, Hanson J, Pieterse CMJ.** 2014. β -Glucosidase BGLU42 is a MYB72-dependent key regulator of rhizobacteria-induced systemic resistance and modulates iron deficiency responses in Arabidopsis roots. *New Phytologist* **204**, 368-379.
- Zhang G, Liu W, Feng Y, Li D, Li K, Sun Q, Zhai L, Wu T, Zhang X, Xu X.** 2020a. Ethylene Response Factors MbERF4 and MbERF72 Suppress Iron Uptake in Woody Apple Plants by Modulating Rhizosphere pH. *Plant and Cell Physiology* **61**, 699-711.
- Zhang H, Li Y, Pu M, Xu P, Liang G, Yu D.** 2020b. Oryza sativa POSITIVE REGULATOR OF IRON DEFICIENCY RESPONSE 2 (OsPRI2) and OsPRI3 are involved in the maintenance of Fe homeostasis. *Plant, cell & environment* **43**, 261-274.
- Zhang H, Li Y, Yao X, Liang G, Yu D.** 2017. Positive regulator of iron homeostasis1, OsPRI1, facilitates iron homeostasis. *Plant Physiology* **175**, 543-554.
- Zhang J, Liu B, Li M, Feng D, Jin H, Wang P, Liu J, Xiong F, Wang J, Wang H-B.** 2015. The bHLH transcription factor bHLH104 interacts with IAA-LEUCINE RESISTANT3 and modulates iron homeostasis in Arabidopsis. *The Plant Cell* **27**, 787-805.
- Zhang L, Nakanishi Itai R, Yamakawa T, Nakanishi H, Nishizawa NK, Kobayashi T.** 2014. The Bowman–Birk Trypsin Inhibitor IBP1 Interacts with and Prevents Degradation of IDEF1 in Rice. *Plant Molecular Biology Reporter* **32**, 841-851.
- Zhang X, Wang C, Zhang Y, Sun Y, Mou Z.** 2012. The Arabidopsis mediator complex subunit16 positively regulates salicylate-mediated systemic acquired resistance and jasmonate/ethylene-induced defense pathways. *The Plant Cell* **24**, 4294-4309.
- Zhao M, Song A, Li P, Chen S, Jiang J, Chen F.** 2014. A bHLH transcription factor regulates iron intake under Fe deficiency in chrysanthemum. *Scientific reports* **4**, 6694.

- Zhao Q, Ren Y-R, Wang Q-J, Wang X-F, You C-X, Hao Y-J.** 2016a. Ubiquitination-related MdbT scaffold proteins target a bHLH transcription factor for iron homeostasis. *Plant Physiology* **172**, 1973-1988.
- Zhao Q, Ren YR, Wang QJ, Yao YX, You CX, Hao YJ.** 2016b. Overexpression of Mdb HLH 104 gene enhances the tolerance to iron deficiency in apple. *Plant biotechnology journal* **14**, 1633-1645.
- Zheng L, Ying Y, Wang L, Wang F, Whelan J, Shou H.** 2010. Identification of a novel iron regulated basic helix-loop-helix protein involved in Fe homeostasis in *Oryza sativa*. *BMC plant biology* **10**, 166.
- Zhu XF, Wu Q, Meng YT, Tao Y, Shen RF.** 2020. AtHAP5A regulates iron translocation in iron-deficient *Arabidopsis thaliana*. *Journal of Integrative Plant Biology* doi.org/10.1111/jipb.12984
- Zhou L-J, Zhang C-L, Zhang R-F, Wang G-L, Li Y-Y, Hao Y-J.** 2019. The SUMO E3 ligase MdSIZ1 targets MdbHLH104 to regulate plasma membrane H⁺-ATPase activity and iron homeostasis. *Plant Physiology* **179**, 88-106.

Table 1. bHLH TFs involved in the regulation of iron homeostasis in plants.

Name	Clade	Species	Interacting proteins	Reference
AtbHLH38*	Ib	<i>A. thaliana</i>	FIT/AtbHLH29, DELLAs	Yuan <i>et al.</i> , 2008; Wang <i>et al.</i> , 2013
AtbHLH39*	Ib	<i>A. thaliana</i>	FIT/AtbHLH29, DELLAs	Yuan <i>et al.</i> , 2008; Wang <i>et al.</i> , 2013
AtbHLH100*	Ib	<i>A. thaliana</i>	FIT/AtbHLH29	Wang <i>et al.</i> , 2013
AtbHLH101*	Ib	<i>A. thaliana</i>	FIT/AtbHLH29	Wang <i>et al.</i> , 2013
FIT/AtbHLH29 *	IIIa	<i>A. thaliana</i>	AtbHLH38, AtbHLH39, AtbHLH100, AtbHLH101, AtbHLH18, AtbHLH19, AtbHLH20, AtbHLH25, BTSL1, BTSL2, CIPK11, DELLAs, EIL1, EIN3, MED16, ZAT12	Colangelo and Guerinot, 2004; Jakoby <i>et al.</i> , 2004; Yuan <i>et al.</i> , 2005
MYC2/AtbHLH6	IIIe	<i>A. thaliana</i>		Cui <i>et al.</i> , 2018
AtbHLH18*	IVa	<i>A. thaliana</i>	FIT/AtbHLH29	Cui <i>et al.</i> , 2018
AtbHLH19*	IVa	<i>A. thaliana</i>	FIT/AtbHLH29	Cui <i>et al.</i> , 2018
AtbHLH20*	IVa	<i>A. thaliana</i>	FIT/AtbHLH29	Cui <i>et al.</i> , 2018
AtbHLH25*	IVa	<i>A. thaliana</i>	FIT/AtbHLH29	Cui <i>et al.</i> , 2018
AtbHLH11	IVb	<i>A. thaliana</i>	IDT1/AtbHLH34, AtbHLH104, ILR3/AtbHLH105, AtbHLH115	Tanabe <i>et al.</i> , 2019
PYE/AtbHLH47	IVb	<i>A. thaliana</i>	ILR3/AtbHLH105, AtbHLH115	Long <i>et al.</i> , 2010
URI/AtbHLH121	IVb	<i>A. thaliana</i>	IDT1/AtbHLH34, AtbHLH104, ILR3/AtbHLH105, AtbHLH115	Gao <i>et al.</i> , 2020b; Kim <i>et al.</i> , 2019; Lei <i>et al.</i> , 2020
IDT1/AtbHLH34	IVc	<i>A. thaliana</i>	IDT1/AtbHLH34, AtbHLH104,	Li <i>et al.</i> , 2016

			ILR3/AtbHLH105, AtbHLH115, AtbHLH11, URI/bHLH121	
AtbHLH104	IVc	<i>A. thaliana</i>	IDT1/AtbHLH34, AtbHLH104, ILR3/AtbHLH105, AtbHLH115, AtbHLH11, URI/ bHLH121	Zhang <i>et al.</i> , 2015
ILR3/AtbHLH105	IVc	<i>A. thaliana</i>	IDT1/AtbHLH34, AtbHLH104, ILR3/AtbHLH105, AtbHLH115, AtbHLH11, PYE/ bHLH47, URI/bHLH121, BTS	Zhang <i>et al.</i> , 2015
AtbHLH115	IVc	<i>A. thaliana</i>	IDT1/AtbHLH34, AtbHLH104, ILR3/AtbHLH105, AtbHLH115, AtbHLH11, PYE/ bHLH47, URI/bHLH121, BTS	Liang <i>et al.</i> , 2015
OsIRO2/OsbHLH56	Ib	<i>O. sativa</i>	OsFIT/OsbHLH156	Ogo <i>et al.</i> , 2006
OsIRO3/OsbHLH63	IVb	<i>O. sativa</i>		Zheng <i>et al.</i> , 2010
OsFIT/OsbHLH156	IIIa	<i>O. sativa</i>	OsIRO2/OsbHLH56	Liang <i>et al.</i> , 2020; Wang <i>et al.</i> , 2020
OsPRI1/OsbHLH60	IVc	<i>O. sativa</i>	OsHRZ1	Zhang <i>et al.</i> , 2017
OsPRI2/OsbHLH58	IVc	<i>O. sativa</i>	OsHRZ1	Kobayashi <i>et al.</i> , 2019
OsPRI3/OsbHLH59	IVc	<i>O. sativa</i>	OsHRZ1	Kobayashi <i>et al.</i> , 2019
OsbHLH133	VIIIc	<i>O. sativa</i>		Wang <i>et al.</i> , 2013
GmbHLH57	IIIa	<i>G. max</i>	GmbHLH300	Li <i>et al.</i> , 2018
GmbHLH300	Ib	<i>G. max</i>	GmbHLH57	Li <i>et al.</i> , 2018
FER	IIIa	<i>S. lycopersicum</i>	SlbHLH68	Ling <i>et al.</i> , 2002
SlbHLH68	Ib	<i>S. lycopersicum</i>	FER	Du <i>et al.</i> , 2015
MxIRO2	Ib	<i>M. xiaojinensis</i>		Yin <i>et al.</i> , 2013
MxFIT	IIIa	<i>M. xiaojinensis</i>	MxERF4	Yin <i>et al.</i> , 2014
PtFIT	IIIa	<i>P. tremula</i>		Huang and Dai, 2015
PtIRO	Ib	<i>P. tremula</i>		Huang and Dai, 2015
SAT1/MdbHLH18	IVa	<i>M. domestica</i>	MdMYB58	Wang <i>et al.</i> , 2018

MdbHLH104	IVc	<i>M. domestica</i>	MdBT1, MdBT2, MdbHLH104, MdbHLH105, MdbHLH115, MdbHLH11, MdbHLH121, MdPYE	Zhao <i>et al.</i> , 2016b
NtbHLH1	IVb	<i>N. tabacum</i>		Li <i>et al.</i> , 2020
CmbHLH1	IVc	<i>C. morifolium</i>		Zhao <i>et al.</i> , 2014
FEFE/CmbHLH38	Ib	<i>C. melo</i>	CmFIT	Ramamurthy and Waters, 2017
GmORG3	Ib	<i>G. max</i>		Xu <i>et al.</i> , 2017

*: for details on posttranslational regulation of FIT activity, see Wu and Ling, 2020 and Schwarz and Bauer, 2020.

Table 2. TFs other than bHLHs involved in the regulation of iron homeostasis in plants.

TF family	Gene name	Species	Interacting proteins	Reference
ABI3/VP1	IDEF1	<i>O. sativa</i>	IBP1	Kobayashi <i>et al.</i> , 2007
ARF	OsARF12	<i>O. sativa</i>		Qi <i>et al.</i> , 2012
ARF	OsARF16	<i>O. sativa</i>		Shen <i>et al.</i> , 2016
C2H2	ZAT12*	<i>A. thaliana</i>	FIT/bHLH29	Le <i>et al.</i> , 2016
EIL	EIN3*	<i>A. thaliana</i>	FIT/bHLH29	Lingam <i>et al.</i> , 2011
EIL	EIL1*	<i>A. thaliana</i>	FIT/bHLH29	Lingam <i>et al.</i> , 2011
ERF	ERF4	<i>A. thaliana</i>		Liu <i>et al.</i> , 2017a
ERF	ERF72	<i>A. thaliana</i>		Liu <i>et al.</i> , 2017b
ERF	ERF95	<i>A. thaliana</i>		Sun <i>et al.</i> , 2020
ERF	MxERF4	<i>M. xiaojinensis</i>	MxFIT	Liu <i>et al.</i> , 2018
ERF	MbERF4	<i>M. baccata</i>	MbERF72	Zhang <i>et al.</i> , 2020a
ERF	MbERF72	<i>M. baccata</i>	MbERF4	Zhang <i>et al.</i> , 2020a
MYB (R2R3)	MYB10	<i>A. thaliana</i>		Palmer <i>et al.</i> , 2013
MYB (R2R3)	MYB28	<i>A. thaliana</i>		Coletto <i>et al.</i> , 2020
MYB (R2R3)	MYB29	<i>A. thaliana</i>		Coletto <i>et al.</i> , 2020
MYB (R2R3)	MYB72	<i>A. thaliana</i>		Palmer <i>et al.</i> , 2013
MYB (R2R3)	MdMYB58	<i>M. domestica</i>	MdSAT1/MdbHLH18	Wang <i>et al.</i> , 2018
MYB (R2R3)	MxMYB1	<i>M. xiaojinensis</i>		Shen <i>et al.</i> , 2008
NAC	IDEF2	<i>O. sativa</i>		Ogo <i>et al.</i> , 2008
NF-YC	HAP5A/NF-YC1	<i>A. thaliana</i>		Zhu <i>et al.</i> , 2020
WRKY	WRKY46	<i>A. thaliana</i>		Yan <i>et al.</i> , 2016

*: for details on posttranslational regulation of FIT activity, see Wu and Ling. 2020 and Schwarz and Bauer, 2020.

Figure legends

Figure 1. Schematic diagram of iron uptake strategies in Arabidopsis and rice. In Arabidopsis, AHA2 secretes protons into the rhizosphere to increase Fe^{3+} solubility; PDR9 secretes Fe-mobilizing coumarins (*i.e.* fraxetin and sideretin) to mobilize and chelate Fe^{3+} . Fe^{3+} is reduced into Fe^{2+} and is subsequently transported into the root epidermal cells by FRO2 and IRT1, respectively. AHA2, FRO2 and IRT1 form a protein complex optimizing Fe acquisition by creating a local environment with low pH and high Fe^{2+} concentration. Rice biosynthesizes and secretes DMA (2-deoxy-mugineic acid) to chelate Fe^{3+} . Fe^{3+} -DMA complexes are transported into root cells via YSL15 and YSL16. In addition, rice also uptakes Fe^{2+} from the soil through the activity of two Fe^{2+} transporters, OsIRT1 and OsIRT2 under waterlogged soil condition. The secretion of CA and PCA via the PEZ2 phenolics efflux transporter participates to Fe^{3+} mobilization and reduction into Fe^{2+} . AHA2: H^+ -ATPase 2; PDR9: PLEIOTROPIC DRUG RESISTANCE 9/ATP-BINDING CASSETTE G37; FRO2: FERRIC REDUCTION OXIDASE 2; IRT1: IRON-REGULATED TRANSPORTER 1, F6'H1: FERULOYL CoA 6' HYDROXYLASE 1; S8H: SCPOLETIN 8-HYDROXYLASE; CYP82C4: FRAXETIN 5-HYDROXYLASE; TOM1: TRANSPORTER OF MUGINEIC ACID 1; YSL15/16: YELLOW STRIPE-LIKE 15/16; OsIRT1/2: Rice IRON-REGULATED TRANSPORTER 1/2; PEZ2: PHENOLICS EFFLUX ZERO 2; OsNAS: NICOTIANAMINE SYNTHASE; OsNAAT: NICOTIANAMINE AMINOTRANSFERASE; OsDMAS: DEOXYMUGINEIC ACID SYNTHASE; SAM: S-adenosyl methionine; NA: nicotianamine; CA: caffeic acid; PCA: protocatechuic acid.

Figure 2. Phylogenetic tree of bHLH TFs involved in the regulation of iron homeostasis in plants. The tree was constructed by using the neighbor-joining method with the MEGA software (v10.0.5) using full-length bHLH amino acid sequences. The bootstrap analysis was carried out with 1000 replicates. Sequences were aligned prior to the construction of the phylogenetic tree using the ClustalX software (v2.0.11). The different bHLH clades are designated as previously reported (Heim *et al.*, 2003). Species abbreviations: At, *Arabidopsis thaliana*; Md, *Malus domestica*; Nt, *Nicotiana tabacum*;

Os, *Oryza sativa*; Sl, *Solanum lycopersicum*; Gm, *Glycine max*; Mx, *Malus xiaojinensis*; Cm, (CmbHLH1), *Chrysanthemum morifolium*; Cm (CmbHLH38), *Cucumis melo*; Pt, *Populus tremula*.

Figure 3. bHLH TFs network that regulates iron homeostasis in Arabidopsis.

Question marks indicate protein complexes for which transcriptional activities have not yet been clearly demonstrated. The colour code for the bHLH TFs refers to the clades described Figure 2. Non bHLH proteins are in grey. EIN3 and EIL1 favor FIT stability whereas MED16 and MED25 mediate the FIT/bHLH29-dependent iron deficiency responses (for additional details on posttranslational regulation of FIT, see Kobayashi, 2019; Schwarz and Bauer, 2020; Spielmann and Vert, 2020; Wu and Ling, 2019). EIN3: ETHYLENE INSENSITIVE 3; EIL1: ETHYLENE INSENSITIVE 3-LIKE1; FBP: FIT-BINDING PROTEIN; MED16 AND 25: MEDIATOR SUBUNIT 16 and 25; ZAT12: ZINC FINGER OF ARABIDOPSIS THALIANA 12.

Figure 4. bHLH TFs network that regulates iron homeostasis in rice. Question mark indicates the putative degradation mechanism for OsPRI1 and OsPRI2 via OsHRZ2. The colour code for the bHLH TFs refers to the clades described Figure 2. Non bHLH proteins are in grey. IBP1.1: IDEF1-BINDING PROTEINS 1.1.

Figure 1. Schematic diagram of iron uptake strategies in Arabidopsis and rice. In Arabidopsis, AHA2 secretes protons into the rhizosphere to increase Fe^{3+} solubility; PDR9 secretes Fe-mobilizing coumarins (i.e. fraxetin and sideretin) to mobilize and chelate Fe^{3+} . Fe^{3+} is reduced into Fe^{2+} and is subsequently transported into the root epidermal cells by FRO2 and IRT1, respectively. AHA2, FRO2 and IRT1 form a protein complex optimizing Fe acquisition by creating a local environment with low pH and high Fe^{2+} concentration. Rice biosynthesizes and secretes DMA to chelate Fe^{3+} . Fe^{3+} -DMA complexes are transported into root cells via YSL15 and YSL16. In addition, rice also uptake Fe^{2+} from the soil through the activity of two Fe^{2+} transporters, OsIRT1 and OsIRT2 under waterlogged soil condition. The secretion of CA and PCA via the PEZ2 phenolics efflux transporter participates to Fe^{3+} mobilization and reduction into Fe^{2+} . AHA2: H^+ -ATPase 2; PDR9: PLEIOTROPIC DRUG RESISTANCE 9/ATP-BINDING CASSETTE G37; FRO2: FERRIC REDUCTION OXIDASE 2; IRT1: IRON-REGULATED TRANSPORTER 1, F6'H1: FERULOYL CoA 6' HYDROXYLASE 1; S8H: SCOPLETIN 8-HYDROXYLASE; CYP82C4: FRAXETIN 5-HYDROXYLASE; TOM1: TRANSPORTER OF MUGINEIC ACID 1; YSL15/16: YELLOW STRIPE-LIKE 15/16; OsIRT1/2: RICE IRON-REGULATED TRANSPORTER 1/2; PEZ2: PHENOLICS EFFLUX ZERO 2; OsNAS: NICOTIANAMINE SYNTHASE; OsNAAT: NICOTIANAMINE AMINOTRANSFERASE; OsDMAS: DEOXYMUGINEIC ACID SYNTHASE; SAM: S-adenosyl methionine; NA: nicotianamine; CA: caffeic acid; PCA: protocatechuic acid.

Figure 2. Phylogenetic tree of bHLH TFs involved in the regulation of iron homeostasis in plants. The tree was constructed by using the neighbor-joining method with the MEGA software (v10.0.5) using full-length bHLH amino acid sequences. The bootstrap analysis was carried out with 1000 replicates. Sequences were aligned, prior to the construction of the phylogenetic tree, using the ClustalX software (v2.0.11). The different bHLH clades are designated as previously reported (Heim et al., 2003). Species abbreviations: At, *Arabidopsis thaliana*; Md, *Malus domestica*; Nt, *Nicotiana tabacum*; Os, *Oryza sativa*; Sl, *Solanum lycopersicum*; Gm, *Glycine max*; Mx, *Malus xiaojinensis*; Cm, (CmbHLH1), *Chrysanthemum morifolium*; Cm (CmbHLH38), *Cucumis melo*; Pt, *Populus tremula*.

Figure 3. bHLH TFs network that regulates iron homeostasis in Arabidopsis. Question marks indicate protein complexes for which transcriptional activities have not yet been clearly demonstrated. The color code for the bHLH TFs refers to the clades described Figure 2. EIN3 and EIL1 favor FIT stability whereas MED16 and MED25 mediate the FIT/bHLH29-dependent iron deficiency responses (for additional details on posttranslational regulation of FIT see Kobayashi, 2019; Schwarz and Bauer, 2020; Spielmann and Vert, 2020; Wu and Ling, 2019). Non bHLH proteins are in grey. EIN3: ETHYLENE INSENSITIVE 3; EIL1: ETHYLENE INSENSITIVE 3-LIKE1; MED16 and 25: FBP: FIT-BINDING PROTEIN; MED16 AND 25: MEDIATOR SUBUNIT 16 and 25; ZAT12: ZINC FINGER OF ARABIDOPSIS THALIANA 12.

Downloaded from <https://academic.oup.com/jxb/advance-article/doi/10.1093/jxb/eraa556/6007775> by guest on 30 November 2020