



HAL
open science

Odor-induced saltiness enhancement: Insights into the brain chronometry of flavor perception

Charlotte Sinding, Henri Thibault, Thomas Hummel, Thierry Thomas-Danguin

► To cite this version:

Charlotte Sinding, Henri Thibault, Thomas Hummel, Thierry Thomas-Danguin. Odor-induced saltiness enhancement: Insights into the brain chronometry of flavor perception. *Neuroscience*, 2021, 452, pp.126-137. 10.1016/j.neuroscience.2020.10.029 . hal-03031862

HAL Id: hal-03031862

<https://hal.inrae.fr/hal-03031862>

Submitted on 30 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH ARTICLE

C. Sinding et al. / Neuroscience xxx (2020) xxx–xxx

Odor-Induced Saltiness Enhancement: Insights Into The Brain Chronometry Of Flavor Perception

Charlotte Sinding,^{a*} Henri Thibault,^a Thomas Hummel^b and Thierry Thomas-Danguin^a

^a Centre des Sciences du Goût et de l'Alimentation, AgroSup Dijon, CNRS, INRAE, Université Bourgogne Franche-Comté, F-21000 Dijon, France

^b Smell & Taste Clinic, Department of Otorhinolaryngology, TU Dresden, Dresden, Germany

Abstract—Flavor perception results from the integration of at least odor and taste. Evidence for such integration is that odors can have taste properties (odor-induced taste). Most brain areas involved in flavor perception are high-level areas; however, primary gustatory and olfactory areas also show activations in response to a combination of odor and taste. While the regions involved in flavor perception are now quite well identified, the network's organization is not yet understood. Using a close to real salty soup model with electroencephalography brain recording, we evaluated whether odor-induced saltiness enhancement would result in differences of amplitude and/or latency in late cognitive P3 peak mostly and/or in P1 early sensory peak. Three target solutions were created from the same base of green-pea soup: i) with a “usual” salt concentration (PPS2), ii) with “reduced” salt (PPS1: –50%), and iii) with reduced salt and a “beef stock” odor (PPS1B). Sensory data showed that the beef odor produced saltiness enhancement in PPS1B in comparison to PPS1. As the main EEG result, the late cognitive P3 peak was delayed by 25 ms in the odor-added solution PPS1B compared to PPS1. The odor alone did not explain this peak amplitude and higher latency in the P3 peak. These results support the classical view that high-level integratory areas process odor–taste interactions with potential top-down effects on primary sensory regions.

© 2020 IBRO. Published by Elsevier Ltd. All rights reserved.

Key words: Olfaction, Taste, Integration, Perception, Food, Electroencephalography.

INTRODUCTION

We experience food as a unitary perception, which we commonly call “taste”. This common “taste” is actually a holistic perception of at least olfactory and gustatory inputs, called “flavor perception”. Odor-induced taste enhancement (OITE) is a phenomenon that derives from the integration of taste and odor into flavor perception. For example, it was shown that a strawberry odor could increase the sweetness of a whipped-cream with sucrose. This result was first highlighted by Frank and Byram (1988). They also defined a fundamental principle of OITE, namely that only congruent odors and tastes would produce OITE, therefore pointing at the role of experience in shaping OITE. Indeed congruent, familiar, and complex flavor mixtures -which are more prone to be perceived as configural units- are more effective in producing OITE (Prescott et al., 2004; Small and Prescott, 2005; Labbe et al., 2006). Several independent labs have later replicated this finding and further demon-

strated odor-induced taste enhancement of other tastes (Frank and Byram, 1988; Schifferstein and Verlegh, 1996; Sakai et al., 2001; Djordjevic et al., 2004; Prescott et al., 2004; Lawrence et al., 2009; Wang et al., 2019). OITE is, therefore, a reliable phenomenon. Other odor–taste interactions have also been established, such as the taste-induced odor enhancement (i.e., the reverse effect of OITE) (Lim et al., 2014; Linscott and Lim, 2016). In our study and the discussion of the results, we focused on the odor-induced saltiness enhancement only.

Most studies on OITE used water with sugar or salt and aroma, which produced non-ecologically relevant, unfamiliar and likely unpleasant perceptions (Prescott et al., 2004; Welge-Lüssen et al., 2005; Marshall et al., 2006; Prescott and Murphy, 2009; Welge-Lussen et al., 2009; Lim and Johnson, 2011, 2012; Seo et al., 2013). To overcome this issue, one can use close-to-real food models, which produce more familiar and holistic food representations. It may also facilitate the OITE with appropriate congruent aroma and smooth out significant hedonic variations that could mask subtle integration mechanisms (Prescott, 1999; Small, 2012; Mroczko-Wąsowicz, 2016; Thomas-Danguin et al., 2016). Other studies used more complex and familiar food models.

*Corresponding author.

E-mail address: charlotte.sinding@inrae.fr (C. Sinding).

Abbreviations: EEG, Electroencephalography; ERPs, event-related potentials; fMRI, functional magnetic resonance imaging; OISE, odor-induced saltiness enhancement.

For example, ethyl 2-methyl butanoate was used in a fruit juice to enhance sweet perception (Barba et al., 2018). In another study, the authors used a sardine aroma to significantly enhance saltiness in a cheese model (Syarifuddin et al., 2016). In the present study, we studied odor-induced saltiness enhancement (OISE). A salty food model has been designed from a green-pea soup base, which was chosen for its composition with a negligible quantity of salt and an easily identifiable odor component. Five conditions with different salt and aroma quantities were selected to produce OISE according to previous results (Lawrence et al., 2009; Nasri et al., 2013). The first condition was the soup added with a standard (usual) level of salt (6.25 g/L), to record the most familiar level of saltiness in this kind of product and to test whether OISE could reinforce saltiness up to a “normal” saltiness intensity. The second solution was 50% salt reduced. The third condition, which is the target beverage, was reduced in salt (50%) and supplemented with a “beef stock” odor chosen for its potential to increase saltiness perception. Finally, two controls were tested, the base soup alone and the base soup with the odor component, to test the effect of added odor in the food model.

Endogenous mechanisms produce OITE in the brain. Several functional magnetic resonance imaging (fMRI) studies have investigated brain areas involved in flavor perception, leading to the identification of a relatively consensual flavor network (Rolls and Baylis, 1994; Rolls, 1997; De Araujo et al., 2003; Small and Prescott, 2005; Seo et al., 2013; Seubert et al., 2015). In these studies, supra-additive activations for the odorant-tastant mixture were found in high-level areas, in the orbitofrontal cortex, the dorsal mid-insula, and the perirhinal cortex (De Araujo et al., 2003; Seo et al., 2013; Small et al., 2013; Seubert et al., 2015). However, odor–taste convergence was also found in the primary gustatory cortex, more precisely in the anterior insula and frontal operculum (De Araujo et al., 2003; Seubert et al., 2015). Regarding these fMRI results, two views exist i) one consists in a hierarchical integration, starting with a parallel unimodal encoding of odor and taste in their respective cortices and further elaborated by higher-order unimodal zones, before converging onto multisensory integrative areas to form the flavor perception; ii) while the second proposed that odor and taste are already integrated into primary olfactory and gustatory cortices (Small and Prescott, 2005; Verhagen and Engelen, 2006; Verhagen, 2007; Prescott, 2012; Small et al., 2013).

To understand whether odor and taste already interact in the primary cortices or later in higher cortices, we need to study the chronometry of odor–taste integration and interaction. Electroencephalography (EEG) is of particular interest to gain insights into these questions. It permits quantitative measures of global brain activations with a resolution of milliseconds. Olfactory-gustatory event-related potentials (ERPs) give access to the chronometry of interactions between gustatory and olfactory cortices. To do so, one should select appropriate stimuli, which permit to isolate variables of interest (e.g., real saltiness or induced saltiness). An

event-related potential (ERP) is a sequence of brain components identified by the maximum amplitudes of a series of positive and negative peaks, from P1 the earliest to P3 the latest measurable. ERP reflects the different steps of information processing in the cortex. The peak amplitude and latency provide a quantitative measure of the intensity and/or amount of neurons discharging in a synchronized way in response to the stimulus provided at t0. The early P1 peak mainly occurs in primary sensory areas and represents the processing of sensory and chemically related properties of food. The late P3 peak occurs mostly in high integratory and cognitive areas and illustrates endogenous processing such as emotional, semantic, decisional, and top-down mechanisms towards primary regions.

While an extensive literature exists on food-related visual event-related potentials (ERPs) (for review, see Carbine et al., 2018), very few studies were based on the senses directly involved in flavor perception: olfaction and gustation. To the best of our knowledge, no EEG studies showed the brain mechanisms of odor-induced taste enhancement. However, Welge-Lüssen et al. (2005), Welge-Lüssen et al. (2009) designed two studies to show the effect of taste (sucrose or lemon pulp) on odor (vanilla) or trigeminal perception (elicited by CO₂) respectively. In these studies, participants were sucking on a taste dispenser when an odor was sent orthonasally (Welge-Lüssen et al., 2005) or retronasally (Welge-Lüssen et al., 2009) with an olfactometer. This moment corresponded to the start of the ERP, which therefore highlighted the odor processing modulated by the taste. Although sensory results did not show odor or taste enhancement, ERPs tended to higher amplitude and reduced early and late peak latencies (P1 and P3), only when the taste matched the odor. These two peaks represent the earliest and latest observable brain mechanisms of the evoked potential measured with EEG. Therefore, Welge-Lüssen’s studies showed that taste sped up the processing of a congruent odor from the very first processing mechanisms (P1 peak). Although these results did not permit observing supra-additive effects for a flavor mixture compared to its odorant-tastant components, they were interestingly discussed in terms of priming. To observe supra-additive effects, one should synchronously stimulate the olfactory and gustatory cortices and compare activation for the mixture to the single components. Recent electrophysiological results in animal reconsidered the classical view of late odor–taste integration. They indeed showed activations in a region of the primary olfactory cortex, i.e., the piriform cortex, in response to sucrose (sweet taste), which the authors considered early interactions (Maier et al., 2012, 2015; Maier, 2017). These results, therefore, challenge the classical theory of late brain interactions between odor and taste and suggest that primary olfactory and gustatory areas may interact as early as the primary EEG peaks such as P1 (100–200 ms).

Therefore, we addressed the chronometry of the integration of odor and taste into flavor perception, in

174 humans, by studying the chronometry of brain
 175 mechanisms leading to OISE. The classical view, which
 176 consisted of a hierarchical integration of flavor, from
 177 primary gustatory and olfactory areas to secondary or
 178 tertiary integratory cortices, has been further expanded
 179 to explain OITE (Verhagen and Engelen, 2006;
 180 Verhagen, 2007; Small, 2008; Prescott, 2012). Following
 181 the integration of odor and taste into the flavor, top-down
 182 feedback may control activations in gustatory areas pro-
 183 ducing an increased endogenous perception of saltiness
 184 intensity. Following this reasoning, the odor-induced salti-
 185 ness enhancement should be observed only on the ERP's
 186 late components. Therefore, we hypothesized that differ-
 187 ences of amplitude and/or latency could be observed
 188 mostly on the latest peak of olfactory-gustatory ERP
 189 (the late P3 peak) and not on the P1 peak involving brain
 190 circuits responding to exogenous properties of the food
 191 such as tastant concentration. In the study, we did not
 192 address whether retronasal odor stimulation is necessary
 193 for the supra-additivity of the flavor solution. Still, to avoid
 194 any potential bias, we used only retronasal odor percep-
 195 tion. Because of the presumed importance of oral referral
 196 in flavor perception (Small, 2008; Spence, 2016), which is
 197 supported by EEG results (Welge-Lüssen et al., 2009;
 198 Masaoka et al., 2010), participants should be stimulated
 199 in-mouth so that aromas could be perceived through the
 200 retronasal route. However, the aroma perception is sup-
 201 posed to be maximal when participants are swallowing.
 202 A dedicated paradigm was designed to account for the
 203 swallowing artifacts and the need for synchrony between
 204 odor and taste perceptions.

205 EXPERIMENTAL PROCEDURES

206 Participants

207 Twenty-one participants naïve to olfactory and gustatory
 208 testing were recruited (18–30 years old, 15 women).
 209 Data from 8 participants were discarded because of
 210 their low number of epochs after artifact rejection (less
 211 than 6 epochs in at least 2 stimulus conditions). The
 212 stimulation of participants in-mouth during EEG
 213 recording is tricky due to tongue and jaw movements
 214 during stimulation, which induce many artifacts. Power
 215 analysis (GPower) showed that 13 participants were
 216 sufficient to have adequate power on the amplitude and
 217 latencies of peaks (power = 0.82). Participants were
 218 asked not to drink or eat, at least 1 h30 min before the
 219 test sessions, to minimize food exposure. All
 220 participants were right-handed (self-reported) and
 221 normosmic (European Test of Olfactory Capabilities,
 222 ETOC; Thomas-Danguin et al., 2003). The experimental
 223 procedure was explained to each participant before
 224 recruitment and again before each test session. Partici-
 225 pants signed an informed consent form to participate in
 226 the study. They received 10€ compensation for each hour
 227 spent performing test sessions. The study was conducted
 228 following the Helsinki Declaration and was validated by
 229 the ethics committee CPP EST 1N°2016/62 (ID RCB:
 230 2016-A01732-49).

Solutions

231 Three stock solutions were prepared: (1) the base
 232 solution was a green-pea soup extracted from an
 233 unsalted green-pea puree (Nestlé®, green pea
 234 NaturNes). The puree was centrifuged at 15,000 RPM
 235 for 20 min at 20 °C (rotor JLA 16–250, Beckman
 236 Coulter), and the supernatant, which contained
 237 everything but the non-soluble particles, was collected
 238 and stored at –20 °C. (2) A salty solution was prepared
 239 at 12.5 g/L NaCl in Evian® water (Danone, France). (3)
 240 The aromatic solution (B) was prepared at 500 ppm of
 241 Beef Bouillon Flavor (YF 555,768 CB, Firmenich) in
 242 Evian water. Evian water was chosen because, in Dijon,
 243 it is the bottled water perceived as the most neutral
 244 (Teillet et al., 2010). Solutions 2 and 3 were prepared
 245 24 h before the test and kept at 4 °C. The test's day, the
 246 base solution was defrosted in the microwave for 5 min-
 247 utes (defrost position). All solutions were then heated at
 248 35 °C (all tubes of the gustometer are water bath) in the
 249 gustometer to match the buccal temperature, 1 h before
 250 the test. Five target solutions were then mixed using the
 251 gustometer: PP (50% base solution + 50% Evian Water),
 252 PPB (50% base solution + 25% aromatic solution
 253 + 25% Evian water), PPS1 (50% base solution + 25%
 254 salty solution + 25% Evian water), PPS1B (50% base
 255 solution + 25% salty solution + 25% aromatic solution),
 256 PPS2 (50% base solution + 50% salty solution). Evian
 257 water, PP, and PPB contained negligible levels of salt
 258 (maximum 0.25 g/L), which are lower than the standard
 259 detection threshold in water (0.58 g/L) (Bartoshuk,
 260 1974). Furthermore, B contained minimal salt (0.03 g/L).
 261

262 Experimental procedure

263 The entire experiment took place in a ventilated air-
 264 conditioned room (23 °C), dedicated to EEG recording,
 265 with controlled light and acoustic insulation. The
 266 procedure was planned over two sessions: 1) a training
 267 session and 2) an EEG recording session, spaced by a
 268 maximum of 8 days, to keep the training effective.

269 Training session

270 Before the training session, participants were tested for
 271 their general olfactory abilities using the ETOC
 272 (Thomas-Danguin et al., 2003). This test consists of an
 273 odor supra-threshold detection task combined with an
 274 identification task for 16 different odors using a 4-
 275 alternatives forced-choice procedure (1 vial odorized
 276 among 4, then 1 correct odor descriptor among 4). The
 277 minimum score required to consider normal olfactory abil-
 278 ities was 12 (out of 16). All participants succeeded in the
 279 ETOC test.

280 The training session's objective was twofold: 1) to
 281 familiarize participants with the gustometer and the
 282 testing room and 2) train participants to breathe with the
 283 velum opened. Solutions were delivered with a GU002
 284 gustometer (Burghart, Wedel, Germany). Initially, the
 285 gustometer is designed to spray raw taste solutions on
 286 the extended tongue; however, we delivered solutions
 287 directly in the mouth to produce the retronasal

perception in the present study. To avoid movements that could cause artifacts during EEG recordings, excess solution and saliva were slowly and continuously withdrawn from the mouth, and participants did not need to swallow. A spray head was hand manufactured specifically for each participant to fit the mouth cavity (modification of the Burghart spray head and fixation on a piece of a hygoformic® Orsing tubing). The solutions were sprayed as a thin mist over a large part of the tongue, permitting homogeneous taste perception over the tongue and odorant diffusion. It also reduces somatosensory and motor artifacts that may occur when the participant has to drink from a glass or even when drops are delivered on the tongue. Furthermore, a salivary ejector head (hygoformic® Orsing) connected to a peristaltic pump permitted removing excess solution and saliva from the mouth to minimize swallowing artifacts during EEG recordings. Participants kept both tubes (external diameter: 4 mm) in the mouth by gently clenching the teeth around them. Participants were asked not to swallow during the sessions.

To trigger as much as possible odor and taste perception in a synchronous way, we trained participants during a dedicated session to inhale through the mouth and exhale through the nose while maintaining an open velum throughout the recording session (free circulation of aromas to the nasal cavity). The combination of in-mouth stimulation for retronasal perception and the open-velum breathing favored the passage of a maximum of odorant volatiles into the nasal cavity. It allowed an increased synchronization between odor and taste perception compared to velum closed (Buettnner et al., 2002; Bonny et al., 2017). The participants were monitored with a breathing sensor (Burghart) coupled with an oscilloscope to receive feedback on their breathing during training. They had to maintain a constant amplitude and regular frequency of the sinusoid for 3 min. Finally, they were trained to maintain breathing while receiving in-mouth stimulations with each solution of interest (5 repeated stimulations for each solution).

EEG recording sessions

Participants started the EEG session by training again on the opened-velum breathing technique (3 min opened-velum breathing). EEG electrodes (Ag/AgCl) were fixed on the head with a conductive paste (EC2 electrode Cream, Natus®) after cleaning the skin (Everi, Spes medical®). Five electrodes were fixed on Fz, Cz, C3, C4, and Pz following a 10/20 system. Ground electrodes were positioned on the mastoids and reference electrodes on the ear lobes. One electrode was fixed above the right eyebrow to record vertical blinking artifacts. Impedance was kept below 30 kΩ. Participants were not notified of the stimulation. They received a flow of Evian water interrupted by air (400 ms water followed by 400 ms air) to limit the quantity of liquid in the mouth. Every 16 or 20 s, they received a target stimulation (400 ms); brain activity was recorded only for target stimulations (500 ms before and 1500 ms after the start of the stimulus). Air pressure

was fixed at 850 mbar. This design permitted fast habituation to somatosensory stimuli (Plattig, 1989). Participants were stimulated 40 times in a row for each of the 5 solutions. Every 20 pulses, a 2 minutes break was made for the participants to relax. Solutions were presented in a counterbalanced order. To maintain a relatively low level of attention during the task, participants had to perform a tracking task (keeping a dot inside a slowly moving square). The game and the sensory evaluation questionnaire were displayed on a monitor (GustOlf custom-made software, adapted from “Tracking performance” software, from the Smell and Taste Clinic Dresden). The tracking task stopped one second after the stimulation, and participants were prompted to rate saltiness intensity after each trial on an unstructured visual analog scale anchored on the left with “low intensity” (corresponding to 0) and on the right side of the scale with “high intensity” (corresponding to 100). To control for noise, participants were listening to “brown noise” through earphones. Brain activity was sampled at 1000 Hz using an EEG Burghart system (OL026) (analog high pass filter 1st order: 0.072 Hz, analog low-pass filter 3rd order: 186 Hz). The recording was triggered by the gustometer and started 500 ms before stimulation and ended 1500 ms after stimulation.

Data analysis

Normality and homogeneity of variances were checked with QQplot and residuals vs. fitted values plots. Furthermore, the homogeneities of variances were confirmed for all models tested (Levene test, $p > 0.35$).

Sensory data analysis

Saltiness intensity was rated for each of the forty stimulations delivered during the EEG recording (R software, R package: nlme (Pinheiro et al., 2015)). A first linear mixed model tested the effect of repetitions to measure habituation to solution in the course of the recording (*repetitions*solutions*, with *participants* as a random factor), no interaction between solutions and repetitions, nor the main effect of repetitions were significant ($p < 0.05$). Therefore, in a second model, the variability of ratings across *participants* was modeled as a random factor, and *repetitions* were nested within each participant, thus taking into account repeated measures, while *solutions* were modeled as fixed factors. Finally, post-hoc tests were performed, pairwise comparisons between solutions were computed with a Tukey test, and p -values were corrected for multiple testing with the false discovery rate (FDR, $p_{corr} < 0.05$).

EEG preprocessing

Data were preprocessed in Letswave (open-source MATLAB EEG signal processing toolbox NOCIONs, Institute of Neuroscience, Université Catholique de Louvain). They were first filtered (Butterworth, bandpass filter: 0.01–30 Hz, filter order 4) and baseline corrected [$y_i = x_i - \text{mean}(b_i)$] using the 500 ms recorded before the stimulation. Epochs contaminated by blinking artifacts

were removed (amplitude criteria $> 90 \mu\text{V}$). Finally, epochs with large alpha waves for most of the epoch duration were visually rejected. The mean number of epochs accepted was 21 ± 7 . After computing the mean across epochs for each participant, a Grand Mean was calculated by weighting each participant's mean by the number of epochs accepted. This weighting, suggested by Mouraux (2015), allowed taking into account the number of epochs included in the individual means to reduce the noise effect on the Grand-Mean and linear mixed models.

EEG RESULTS ANALYSIS

Peak amplitudes and latencies were measured with ERPLab implemented in the EEGlab toolbox from MATLAB (Lopez-Calderon and Luck, 2014). We started the definition of time windows from the literature using gustatory ERPs for P1, which appeared between 120–180 ms (Mizoguchi et al., 2002; Franken et al., 2011), and on olfactory or gustatory ERPs for P3, which appeared between 550–750 ms (Pause et al., 1996; Welge-Lüssen et al., 2005; 2009; Franken et al., 2011; Huart et al., 2012). We used relatively small windows (windows with a minimum of 100 ms and a maximum of 150 ms) to avoid overlaps between peaks. Local peak amplitude was searched in each time window within averages of 10 points on each local peak side. Local peak amplitude is defined as having a greater voltage than the average of the n number points on either side (Luck, 2014). When no peak was found, the NA value (non-applicable) was used. Time windows were checked and adapted to have as few as possible NA values on Cz, Fz, and Pz electrodes. Following these criteria, the P1 peak was analyzed within a 100–200 ms time window, and the P3 peak was analyzed within a 560–710 ms time window.

Before peak analysis, following Luck recommendations (Luck, 2014), data were filtered once more using a mild low-pass filter (half amplitude cutoff of 30 Hz, slope of 12 dB/octave). Finally, local peak amplitude and latency were measured for each participant at each electrode position. Data were analyzed with a linear mixed model (R package: nlme, lme function (Pinheiro et al., 2015)). The factor *participant* was modeled as a random factor, while solutions and electrodes were modeled as fixed factors. The interaction between solution and electrodes was also tested to highlight electrodes that may behave differently regarding solutions. The estimation of variance components followed the method of Restricted Maximum Likelihood (ReML) estimation. The variance of the response variables (amplitude or latency) measured on the peak of average ERP per participant and condition were weighted by the number of epochs accepted for each participant to reduce the impact of noisy mean, to correspond to grand-average ERP. Because no interactions between electrodes and solutions were found (cf. Results), electrodes that did not significantly differ were included in the models to highlight global brain responses and restrict the analysis to major effects. Only electrodes that did not vary from the elec-

trode with the highest amplitude were analyzed. Finally, the effects of electrodes (used as a repeated measure) and solutions were tested without the interaction effect. Pairwise, Tukey's tests for defined contrasts were computed in cases of significant fixed effects. The contrasts were PPS1B vs. PPS1, PPS1B vs. PPS2, PPS2 vs. PPS1, PP vs. PPB, salted solutions vs. unsalted solutions, solutions with beef stock aroma vs. solutions without beef stock aroma. Pearson correlation was tested between P3 amplitude and the saltiness intensity evaluated for all participants and all conditions. Linear effects and correlation were considered significant at $p < 0.05$. Pairwise comparisons were FDR corrected for multiple comparisons ($p_{\text{corr}} < 0.05$).

RESULTS

Sensory results

Intensity evaluations did not decrease in the course of the 40 repetitions ($F [39, 2548] = 0.19, p = 1$). Moreover, no interaction between solutions and repetitions ($F [156, 2388] = 0.74, p = 0.99$), nor main effect of repetitions ($F [156, 2388] = 0.46, p = 0.99$) was significant. These results highlighted the lack of habituation to the different solutions in the course of the 40 repetitions.

Saltiness intensity differed significantly between solutions ($F [4, 2076] = 939.21, p < 0.0001$, sd random effects = 18.18). Sensory results revealed a small odor-induced saltiness enhancement (OISE) by the “beef stock” aroma in the salt-reduced green-pea soup (PPS1B). PPS1B (mean of intensity ratings of 40 repetitions across participants: $M = 63.60$, standard error of the mean: $SEM = 0.86$) was perceived as slightly more salty than PPS1 ($M = 60.53$, $SEM = 0.96$; $z = 2.72$, $p_{\text{corr}} = 0.007$, estimate = 3.06, sd = 1.13.), although not as salty as PPS2 ($M = 72.80$, $SEM = 0.81$; $z = 8.16$, $p_{\text{corr}} < 0.0001$, estimate = 9.20, sd = 1.13). All salt-added solutions (PPS1, PPS1B and PPS2) were perceived as more salty than the control solutions (PP, $M = 17.87$, $SEM = 0.70$, and PPB, $M = 26.42$, $SEM = 1.05$) ($p_{\text{corr}} < 0.00001$). PPB was rated as more salty than PP ($z = 7.59$, $p_{\text{corr}} < 0.0001$, estimate = 8.56, sd = 1.13) which support the odor-induced salty taste of the odor since no salt was added in this sample.

EEG results

No interaction was found between *electrode position* and *solution* for any of the peaks (P1, P3) regarding amplitude or latency (all comparisons: $F [16, 282:288] < 1.06, p > 0.39$). Therefore, this interaction was never included in the models. Because no interactions occurred, electrodes that did not significantly differ were included in the models to highlight global brain responses and restrict the analysis to major effects. P3 and P1 peaks were analyzed to determine whether odor-induced saltiness enhancement could be observed in early and/or later brain processing (Fig. 1).




Fig 1. Event-related potentials for all solutions and all electrodes. The curves are Grand Mean weighted by the number of epochs finally accepted for each participant (amplitude (μV) as a function of time (ms) for each solution). The solutions were green-pea soup (PP), PP with beef stock aroma (PPB), PP with a reduced level of salt (3.125 g/L) (PPS1), PPS1 with beef stock aroma (PPS1B) and PP with a usual level of salt (6.25 g/L) (PPS2). The five recorded electrodes are shown (Cz, C3, C4, Fz and Pz). To improve graphical representation of ERPs, grand means were filtered with a Butterworth low-pass filter at 20 Hz and baseline was corrected.

P1 peak

There was a significant effect of channel on amplitude ($F [4, 298] = 4.52, p = 0.002$). Fz presented the highest P1 mean amplitude ($M = 3.98, \text{SEM} = 0.33$) and did not significantly differ from Cz ($z = 1.45, \text{pcorr} = 0.26$) but differed from Pz, C4 and C3 (for all comparisons: $p < 0.02$). Therefore, Fz and Cz were included as repeated measures in the following analyses.

There was no significant difference in the amplitude of P1 between solutions ($F [4, 98] = 1.22, p = 0.31, \text{sd random effects} = 10.14$) but the latency of P1 differed ($F[4, 98] = 3.31, p = 0.01, \text{sd random effects} = 108.6029$) (Fig. 2). P1 for PPS2 ($M = 143 \pm 5 \text{ ms}$) appeared earlier than PPS1B ($M = 164 \pm 4 \text{ ms}$) and PPB ($M = 163 \pm 5 \text{ ms}$). Indeed, P1 for PPS2 appeared respectively 21 ms ($z = -3.22, p = 0.01, \text{estimate} = 21, \text{sd} = 6.45$) and 20 ms ($z = -3.05, p = 0.01, \text{estimate} = 20, \text{sd} = 6.44$) earlier than PPS1B and PPB, which means that P1 for PPS2 appeared earlier than the two “aromatic” solutions (beef stock odor). Latency did not differ between PPS1B, PPS1, PPB and PP ($p > 0.13, \text{estimates} < 13.6, \text{sd} < 6.99$).

P3 peak

There was a significant effect of electrode position on the amplitude of P3 peak ($F [4, 287] = 2.45, p = 0.05$). Cz presented the highest mean amplitude of P3 ($M = 5.03, \text{SEM} = 0.53$) and did not significantly differ from the Fz, C4 and Pz electrodes ($p > 0.75$) but differed from C3 ($z = 2.77, p = 0.05$). Therefore, Cz, Fz, C4, and Pz were included as repeated measures in the analysis.

A significant difference was found on the amplitude of P3 peak between solutions ($F [4, 204] = 18.52, p < 0.0001, \text{sd random effects} = 12.76$). PPS1, PPS1B and PPS2 induced significantly larger P3 peaks than PP and PPB (contrast between unsalted and salted solutions, $z = 8.443, \text{pcorr} < 0.0001, \text{estimate} = 9.02, \text{sd} = 1.07$, Fig. 3A). A weak but significant positive correlation was found between saltiness intensity ratings and P3 peak amplitude ($z = 3.05, p = 0.002, \text{tau} = 0.26$) (Fig. 4). The P3 peak amplitudes for PP and PPB did not differ from a null amplitude ($\text{pcorr} > 0.39, \text{estimates} < 0.16, \text{sd} < 0.55$). There was no difference of P3 amplitude between PPS1, PPS1B and PPS2 ($\text{pcorr} > 0.30, \text{estimates} < 0.69, \text{sd} < 0.56$).

There was a significant difference on latency of P3 between the tested solutions ($F[2, 102] = 9.05, p = 0.0003, \text{sd random effects} = 152.85$). The P3 peak appeared later for PPS1B ($M = 662 \pm 6 \text{ ms}$) compared to PPS1 ($M = 637 \pm 5 \text{ ms}, z = 3.93, \text{pcorr} = 0.0003, \text{estimate} = 26.3, \text{sd} = 6.70$) and to PPS2 ($M = 642 \pm 4 \text{ ms}, z = -3.24, \text{pcorr} = 0.002, \text{estimate} = 20.58, \text{sd} = 6.36$) (Fig. 3B). In summary, the P3 peak was significantly different from zero amplitude only for the solutions PPS1, PPS1B and PPS2. The P3 peak, in response to PPS1B, was delayed of 25 ms and 20 ms compared with PPS1 and PPS2 respectively. Finally, the amplitude of the P3 peak increased as a function of the saltiness intensity rated.

518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576




Fig 2. P1 peak amplitude and latency. Weighted mean P1 amplitudes (A) and latencies (B) (\pm CI95%) for electrodes Fz and Cz for each solution: green-pea soup (PP), PP with beef stock aroma (PPB), PP with a reduced level of salt (3.125 g/L) (PPS1), PPS1 with beef stock aroma (PPS1B) and PP with a usual level of salt (6.25 g/L) (PPS2).




Fig 3. P3 peak amplitude and latency. Weighted mean P3 peak amplitudes (A) and latencies (B) (\pm CI95%) for electrodes Fz, Cz, C4, and Pz for each solution: green-pea soup (PP), PP with beef stock aroma (PPB), PP with a reduced level of salt (3.125 g/L) (PPS1), PPS1 with beef stock aroma (PPS1B) and PP with a usual level of salt (6.25 g/L) (PPS2). A P3 peak was not found for PP and PPB. Therefore, mean latencies were calculated only for PPS1B, PPS1, and PPS2.

DISCUSSION

The objective of the study was to highlight the brain chronometry of odor–taste integration using OISE. Our hypothesis, based on the classical view of odor–taste integration, was that differences of amplitude and/or latencies between solutions with and without OISE would be observed on the late P3 peak of olfactory-gustatory ERP, but no difference would appear on the early P1 peak. The sensory results showed a significant OISE using the “beef stock” aroma (PPS1B vs. PPS1) in the green pea soup. The ERP results showed an increased latency on the P3 peak with the same solutions (PPS1B vs. PPS1). No difference in amplitude or latency was observed on the P1 peak between PPS1B and PPS1. Therefore, our results support the hypothesis of late brain integration in the high cognitive areas as proposed in the classical view of flavor perception (Verhagen and Engelen, 2006; Verhagen, 2007; Small, 2008; Prescott, 2012).

The ERPs measured in our study mainly reflect the gustatory component (specifically the saltiness processing) of the solutions, and its modulation by the olfactory component. Indeed the grand-averages showed ERPs with proper peaks, well-differentiated from noise, for the salted solutions and small ones for the control solutions apart from the P1 peak (Fig. 2). These control solutions (PP and PPB) were mainly odorant and had a poor taste. Furthermore, as the P3 peak correlated to the saltiness intensity (Fig. 4), it also supports that the ERPs represent mainly the gustatory brain activations and their modulation by the olfactory component. Therefore, any effect observed on P1 or P3 peaks will be either explained by the concentrations of salt or by the modulation of the gustatory processing by the olfactory one (i.e., odor-induced saltiness enhancement). This modulation

577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637




Fig 4. Correlation between the mean saltiness intensity ratings and the amplitude (μV) of the P3 peak at the Cz electrode. Each data point is the evaluation of one solution evaluated 40 times by one participant (13 participants and 5 solutions). The amplitude of the P3 peak increased as a function of the saltiness intensity. Kendall correlation: $z = 3.05$, $p = 0.002$, $\tau = 0.26$.

638 can either intervene early in the processing by direct
639 interactions between primary olfactory and gustatory
640 areas (emerging view) or occur later in the processing
641 through top-down pathways (classical view). Because
642 the modulation observed appeared on the P3 peak and
643 not on P1, the results confirmed OISE's classical view.

644 We observed a higher salty taste intensity in PPB
645 compared to PP. However, the effect in PPB likely
646 implied an odor-induced saltiness perception (OISP),
647 but no enhancement (OISE) as salt was not present in
648 the solution. Because gustatory responses mainly drive
649 the ERPs here, although we observed a perceptual
650 difference between PP and PPB (OISP), we did not
651 observe brain correlates of such sensory response. It
652 could be that OISP and OISE involve different brain
653 circuits or that the number of neurons involved by the
654 OISP may not be sufficient to be visible in ERP, on the
655 contrary to OISE. The sensory effects may seem
656 contradictory, but the low range of salt concentration
657 may explain this result. The OISE effect is statistically
658 significant in the PPS1B solution, but the OISE
659 appeared small and lower than the OISP (Fig. 1). The
660 enhancement of salty taste induced by an odor is
661 usually observed in the range of 10 to 30% (Lawrence
662 et al., 2009). In the present study, the percentage of
663 enhancement (5%) might have been flattened because
664 of the evaluation of no-salt-added samples (PP and
665 PPB) in the sample set. Indeed, these two samples' salti-
666 ness intensity was likely low compared to the three other
667 salt-containing samples, which could have reduced the
668 discrimination between PPS1, PPS1B, and PPS2. This

669 hypothesis is supported by the
670 unexpected small difference
671 between PPS1 and PPS2 that
672 contained double salt.

673 One could think that rating
674 saltiness intensity only at the end
675 of each epoch recording would
676 involve a dumping effect. A
677 dumping effect occurs when "a
678 salient attribute is not included
679 on a ballot; the opinion about
680 that aspect of the product may
681 then be displaced onto another,
682 sometimes inappropriate scale.
683 In other words, if consumers are
684 not given a rating scale in which
685 they can voice some important
686 opinion about a product, they will
687 'dump' this perception onto some
688 other available rating scale or
689 question" (Clark and Lawless,
690 1994). Dumping is usually consid-
691 ered as a bias in sensory rating
692 because it could lead to perceived
693 intensity overrating. However,
694 several recent studies reconsider
695 the dumping effect not as bias
696 but as a proof of odor–taste inte-
697 gration, because flavor perception
698 is by nature configural (Prescott,
699 2012; Onuma et al., 2018). There-

700 fore, when participants have no clue about the elements
701 (odor and taste), the configuration (flavor) is attended,
702 and the OITE occurs. The configural perception is a typi-
703 cal integratory perception described in the significant
704 contribution of the "unique cue theory" (Rescorla, 1972,
705 1973). This theory was further developed by Pearce
706 (2002) and finally demonstrated in the context of odor-
707 odor and odor–taste configural perceptions (Le Berre
708 et al., 2008, 2010; Sinding et al., 2011; White et al., 2020).

709 P1 peak

710 P1 peak, in chemosensory studies (70–302 ms), is
711 associated to brain circuits responding to exogenous
712 properties of the food such as tastant concentration
713 (Funakoshi and Kawamura, 1971; Kobayakawa et al.,
714 1999, 2007; Mizoguchi et al., 2002; Ohla et al., 2010).
715 Here, we did not find early changes in the brain process-
716 ing of PPS1B compared to PPS1 when considering ampli-
717 tude or latencies of the P1 peak. Therefore, no effect
718 possibly linked to OISE could be observed on the P1 peak
719 amplitude. We found a delay on the P1 peak, between
720 PPS2 vs. PPS1B and PPB. These contrasts could not
721 account for the OISE phenomenon, which would suppose
722 a difference between PPS1 and PPS1B. Both PPS1B and
723 PPB solutions contained a "beef stock aroma" and were
724 delayed compared to the most salted solution PPS2. This
725 polarization of the two types of solutions (high salt solution
726 vs. high odorant solutions) regarding their P1 latencies
727 could highlight the differential processing between salient
728 gustatory solution (PPS2) and salient olfactory solutions

(PPB and PPS1B). In the literature, it is reported that olfactory ERPs have a later P1 peak than gustatory ERP. In studies that showed a P1 peak in response to olfactory stimulation, this peak appeared between 200–280 ms (Tateyama et al., 1998; Iannilli et al., 2013), while it appeared at 120–140 ms for gustatory stimulation (Mizoguchi et al., 2002; Ohla et al., 2009, 2010; Jacquin-Piques et al., 2015).

Significant differences in P1 amplitude regarding salt concentration in PP, PPS1, and PPS2 were not found, although these solutions were perceived as significantly different. This result is inconsistent with Kobayakawa and colleagues (Kobayakawa et al., 1996, 1999, 2008). It is generally assumed that P1 amplitude is related to tastant concentration. Three studies from two independent labs showed that P1 latency is more likely to reflect a change in tastant concentration, with lower latencies for higher concentrations (Funakoshi and Kawamura, 1971; Kobal, 1985; Tateyama et al., 1998). In our study, the solution with the highest concentration of salt (PPS2) was processed with the lowest latency, but the result is not significant. Altogether, these results could be explained by the concentrations of salt used in our study, which are not sufficiently extreme to show the expected latency difference. Studies that tackle the brain peaks associated with the perception of tastant concentrations use steps of concentrations 5 times larger than those used in our study. In Kobayakawa et al. (2008), differences of P1 amplitude were observed between 100 mM (5.84 g/L) and 1 M (58.4 g/L), or between 30 mM (1.752 g/L) and 300 (17.52 g/L), but not between 100 mM and 300 mM. In our study, salt concentrations were in line with a usual soup that is 6.25 g/L for PPS2 and 3 g/L for PPS1 (soup with reduced-salt level). Likely, the low, but food relevant, salt concentrations used in our study could explain why we did not observe significant differences in either amplitude or latency for P1 peak as a function of salt levels.

767 **P3 latency might be a marker of odor-induced taste** 768 **enhancement**

769 P3 peak might be linked to cognitive processing diversity,
770 including emotions integration, attention allocation, and
771 working memory, which have in common their endogenous
772 origin. Furthermore, the P3 peak is elicited
773 by multiple intracerebral generators revealing its
774 integratory component (Picton, 1992; Li et al., 2015). In
775 our study, the P3 peak was significantly delayed by
776 20 ms or more in response to PPS1B compared to
777 PPS1 and PPS2. Moreover, the brain processing of
778 PPB did not present a significant P3 peak; therefore, we
779 could exclude an impact of the odor component on P3,
780 and we could then directly compare PPS1B to PPS1
781 and PPS2. We can also exclude an earlier origin of this
782 delay, as no significant delay was present at the early
783 stages of brain processing (only 7 ms separated P1 peaks
784 for PPS1B compared with PPS1). Although the interpreta-
785 tion of latencies, in terms of neuronal activity, is challeng-
786 ing, we might consider this delay as evidence for a higher
787 number of synapses involved in processing the sensory
788 information for solutions presenting an odor-induced taste

enhancement (for review, see Woodman, 2010). This
result would comply with the classical view of flavor inte-
gration proposed by Verhagen (Verhagen and Engelen,
2006; Verhagen, 2007). In this theory, inspired by visual
studies, downstream areas activate heteromodal integra-
tion areas that then loop back to the primary sensory
areas to form a refined activation pattern. The activation
of these associative areas, and the back projections to
primary gustatory regions, may explain the higher latency
observed for the odor-induced taste enhancement solu-
tion. Heteromodal multisensory processing areas, such
as the superior temporal sulcus (Calvert, 2001; Calvert
and Thesen, 2004), were found to be activated between
518–730 ms after food odor stimulation, with magnetoen-
cephalography (MEG) (Kettenmann et al., 1997). This
timing is consistent with the latency of the P3 peak iden-
tified in the present study.

806 **P3 peak amplitude might be a marker of conscious** 807 **perception of the saltiness intensity**

808 An intriguing result is that the amplitude of the P3 peak
809 was significantly different from the null amplitude only in
810 solutions with added-salt. Furthermore, looking at
811 individual results, we found a small but significant
812 correlation of this peak amplitude with salty taste
813 intensity ratings. Several explanations may account for
814 these results and may rely on different cognitive steps
815 involved in the intensity rating task. These steps may be
816 decomposed: i) non-conscious salt concentration
817 processing, ii) conscious representation of saltiness
818 intensity, and iii) evaluation of/decision on saltiness
819 intensity. The difference in P3 amplitude did not
820 correspond to the first step, as we did not find a
821 significant difference in amplitude between PPS1 and
822 PPS2. These results could be linked to the evaluation/
823 decision on saltiness intensity rating, which occurred at
824 least 320 ms after the P3 peak. As the participants were
825 likely expecting to evaluate the saltiness intensity after
826 each stimulation, the P3 peak could reflect their
827 anticipatory evaluation. However, in this case, P3 would
828 have also been observed for PP and PPB solutions.
829 Finally, suppose these results are neither linked to “the
830 non-conscious salt concentration representation” nor “to
831 the evaluation/decision on the saltiness intensity”. In
832 that case, it could be an intermediate state: “a
833 conscious representation of the saltiness intensity”.
834 Notably, the expectancy of the intensity scale could
835 favor the conscious representation of the saltiness
836 intensity. This late conscious perception has been
837 shown in visual attentional blink studies (Sergent et al.,
838 2005). This task permitted us to decipher the peaks
839 involved in non-conscious processing versus conscious
840 one. Although such results were not yet shown in taste
841 sensory modality, we suggest that P3 might represent
842 the conscious perception of the saltiness intensity.

843 **The classical view of odor–taste integration** 844 **supported by our results**

845 Overall, our results support the odor–taste integration
846 theoretical framework proposed by Verhagen (Verhagen

and Engelen, 2006; Verhagen, 2007), Prescott (2012), and Small (Small, 2008). In this framework, the integration of odor and taste occurs in high-level brain areas (OFC, perirhinal, and dorsal mid insula), as brain correlations of odor-induced saltiness enhancement were found only at the later stages of brain processing. More recently, Small and colleagues (Small et al., 2013) proposed an emerging model of odor–taste integration relying on animal studies and fMRI human studies (De Araujo et al., 2003; Maier et al., 2012, 2015; Seubert et al., 2015; Maier, 2017). They showed early interactions between primary olfactory and gustatory regions. They proposed: i) a densely connected system between olfactory and gustatory areas and ii) that flavor might be already integrated into primary chemosensory cortices. Although our study did not refute the dense connections between primary olfactory and gustatory cortices, it did not show early modulation in response to odor and taste presented together. However, we cannot completely exclude that our EEG design was not powered enough to detect early changes.

Our results provide the first insight into the brain chronometry of odor–taste integration, focusing on salty flavor. We found that olfactory-gustatory interactions mainly occur in the late brain processing of sensory information carried by a close-to-real food solution. We have developed an adequate stimulation method to understand the chronometry of odor–taste integration in the flavor system. Other converging results on similar questions would be necessary to comprehend flavor perception and underlying brain mechanisms further. The enhancing effect was small in our solution, likely due to the type of evaluation (intensity scales) and context (during EEG recording). Therefore, our results' reproducibility should be tested with a more reliable sensory evaluation method and several food models. Due to the long recording time, we could neither test several solutions presenting an odor-induced taste enhancement nor control solutions without such an effect. A study comparing enhancing and non-enhancing aroma would be of interest.

UNCITED REFERENCES

Geisler and Murphy (2000), Kobal et al. (1990), Lascano et al. (2010), Livermore et al. (1992), Lundstrom et al. (2006), Murphy et al. (2000), Murphy et al. (1998), Pause and Krauel (2000), , Spence et al. (2015), Stevenson et al. (2011), Thesen and Murphy (2002).

ACKNOWLEDGMENTS

This work was supported by a French INRA grant (ANS-COGITO). We would like to thank Jacques Maratray for setting up the GustOlf sensory evaluation software.

REFERENCES

Barba C, Beno N, Guichard E, Thomas-Danguin T (2018) Selecting odorant compounds to enhance sweet flavor perception by gas chromatography/olfactometry-associated taste (GC/O-AT). *Food*

- Chem 257:172–181. <https://doi.org/10.1016/j.foodchem.2018.02.152>. 902
903
- Bartoshuk LM (1974) NaCl thresholds in man: Thresholds for water taste or NaCl taste? *J Comp Physiol Psychol* 87:310–325. 904
905
- Bonny JM, Sinding C, Thomas-Danguin T (2017). In: modern magnetic resonance. Cham: Springer International Publishing. 906
p. 1–20. DOI: 10.1007/978-3-319-28275-6_132-1. 907
908
- Buettner A, Beer A, Hannig C, Settles M, Schieberle P (2002) Physiological and analytical studies on flavor perception dynamics as induced by the eating and swallowing process. *Food Qual Prefer* 13(7-8):497–504. [https://doi.org/10.1016/S0950-3293\(02\)00052-6](https://doi.org/10.1016/S0950-3293(02)00052-6). 909
910
911
912
913
- Calvert GA (2001) Crossmodal processing in the human brain: insights from functional neuroimaging studies. *Cereb Cortex* 11:1110–1123. <https://doi.org/10.1093/cercor/11.12.1110>. 914
915
- Calvert GA, Thesen T (2004) Multisensory integration: methodological approaches and emerging principles in the human brain. *Journal of Physiology-Paris* 98(1-3):191–205. <https://doi.org/10.1016/j.jphysparis.2004.03.018>. 916
917
918
919
920
- Carbone KA, Rodeback R, Modersitzki E, Miner M, LeCheminant JD, Larson MJ (2018) The utility of event-related potentials (ERPs) in understanding food-related cognition: A systematic review and recommendations. *Appetite* 128:58–78. <https://doi.org/10.1016/j.appet.2018.05.135>. 921
922
923
924
925
- Clark CC, Lawless HT (1994) Limiting response alternatives in time-intensity scaling: an examination of the halo-dumping effect. *Chem Senses* 19(6):583–594. <https://doi.org/10.1093/chemse/19.6.583>. 926
927
928
929
- de Araujo IET, Rolls ET, Kringelbach ML, McGlone F, Phillips N (2003) Taste-olfactory convergence, and the representation of the pleasantness of flavour, in the human brain. *Eur J Neurosci* 18(7):2059–2068. <https://doi.org/10.1046/j.1460-9568.2003.02915.x>. 930
931
932
933
934
- Djordjevic J, Zatorre RJ, Jones-Gotman M (2004) Odor-induced changes in taste perception. *Exp Brain Res* 159(3):405–408. <https://doi.org/10.1007/s00221-004-2103-y>. 935
936
937
- Frank RA, Byram J (1988) Taste–smell interactions are tastant and odorant dependent. *Chem Senses* 13(3):445–455. <https://doi.org/10.1093/chemse/13.3.445>. 938
939
940
- Franken IHA, Huijding J, Nijs IMT, van Strien JW (2011) Electrophysiology of appetitive taste and appetitive taste conditioning in humans. *Biol Psychol* 86(3):273–278. <https://doi.org/10.1016/j.biopsycho.2010.12.008>. 941
942
943
944
- Funakoshi M, Kawamura Y (1971) Summated cerebral evoked responses to taste stimuli in man. *Electroencephalogr Clin Neurophysiol* 30(3):205–209. [https://doi.org/10.1016/0013-4694\(71\)90055-1](https://doi.org/10.1016/0013-4694(71)90055-1). 945
946
947
948
- Geisler MW, Murphy C (2000) Event-related brain potentials to attended and ignored olfactory and trigeminal stimuli. *Int J Psychophysiol* 37(3):309–315. [https://doi.org/10.1016/S0167-8760\(00\)00111-2](https://doi.org/10.1016/S0167-8760(00)00111-2). 949
950
951
952
- Huart C, Legrain V, Hummel T, Rombaux P, Mouraux A (et) Time-frequency analysis of chemosensory event-related potentials to characterize the cortical representation of odors in humans. *PLoS ONE* 7:e33221. <https://doi.org/10.1371/journal.pone.0033221>. 953
954
955
956
- Iannilli E, Wiens S, Arshamian A, Seo H-S (2013) A spatiotemporal comparison between olfactory and trigeminal event-related potentials. *NeuroImage* 77:254–261. <https://doi.org/10.1016/j.neuroimage.2012.12.057>. 957
958
959
960
- Jacquin-Piques A, Gaudillat S, Mouillot T, Gigot V, Meillon S, Leloup C, Penicaud L, Brondel L (2015) Prandial states modify the reactivity of the gustatory cortex using gustatory evoked potentials in humans. *Front Neurosci* 9:490. <https://doi.org/10.3389/fnins.2015.00490>. 961
962
963
964
965
- Kettenmann B, Hummel C, Stefan H, Kobal G (1997) Multiple olfactory activity in the human neocortex identified by magnetic source imaging. *Chem Senses* 22(5):493–502. <https://doi.org/10.1093/chemse/22.5.493>. 966
967
968
969
- Kobal G (1985) Gustatory evoked potentials in man. *Electroencephalogr Clin Neurophysiol/Evoked Poten Sect* 62(6):449–454. [https://doi.org/10.1016/0168-5597\(85\)90055-3](https://doi.org/10.1016/0168-5597(85)90055-3). 970
971
972

- 973 Kobal G, Hummel C, Nuernberg B, Brune K (1990) Effects of
974 pentazocine and acetylsalicylic acid on pain-rating, pain-related
975 evoked potentials and vigilance in relationship to pharmacokinetic
976 parameters. *Agents Actions* 29:342–359.
- 977 Kobayakawa Ko, Kobayakawa R, Matsumoto H, Oka Y, Imai T, Ikawa
978 M, et al. (2007) Innate versus learned odour processing in the
979 mouse olfactory bulb. *Nature* 450(7169):503–508. <https://doi.org/10.1038/nature06281>.
- 980
981 Kobayakawa T, Endo H, Ayabe-Kanamura S, Kumagai T, Yamaguchi
982 Y, Kikuchi Y, Takeda T, Saito S, Ogawa H (1996) The primary
983 gustatory area in human cerebral cortex studied by
984 magnetoencephalography. *Neurosci Lett* 212(3):155–158.
985 [https://doi.org/10.1016/0304-3940\(96\)12798-1](https://doi.org/10.1016/0304-3940(96)12798-1).
- 986 Kobayakawa T, Ogawa H, Kaneda H, Ayabe-Kanamura S, Saito S,
987 Endo H, Saito S (1999) Spatio-temporal analysis of cortical
988 activity evoked by gustatory stimulation in humans. *Chem.*
989 *Senses* 24:201–209. <https://doi.org/10.1093/chemse/24.2.201>.
- 990 Kobayakawa T, Saito S, Gotow N, Ogawa H (2008) Representation
991 of salty taste stimulus concentrations in the primary gustatory
992 area in humans. *Chem Percept* 1(4):227–234. <https://doi.org/10.1007/s12078-008-9030-4>.
- 993
994 Labbe D, Damevin L, Vaccher C, Morgenege C, Martin N (2006)
995 Modulation of perceived taste by olfaction in familiar and
996 unfamiliar beverages. *Food Quality Preference* 17(7-
997 8):582–589. <https://doi.org/10.1016/j.foodqual.2006.04.006>.
- 998 Lascano AM, Hummel T, Lacroix J-S, Landis BN, Michel CM (2010)
999 Spatio-temporal dynamics of olfactory processing in the human
1000 brain: an event-related source imaging study. *Neuroscience* 167
1001 (3):700–708. <https://doi.org/10.1016/j.neuroscience.2010.02.013>.
- 1002 Lawrence G, Salles C, Septier C, Busch J, Thomas-Danguin T (2009)
1003 Odour-taste interactions: A way to enhance saltiness in low-salt
1004 content solutions. *Food Quality Preference* 20(3):241–248.
1005 <https://doi.org/10.1016/j.foodqual.2008.10.004>.
- 1006 Le Berre E, Jarmuzek E, Béno N, Etiévant P, Prescott J, Thomas-
1007 Danguin T (2010) Learning influences the perception of odour
1008 mixtures. *Chem. Percept.* 3(3-4):156–166. <https://doi.org/10.1007/s12078-010-9076-y>.
- 1009
1010 Le Berre E, Thomas-Danguin T, Béno N, Coureaud G, Etiévant P,
1011 Prescott J (2008) Perceptual processing strategy and exposure
1012 influence the perception of odour mixtures. *Chem Senses*
1013 33:193–199. <https://doi.org/10.1093/chemse/bjm080>.
- 1014 Li F, Liu T, Wang F, Li He, Gong D, Zhang R, et al. (2015)
1015 Relationships between the resting-state network and the P3:
1016 Evidence from a scalp EEG study. *Sci Rep* 5(1). <https://doi.org/10.1038/srep15129>.
- 1017
1018 Lim J, Fujimaru T, Linscott TD (2014) The role of congruency in
1019 taste-odor interactions. *Food Quality Preference* 34:5–13. <https://doi.org/10.1016/j.foodqual.2013.12.002>.
- 1020
1021 Lim J, Johnson MB (2011) Potential mechanisms of retronasal odor
1022 referral to the mouth. *Chem Senses* 36(3):283–289. <https://doi.org/10.1093/chemse/bjq125>.
- 1023
1024 Lim J, Johnson MB (2012) The role of congruency in retronasal odor
1025 referral to the mouth. *Chem Senses* 37(6):515–522. <https://doi.org/10.1093/chemse/bjs003>.
- 1026
1027 Linscott TD, Lim J (2016) Retronasal odor enhancement by salty and
1028 umami tastes. *Food Quality Preference* 48:1–10. <https://doi.org/10.1016/j.foodqual.2015.08.004>.
- 1029
1030 Livermore A, Hummel T, Kobal G (1992) Chemosensory event-
1031 related potentials in the investigation of interactions between the
1032 olfactory and the somatosensory (trigeminal) systems.
1033 *Electroencephalogr Clin Neurophysiol* 83(3):201–210. [https://doi.org/10.1016/0013-4694\(92\)90145-8](https://doi.org/10.1016/0013-4694(92)90145-8).
- 1034
1035 Lopez-Calderon J, Luck SJ (2014) ERPLAB: an open-source toolbox
1036 for the analysis of event-related potentials. *Front Human Neurosci*
1037 8:213. <https://doi.org/10.3389/fnhum.2014.00213>.
- 1038
1039 Luck SJ (2014) An introduction to the event-related potential
1040 technique (Gazzaniga MS, ed). Massachusetts: The MIT Press.
- 1041
1042 Lundstrom JN, Seven S, Olsson MJ, Schaal B, Hummel T (2006)
1043 Olfactory event-related potentials reflect individual differences in
1044 odor valence perception. *Chem Senses* 31(8):705–711. <https://doi.org/10.1093/chemse/bjl012>.
- 1045
1046 Maier J, Blankenship M, Li J, Katz D (2015) A multisensory network
1047 for olfactory processing. *Curr Biol* 25(20):2642–2650. <https://doi.org/10.1016/j.cub.2015.08.060>.
- 1048
1049 Maier JX (2017) Single-neuron responses to intraoral delivery of odor
1050 solutions in primary olfactory and gustatory cortex. *J Neurophysiol*
1051 117(3):1293–1304. <https://doi.org/10.1152/jn.00802.2016>.
- 1052
1053 Maier JX, Wachowiak M, Katz DB (2012) Chemosensory
1054 convergence on primary olfactory cortex. *J Neurosci* 32
1055 (48):17037–17047. <https://doi.org/10.1523/JNEUROSCI.3540-12.2012>.
- 1056
1057 Marshall K, Laing DG, Jinks AL, Hutchinson I (2006) The capacity of
1058 humans to identify components in complex odor-taste mixtures.
1059 *Chem Senses* 31:539–545. <https://doi.org/10.1093/chemse/bji058>.
- 1060
1061 Masaoka Y, Satoh H, Akai L, Homma I (2010) Expiration: The
1062 moment we experience retronasal olfaction in flavor. *Neurosci*
1063 *Lett* 473(2):92–96. <https://doi.org/10.1016/j.neulet.2010.02.024>.
- 1064
1065 Mizoguchi C, Kobayakawa T, Saito S, Ogawa H (2002) Gustatory
1066 evoked cortical activity in humans studied by simultaneous EEG
1067 and MEG recording. *Chem Senses* 27:629–634. <https://doi.org/10.1093/chemse/27.7.629>.
- 1068
1069 Mouraux A (2015) Average. *NOCIONS/letswave6:1*.
- 1070
1071 Mroczko-Wąsowicz A (2016) The unity and complexity of flavour
1072 perception. In: *Proceedings of the 1st workshop on multi-sensorial*
1073 *approaches to human-food interaction - MHFI '16*. New York, New
1074 York, USA: ACM Press. p. 1–4. DOI: 10.1145/3007577.3007583.
- 1075
1076 Murphy C, Morgan CD, Geisler MW, Wetter S, Covington JW,
1077 Madowitz MD, Nordin S, Polich JM (2000) Olfactory event-related
1078 potentials and aging: normative data. *Int J Psychophysiol* 36
1079 (2):133–145. [https://doi.org/10.1016/S0167-8760\(99\)00107-5](https://doi.org/10.1016/S0167-8760(99)00107-5).
- 1080
1081 Murphy C, Wetter S, Morgan CD, Ellison DW, Geisler MW (1998) Age
1082 effects on central nervous system activity reflected in the olfactory
1083 event-related potential: evidence for decline in middle age. *Annals*
1084 *NY Acad Sci* 855:598–607. <https://doi.org/10.1111/j.1749-6632.1998.tb10630.x>.
- 1085
1086 Nasri N, Septier C, Beno N, Salles C, Thomas-Danguin T (2013)
1087 Enhancing salty taste through odour-taste-taste interactions:
1088 Influence of odour intensity and salty tastants' nature. *Food*
1089 *Quality Preference* 28(1):134–140. <https://doi.org/10.1016/j.foodqual.2012.07.004>.
- 1090
1091 Ohla K, Hudry J, le Coutre J (2009) The cortical chronometry of
1092 electrogustatory event-related potentials. *Brain Topogr* 22
1093 (2):73–82. <https://doi.org/10.1007/s10548-009-0076-7>.
- 1094
1095 Ohla K, Toepel U, le Coutre J, Hudry J (2010) Electrical neuroimaging
1096 reveals intensity-dependent activation of human cortical gustatory
1097 and somatosensory areas by electric taste. *Biolo Psychol* 85
1098 (3):446–455. <https://doi.org/10.1016/j.biopsycho.2010.09.007>.
- 1099
1100 Onuma T, Maruyama H, Sakai N (2018) Enhancement of saltiness
1101 perception by monosodium glutamate taste and soy sauce odor: a
1102 near-infrared spectroscopy study. *Chem. Senses* 43:151–167.
1103 <https://doi.org/10.1093/chemse/bix084>.
- 1104
1105 Pause BM, Krauel K (2000) Chemosensory event-related potentials
1106 (CSERP) as a key to the psychology of odors. *Int J Psychophysiol*
1107 36(2):105–122. [https://doi.org/10.1016/S0167-8760\(99\)00105-1](https://doi.org/10.1016/S0167-8760(99)00105-1).
- 1108
1109 Pause BM, Sojka B, Krauel K, Ferstl R (1996) The nature of the late
1110 positive complex within the olfactory event-related potential
1111 (OERP). *Psychophysiology* 33(4):376–384. <https://doi.org/10.1111/j.1469-8986.1996.tb01062.x>.
- 1112
1113 Pearce JM (2002) Evaluation and development of a connectionist
1114 theory of configural learning. *Anim Learn Behav* 30(2):73–95.
1115 <https://doi.org/10.3758/BF03192911>.
- 1116
1117 Picton TW (1992) The P300 wave of the human event-related
1118 potential. *J Clin Neurophysiol* 9(4):456–479. <https://doi.org/10.1097/00004691-199210000-00002>.
- 1119
1120 Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team (2015) nlme:
1121 Linear and Nonlinear Mixed Effects Models. R Packag version 31-
1122 120 <http://CRA>.
- 1123
1124 Plattig K-H (1989) Electrophysiology of taste and smell. *Clin Phys*
1125 *Physiol Measur* 10(2):91–125. <https://doi.org/10.1088/0143-0815/10/2/001>.

- 1114 Prescott J (1999) Flavour as a psychological construct: implications
1115 for perceiving and measuring the sensory qualities of foods. *Food*
1116 *Qual Prefer* 10(4-5):349–356. [https://doi.org/10.1016/S0950-](https://doi.org/10.1016/S0950-3293(98)00048-2)
1117 [3293\(98\)00048-2](https://doi.org/10.1016/S0950-3293(98)00048-2).
- 1118 Prescott J (2012) Multimodal chemosensory interactions and
1119 perception of flavor. In: Murray M, Wallace M, editors. *The*
1120 *neural bases of multisensory processes*. Boca Raton (FL): CRC
1121 Press/Taylor & Francis.
- 1122 Prescott J, Johnstone V, Francis J (2004) Odor-taste interactions:
1123 effects of attentional strategies during exposure. *Chem Senses*
1124 29:331–340. <https://doi.org/10.1093/chemse/bjh036>.
- 1125 Prescott J, Murphy S (2009) Inhibition of evaluative and perceptual
1126 odour–taste learning by attention to the stimulus elements.
1127 *Quarterly Journal of Experimental Psychology* 62
1128 (11):2133–2140. <https://doi.org/10.1080/17470210903031169>.
- 1129 Rescorla RA (1972) Configural conditioning in discrete-trial bar
1130 pressing. *J Comp Physiol Psychol* 79:307.
- 1131 Rescorla RA (1973) Evidence for “unique stimulus” account of
1132 configural conditioning. *J Comp Physiol Psychol* 85(2):331–338.
1133 <https://doi.org/10.1037/h0035046>.
- 1134 Rolls ET, Baylis LL (1994) Gustatory, olfactory, and visual
1135 convergence within the primate orbitofrontal cortex. *J Neurosci*
1136 14(9):5437–5452. [https://doi.org/10.1523/JNEUROSCI.14-09-](https://doi.org/10.1523/JNEUROSCI.14-09-05437.1994)
1137 [05437.1994](https://doi.org/10.1523/JNEUROSCI.14-09-05437.1994).
- 1138 Rolls ET (1997) Taste and olfactory processing in the brain and its
1139 relation to the control of eating. *Crit Rev Neurobiol* 11(4):263–287.
1140 <https://doi.org/10.1615/CritRevNeurobiol.v11.i4.20>.
- 1141 Sakai N, Kobayakawa T, Gotow N, Saito S, Imada S (2001)
1142 Enhancement of sweetness ratings of aspartame by a vanilla
1143 odor presented either by orthonasal or retronasal routes. *Percept*
1144 *Mot Skills* 92(3_suppl):1002–1008. [https://doi.org/10.2466/](https://doi.org/10.2466/pms.2001.92.3c.1002)
1145 [pms.2001.92.3c.1002](https://doi.org/10.2466/pms.2001.92.3c.1002).
- 1146 Schifferstein HNJ, Verlegh PWJ (1996) The role of congruency and
1147 pleasantness in odor-induced taste enhancement. *Acta Psychol*
1148 94(1):87–105. [https://doi.org/10.1016/0001-6918\(95\)00040-2](https://doi.org/10.1016/0001-6918(95)00040-2).
- 1149 Seo H-S, Iannilli E, Hummel C, Okazaki Y, Buschhüter D, Gerber J,
1150 Krammer GE, van Lengerich B, Hummel T (2013) A salty-
1151 congruent odor enhances saltiness: Functional magnetic
1152 resonance imaging study. *Hum Brain Mapp* 34(1):62–76. [https://](https://doi.org/10.1002/hbm.21414)
1153 doi.org/10.1002/hbm.21414.
- 1154 Sergent C, Baillet S, Dehaene S (2005) Timing of the brain events
1155 underlying access to consciousness during the attentional blink.
1156 *Nat Neurosci* 8(10):1391–1400. <https://doi.org/10.1038/nn1549>.
- 1157 Seubert J, Ohla K, Yokomukai Y, Kellermann T, Lundström JN (2015)
1158 Superadditive opercular activation to food flavor is mediated by
1159 enhanced temporal and limbic coupling: Food Flavor Enhances
1160 Temporal-Limbic Coupling. *Hum Brain Mapp* 36(5):1662–1676.
1161 <https://doi.org/10.1002/hbm.22728>.
- 1162 Sinding C, Thomas-Danguin T, Crepeaux G, Schaal B, Coureaud G
1163 (2011) Experience influences elemental and configural perception
1164 of certain binary odour mixtures in newborn rabbits. *J Exp Biol* 214
1165 (24):4171–4178. <https://doi.org/10.1242/jeb.063610>.
- 1166 Small DM (2008) Flavor and the formation of category-specific
1167 processing in olfaction. *Chem. Percept.* 1(2):136–146. [https://doi.](https://doi.org/10.1007/s12078-008-9015-3)
1168 [org/10.1007/s12078-008-9015-3](https://doi.org/10.1007/s12078-008-9015-3).
- 1169 Small DM (2012) Flavor is in the brain. *Physiol Behav* 107
1170 (4):540–552. <https://doi.org/10.1016/j.physbeh.2012.04.011>.
- 1171 Small DM, Prescott J (2005) Odor/taste integration and the
1172 perception of flavor. *Exp Brain Res* 166(3-4):345–357. [https://](https://doi.org/10.1007/s00221-005-2376-9)
1173 doi.org/10.1007/s00221-005-2376-9.
- 1174 Small DM, Veldhuizen MG, Green B (2013) Sensory neuroscience:
1175 Taste responses in primary olfactory cortex. *Curr Biol* 23:
1176 R157–R159. <https://doi.org/10.1016/j.cub.2012.12.036>.
- 1177 Spence C (2016) Oral referral: On the mislocalization of odours to the
1178 mouth. *Food Qual Prefer* 50:117–128. [https://doi.org/10.1016/](https://doi.org/10.1016/j.foodqual.2016.02.006)
1179 [j.foodqual.2016.02.006](https://doi.org/10.1016/j.foodqual.2016.02.006).
- 1180 Spence C, Smith B, Auvray M (2015) Confusing tastes and flavours.
1181 In: Stokes D, Matthen M, Biggs S, editors. *Perception and its*
1182 *modalities*. Oxford, UK: Oxford University Press. p. 247–274.
- 1183 Stevenson RJ, Mahmut MK, Oaten MJ (2011) The role of attention in
1184 the localization of odors to the mouth. *Atten Percept Psychophys*
1185 73(1):247–258. <https://doi.org/10.3758/s13414-010-0013-6>.
- 1186 Syarifuddin A, Septier C, Salles C, Thomas-Danguin T (2016)
1187 Reducing salt and fat while maintaining taste: An approach on a
1188 model food system. *Food Qual Prefer* 48:59–69. [https://doi.org/](https://doi.org/10.1016/j.foodqual.2015.08.009)
1189 [10.1016/j.foodqual.2015.08.009](https://doi.org/10.1016/j.foodqual.2015.08.009).
- 1190 Tateyama T, Hummel T, Roscher S, Post H, Kobal G (1998) Relation
1191 of olfactory event-related potentials to changes in stimulus
1192 concentration. *Electroencephalogr Clin Neurophysiol/Evoked*
1193 *Potent Sect* 108(5):449–455. [https://doi.org/10.1016/S0168-](https://doi.org/10.1016/S0168-5597(98)00022-7)
1194 [5597\(98\)00022-7](https://doi.org/10.1016/S0168-5597(98)00022-7).
- 1195 Teillet E, Urbano C, Cordelle S, Schlich P (2010) Consumer
1196 perception and preference of bottled and tap water. *J Sens Stud*
1197 25:463–480. <https://doi.org/10.1111/j.1745-459X.2010.00280.x>.
- 1198 Thesen T, Murphy C (2002) Reliability analysis of event-related brain
1199 potentials to olfactory stimuli. *Psychophysiology* 39(6):733–738.
1200 <https://doi.org/10.1111/1469-8986.3960733>.
- 1201 Thomas-Danguin T, Rouby C, Sicard G, Vigoroux M, Barkat S, Brun
1202 V... Rousseau F (2003) ETOC: A short olfactory test culturally
1203 validated across Europe. *Chem Senses* 28:E60.
- 1204 Thomas-Danguin T, Sinding C, Tournier C, Saint-Eve A (2016)
1205 Multimodal interactions. In: Etiévant P, Guichard E, Salles C,
1206 Voilley A, editors. *Flavor from food to behaviors, wellbeing and*
1207 *health*. Woodhead Publishing Series In Food Science,
1208 *Technology And Nutrition*. p. 430.
- 1209 Verhagen JV (2007) The neurocognitive bases of human multimodal
1210 food perception: Consciousness. *Brain Res Rev* 53(2):271–286.
1211 <https://doi.org/10.1016/j.brainresrev.2006.09.002>.
- 1212 Verhagen JV, Engelen L (2006) The neurocognitive bases of human
1213 multimodal food perception: Sensory integration. *Neurosci*
1214 *Biobehav Rev* 30(5):613–650. [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.neubiorev.2005.11.003)
1215 [neubiorev.2005.11.003](https://doi.org/10.1016/j.neubiorev.2005.11.003).
- 1216 Wang G, Bakke AJ, Hayes JE, Hopfer H (2019) Demonstrating cross-
1217 modal enhancement in a real food with a modified ABX test. *Food*
1218 *Qual Prefer* 77:206–213. [https://doi.org/10.1016/](https://doi.org/10.1016/j.foodqual.2019.05.007)
1219 [j.foodqual.2019.05.007](https://doi.org/10.1016/j.foodqual.2019.05.007).
- 1220 Welge-Lüssen A, Drago J, Wolfensberger M, Hummel T (2005)
1221 Gustatory stimulation influences the processing of intranasal
1222 stimuli. *Brain Res* 1038(1):69–75. [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.brainres.2005.01.011)
1223 [brainres.2005.01.011](https://doi.org/10.1016/j.brainres.2005.01.011).
- 1224 Welge-Lussen A, Ebnother M, Wolfensberger M, Hummel T (2009)
1225 Swallowing is differentially influenced by retronasal compared
1226 with orthonasal stimulation in combination with gustatory stimuli.
1227 *Chem Senses* 34(6):499–502. [https://doi.org/10.1093/chemse/](https://doi.org/10.1093/chemse/bjp024)
1228 [bjp024](https://doi.org/10.1093/chemse/bjp024).
- 1229 Welge-Lüssen A, Husner A, Wolfensberger M, Hummel T (2009)
1230 Influence of simultaneous gustatory stimuli on orthonasal and
1231 retronasal olfaction. *Neurosci Lett* 454(2):124–128. [https://doi.](https://doi.org/10.1016/j.neulet.2009.03.002)
1232 [org/10.1016/j.neulet.2009.03.002](https://doi.org/10.1016/j.neulet.2009.03.002).
- 1233 White TL, Thomas-Danguin T, Olofsson JK, Zucco GM, Prescott J
1234 (2020) Thought for food: Cognitive influences on chemosensory
1235 perceptions and preferences. *Food Qual Prefer* 79:103776.
1236 <https://doi.org/10.1016/j.foodqual.2019.103776>.
- 1237 Woodman GF (2010) A brief introduction to the use of event-related
1238 potentials in studies of perception and attention. *Attenti Percept*
1239 *Psychophys* 72(8):2031–2046. [https://doi.org/10.3758/](https://doi.org/10.3758/BF03196680)
1240 [BF03196680](https://doi.org/10.3758/BF03196680).

(Received 9 September 2019, Accepted 26 October 2020)
(Available online xxx)