



HAL
open science

Iron deficiency and anemia in adolescent girls consuming predominantly plant-based diets in rural Ethiopia

Yohannes Seyoum, Christèle Humblot, Gaël Nicolas, Muriel Thomas, Kaleab Baye

► **To cite this version:**

Yohannes Seyoum, Christèle Humblot, Gaël Nicolas, Muriel Thomas, Kaleab Baye. Iron deficiency and anemia in adolescent girls consuming predominantly plant-based diets in rural Ethiopia. *Scientific Reports*, 2019, 9, <10.1038/s41598-019-53836-5>. <hal-03033106>

HAL Id: hal-03033106

<https://hal.inrae.fr/hal-03033106v1>

Submitted on 2 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

OPEN

Iron deficiency and anemia in adolescent girls consuming predominantly plant-based diets in rural Ethiopia

Yohannes Seyoum^{1,2,6}, Christèle Humblot², Gaël Nicolas^{3,4,5} , Muriel Thomas⁶ & Kaleab Baye^{1*}

Rapid physical growth and the onset of menstruation during adolescence can increase the risk of iron deficiency (ID) and related adverse effects. However, little is known about the risk of anemia and ID among adolescent girls in Ethiopia. Therefore, we aimed to determine the prevalence of ID, low iron stores, and anemia and characterize selected risk factors in Huruta, Arsi Zone, Oromia Region, Ethiopia. A cross-sectional study was conducted among non-pregnant adolescent girls (15–19 years of age; $n = 257$). Data on household socio-demographic characteristics, anthropometric measurements, and women's dietary diversity score (WDDS) were collected. Hemoglobin (Hb) and serum ferritin (SF), C-reactive protein (CRP), and α -1-acid-glycoprotein (AGP) concentrations were measured. Diets were predominantly plant-based, with a low consumption of animal source foods, fruits, and dark-green leafy vegetables. Only 4% of the adolescent girls had adequate dietary diversity ($WDDS \geq 5$), and 35% were underweight. The prevalence of anemia (Hb < 11 g/dL, 8.7%) and clinical ID (SF < 15 μ g/L, 8.7%) was low, but 41% had marginal iron stores (SF < 50 μ g/L). The low prevalence of ID, despite a predominantly plant-based diet is atypical and calls for adapted strategies to address low iron stores in this and other similar settings of Ethiopia.

Anemia affects a staggering two billion people worldwide¹. Women of reproductive age in low- and middle-income countries (LMIC) are disproportionately affected. The etiology of anemia is multi-faceted and includes micronutrient deficiencies, chronic infections, and inherited hemoglobin disorders². It has long been assumed that up to 50% of the anemia burden is related to iron deficiency (ID). However, recent studies have shown causal factors to be highly heterogeneous and suggested that the burden of anemia related to ID may be lower than previously assumed³.

ID is associated with impaired physical work capacity, cognitive function, and reproductive physiology and poor pregnancy outcomes⁴. To date, most initiatives that prevent or correct iron deficiency/anemia have focused on the first 1,000 days of life from conception to the second birthday of the child. However, growing evidence suggests that maternal nutrition before conception is as equally important for preventing adverse pregnancy outcomes⁵. For example, pre-conception iron-deficiency anemia (IDA) is associated with reduced infant growth and an increased risk of adverse pregnancy outcomes⁶.

The rapid growth that occurs during adolescence, the onset of menstruation, and the consumption of predominantly plant-based diets with low bioavailable iron, all contribute to the depletion of iron stores that substantially increase the susceptibility of adolescents to IDA and related adverse outcomes in future pregnancies⁷. Consequently, WHO is promoting weekly iron-folic acid supplementation programs in populations in which the prevalence of anemia in non-pregnant women of reproductive age is $\geq 20\%$ ⁸. Although there are increasing

¹Center for Food Science and Nutrition, College of Natural and Computational Sciences, Addis Ababa University, Addis Ababa, PO Box 1176, Ethiopia. ²IRD, UMR 204 Food and Nutrition Research in the Global South (NUTRIPASS), IRD/University of Montpellier/SupAgro, Montpellier, France. ³INSERM UMR1149, CNRS ERL 8252, Centre de Recherche sur l'inflammation, Université Paris Diderot, Site Bichat, Sorbonne Paris Cité, 75018, Paris, France.

⁴Assistance Publique-Hôpitaux de Paris, Centre Français des Porphyries, Hôpital Louis Mourier, 92701, Colombes, Cedex, France. ⁵Laboratory of Excellence, GR-Ex, 75015, Paris, France. ⁶Micalis Institute, INRA, AgroParisTech, Université Paris-Saclay, 78350, Jouy-en-Josas, France. *email: kaleabbaye@gmail.com

	Proportion(%) or mean \pm SD
Household size (mean \pm SD)	6.0 \pm 2.0
Male-headed households	75.9
Household head education	
No formal education	23.4
Primary	48.0
Secondary	22.2
University or college diploma and degree	6.4
Household income source	
Wage income	5.9
Pension	1.2
Salary	10.2
Business	9.0
Agricultural production	73.7
Source of drinking water	
Improved source	95.4
Non-improved source	4.6
Type of toilet facility	
Improved facility	88.7
Non-improved facility	11.3
Access to electricity	50.0
Source of cooking fuel [§]	
Electricity	17.6
Charcoal	78.0
Wood	84.7
Straw/Shrubs/Grass	3.5
Animal dung	3.9
Fruits and vegetable cultivation	49.8
Use of cultivated fruits and vegetables (out of 49.8%)	
For sale	20.3
For food use	79.7
Ownership of agricultural land	83.1
Ownership of livestock	71.5
Mean age of participants (years \pm SD)	16 \pm 1.1
Respondent's education	
Primary	77.6
Secondary	22.4

Table 1. Household and respondents' characteristics. SD, standard deviation; [§]multiple source of cooking fuel can be used by households; hence, options are not mutually exclusive.

number of programs introducing intermittent (weekly) iron-folic acid supplementation for adolescent girls, there is little information on the prevalence of anemia, iron deficiency, and low iron stores for this population in many LMIC, including Ethiopia. Such information is critical for the design and implementation of effective intervention that maximizes benefits while minimizing possible adverse effects.

We aimed to determine the prevalence of anemia, iron deficiency, and low iron stores in adolescent girls (15 to 19 years old), using a cross-sectional design, in the Arsi Zone, Oromia Region, Ethiopia, where diets are predominantly plant-based.

Results

Characteristics of the study population. Among the 257 adolescent girls (age 15 to 19 years), 78% were enrolled in primary school (Table 1). The average household size was six and they were mostly male-headed (76%). Nearly all households had access to public tap water (96%) and 89% had a toilet facility (pit latrine with slab) on their premises. Half of the households had access to electricity. Wood was the primary source of energy for cooking (84%). Approximately, 83% of the households owned agricultural land and 71% livestock. Accordingly, agricultural production was the main source of income for their families.

Anthropometric characteristics and food consumption. Approximately a third of the adolescent girls (35%) were underweight (BMI <18.5; Table 2). Only 4% consumed at least five food groups, which is considered to be the minimum cut-off for the WDDS. The mean number of food groups consumed was three.

	Proportion (%) or mean \pm SD
Weight (kg)	47.6 \pm 6.6
Height (cm)	155.4 \pm 6.0
Body mass index (BMI)	
Underweight (<18.5)	34.6%
Overweight (25–29.99)	3.5%
WDDS	3.2 \pm 0.6
WDDS category	
Adequate (\geq 5)	4.3%
Inadequate (<5)	95.7%
Hemoglobin (g/dL)	14.2 (1.0)

Table 2. Anthropometry, hemoglobin level, and women's dietary diversity score (WDDS) of adolescent girls. SD, standard deviation.




Figure 1. Food groups consumed according to the women's dietary diversity score (WDDS) classification (n = 235). VA, vitamin A; ASF, animal-source foods.




Figure 2. Prevalence of anemia, infection, and iron deficiency (adjusted/unadjusted for the presence/duration of infection/inflammation) in adolescent girls. ID, iron deficiency.

Further investigation into the food groups showed that the diet of the girls was mainly composed of grains and tubers (100%), other vegetables (mainly onions; 99%), and pulses (84%; Fig. 1). The proportion of adolescents that consumed nuts and seeds, vitamin A-rich fruits and vegetables, dark green leafy vegetables, animal-source foods, and dairy products was very low.

Anemia, infection and iron deficiency. Despite a significant proportion of the study population being underweight and the low consumption of animal-source foods (ASF), the prevalence of anemia (8.7%) was very low, with no severe anemia, but 0.8% moderate and 7.9% mild anemia (Fig. 2).

The prevalence of clinical iron deficiency after adjusting for inflammation/infection based on serum CRP and AGP levels was low (8.7%; SF <15 μ g/L), but 41% had marginal iron stores (SF <50 μ g/L). The prevalence of infection in the area was 17%, of which 0.8%, 14.8%, and 1.5% were in incubation, early-convalescence, and

late-convalescence phase, respectively. All of the anemic subjects were also iron deficient. Furthermore, correction of the ferritin values for inflammation did not significantly affect the estimate of the prevalence of iron deficiency (+0.4%) in this study.

Discussion

We investigated the dietary diversity, prevalence of iron deficiency, and frequency of low iron stores of adolescent girls in rural Ethiopia. Diets were predominantly plant-based with little or no consumption of ASFs, fruits, or dark-green leafy vegetables. A considerable proportion of the adolescent girls were underweight, but surprisingly the prevalence of anemia and iron deficiency was low, with no cases of IDA or severe anemia.

The dietary diversity score was very low and was characterized by a very low proportion of adolescent girls consuming ASFs, fruits, and vegetables. This is in accordance with findings from the Ethiopian National Food Consumption Survey⁹. Such predominantly plant-based diets are poor sources of bioavailable iron, partially because of the high content of iron-absorption inhibitors, such as phytate and polyphenols^{10,11}. Such diets, when compounded with the low consumption of fruit, which contributes to the intake of iron absorption enhancers, such as ascorbic acid, can increase the risk of IDA¹⁰.

Surprisingly, the prevalence of anemia and iron deficiency in this study was, based on the WHO classification, found to be a mild public health concern (8%) and was even lower than the national average (20%) for adolescents of 15 to 19 years of age¹². This confirms the need for sub-national estimates and geospatial characterizations of anemia and its etiology. The relatively low prevalence of ID and anemia is in line with findings among more vulnerable age groups like young children and pregnant women in Ethiopia^{13,14}. This is attributed to high iron intake, partially from soil contamination of major food grains^{9,15}. Such soil contamination, attributed to the traditional threshing of grain under the hooves of cattle, was recently reported to contribute to the hemoglobin regeneration of iron-depleted rats¹⁶. However, there is limited knowledge on the bioavailability of iron from soil contamination and its interactions with the absorption of other nutrients and this subject warrants further investigation in humans¹⁷.

Although we observed no clinical iron deficiency, approximately half of the adolescent girls had marginal iron stores. As adolescent girls are future mothers and approximately 21 million adolescent girls in developing countries are expected to become pregnant every year, appropriate intervention strategies that restore iron stores are needed¹⁸. This is even more important in settings, such as rural Ethiopia, where pregnant women attend antenatal care late and are thus not able to start iron-folic acid supplementation sufficiently early¹⁹. Furthermore, a significant proportion of adolescent girls were underweight, suggesting that wider nutritional intervention that promotes dietary diversity and appropriate energy-balance is needed to reduce the risk of preterm birth and low birth weight²⁰.

This study has a number of limitations that needs to be accounted when interpreting our findings: first, the cross-sectional design does not allow causal inference regarding the etiology of anemia and iron deficiency; second, although a battery of iron-related biomarkers were analysed, because of financial and technical constraints we were not able to analyze all nutrients whose deficiencies may be related to anemia. Ideally, analyses of both serum ferritin and sTfR would have provided the most effective assessment of the population's iron status. This is because sTfR indicates the balance between cellular iron requirements and iron supply, while serum ferritin provides information on the iron stores. Unlike serum ferritin, sTfR is little affected by infection and inflammation and may also indicate the severity of iron insufficiency when iron stores are depleted and no other cause of erythropoiesis exists. However, given the low prevalence of depleted iron stores (SF <15 µg/L) and the correction of SF values for inflammation and infection by measuring both AGP and CRP, our assessment of iron status based on SF alone can still be considered accurate.

Notwithstanding the above limitations, this study contributes to the limited body of evidence on the prevalence of iron deficiency and low iron stores in adolescent girls in rural Ethiopia. In light of this finding, the most appropriate regimen of iron-folic acid supplementation that maximizes benefit while preventing adverse effects warrants investigation. Improving dietary quality through the promotion of diversified diet is not only critical for promoting healthy dietary patterns in the long-term, but also can support adolescents to enter pregnancy with adequate iron stores to support fetal development and their wellbeing.

Methods

Study area. The study was conducted in Huruta district, in the Oromia Region of Southeast Ethiopia, situated at 80°09' north and 39°21' east with an average elevation of ~1,978 m²¹. The study site was purposively selected for the baseline of a larger study that aimed to understand the etiology of anemia. According to the population census report of the CSA²¹, it has a total population of 15,298 of whom 7,463 are male and 7,835 female in its two *kebeles* (lowest administrative unit). Wheat is the major crop produced in the area. Cereals (barley, teff, and maize), legumes (horse beans and field peas), and various types of oil seeds are also cultivated. Furthermore, livestock, such as cattle, sheep, goats, and poultry are commonly raised in the area²².

Study design and sampling. A cross-sectional study was conducted among adolescent girls (n = 257) aged 15 to 19 years. The sample size was calculated using a single-proportion formula for a finite population using OpenEpi (<https://www.openepi.com/SampleSize/SSPropor.htm>). A population size (total number of females) of 7,835, 20% prevalence of anemia for the Oromia Region²³, 95% confidence level, and a design effect of 1.0 (because of random sampling) were taken into consideration for sample size calculation. The required sample size was augmented to account for 8% non-response rate. All participants were apparently healthy, with no chronic diseases, were not pregnant, and did not have any children. A listing of all the eligible adolescent girls was conducted with the help of the health extension workers. The study participants were then selected randomly from the listing of the two *kebeles* of Hururta.

Ethical considerations. Ethical approval was obtained from the Ethics Review Committee of the College of Natural and Computational Sciences of Addis Ababa University (CNSDO/100/09/2016) and the Oromia Regional Health Bureau (AHIQTFWO/408). Informed consent was obtained after a detailed explanation was given about the objectives and principles of the study. Informed consent/assent was obtained from parents and/or legal guardians for those under 18 years of age and from adolescents only when they were above 18. All experiments were performed in accordance with relevant guidelines and regulations.

Socio-demographic and anthropometric measurements. Data on socio-demographic characteristics was collected through interviews administered using a pre-tested questionnaire that included questions about the educational status of the family, household composition, available facilities in the household, and access to healthcare. The weight and height of each respondent were measured at the beginning of the survey using calibrated equipment, with the subjects wearing light clothing and no shoes. Then, the body mass index (BMI) was calculated as the weight in kilograms divided by the square of the height in meters (kg/m^2). Accordingly, BMI values were categorized as underweight (<18.5), normal ($18.5\text{--}24.99$), and overweight (>25)²⁴.

Women's dietary diversity score (WDDS). The WDDS was estimated using open-ended qualitative 24-hour recall. All foods eaten, inside and/or outside the home, were recorded with frequent probing. A detailed list of all ingredients of the dishes, snacks, or other foods consumed was generated to enable the classification of mixed dishes. The foods consumed were then categorized into the ten food groups that constitute the women's dietary diversity score²⁵: (1) grains, white roots and tubers, and plantains (starchy staples); (2) pulses (beans, peas, and lentils); (3) nuts and seeds; (4) vitamin A-rich fruits and vegetables; (5) dark green leafy vegetables; (6) other vegetables; (7) other fruits; (8) meat, poultry, and fish; (9) dairy; and (10) eggs. Finally, the WDDS (0–10) was calculated by summing the number of food groups consumed in the previous 24 h²⁶.

Blood sample collection and hemoglobin determination. Non-fasting venous blood samples (~4 mL) were collected in BD Vacutainer[®], Serum Separation Tubes (SST[™]) by experienced phlebotomists. A few drops of blood were immediately taken to measure hemoglobin (Hb) using the HemoCue[®] Hb 201+ system²⁷. Then, the remaining sample was immediately centrifuged and aliquoted into 2-mL Eppendorf Tubes[®] that were stored at -20°C until further analysis.

Serum ferritin (SF), C-reactive protein (CRP), and α -1 acid-glycoprotein (AGP) analyses. SF was analyzed using a fully automated clinical analyzer electrochemiluminescence immunoassay (ECLIA, Elecsys[®] 2010 analyzer Cobas e 411; Roche Diagnostics GmbH, Mannheim, Germany). CRP and AGP were determined by immunoturbidimetric methods with a clinical chemistry analyzer (Cobas Integra 400 system; Roche Diagnostic GmbH) at the Ethiopian Public Health Institute. The coefficient of variation (inter-/intra-assay) was: hemoglobin (2.1%/ 4.5%), serum ferritin (3.1%/ 3.9%), CRP (1.2%/ 2.4%) and AGP (0.8%/ 1.6%).

Deficiency cut-offs. The Hb values were adjusted for altitude and anemia was classified as mild (11–11.9 g/dL), moderate (8–10.9 g/dL), or severe (<8 g/dL)²⁸. The presence and type of inflammation were determined using AGP and CRP values. Inflammation was considered to be at the incubation stage for CRP >5 mg/L and AGP ≤ 1 g/L, early convalescence for CRP >5 mg/L and AGP >1 g/L, and late convalescence for CRP ≤ 5 mg/L and AGP >1 g/L²⁹. SF was corrected for inflammation by multiplying by 0.77 (incubation), 0.53 (early convalescence), or 0.75 (late convalescence). ID was defined as SF <15 $\mu\text{g}/\text{L}$ and marginal iron stores as SF <50 $\mu\text{g}/\text{L}$. IDA was defined as a combination of anemia and ID³⁰.

Statistical analyses. Statistical analyses of the data were performed using SPSS for windows (IBM SPSS, Inc version 20). Descriptive statistics, such as the frequency (%), means, and standard deviations, are presented. The normal distribution of data was assessed using the Shapiro-Wilk test. P-values < 0.05 were considered statistically significant.

Received: 12 May 2019; Accepted: 5 November 2019;

Published online: 21 November 2019

References

- Benoist, B. d., McLean, E., Egl, I. & Cogswell, M. Worldwide prevalence of anaemia 1993–2005: WHO global database on anaemia. *Worldwide prevalence of anaemia 1993–2005: WHO global database on anaemia*. (2008).
- Kassebaum, N. J. *et al.* A systematic analysis of global anemia burden from 1990 to 2010. *Blood* **123**, 615–624 (2014).
- Petry, N. *et al.* The proportion of anemia associated with iron deficiency in low, medium, and high human development index countries: a systematic analysis of national surveys. *Nutrients* **8**, 693 (2016).
- Camaschella, C. Iron deficiency: new insights into diagnosis and treatment. *ASH Education Program Book* **2015**, 8–13 (2015).
- Patton, G. C. *et al.* Our future: a Lancet commission on adolescent health and wellbeing. *The Lancet* **387**, 2423–2478 (2016).
- Ronnenberg, A. G. *et al.* Preconception hemoglobin and ferritin concentrations are associated with pregnancy outcome in a prospective cohort of Chinese women. *The Journal of nutrition* **134**, 2586–2591 (2004).
- Beard, J. L. Iron requirements in adolescent females. *The Journal of nutrition* **130**, 440S–442S (2000).
- WHO. Guideline: intermittent iron and folic acid supplementation in non-anaemic pregnant women (2011).
- EPHI. Ethiopian National Food Consumption Survey. (Ethiopian Public Health Institute, Addis Ababa, Ethiopia, 2013).
- Hurrell, R. & Egli, I. Iron bioavailability and dietary reference values. *The American Journal of Clinical Nutrition* **91**, 1461S–1467S, <https://doi.org/10.3945/ajcn.2010.28674F> (2010).
- Petry, N., Egli, I., Zeder, C., Walczyk, T. & Hurrell, R. Polyphenols and Phytic Acid Contribute to the Low Iron Bioavailability from Common Beans in Young Women. *The Journal of Nutrition* **140**, 1977–1982, <https://doi.org/10.3945/jn.110.125369> (2010).
- ICF, C. a. Ethiopia Demographic and Health Survey (Central Statistical Agency and ICF, Addis Ababa, Ethiopia and Rockville, Maryland, USA, 2016).

13. Gashu, D. *et al.* Ethiopian pre-school children consuming a predominantly unrefined plant-based diet have low prevalence of iron-deficiency anaemia. *Public health nutrition* **19**, 1834–1841 (2016).
14. Gebreegziabher, T. & Stoecker, B. J. Iron deficiency was not the major cause of anemia in rural women of reproductive age in Sidama zone, southern Ethiopia: A cross-sectional study. *PloS one* **12**, e0184742 (2017).
15. Baye, K., Mouquet-Rivier, C., Icard-Vernière, C., Picq, C. & Guyot, J. P. Changes in mineral absorption inhibitors consequent to fermentation of Ethiopian injera: implications for predicted iron bioavailability and bioaccessibility. *International Journal of Food Science & Technology* **49**, 174–180 (2014).
16. Guja, H. & Baye, K. Extrinsic iron from soil contributes to Hb regeneration of anaemic rats: implications for foods contaminated with soil iron. *British Journal of Nutrition* **119**, 880–886 (2018).
17. Gibson, R. S. *et al.* Dietary iron intakes based on food composition data may underestimate the contribution of potentially exchangeable contaminant iron from soil. *Journal of Food Composition and Analysis* **40**, 19–23 (2015).
18. Darroch, J. E., Woog, V., Bankole, A. & Ashford, L. S. Adding it up: Costs and benefits of meeting the contraceptive needs of adolescents (2016).
19. Gebremedhin, S., Samuel, A., Mamo, G., Moges, T. & Assefa, T. Coverage, compliance and factors associated with utilization of iron supplementation during pregnancy in eight rural districts of Ethiopia: a cross-sectional study. *BMC public Health* **14**, 607 (2014).
20. Zerfu, T. A., Umata, M. & Baye, K. Dietary diversity during pregnancy is associated with reduced risk of maternal anemia, preterm delivery, and low birth weight in a prospective cohort study in rural Ethiopia. *The American journal of clinical nutrition* **103**, 1482–1488 (2016).
21. CSA. Summary and statistical report of the 2007 population and housing census. *Addis Ababa, Ethiopia: Federal democratic republic of Ethiopia population census commission*, 1–10 (2007).
22. Mekonnen, Z. Community opinion, marketing and current debates on eucalyptus in Huruta District, Arsi Zone of Oromia Region, Ethiopia. *Eucalyptus Species Management, History, Status and Trends in Ethiopia*, 15–17 (2010).
23. CSA, I. Ethiopia demographic and health survey 2011. (2012).
24. WHO. *BMI classification Global Database on Body Mass Index*, 2006).
25. Martin-Prével, Y. *et al.* (FAO, 2015).
26. FAO. (Food and Agriculture Organization of the United Nations Rome, Italy, 2011).
27. HemoCue. *HemoCue® Hb 201+ System*, <https://www.hemocue.com/en/solutions/hematology/hemocue-hb-201plus-system> (2019).
28. WHO. (World Health Organization, Geneva, Switzerland, 2011).
29. Thurnham, D. I., Northrop-Clewes, C. A. & Knowles, J. The Use of Adjustment Factors to Address the Impact of Inflammation on Vitamin A and Iron Status in Humans–3. *The Journal of nutrition* **145**, 1137S–1143S (2015).
30. WHO. (World Health Organization, VMNIS | Vitamin and Mineral Nutrition Information System, 2011).

Acknowledgements

This research is supported by a grant from the French Embassy in Ethiopia. We would like to thank the Ethiopian Public Health Institute for serum ferritin and inflammation/infection markers analyses.

Author contributions

Y.S., C.H. and K.B. designed the study. Y.S. implemented the fieldwork, analyzed the data, and wrote the first draft. C.H., G.N., M.T. and K.B. contributed to the interpretation of the results and critically reviewed the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to K.B.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019