



HAL
open science

Sixty years from the first disease description, a novel badnavirus associated with chestnut mosaic disease

Armelle Marais, Sergio Murolo, Chantal Faure, Yoann Brans, Clement Larue, François Maclot, Sébastien Massart, Michela Chiumenti, Angelantonio Minafra, Gianfranco Romanazzi, et al.

► To cite this version:

Armelle Marais, Sergio Murolo, Chantal Faure, Yoann Brans, Clement Larue, et al.. Sixty years from the first disease description, a novel badnavirus associated with chestnut mosaic disease. *Phytopathology*, 2021, 111 (6), pp.1051-1058. 10.1094/PHYTO-09-20-0420-R . hal-03036205

HAL Id: hal-03036205

<https://hal.inrae.fr/hal-03036205>

Submitted on 6 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1 **Sixty years from the first disease description, a novel badnavirus associated with**
2 **chestnut mosaic disease**

3

4 Armelle Marais*†¹, Sergio Murolo†², Chantal Faure¹, Yoann Brans³, Clément Larue^{4,5}, François
5 Maclot⁶, Sébastien Massart⁶, Michela Chiumenti⁷, Angelantonio Minafra⁷, Gianfranco Romanazzi²,
6 Marie Lefebvre¹, Teresa Barreneche¹, Cécile Robin⁴, Rémy J. Petit⁴, Thierry Candresse¹

7

8 ¹ Univ. Bordeaux, INRAE, UMR BFP, Villenave d'Ornon, France

9 ² Department Agricultural, Food and Environmental Sciences, UNIVPM, Ancona, Italy

10 ³ Laboratoire de Virologie et de Biologie moléculaire, CTIFL, Prignonrieux, France

11 ⁴ Univ. Bordeaux, INRAE, UMR Biogeco, Cestas, France

12 ⁵ INVENIO, Maison Jeannette, Douville, France

13 ⁶ Plant Pathology Laboratory, TERRA-Gembloux Agro-BioTech, University of Liège, Belgium

14 ⁷ CNR Institute for Sustainable Plant Protection —Bari, Italy

15

16 * Corresponding author: A. Marais; Email: armelle.marais-colombel@inrae.fr

17 † AM and SM contributed equally to this work

18

19 **Keywords: High throughput sequencing, *Castanea* sp., etiology, viral disease, *Badnavirus*,**
20 ***Caulimoviridae***

21

22 The nucleotide sequences reported here have been deposited in GenBank under the accession
23 numbers MT261366, MT269853, MT270664-MT270682, and MT339503-MT339590.

24

25 **ABSTRACT**

26 Although the chestnut mosaic disease (ChMD) was described several decades ago, its etiology is still
27 not elucidated. Here, using classical approaches in combination with high throughput sequencing
28 (HTS) techniques, we identify a novel *Badnavirus* that is a strong etiological candidate for ChMD.
29 Two disease sources from Italy and France were submitted to HTS-based viral indexing. Total RNAs
30 were extracted, ribodepleted and sequenced on an Illumina NextSeq500 (2x150 or 2x 75 nt). In each
31 source, we identified a single contig of about 7.2 kilobases that corresponds to a complete circular
32 viral genome and shares homologies with various badnaviruses. The genomes of the two isolates
33 have an average nucleotide identity of 90.5% with a typical badnaviral genome organization
34 comprising three open reading frames. Phylogenetic analyses and sequence comparisons show that
35 this virus is a novel species for which we propose the name *Chestnut mosaic virus* (ChMV). Using a
36 newly developed molecular detection test, we systematically detected the virus in symptomatic
37 graft-inoculated indicator plants (chestnut and American oak), as well in chestnut trees presenting
38 typical ChMD symptoms in the field (100% and 87% in France and Italy surveys, respectively).
39 Datamining of publicly available chestnut SRA transcriptomic data allowed the reconstruction of two
40 additional complete ChMV genomes from two *Castanea mollissima* sources from the USA, as well
41 as ChMV detection in *C. dentata* from the USA. Preliminary epidemiological studies, performed in
42 France and in Central Eastern Italy, showed that ChMV has a high incidence in some commercial
43 orchards, with a low within-orchard genetic diversity.

44

45

46 European chestnut (*Castanea sativa* Mill.) has a long-standing tradition of cultivation in many
47 European countries. It is an important species, both economically as a source of timber and fruit
48 and ecologically through the multiple ecosystemic services it provides. In Europe, chestnut covers
49 about 2.5 million hectares, mainly concentrated in France, Italy, Spain, Portugal, Switzerland, the
50 Balkan regions, and Southern England (Conedera et al. 2016). Chestnut (*Castanea* spp.) can be
51 heavily affected by various pathogens. The most detrimental are caused by fungal-like organisms
52 (Oomycetes) and fungi such as *Phytophthora cambivora* Petri and *P. cinnamomic* Rands., the agents
53 of ink disease, or *Cryphonectria parasitica*, the causal agent of chestnut blight, which all provoke
54 disorders that can lead to tree mortality (Prospero et al. 2012; Rigling and Prospero, 2018). In Italy,
55 Gualaccini (1958) described a chestnut disease associated with viral symptoms (mosaic, shoots with
56 asymmetric leaf blade deformation) which was again reported in Campania during the 80s,
57 (Ragozzino and Lahoz 1986), and in the Marche region (central eastern Italy) in 2000 (Antonaroli
58 and Perna; 2000). In France, the disease was first identified circa 1987 on cultivars of *C. sativa* x *C.*
59 *crenata* hybrids from commercial orchards located in the south-west of the country. Desvignes
60 (1999b) made a more detailed description of the symptoms, which are characterized by necrotic
61 lesions in the bark and wood that turns into cankers, chlorotic lesions and yellow stripes on leaf
62 veins and partial limb atrophy, and called this disease Chestnut Mosaic Disease (ChMD). This disease
63 can heavily affect the production of both young and secular trees (Antonaroli and Perna, 2000). It
64 has also been reported in Japan and Hungary (Shimada, 1962; Horvath et al. 1975). Even if its
65 etiology has long remained unknown, researchers hypothesized that the causal agent of ChMD
66 could be a virus, introduced in Europe between 1940 and 1960 when a number of *C. crenata*
67 cultivars were imported from Japan for breeding purposes. Investigations in France and Italy
68 established that the causal agent can be eliminated by thermotherapy, is aphid-transmissible, and
69 is graft-transmissible to *Castanea* and *Quercus* species in which it may elicit symptoms (Desvignes

70 and Lecoq, 1995; Desvignes, 1999b; Vettraino et al. 2005). The susceptibility to the ChMD agent of
71 *Castanea* species/cultivars has been evaluated in several studies (Desvignes, 1992; 1999b;
72 Desvignes and Lecoq, 1995). Three categories of cultivars could thus be defined from tolerant to
73 moderately and fully susceptible. Graft incompatibility was also observed when cultivars of different
74 susceptibilities are assembled by grafting. Most of the *C. sativa* cultivars and hybrids are tolerant to
75 ChMD, although some well-known French hybrids like 'Maraval' (Ca 74) are fully susceptible, and
76 used for indexing purposes to detect the ChMD agent in tolerant cultivars (Desvignes and Lecoq,
77 1995).

78 In the last decade, a number of studies have highlighted the potential of non-targeted molecular
79 diagnostics based on high-throughput sequencing (HTS) to elucidate the etiology of viral plant
80 diseases and to provide viral sequence data from which rapid diagnostic molecular assays can be
81 developed (Martin et al. 2016; Villamor et al. 2019). Since 2009, HTS combined with bioinformatics
82 have been used for the discovery, characterization, and *de novo* assembly of the genome of known
83 and novel plant viruses and viroids (Rott et al. 2017; Kreuze et al. 2009). This has accelerated the
84 application of HTS technologies in the field diagnostic of diseases (Massart et al. 2014), and in
85 quarantine regulations (Martin et al. 2016; Massart et al. 2017).

86 Badnaviruses are plant pararetroviruses belonging to the family *Caulimoviridae* that have
87 emerged as serious pathogens causing severe yield losses in a wide range of economically important
88 crops all over the world (Bhat et al. 2016). The genome of badnaviruses is composed of a non-
89 covalently closed, circular double-stranded DNA (ranging from 7.2 to 9.2 kbp) and is encapsidated
90 in bacilliform virions. This genome typically harbors three open reading frames (ORFs) encoding,
91 respectively, a protein of unknown function, the virion-associated protein (VAP), and a polyprotein
92 containing functional and structural domains [movement protein (MP), coat protein (CP), aspartic
93 protease (AP), reverse-transcriptase (RT) and RNase H]] (Hohn and Rothnie, 2013; Bhat et al. 2016).

94 Badnaviruses can also be present as integrated sequences in some host plant genomes (endogenous
95 badnaviruses) (Staginnus et al. 2009; Bhat et al. 2016). The contribution of these integrated
96 sequences to host and virus evolution is still poorly understood (Geering et al. 2014).

97 Given the very limited knowledge on the etiology of ChMD, and based on previously published
98 studies (Desvignes, 1992; 1999a, 1999b; Desvignes and Lecoq, 1995; Desvignes and Cornaggia,
99 1996), we investigated the hypothesis that a virus might be involved in this disease. Combining HTS-
100 based viral indexing and classical approaches, we report here the complete genome sequence of a
101 novel badnavirus species for which the name *Chestnut mosaic virus* (ChMV) is proposed. We further
102 show that there is a strict correlation between the presence of the virus and the appearance of
103 typical ChMD symptoms in various graft-inoculated indicator plants. Preliminary epidemiological
104 studies carried out in Italy and in France reveal that the virus can have high incidence in some
105 orchards, and, as expected, can be associated with symptomatic or asymptomatic infections.

106

107

MATERIALS AND METHODS

108 **Plant samples and virus isolates.** Virus isolates included in this study are listed in Supplementary
109 Table S1. Isolate LC1224H is originated from a red oak (*Quercus rubra*) artificially inoculated in 1992
110 with a chestnut mosaic source from a hybrid *Castanea sativa* x *Castanea crenata* included in a
111 French breeding program. Leaves of grafted oaks displayed typical symptoms characterized by
112 chlorotic mottle, yellow veins, and mosaic (Desvignes and Lecoq 1995) (Figure 1A). Isolate
113 FRlc1224A was derived from the same source, and is the result of a back-inoculation by grafting of
114 LC1224H to the natural chestnut hybrid Maraval (Ca 74; *C. crenata* x *C. sativa*) indicator (Desvignes
115 et al. 1992). Isolate LC1224F originated from a Maraval indicator inoculated by aphid transmission
116 from an initial ChMD source in a *C. crenata* x *C. sativa* French hybrid (Desvignes and Cornaggia,
117 1996). The LCA552 and LCA584 isolates were collected from *C. sativa* trees in France in 2009 and

118 2018, while the T32018 disease source was isolated from a French hybrid *C. crenata* x *C. sativa* in
119 2018. All of these isolates have been held and propagated on 'Maraval' indicator plants at the CTIFL
120 virology laboratory (Lanxade, France).

121 In the framework of a survey carried out in Italian chestnut orchards to monitor chestnut blight
122 (Acquasanta Terme (AP), locality Umito, Italy) (Murolo et al. 2018), typical leaf symptoms of ChMD
123 were recorded in 2016. Six symptomatic plants were collected, pooled (10 -15 symptomatic shoots)
124 and included in the HTS analysis (ITumito39 source).

125 In order to evaluate the incidence of ChMV, chestnut trees from INRAE chestnut biological
126 resource center (<https://www6.bordeaux-aquitaine.inrae.fr/biogeco/Ressources>) were sampled.
127 This orchard is located on the Villenave d'Ornon INRAE center (France) with trees distributed in
128 three plots (A, E, or Port, Table S1). A total of 43 *C. sativa*, 14 *C. mollissima*, six *C. crenata* and 32
129 hybrid chestnut trees were sampled, corresponding to a total of 38 symptomatic trees with typical
130 ChMD symptoms, 47 asymptomatic trees, and 10 trees with dubious or atypical symptoms. In
131 addition, in the Central eastern Italy Marche region, leaves from 60 symptomatic and from 10
132 asymptomatic grafted *C. sativa* cv. Marrone trees of different ages were collected in a commercial
133 chestnut orchard (Plot I, Table S1).

134 Isolates FRlc1224A and ITumito39 were used for the HTS analysis, whereas all other samples were
135 included either in the incidence analysis or in the causal relationship analysis (Table S1).

136 **Total RNA extraction and RNA-Seq analysis.** Symptomatic leaves from a 'Maraval' indicator
137 (FRlc1224A) were collected and used to extract total RNAs according to the protocol described by
138 Reid et al. (2006). For the Italian material, total RNAs were extracted from symptomatic leaves
139 according to the protocol described by Gambino et al. (2008). Total RNAs were then submitted to a
140 DNase treatment following the manufacturer's recommendations (Fisher Scientific, Illkirch, France).
141 Ribosomal RNAs were removed using a RiboMinus Plant Kit for RNA-Seq (Invitrogen, Fisher

142 Scientific, Illkirch, France) before cDNA library synthesis with the Illumina TruSeq Stranded RNA
143 library Prep kit (Illumina, Inc., San Diego, CA) and sequenced on an Illumina NextSeq500 (2x150 nt
144 or 2x75 nt) in a multiplexed format (GIGA-Genomics facility, Université de Liège, Belgium).

145 **Bioinformatic analysis.** Primary quality analyses were performed using Geneious Prime
146 2019.2.1 Software (<https://www.geneious.com>). *De novo* assemblies of quality filtered reads were
147 performed using Velvet (Zerbino and Birney, 2008), Geneious R 11 (<https://www.geneious.com>),
148 and Spades (Bankevich et al. 2012), or using the CLC genomics workbench 8.0
149 <http://www.clcbio.com>). Contigs were annotated by BlastN and BlastX comparisons with nucleotide
150 and non-redundant protein GenBank databases, respectively. Blast results were screened using e-
151 value thresholds of 10^{-6} and 10^{-4} for BlastN and BlastX, respectively. Publicly available chestnut RNA-
152 Seq transcriptomic data were retrieved from the NCBI Sequence Read Archive (SRA) and
153 downloaded reads were mapped against the sequence of the FRlc1224A isolate using CLC Genomics
154 Workbench 11.0. When needed, *de novo* assembly and contig annotations were also performed as
155 described above.

156 **Total DNA extraction and PCR confirmation of genome completeness and circularity.** In order
157 to verify both the completeness of the assembled genome sequences and genome circularity, pairs
158 of specific outward-facing primers were designed for each isolate (Ch-Bad-6976F/ Ch-Bad-252R for
159 the isolate FRlc1224A and Bad-Ch-6481F/Bad-Ch-325R for the isolate ITumito39, Table S2). Leaf
160 tissues (0.5 g) were pulverized in liquid nitrogen and total DNA was extracted in CTAB buffer (2%
161 cetyl trimethylammonium bromide, 100 mM Tris-HCl, 1.4 M NaCl, 20 mM EDTA), adding 3%
162 polyvinyl pyrrolidone 40, and 0.5% sodium metabisulfite (Doyle and Doyle, 1990). Finally, the DNA
163 pellets were resuspended in 50 μ l sterile water. PCRs were performed in a 50 μ l reaction volume
164 containing 10 mM Tris-HCl (pH 8.5), 2 mM $MgCl_2$, 50 mM KCl, 0.2 mM dNTPs, forward and reverse
165 primers at 1 μ M each, and 1.25U of Dream Taq (ThermoFisher) using 50 ng of the template. After an

166 initial denaturation step at 95°C for 4 min, 40 or 35 cycles, respectively, were set at 94°C for 30 sec,
167 60°C (Ch-Bad-6976F/ Ch-Bad-252R) or 55°C (Bad-Ch-6481F/325R) for 30 sec, and 72°C for 90 sec,
168 followed by a final extension step of 10 min at 72°C. PCR amplification products were sequenced on
169 both strands (GATC, Eurofins, Ebersberg, Germany).

170 **ChMV molecular detection and variant analysis by PCR.** For the molecular detection of ChMV,
171 two sets of primers were designed in conserved regions of ORF3 designed using the sequences of
172 isolates FRlc1224A and ITumito39. One primer pair (Ch-Bad-1466F/Ch-Bad-1800R, Table S2) allows
173 the amplification of a genomic region (335 nt) in the MP domain (Figure 2), whereas the second pair
174 (Ch-Bad-5860F/Ch-Bad-6109R, Table S2) amplifies a 232-nt fragment in the RH domain (Figure 2).
175 An aliquot of 25 ng of total DNA was used for the PCR assays in a 50 µl volume containing 10 mM
176 Tris-HCl (pH 8.5), 2 mM MgCl₂, 50 mM KCl, 0.2 mM dNTPs, forward and reverse primers at 1 µM
177 each, and either 1.25U of DreamTaq or 1U of GoTaq. After an initial denaturation step at 95°C for 4
178 min, 35 cycles were set at 94°C for 30 sec, 56°C for 30 sec, and 72°C for 90 sec, followed by a final
179 extension step of 10 min at 72°C. Amplicons were analyzed by electrophoresis on 1.5% agarose gel
180 and were directly sequenced on both strands (GATC).

181 Possible phytoplasma infection was evaluated using primer pair P1/P7 (Deng and Hiruki. 1991;
182 Smart et al. 1996) and, in nested PCR, primers R16F2n/R2 (Gundersen and Lee 1996).

183 **Sequence and phylogenetic analyses.** The full-length genomes were analyzed by ORF Finder
184 (<http://www.ncbi.nlm.nih.gov/projects/gorf/>) to identify putative ORFs in the viral genome.
185 Deduced amino acid (aa) sequences were analyzed for conserved protein domains gathered in
186 Conserved Domains Database (CDD) (<http://www.ncbi.nlm.nih.gov/structure/cdd.shtml>) and
187 theoretical molecular weights were calculated using ExpASY (http://web.expasy.org/compute_pi/).
188 Multiple alignments of nucleotide (nt) or amino acid (aa) sequences were performed using the
189 ClustalW program (Thompson et al. 1994) implemented in MEGA version 7.0 (Kumar et al. 2016).

190 Genetic distances (p-distances using strict nt or aa identity) were calculated using MEGA 7.0.
191 Phylogenetic trees were reconstructed using the neighbor-joining method implemented in MEGA
192 7.0 and robustness of nodes was assessed from 1,000 bootstrap resamplings.

193

RESULTS

194 **Determination of the complete genome sequence of a novel badnavirus from two chestnut**
195 **disease sources.** Two ChMD sources were included in the HTS analysis. The French source
196 (FRlc1224A) showed typical ChMD symptoms, with leaf deformation, yellow veins and chlorotic
197 diffuse mottling (Figure 1B) and the Italian source (ITumito39) is a mixture of six plants showing
198 intensive vein banding and leaf blade deformation (Figure 1C). High-throughput sequencing of
199 ribodepleted RNAs extracted from the sources FRlc1224A and ITumito39 yielded a total of
200 10,737,052 reads and 4,135,330 reads, respectively. *De novo* assembly and Blast annotation allowed
201 for the identification of a single long contig with significant homology with badnaviruses. These
202 contigs were respectively 7,264 and 7,214 bp long and showed short terminal redundancies,
203 consistent with the structure of the long RNA transcript involved in the replication of badnaviruses
204 (Teycheney et al. 2020) and suggesting they represented the full coverage of a circular badnaviral
205 genome. A total of 39,657 reads were integrated in the FRlc1224A contig, representing 0.37% of
206 total reads, with a mean coverage depth of 795X, whereas 611 reads (0.015% of total reads) were
207 integrated in the ITumito39 contig, with a mean coverage depth of 14.4X. The circularity and
208 completion of the DNA genome sequence of each isolate were validated by a PCR on purified DNA
209 extracted from the host plants and using specific outward-facing primers designed from the contig
210 sequences. The respective 436 nt- and 1,007 nt fragments were amplified and sequenced,
211 confirming DNA genome completeness and circularity (data not shown). The assembled sequences

212 have been deposited in GenBank under accession numbers MT269853 and MT261366, respectively.
213 No other plant virus was detected in the two datasets during the Blast annotation of contigs.

214 **Genome organization of chestnut mosaic virus and determination of its phylogenetic**
215 **relationships.** The badnaviral genomes characterized independently from the French and Italian
216 ChMD sources are respectively 7,160 bp and 7,161 bp long, within the range of badnavirus genome
217 sizes (Teycheney et al. 2020). The genomic organization is the same for both isolates, comprising
218 three open reading frames (ORFs) encoded on the positive strand (Figure 2), and is typical for
219 badnaviruses (Teycheney et al. 2020). The ORF1 (nt 245-751, numbering according to the isolate
220 FRlc1224A sequence) encodes a protein of 169 aa (19.8 kDa), the ORF2 (nt 751-1161) encodes a
221 137-aa protein (15 kDa), and the third ORF (nt 1,163-6,721) encodes a polyprotein of 1,853 aa (211.7
222 kDa) with five conserved protein domains (Figure 2): a viral movement protein (MP, cl03100), a zinc-
223 binding motif (ZnF, pfam00098), a retroviral aspartyl protease domain (RVP, pfam00077), a reverse
224 transcriptase domain (RT, cd01647) and a ribonuclease H domain (RH, cl14782). The two “Cys”
225 motives (C-X₂-C-X₄-H-X₄-C, and C-X₂-C-X₁₁-C-X₂-C-X₄-C-X₂-C) usually found in the coat protein of
226 badnaviruses (Bath et al. 2016) were also detected in the ORF3-deduced protein, between aa
227 positions 777-790 and 902-928.

228 Both isolates are closely related, with an overall 90.5% nt identity. The three indels observed
229 between the two sequences are located in the intergenic region, the isolate ITumito39 ended up
230 being one nucleotide longer. The three ORFs have the same sizes, are strictly colinear and the
231 encoded proteins share respectively 95.2% (ORF1), 95.5% (ORF2) and 94.8% (ORF3) aa identity.

232 To characterize the phylogenetic relationships and taxonomic position of the chestnut
233 badnavirus, a phylogenetic tree was reconstructed using an alignment of full genome nucleotide
234 sequences of *Badnavirus* genus members, with the rice tungro bacilliform virus used as an outgroup
235 (Figure 3). Both isolates cluster in group 3 defined by Wang et al. (2014), together with gooseberry

236 vein banding virus (GVBV), rubus yellow net virus (RYNV), grapevine vein clearing virus (GVCV), birch
237 leafroll-associated virus (BLRaV), wisteria badnavirus 1 (WBV1), and pagoda yellow mosaic-
238 associated virus (PYMaV) (Figure 3). Nevertheless, they are clearly distant from all of these species,
239 defining a novel branch, supported by a 99% bootstrap value (Figure 3). Tree topology was similar
240 when using an alignment of representative badnaviral ORF3 protein sequences (Figure S1). To
241 confirm these analyses, pairwise comparisons of genome sequences showed that the isolate
242 FRlc1224A has only weak identity levels with representative members of the genus *Badnavirus*,
243 comprised between 42.1% nt identity (sugarcane bacilliform IM virus; 42.5% for the isolate
244 ITumito39) and 50.9% (WBV1; 50.8% for the isolate ITumito39). The same tendency is observed
245 when considering the genome proteins. The ORF1-encoded protein shows only weak homology with
246 the corresponding proteins of WBV1 (27.8% aa identity) and PYMAV (26.1% aa identity), and the
247 ORF2-encoded protein shares only 33.1% aa identity with the corresponding protein of the most
248 closely related virus, WBV1. The polyprotein encoded by ORF3 shares 49.5% aa identity with the
249 corresponding protein of the closest relative, PYMAV. Using the ORF3 region (RT and RH domain)
250 used for taxonomical discrimination in the family *Caulimoviridae* (Teycheney et al. 2020), the
251 FRlc1224A isolate shows between 64% (with GVBV) and 68.4% (with BLRaV) nt identity (Table 1),
252 which is below the 80% nt identity value used as the species demarcation threshold in the family.
253 Therefore, this virus represents a novel species in the family *Caulimoviridae*, for which we propose
254 the name *Chestnut mosaic virus* (ChMV). In the same taxonomically informative region, the isolates
255 FRlc1224A and ITumito39 share 91.9% nt identity (97.8% aa identity), indicating that they belong to
256 the same viral species (Table 1).

257 **Identification of ChMV in publicly available chestnut HTS data.** The datamining of chestnut
258 HTS data from various chestnut sources publicly available at GenBank [EST sequences, whole
259 genome assembly, RNA-Seq and Genotyping-by-sequencing (GBS) reads available as Sequence Read

260 Archives (SRA)] allowed the identification of ChMV in several of those datasets (Table S3). In
261 particular, two complete genomes were obtained from datasets involving *C. mollissima* cv. Vanuxem
262 in the USA, one from the whole genome assembly (JRKL01079565) and the other by *de novo*
263 assembly of RNA-Seq data (SRX4015368) with 99.2% and 97.4% nt identity, respectively, with the
264 FRlc1224A isolate over the whole genome (Figure 4). In addition, partial ChMV genome assemblies
265 of >3 kbp could be obtained from a range of other datasets generated in the USA or in China from
266 *C. mollissima* (Table S3), all of which showed significant relatedness with the FRlc1224A sequence
267 as shown by a phylogenetic tree reconstructed using nucleotide alignments of concatenated ChMV
268 sequences retrieved from the various datasets (Figure 4). In addition, partial ChMV genomes could
269 also be reconstructed from two datasets obtained from *C. dentata* in the USA. Interestingly, one of
270 these two *C. dentata* isolate sequences shows closest relationship with ITumito39 sequence (Figure
271 4) with only 89.2% nt identity with the isolate FRlc1224A as compared to 93.9% nt identity with
272 ITumito39. The second isolate of *C. dentata* appears to be equally related to the FRlc1224A and
273 ITumito39 isolates, with 90.9% and 90.6% nt identity, respectively.

274 **Incidence and genetic variability of ChMV in France and Italy.** The incidence and genetic
275 variability of ChMV were investigated by analyzing two genomic regions of ORF3, one 335-nt long
276 located in the MP domain amplified using primer pair Ch-Bad-1466F/Ch-Bad-1800R and the other
277 232-nt long in the RNase H domain and amplified with primer pair Ch-Bad-5860F/Ch-Bad-6109R
278 (Table S2, and Figure 2). The two primer pairs were designed to be able to detect both isolates
279 FRlc1224A and ITumito39. In Italy, a total of 70 *C. sativa* cv Marrone samples were collected in the
280 same location, while in France, 95 chestnut accessions belonging to three different *Castanea* species
281 or hybrids were sampled in three plots (A, Port, E). Both symptomatic and asymptomatic samples
282 were collected, as well as some samples with atypical or dubious symptoms. Globally, ChMV was
283 frequent in the surveyed plots, with 57/70 (81.5%) infected *C. sativa* samples in Italy and 65/95 trees

284 (68%) in France (Table 2). In the Italian orchard, half of the asymptomatic trees were found to be
285 infected by ChMV, compared to 87% of the symptomatic ones (Table 2). None of the analyzed
286 samples were found positive using a phytoplasma-specific PCR assay. In the French collection, the
287 virus was detected in 100% (38/38) of the trees showing typical ChMD symptoms, and in 49%
288 (23/47) of the asymptomatic trees, including two trees that were symptomless but showed strong
289 symptoms on rootstock off-shoots (Figure S2). ChMV was also detected in four out of the 10 trees
290 showing atypical/doubtful symptoms.

291 The genetic variability of ChMV was evaluated by analyzing the sequences of the two PCR
292 amplicons generated for the incidence survey. Considering the relative homogeneity of the origin
293 of the Italian samples, the number of samples included in this analysis was limited to 13 (four from
294 asymptomatic trees, and nine symptomatic ones) (Table S1). In total, the final dataset consisted in
295 53 isolates for which the sequence of the two genomic regions were available (49 from the incidence
296 survey and four from independent ChMD sources held in collection at CTIFL, see below). As
297 illustrated by the unrooted neighbor-joining tree reconstructed from the alignments of RT-RnaseH
298 domain nucleotide sequences (Figure S3A), ChMV diversity is structured into two clusters, defined
299 by the geographical origin of the samples (Italy and France). The sequences determined from the
300 four independent French disease sources (FR1c1224A, T30218, LCA552, LCA584) belong to the same
301 «French» cluster. Overall, the level of genetic diversity is very low in this genomic region, with an
302 average pairwise nt divergence (diversity) of 2.2% +/- 0.5%. This value is even lower when
303 considering the intra-group diversity, as 0.2% +/- 0.1% within the French cluster and 0.1% +/- 0.1%
304 within the Italian ones. In contrast, the inter-group diversity reaches 6.3% +/- 1.5%, confirming the
305 existence of two geographical clusters. The same trends are observed when analyzing the genomic
306 region located in the MP domain (Figure S3B). The same geographical clustering could be observed,
307 with the exceptions of three French isolates that seem to be more closely related to the Italian

308 cluster (Figure S3B). Another French isolate, 20971-E remains isolated and does not fit in either
309 group. The average nt divergence in this region is slightly higher than in the RT-RNaseH region, (5.8%
310 +/- 0.7%), and the inter-group diversity reaches a value of 13.4% +/- 1.8%, as compared to the 6.3%
311 value for the other region.

312

DISCUSSION

313 Since the seminal work of Desvignes and collaborators in the 1990s, it has been acknowledged
314 that the agent responsible for ChMD is most likely a thermosensitive, graft-transmissible virus that
315 can be transmitted experimentally and probably naturally by the aphid *Myzocallis castanicola*
316 Desvignes, 1992; 1999a; 1999b; Desvignes and Lecoq, 1995; Desvignes and Cornaggia, 1996).
317 Depending on the chestnut genotype, this infection can either be asymptomatic or can result in the
318 expression of severe and conspicuous ChMD symptoms. In chestnut orchards in the Marche region
319 (Italy), both young and mature plants were shown to be affected, significantly decreasing chestnut
320 production. Symptoms have also been observed in some *Quercus* species following experimental
321 graft inoculation. To date, however, the causal agent remains to be identified.

322 Here, by using HTS-based viral indexing, we were able to identify and characterize, in two
323 independent ChMD sources, two isolates of the same novel virus. Phylogenetic and sequence
324 analyses showed that this virus belongs to the genus *Badnavirus*, in the family *Caulimoviridae*, and
325 could be considered as a new species, for which the name *Chestnut mosaic virus* (ChMV) is
326 proposed. Interestingly, this new virus clusters with a group of badnaviruses that includes RYNV,
327 GVBaV, and GVCV.

328 There is unambiguous evidence that ChMV as reported here is an episomal virus. It was detected
329 in graft-inoculated indicators, and not in non-inoculated control plants of the same variety,
330 demonstrating its graft-transmissibility, a property of episomal viruses. This line of evidence is
331 further reinforced by the detection of ChMV in symptomatic, graft-inoculated indicator *Quercus*

332 plants and, again, not in the corresponding control plants. In parallel, the HTS detection of ChMV
333 from DNase-treated RNAs, the failure to detect ChMV in a range of the surveyed chestnut trees and
334 the sequence diversity identified in ChMV all rule out a scenario in which an endogenous ChMV
335 genome, integrated in the chestnut genome could be responsible for the HTS and PCR results
336 reported here. There was in fact no indication of ChMV in the chestnut genome assembly
337 (JRKL01079565) since no integration borders could be identified and a single contig, representing a
338 complete unintegrated viral genome transcript, was identified. Integration of ChMV as an
339 endogenous viral element (EVE, Bhat et al. 2016) therefore does not appear to be a general genomic
340 feature of chestnut.

341 According to the simplified hierarchical approach proposed by Fox (2020) for assessing causal
342 relationships in plant virology, ChMV appears as a good candidate, if not as the causative agent of
343 ChMD. There are several arguments and experimental evidence supporting this idea. Following HTS
344 analyses, ChMV was the sole virus detected in the French source FRlc1224A, coming from a ChMD
345 source initially involving a *C. sativa* x *C. crenata* hybrid. It was also the sole virus detected in the
346 Italian ChMD source analyzed by HTS. Using molecular detection tests developed in this work, the
347 virus was consistently found in other symptomatic accessions derived from the same diseased
348 source (LC1224H, a *Q. rubra* artificially inoculated and LC1224F, an indicator plant inoculated by
349 aphid transmission; Figure 5). In addition, three other independent chestnut sources shown by
350 biological indexing on the 'Maraval' indicator to be affected by ChMD were found to be infected by
351 ChMV (LCA552, LCA584 and T32018 in Figure 5). There is therefore a correlation between the
352 appearance of ChMD symptoms and the presence of ChMV in the graft-inoculated indicators,
353 supporting the hypothesis of a causal relationship between ChMV infection and ChMD. In total, five
354 independent ChMD sources collected between 1990 and 2018 in two countries (Italy and France)

355 were ChMV positive, satisfying the Bradford-Hill's experimental and consistency criteria (Bradford
356 Hill, 1965; Fox, 2020).

357 Preliminary studies indicate that ChMV is highly prevalent in the analyzed orchards in France
358 and Italy, confirming the earlier results of Desvignes (1999a). In parallel, the identification of ChMV
359 sequences in publicly available HTS data provides a strong indication of the presence of ChMV in *C.*
360 *mollissima* in the USA and in China and in *C. dentata* in the USA. In the surveys, ChMV was not
361 systematically associated with symptomatic infections, although its frequency was systematically
362 higher in symptomatic plants. This result was expected since previous grafting experiments had
363 demonstrated that not all chestnut varieties/species are susceptible to ChMD and develop
364 symptomatic infections (Desvignes and Lecoq, 1995; Desvignes, 1992; 1999b). Biological indexing
365 on the susceptible 'Maraval' indicator has in particular identified latent ChMV infections in many
366 symptomless *C. sativa* varieties or *C. sativa* x *C. crenata* hybrids (Desvignes and Lecoq, 1995;
367 Desvignes, 1992; 1999b). On the other hand, all surveyed symptomatic plants in France were found
368 to harbor the virus, while it was detected in 52/60 (87%) of tested symptomatic Italian trees. The
369 failure to detect ChMV in eight symptomatic Italian trees might reflect sequence variability and an
370 incomplete inclusiveness of the PCR primers used or low or uneven virus accumulation. Indeed,
371 using biological indexing, Desvignes et al. have previously found an uneven distribution of the ChMD
372 agent in infected trees leading to a failure to detect it in parts of some infected trees (Desvignes and
373 Lecoq, 1995; Desvignes et al. 1999b).

374 Taken together, and even though Koch's postulates were not fully verified, the experiments
375 reported here make a very strong case for a role of ChMV as the causal agent of the chestnut mosaic
376 disease. The low ChMV diversity observed in France and Italy are consistent with the scenario of its
377 recent introduction in Europe (Desvignes and Lecoq, 1995), while the genetic separation of the
378 Italian and French clusters is suggestive of separate introduction events. These results and the

379 associated development of molecular tools for the detection of ChMV will help speed up the
380 selection of virus-free mother plants and mitigate the virus spread in new chestnut orchards and
381 layerings. However, many questions remain regarding the variability of symptom intensity in
382 relationship to cultivar susceptibility, ChMV-induced graft incompatibility, the impact of
383 pedoclimatic conditions and of synergic and competitive interferences with other chestnut
384 pathogens, and silvicultural management.

385

386

ACKNOWLEDGMENTS

387 The authors thank the INRAE Vine Experimental Unit of Bordeaux (UEVB) and X. Capdevielle for
388 taking care of chestnut orchards. We would like to thank Ascenzio Santini (Azienda Agricola Laga
389 Nord, Umito, Aquasanta Terme, Italy) for the essential support during the phytosanitary monitoring
390 in chestnut orchards. Thanks to D. Cornaggia and P. Gentit for the information provided on the past
391 work of CTIFL, for the characterization of the symptoms of the disease on biological indicators and
392 the conservation of the isolates. Thanks to Eden Darnige for proofreading the manuscript. Part of
393 this work was based upon COST Action FA1407 (DIVAS), supported by COST (European Cooperation
394 in Science and Technology).

395

396

LITERATURE CITED

397 Antonaroli, R., and Perna, M.R. 2000. Una fitopatia ad eziologia ancora incerta: il giallume di
398 castagno in Emilia Romagna e nelle Marche. *Sherwood* 6:43-46.

399 Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., Lesin, V. M.,
400 Nikolenko, S. I., Pham, S., Prjibelski, A. D., Pyshkin, A. V., Sirotkin, A. V., Vyahhi, N., Tesler, G.,
401 Alekseyev, M. A., and Pevzner, P. A. 2012. SPAdes: a new genome assembly algorithm and its
402 applications to single-cell sequencing. *J. Comput. Biol.* 9:455-477.

- 403 Bhat, A. I., Hohn, T., and Selvarajan, R. 2016. Badnaviruses: the current global scenario. *Viruses*
404 8:177; doi:10.3390/v8060177.
- 405 Bradford Hill, A. 1965. The environment and disease: association or causation? *Proc. R. Soc. Med.*
406 58:295–300.
- 407 Conedera, M., Tinner, W., Krebs, P., de Rigo, D., and Caudullo, G. 2016. *Castanea sativa* in Europe:
408 distribution, habitat, usage and threats. Pages 78-79. in: *European Atlas of Forest Tree Species*.
409 J. San-Miguel-Ayanz, D. de Rigo, G. Caudullo, T. Houston Durrant, A. Mauri eds.
- 410 Deng, S. and Hiruki, C. 1991. Amplification of 16S rRNA genes from culturable and non-culturable
411 mollicutes. *J. Microbiol. Meth.* 14:53-61.
- 412 Desvignes, J. C. 1992. Characterization of the chestnut mosaic. *Acta Hortic.* 309:353-358.
- 413 Desvignes, J. C., and Lecocq, G. 1995. New knowledges on the chestnut mosaic virus disease. *Acta*
414 *Hortic.* 386:578-584.
- 415 Desvignes, J. C., and Cornaggia, D. 1996. Mosaïque du chataignier: Transmission par le puceron
416 *Myzocallis castanicola*. *Phytoma* 481:39-41.
- 417 Desvignes, J. C. 1999a. Mosaïque du chataignier. Pages 179-181. in: *Maladies à virus des arbres*
418 *fruitiers*. Ctifl Ed. France
- 419 Desvignes, J. C. 1999b. Sweet chestnut incompatibility and mosaics caused by the chestnut mosaic
420 virus (ChMV). *Acta Hortic.* 494:451-454.
- 421 Doyle, J. J., and Doyle, J. L. 1990. Isolation of plant DNA from fresh tissue. *Focus* 12:13–15.
- 422 Fox, A. 2020. Reconsidering causal association in plant virology. *Plant Pathol.* doi:
423 10.1111/ppa.13199
- 424 Gambino, G., Perrone, I., and Gribaudo, I. 2008. A rapid and effective method for RNA extraction
425 from different tissues of grapevine and other woody plants. *Phytochem. Anal.* 19:520-525.

- 426 Geering, A. D. W., Maumus, F., Copetti, D., Choisne, N., Zwickl, D. J., Zytnicki, M., McTaggart, A. R.,
427 Scalabrin, S., Vezzulli, S., Wing, R. A., Quesneville, H. and Teycheney, P. Y 2014. Endogenous
428 florendoviruses are major components of plant genomes and hallmarks of virus evolution. *Nat.*
429 *Comm.* 5:5269
- 430 Gualaccini, G. 1958. Una virosi nuova del castagno. *Boll. Staz Patol. Veg. Roma*, 16:67-75.
- 431 Gundersen, D. E. and Lee, I. M. 1996. Ultrasensitive detection of phytoplasmas by nested-PCR assays
432 using two universal primer pairs. *Phytopathol. Mediterr.* 35:144-151.
- 433 Hohn, T., and Rothnie, H. 2013. Plant pararetroviruses: replication and expression. *Curr. Opin. Virol.*
434 3:621-628.
- 435 Horvath, J., Ecke, I., Gal, T., and Dezcery, M. 1975. Demonstration of virus-like particles in sweet
436 chestnut and oak with leaf deformations in Hungary. *Z. Pflkrankh. Pflsdrrntz.* 82:498-502.
- 437 Kreuze, J. F., Perez, A., Untiveros, M., Quispe, D., Fuentes, S., and Barker, I. 2009. Complete viral
438 genome sequence and discovery of novel viruses by deep sequencing of small RNAs: A generic
439 method for diagnosis, discovery and sequencing of viruses. *Virology* 388:1-7.
- 440 Kumar, S., Stecher, G., and Tamura, K. 2016. MEGA 7: Molecular Evolutionary Genetics Analysis
441 version 7 for bigger datasets. *Mol. Biol. Evol.* 33:1870-1874.
- 442 Martin, R. R., Constable, F., and Tzanetakis, I. E. 2016. Quarantine regulations and the impact of
443 modern detection methods. *Annu. Rev. Phytopathol.* 54:189-205.
- 444 Massart, S., Olmos, A., Jijakli, H., and Candresse, T. 2014. Current impact and future directions of
445 high throughput sequencing in plant virus diagnostics. *Virus Res.* 188:90-96.
- 446 Massart, S., Candresse, T. Gil, J., Lacomme, C., Predajna, L., Ravnikar, M., Reynard, J. S., Rumbou, A.,
447 Saldarelli, P., Škorić, D., Vainio, E. J., Valkonen, J. P. T., Vanderschuren, H., Varveri, C., and Wetzel,
448 T. 2017. A framework for the evaluation of biosecurity, commercial, regulatory, and scientific

- 449 impacts of plant viruses and viroids identified by NGS technologies. *Front. Microbiol.* 8:45. doi:
450 10.3389/fmicb.2017.00045
- 451 Murolo, S., De Miccolis Angelini, R. M., Faretra F., and Romanazzi, G. 2018. Phenotypic and
452 molecular investigations on hypovirulent *Cryphonectria parasitica* in Italy. *Plant Dis.* 102:540-
453 545.
- 454 Prospero, S., Vannini, A., and Vettrai, A. M. 2012. Phytophthora on *Castanea sativa* Mill. (sweet
455 chestnut). *JKI Data Sheets* 6, DOI 10.5073/jkidspdd.2012.006
- 456 Ragozzino, A., and Lahoz, E. 1986. Una malattia virus-simile del castagno in provincia di Avellino.
457 *Giornate di studio sul castagno. Soc. Orti. Italiana* 307-311.
- 458 Reid, K. E., Olsson, N., Schlosser, J., Peng, F., and Lund, S. T. 2006. An optimized grapevine RNA
459 isolation procedure and statistical determination of reference genes for real-time RT-PCR during
460 berry development. *BMC Plant Biol.* 6:26-37.
- 461 Rigling, D., and Prospero, S. 2018. *Cryphonectria parasitica*, the causal agent of chestnut blight:
462 invasion history, population biology and disease control. *Mol. Plant Pathol.* 19:7-20.
- 463 Rott, M., Xiang, Y., Boyes, I., Belton, M., Saeed, H., Kesanakurti, P., Hayes, S., Lawrence, T., Birch, C.,
464 Bhagwat, B., and Rast, H. 2017. Application of Next Generation Sequencing for diagnostic testing
465 of tree fruit viruses and viroids. *Plant Dis.* 101:14891499.
- 466 Shimada, S. 1962. Chestnut Yellows. *Pl. Prot. Tokyo*, 16:253-254.
- 467 Smart, C. D., Schneider, B., Blomquist, C. L., Guerra, L. J., Harrison, N. A., Ahrens, U., Lorenz, K. H.,
468 Seemuller, E., and Kirkpatrick, B. C. 1996. Phytoplasma-specific PCR primers based on sequences
469 of 16S-23S rRNA spacer region. *Appl. Environ. Microbiol.* 62:2988-2993.
- 470 Staginnus, C., Iskra-Caruana, M. L., Lockhart, B., Hohn, T., and Richert-Pöggeler, K. R. 2009.
471 Suggestions for a nomenclature of endogenous pararetroviral sequences in plants. *Arch. Virol.*
472 154:1189-1193.

- 473 Teycheney, P. Y., Geering, A. D. W., Dasgupta, I., Hull, R., Kreuze, J.F., Lockhart, B., Muller, E.,
474 Olszewski, N., Pappu, H., Pooggin, M. M., Richert-Pöggeler, K. R., Schoelz, J. E., Seal, S., Stavelone,
475 L., Umber, M. and ICTV Report Consortium. 2020. ICTV Virus Taxonomy Profile: *Caulimoviridae*.
476 J. Gen. Virol. DOI 10.1099/jgv.0.001497
- 477 Thompson, J. D., Higgins, D. G., and Gibson, T. J. 1994. CLUSTALW: Improving the sensitivity of
478 progressive multiple sequence alignment through sequence weighting, position-specific gap,
479 penalties and weight matrix choice. *Nucleic Acids Res.* 22, 4673-4680.
- 480 Vettraino, M., Vannini, A., Flamini, L., Lagnese, R., Pizzichini, L., Talevi, S., and Fulbright, D. W. 2005.
481 A new transmissible symptomology on sweet chestnut in Italy. *Acta Hort.* 693:547-550.
- 482 Villamor, D. E. V., Ho, T., Al Rwahnih, M., Martin, R. R., and Tzanetakis, I. E. 2019. High Throughput
483 Sequencing for plant virus detection and discovery. *Phytopathology* 109:716-725.
- 484 Wang, Y., Cheng, X., Wu, X., Wang, A., and Wu, X. 2014. Characterization of complete genome and
485 small RNA profile of pagoda yellow mosaic virus, a novel badnavirus in China. *Virus Res.* 188:103-
486 108.
- 487 Zerbino, D. R., and Birney, E. 2008. Velvet: algorithms for de novo short read assembly using de
488 Bruijn graphs. *Genome Res.* 18:821-829.

489 **TABLE 1.** Percentage of identity between the ORF3 region encoding the Reverse Transcriptase -
490 Rnase H^a of chestnut mosaic virus (ChMV) isolate FRlc1224A and the corresponding genomic regions
491 of the isolate ITumito39 and of the most closely related members of the genus *Badnavirus*

Virus ^b	Nucleotide identity (%)	Amino acid identity (%)
ChMV ITumito39	91.9%	97.8%
RYNV	65.1%	71.6%
GVBV	64%	68.9%
GVCV	66.8%	72.6%
BLRaV	68.4 %	72.6%
WBV1	68.2%	71.6%
PYMaV	67.7%	71.9%

492 ^a This region is the one typical used for taxonomic discrimination in the family *Caulimoviridae*
493 (Teycheney et al. 2020)

494 ^b Acronyms used: RYNV, rubus yellow net virus; GVBV, gooseberry vein banding virus; GVCV,
495 grapevine vein-clearing virus; BLRaV, birch leafroll-associated virus; WBV1, wisteria badnavirus 1;
496 PYMaV, pagoda yellow mosaic-associated virus.

497

498

499 **TABLE 2.** Number and percentage of chestnut mosaic virus-infected plants regarding the plot, the
500 *Castanea* species sampled, and the symptomatology

Origin of the sampled plants	Infected/total plants (%)	Infected/asympto matic plants (%)	Infected/sympto matic plants (%)	Infected/plants with atypical symptoms (%)
France (overall)	65/95 (68.4%)	23/47 (48.9%)	38/38 (100%)	4/10 (40%)
Plot A	13/23 (56.5%)	6/15 (40%)	6/6 (100%)	1/2 (50%)
Plot E	48/66 (72.7%)	15/28(53.6%)	30/30 (100%)	3/8 (37.5%)
Plot Port	4/6 (66.6%)	2/4 (50%)	2/2 (100%)	na ^b
<i>C. crenata</i>	5/6 (83%)	1/1 (100%)	4/4 (100%)	0/1
<i>C. sativa</i>	30/43 (70%)	11/20 (55%)	18/18 (100%)	1/5 (20%)
<i>C. mollissima</i>	13/14 (93%)	1/2 (50%)	10/10 (100%)	2/2 (100%)
Hybrid ^a	17/32 (53.1%)	10/24 (41.6%)	6/6 (100%)	1/2 (50%)
Italy <i>Castanea</i> <i>sativa</i>	57/70 (81.5%)	5/10 (50%)	52/60 (87%)	na ^b

501 ^a Interspecific hybrids between *C. crenata*, *C. mollissima* and *C. sativa*

502 ^b not applicable

503

504 CAPTIONS FOR FIGURES

505 **Fig. 1. Symptoms of chestnut mosaic disease on various hosts.** (A) Isolate LC1224H: Red oak
506 (*Quercus rubra*) graft-inoculated with a diseased source; (B) Isolate FRlc1224A: 'Maraval' Ca 74
507 graft-inoculated with LC1224H; (C) Isolate ITumito39: symptomatic leaves from cv Marrone grafted
508 onto *Castanea sativa*; (D) non-inoculated *Q. rubra*; (E) non-inoculated 'Maraval' Ca 74; (F)
509 Asymptomatic leaves from cv Marrone grafted onto *C. sativa*.

510

511 **Fig. 2: Schematic representation of the genomic organization of the chestnut mosaic virus.** The
512 tRNA binding site is indicated and defines the position 1 on the genome. The three open reading
513 frames (ORFs) are shown as grey arrows, as well as their position in parentheses. Five conserved
514 motifs are identified in the ORF3 polyprotein: MP, Viral movement protein (pfam01107); ZnF, Zinc
515 finger (pfam00098); RVP, Retroviral aspartyl protease (pfam00077); RT, Reverse transcriptase
516 (cd01647); RH, Ribonuclease H (cd09274)

517

518 **Fig. 3. Phylogenetic tree reconstructed using the complete genome sequences of badnavirus**
519 **members.** Virus names as well as GenBank accession numbers are indicated. The tree was
520 reconstructed using the neighbor-joining method, and randomized bootstrapping was performed
521 to evaluate the statistical significance of branches (1,000 replicates). Bootstrap values above 70%
522 are shown. The scale bar represents 5% nucleotide divergence between sequences. The groups as
523 defined by Wang et al. (2014) are indicated. Chestnut mosaic virus isolates determined in this work
524 are indicated by black triangles. Rice tungro bacilliform virus was used as outgroup.

525

526 **Fig. 4. Unrooted neighbor-joining phylogenetic tree reconstructed from the alignment of**
527 **concatenated nucleotide sequences related to chestnut mosaic virus detected by datamining of**
528 **publicly available transcriptomic chestnut data.** Randomized bootstrapping was performed to
529 evaluate the statistical significance of branches (1,000 replicates). Bootstrap values above 70% are
530 shown. The scale bar represents 10% nucleotide divergence between sequences.

531

532 **Fig. 5. Detection of chestnut mosaic virus in various samples by PCR using primers pairs Ch-**
533 **Bad1466F/1800R (A) and Ch-Bad5860F/6109R (B).** Lane 1: LC1224F; Lane 2: LC1224H; Lane 3:
534 FRlc1224A; Lane 4: T32018; Lane 5: LCA552; Lane 6: LCA584; Lane 7: 'Maraval' Ca 74 non-inoculated
535 plant; Lane 8: *Quercus rubra* non-inoculated plant; Lane 9: no template; L: molecular weight marker.
536 Horizontal bars on the left of the figure indicate the size of the amplification products. The isolates
537 are listed in Table S1.

538

539 **e-EXTRA FIGURE CAPTIONS AND TABLE TITLES**

540

541 **Supplementary Fig. S1. Phylogenetic tree reconstructed using the ORF3-deduced amino acid**
542 **sequences of badnavirus members.** Virus names as well as GenBank accession numbers are
543 indicated. The tree was reconstructed using the neighbor-joining method, and randomized
544 bootstrapping was performed to evaluate the statistical significance of branches (1,000 replicates).
545 Bootstrap values above 70% are shown. The scale bar represents 10% amino acid divergence
546 between sequences. Chestnut mosaic virus isolates determined in this work are indicated by black
547 triangles. Rice tungro bacilliform virus was used as outgroup.

548

549 **Supplementary Fig. S2. Leaves of the sample 20893. (A) rootstock leaves (*Castanea sativa*); (B)**
550 **cultivar leaves**

551

552 **Supplemental Fig. S3. Unrooted neighbor-joining phylogenetic trees reconstructed from the**
553 **alignment of nucleotide sequences of the PCR fragments targeted partial RT-RnaseH domain (A)**
554 **and partial MP domain (B) obtained from a range of chestnut mosaic virus isolates (listed in Table**
555 **S1).** Randomized bootstrapping was performed to evaluate the statistical significance of branches
556 (1,000 replicates). Bootstrap values above 70% are shown. The scale bar represents 5% (A) or 10%
557 (B) nucleotide divergence between sequences.

558

559 **Supplemental Table S1. List of chestnut samples used in the present study together with relevant**
560 **ChMV accession numbers**

561

562 **Supplemental Table S2. Primers used for genome circularity confirmation and for molecular**
563 **detection of chestnut mosaic virus by PCR**

564

565 **Supplemental Table S3. Datamining of publicly available chestnut HTS data for chestnut mosaic**
566 **virus sequences**






Fig. 2: Schematic representation of the genomic organization of the chestnut mosaic virus. The tRNA binding site is indicated and defines the position 1 on the genome. The three open reading frames (ORFs) are shown as grey arrows, as well as their position in parentheses. Five conserved motifs are identified in the ORF3 polyprotein: MP, Viral movement protein (pfam01107); ZnF, Zinc finger (pfam00098); RVP, Retroviral aspartyl protease (pfam00077); RT, Reverse transcriptase (cd01647); RH, Ribonuclease H (cd09274)

338x190mm (96 x 96 DPI)





Fig. 3. Phylogenetic tree reconstructed using the complete genome sequences of badnavirus members. Virus names as well as GenBank accession numbers are indicated. The tree was reconstructed using the neighbor-joining method, and randomized bootstrapping was performed to evaluate the statistical significance of branches (1,000 replicates). Bootstrap values above 70% are shown. The scale bar represents 5% nucleotide divergence between sequences. The groups as defined by Wang et al. (2014) are indicated. Chestnut mosaic virus isolates determined in this work are indicated by black triangles. Rice tungro bacilliform virus was used as outgroup.

190x338mm (96 x 96 DPI)







Fig. 5. Detection of chestnut mosaic virus in various samples by PCR using primers pairs Ch-Bad1466F/1800R (A) and Ch-Bad5860F/6109R (B). Lane 1: LC1224F; Lane 2: LC1224H; Lane 3: FRlc1224A; Lane 4: T32018; Lane 5: LCA552; Lane 6: LCA584; Lane 7: 'Maraval' Ca 74 non-inoculated plant; Lane 8: *Quercus rubra* non-inoculated plant; Lane 9: no template; L: molecular weight marker. Horizontal bars on the left of the figure indicate the size of the amplification products. The isolates are listed in Table S1.

338x190mm (96 x 96 DPI)

Fig. S1



0.1

Fig. S2




Fig. S3A





Fig. S3B



1 **Supplementary Table S1. List of chestnut samples used in the present study together with relevant ChMV**
 2 **accession numbers**

Isolate ID ^a	Chestnut Accession ID	Chestnut Accession name	Species ^b	Rootstock	Country	Plot ^c	Accession numbers ^d	Use ^e
20885	Ca04x Ca03	-	<i>C. crenata</i>	Not graft	France	A	MT339547; MT339503	Incidence
20887	Ca 715	'Merle'	<i>C. sativa</i>	G1. Ca15	France	A	MT339548; MT339504	Incidence
20890	Ca 663	'Trigueira'	<i>C. sativa</i>	unknown	France	Port	MT339549; MT339505	Incidence
20891	Ca 664	'Longal Special'	<i>C. sativa</i>	Ca 74	France	Port	MT339550; MT339506	Incidence
<u>20895</u>	Ca 43	'Vignols'	<i>C. sativa x C. crenata</i>	Ca 116	France	A	MT339551; MT339507	Incidence
20899	Ca 599	'Ibuki'	<i>C. crenata</i>	G1.Ca 02	France	A	MT339552; MT339508	Incidence
20901	Ca 598	'Riheï'	<i>C. crenata x C. mollissima</i>	G1. Ca102	France	A	MT339553; MT339509	Incidence
20902	Ca 75	'Fertil'	<i>C. mollissima</i>	Ca 07	France	A	MT339554; MT339510	Incidence
20904	Tree- A68Ks ⁱ	-	<i>C. mollissima</i>	Not graft	France	A	MT339555; MT339511	Incidence
20907	Ca 118	'Marlhac'	<i>C. sativa x C. crenata</i>	Not graft	France	A	MT339556; MT339512	Incidence
20908	Ca 124	'Maridonne'	<i>C. sativa x C. crenata</i>	G1.Ca116	France	A	MT339557; MT339513	Incidence
20909	Ca 125	'Bouche de Bétizac'	<i>C. sativa x C. crenata</i>	Ca 74	France	A	MT339558; MT339514	Incidence
20910	Ca 860	-	hybrid SC	unknown	France	E	MT339559; MT339515	Incidence
20911	Ca 860	-	hybrid SC	unknown	France	E	MT339560; MT339516	Incidence
20913	Ca 844 ⁱ	-	<i>C. mollissima</i>	Not graft	France	E	MT339561; MT339517	Incidence
20917	Ca 837 ^h	-	<i>C. mollissima</i>	Not graft	France	E	MT339562; MT339518	Incidence
20923	Ca 846 ^h	-	<i>C. mollissima</i>	Not graft	France	E	MT339563; MT339519	Incidence
20925	Ca 741	'Dauphine'	<i>C. sativa</i>	Ca 07	France	E	MT339564; MT339520	Incidence
20926	Ca 665	'Longal'	<i>C. sativa</i>	Ca 07	France	E	MT339565; MT339521	Incidence
20930	Ca 860	-	hybrid SC	unknown	France	E	MT339566; MT339522	Incidence
20932	Ca 564	'Ipharra 16'	<i>C. crenata</i>	Ca 07	France	E	MT339567; MT339523	Incidence
<u>20935</u>	Ca 576	'Sardonne'	<i>C. sativa</i>	Ca 07	France	E	MT339568; MT339524	Incidence
20937	Ca 138	'Marron de Redon'	<i>C. sativa</i>	Ca 07	France	E	MT339569; MT339525	Incidence
20943	unknown	-	<i>C. sativa</i>	Ca 74	France	E	MT339570; MT339526	Incidence
20944	Ca 520	'Montagne'	<i>C. sativa</i>	Ca 07	France	E	MT339571; MT339527	Incidence
20945	Ca 106	'Marron Comballe'	<i>C. sativa</i>	Ca 07	France	E	MT339572; MT339528	Incidence

20947	Ca 111	'Marron de Lyon'	<i>C. sativa</i>	Ca 07	France	E	MT339573; MT339529	Incidence
20951	Ca 126	-	<i>C. sativa x C. crenata</i>	Ca 07	France	E	MT339574; MT339530	Incidence
20956	Ca 105	'Sardonne'	<i>C. sativa</i>	Ca 07	France	E	MT339575; MT339531	Incidence
20959	Ca 03	-	<i>C. crenata</i>	Ca 74	France	E	MT339576; MT339532	Incidence
20960	Ca 127	-	<i>(C. crenata x C. sativa) x C. mollissima</i>	Ca 07	France	E	MT339577; MT339533	Incidence
20961	Ca 127	-	<i>(C. crenata x C. sativa) x C. mollissima</i>	Ca 07	France	E	MT339578; MT339534	Incidence
20964	Ca 112	'Bournette'	<i>C. crenata x C. sativa</i>	Ca 07	France	E	MT339579; MT339535	Incidence
20965	Ca 501	-	<i>C. sativa</i>	Ca 07	France	E	MT339580; MT339536	Incidence
20966	Ca 501	-	<i>C. sativa</i>	Ca 07	France	E	MT339581; MT339537	Incidence
20967	Ca 665	'Longal'	<i>C. sativa</i>	Ca 07	France	E	MT339582; MT339538	Incidence
20968	Ca 48	'Précoce Migoule'	<i>(C. crenata x C. sativa)</i>	Ca 07	France	E	MT339583; MT339539	Incidence
20971	Ca 151	'Bouche Rouge'	<i>C. sativa</i>	Ca 07	France	E	MT339584; MT339540	Incidence
20975	unknown	-	<i>C. sativa x C. crenata</i>	Ca 07	France	E	MT339585; MT339541	Incidence
20889	unknown	-	<i>C. sativa</i>	unknown	France	Port	na	Incidence
20893	Ca 106	'Marron Comballe'	<i>C. sativa</i>	unknown	France	Port	na	Incidence
20898	Ca 105	'Sardonne'	<i>C. sativa</i>	G1. Ca394	France	A	na	Incidence
20900	Ca 501	-	<i>C. sativa</i>	G1.Ca486	France	A	na	Incidence
20903	Ca744	'Quing Zha'	<i>C. mollissima</i>	G1.Moll	France	A	na	Incidence
20912	Ca 845 ^h	-	<i>C. mollissima</i>	Not graft	France	E	na	Incidence
20914	Ca 843 ^h	-	<i>C. mollissima</i>	Not graft	France	E	na	Incidence
20915	Ca 842 ^h	-	<i>C. mollissima</i>	Not graft	France	E	na	Incidence
<u>20918</u>	Ca 838 ^h	-	<i>C. mollissima</i>	Not graft	France	E	na	Incidence
<u>20919</u>	Ca 839 ^h	-	<i>C. mollissima</i>	Not graft	France	E	na	Incidence
20920	Ca 840 ^h	-	<i>C. mollissima</i>	Not graft	France	E	na	Incidence
20921	Ca 841 ^h	-	<i>C. mollissima</i>	Not graft	France	E	na	Incidence
20922	Ca 74	'Maraval'	<i>C. crenata x C. sativa</i>	Not graft	France	E	na	Incidence
20924	Ca 825	'Précoce de Besse'	<i>C. sativa</i>	Ca 07	France	E	na	Incidence
20940	Ca 393	'Marron de Chevanceaux'	<i>C. sativa</i>	Ca 07	France	E	na	Incidence
20941	Ca 460	-	<i>(C. crenata x C. sativa) x (C. crenata x C. sativa)</i>	Ca 74	France	E	na	Incidence
29942	Ca 511	'Marrone di Chiusa Pesio'	<i>C. sativa</i>	Ca 74	France	E	na	Incidence
20948	Ca 116	-	<i>C. sativa x C. crenata</i>	Ca 74	France	E	na	Incidence
20949	Ca 135	'Précoce des Vans'	<i>C. sativa</i>	Ca 74	France	E	na	Incidence
20952	Ca 521	-	<i>C. crenata</i>	Ca 07	France	E	na	Incidence

20955	Ca 399	'Comballe'	<i>C. sativa</i>	Ca 07	France	E	na	Incidence
20962	Ca 639	-	<i>C. sativa</i>	Ca 07	France	E	na	Incidence
20963	Ca 730	'Sauvage Marron'	<i>C. sativa</i>	Ca 07	France	E	na	Incidence
20969	Ca 381	-	<i>C. sativa</i>	Ca 07	France	E	na	Incidence
20970	Ca 381	-	<i>C. sativa</i>	Ca 07	France	E	na	Incidence
20972	Ca 151	'Bouche Rouge'	<i>C. sativa</i>	Ca 07	France	E	na	Incidence
FRIC1224A	Ca 74	'Maraval'	<i>C. crenata x C. sativa</i>	na	France	Ctifl	MT269853 (complete genome)	HTS
LC1224H	na	na	<i>Q. rubra</i>	na	France	Ctifl	MT339586; MT339542	CRA
T32018	Ca 74	Maraval	<i>C. crenata x C. sativa</i>	na	France	Ctifl	MT339587; MT339543	CRA
LCA552	Ca 74	Maraval	<i>C. sativa</i>	na	France	Ctifl	MT339588; MT339544	CRA
LCA584	Ca 74	Maraval	<i>C. sativa</i>	na	France	Ctifl	MT339589; MT339545	CRA
LC1224F	Ca 74	Maraval	<i>C. crenata x C. sativa</i>	na	France	Ctifl	MT339590; MT339546	CRA
ITUmito39	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	MT261366 (complete genome)	HTS
ITCh2	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	MT270674; MT270683	Incidence
ITCh3	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	MT270667 ^f	Incidence
ITCh4	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	MT270668; MT270684	Incidence
ITCh8	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	MT270685 ^g	Incidence
ITCh10	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	MT270669; MT270678	Incidence
ITCh11	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	MT270664; MT270679	Incidence
ITCh12	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	MT270665; MT270680	Incidence
ITCh14	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	MT270666; MT270681	Incidence
ITCh20	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	MT270670; MT270675	Incidence
ITChAs3	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	MT270671; MT270676	Incidence
ITChAs4	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	MT270672 ^f	Incidence
ITChAs5	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	MT270677 ^g	Incidence
ITChAs8	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	MT270673; MT270682	Incidence
ITChAs9	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh1	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh5	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh6	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh7	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh9	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh13	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh14	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh15	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh20	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh21	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence

ITCh22	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh23	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh24	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh25	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh27	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh28	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh29	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh30	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh31	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh32	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh33	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh35	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh36	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh37	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh38	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh39	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh41	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh42	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh43	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh44	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh46	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh47	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh48	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh49	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh50	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh51	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh52	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh54	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh55	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh56	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh58	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh59	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence
ITCh60	unknown	Marrone	<i>C. sativa</i>	na	Italy	I	na	Incidence

^a Isolates from asymptomatic trees are indicated in bold; isolates with doubtful symptoms are underlined; isolates from asymptomatic cultivars with typical symptoms visible on rootstock regrowths are indicated in bold italic.

^b; hybrid SC : hybrid between *C. sativa* and *C. crenata*, regardless the knowledge about which are the female and male parents.

^c : Five plots have been sampled (A, Port, E, I, Ctifl)

^d : Accession numbers are relative to the sequences obtained with both PCR detection assays (Ch-Bad-1466F/Ch-Bad-1800R and Ch-Bad-5860F/Ch-Bad-6109R)

^e: Isolates were included either in the HTS analysis, or in the incidence analysis, or in the causal relationship analysis (CRA)

^f: Only the fragment amplified with Ch-Bad-1466F/Ch-Bad-1800R was sequenced

^g: Only the fragment amplified with Ch-Bad-5860F/Ch-Bad-6109R was sequenced

^h 'Mengshankui' (*C. mollissima* cultivar) seedling

ⁱ *C. mollissima* seedling

Supplemental TABLE S2. Primers used for genome circularity confirmation and for molecular detection of chestnut mosaic virus by PCR

Primer	Nucleotide sequence (5' to 3')	Genome coordinates	Amplicon size	Isolates detected
Ch-Bad-6976F	CCCGAGCCATTTAC ACTTCACAAC	6,976-6,999	436 nt	FRlc1224A
Ch-Bad-252R	TCACTCATCTACCTC ACACGCTC	252-230		
Ch-Bad-6481F	GAAATTGAATTGGA AGGAAGA	6,481-6,501	1,007 nt	ITumito39
Ch-Bad-325R	TCAGATCAGCAAAC TCGAAC	344-325-		
Ch-Bad-1466F	TATCAGCACTACAG TGAACAACC	1,466-1,488	335 nt	
Ch-Bad-1800R	GTCATGACGCAAAC TTGGAA	1,800-1,781		polyvalent
Ch-Bad-5860F	AGTATGTAAATGG GCACCGTTC	5,857-5,878	232 nt	
Ch-Bad-6109R	GTTGATCCATCGCA CTCTTG	6,109-6,090		polyvalent

1 **Supplemental TABLE S3.** Datamining of publicly available chestnut HTS data for chestnut mosaic
 2 virus sequences

Type of data ^a	Dataset	<i>Castanea</i> species/cultivar	Country	Length of assembled sequence (nt)	% mapped reads ^b	% nt identity with FRlc1224A
EST	GO917001	<i>C. mollissima</i> , BX3, clone Vanuxum	USA	436	na	99%
WGA	JRKL01079565	<i>C. mollissima</i> , cv Vanuxem	USA	7,164 (full length)	na	99%
RNA-Seq	SRX4015368	<i>C. mollissima</i>	USA	7,161 (full length)	0.004%	97.4%
RNA-Seq	SRX001805	<i>C. mollissima</i> , cv Vanuxem	USA	5,889 (scaffold)	0.04%	95-100%
GBS	SRX5144434	<i>C. mollissima</i>	China	5,933 (scaffold)	0.06%	98.7% (average)
GSB	SRX5144449	<i>C. mollissima</i> , clone Vanuxem	China	6,396 (scaffold)	0.04%	97.5% (average)
GBS	SRX5145051	<i>C. mollissima</i> , cv Vanuxem	China	5,873 (scaffold)	0.02%	99% (average)
GBS	SRX5145050	<i>C. mollissima</i>	China	6,680 (scaffold)	0.01%	99% (average)
GBS	SRX51445044	<i>C. mollissima</i>	China	6,183 (scaffold)	0.01%	97.4% (average)
GBS	SRX51444621	<i>C. mollissima</i>	China	5,875 (scaffold)	0.02%	95.6% (average)
GBS	SRX51444448	<i>C. mollissima</i>	China	2,948 (scaffold)	0.04%	99% (average)
GBS	SRX51444446	<i>C. mollissima</i>	China	5,873 (scaffold)	0.04%	99% (average)

GBS	SRX5825095	<i>C. dentata</i>	USA	1,857 (scaffold)	0.0036%	89.2% (average)
GBS	SRX5145077	<i>C. dentata</i>	USA	6,068 (scaffold)	0.01%	90.9% (average)

3 ^a EST: expressed sequence tags; WGA: whole genome assembly; GBS: genotyping by sequencing

4 ^b Reads mapped to the genomic sequence of chestnut mosaic virus (French isolate)

