



HAL
open science

Interacting effects of land use type, soil microbes and plant traits on aggregate stability

Luis Merino-Martín, Alexia Stokes, Hyun Gweon, Lur Moragues-Saitua, Siobhan Staunton, Claude Plassard, Anna Oliver, Yves Le Bissonnais, Robert Griffiths

► To cite this version:

Luis Merino-Martín, Alexia Stokes, Hyun Gweon, Lur Moragues-Saitua, Siobhan Staunton, et al.. Interacting effects of land use type, soil microbes and plant traits on aggregate stability. *Soil Biology and Biochemistry*, 2021, 154, pp.108072. 10.1016/j.soilbio.2020.108072 . hal-03040299

HAL Id: hal-03040299

<https://hal.inrae.fr/hal-03040299>

Submitted on 18 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Interacting effects of land use type, soil microbes and plant traits on aggregate stability

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Merino-Martín, L., Stokes, A., Gweon, H. S. ORCID: <https://orcid.org/0000-0002-6218-6301>, Moragues-Saitua, L., Staunton, S., Plassard, C., Oliver, A., Le Bissonnais, Y. and Griffiths, R. I. (2020) Interacting effects of land use type, soil microbes and plant traits on aggregate stability. *Soil Biology & Biochemistry*. 108072. ISSN 0038-0717 doi: <https://doi.org/10.1016/j.soilbio.2020.108072> Available at <https://centaur.reading.ac.uk/94779/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.1016/j.soilbio.2020.108072>

To link to this article DOI: <http://dx.doi.org/10.1016/j.soilbio.2020.108072>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 **Interacting effects of land use type, soil microbes and plant traits on aggregate stability**

2

3 **Author list / address list:**

4 Luis Merino-Martín^{1,2,3}, Alexia Stokes¹, Hyun S. Gweon², Lur Moragues-Saitua⁴, Siobhan Staunton⁵,
5 Claude Plassard⁵, Anna Oliver², Yves Le Bissonnais⁶, Robert I. Griffiths².

6 ¹ *Amap, University Montpellier, INRAE, Cirad, Cnrs, Ird, Montpellier, France.*

7 ² *Centre for Ecology & Hydrology, Crowmarsh Gifford, Wallingford, Oxfordshire, United Kingdom.*

8 ³ *CEFE, Univ Montpellier, CNRS, Univ Paul Valéry Montpellier 3, EPHE, IRD, Montpellier, France.*

9 ⁴ *Department of Conservation of Natural Resources, NEIKER-Tecnalia, Derio, Spain.*

10 ⁵ *Eco&Sols, INRAE, CIRAD, IRD, Institut Agro, Univ Montpellier, Montpellier, France.*

11 ⁶ *LISAH, INRAE, IRD, Institut Agro, University Montpellier, Montpellier, France.*

12 *Corresponding author. Tel.: +33(0)789288168. E-mail address: luismerinomartin@gmail.com (L.
13 Merino-Martín).

14

15 For submission to *Soil Biology and Biochemistry*.

16 **Article type:** Full paper.

17 **Word count:**

Total word count (excluding summary, highlights, references and legends):	10329	No. of figures:	8
Summary:	427	No. of Tables:	6
Introduction:	1396	No of Supporting Information files:	6 tables 5 figures 1 research note
Materials and methods:	4497		
Results:	2013		
Discussion:	1989		
Conclusions:	325		
Acknowledgements:	96		

18

19

20 **Summary**

21 Soil aggregates are critical to soil functionality, but there remain many uncertainties with respect to
22 the role of biotic factors in forming aggregates. Understanding the interacting effects of soil, land use
23 type, vegetation and microbial communities is a major challenge that needs assessment in both field
24 and controlled laboratory conditions, as well as in bulk and rhizosphere soils. To address these effects
25 and their feedbacks, we first examined the influence of soil, root and litter characteristics along a land
26 use gradient (ancient woodland, secondary woodland, grassland, pasture and arable land) on microbial
27 community structure (in both bulk and rhizosphere soil), as well as on aggregate stability. Then, we
28 performed an inoculation experiment where we extracted soil columns from the arable and secondary
29 woodland and used a third unstructured loamy soil as a control. We sterilized these three soils to
30 remove microbial communities, and then either inoculated the tops of sterilized soil columns with soil
31 from the secondary woodland or the arable field sites. Control columns of all soil types were not
32 inoculated. In a fully-crossed design, we planted two species possessing distinct root system
33 morphological traits: *Brachypodium sylvaticum* (fibrous system with many thin and fine roots) and
34 *Urtica dioica* (taproot system with few fine roots). After four months, microbial communities (in bulk
35 and rhizospheric soil) and aggregate stability were measured, along with root traits. In both the field
36 and laboratory experiments, bacterial (16S) and fungal (ITS) biodiversity was determined using high
37 throughput sequencing. In the field study we found that: i) there were strong relationships between
38 aggregate stability and microbial community composition that were driven by land use, ii) the
39 relationship between aggregate stability along the land use gradient and the trophic nature of bacterial
40 communities was not significant, but that certain soil, root and litter parameters shaped bacterial
41 phyla, with oligotrophic bacteria conditioned by the rhizosphere niche, and copiotrophic phyla more
42 dependent on bulk soil conditions, iii) land use gradient (from woodland to arable), reduced the
43 relative abundance of saprotrophic and ectomycorrhizal fungi with an increase in the relative
44 abundance of Ascomycota and a reduction in the relative abundance of Basidiomycota. In the
45 laboratory experiment we found that: i) the inoculation of sterilized soils with soils from the field
46 significantly increased aggregate stability in control soil that was initially poorly structured, ii) the
47 effects of inoculation on aggregate stability were similar when either secondary woodland or arable

48 soils were used as inoculums and iii) these effects were affected significantly by root length density.
49 Our results show that microbial communities influence soil structure and that bacterial communities
50 are intimately associated to rhizospheric conditions and root traits (of which root length density was
51 the most pertinent).

52 **Key words:** bacteria, fungi, glomalin, ergosterol, ITS, root traits, root systems, soil structure, 16S.

53 **1. Introduction**

54 Soil structure is an essential factor driving the success of many ecosystem services worldwide
55 (Adhikari and Hartemink, 2016), but most soils have been severely degraded in recent decades, due
56 largely to anthropogenic pressures (Jie et al., 2002; Peng et al., 2015). Therefore, the capacity of soil
57 to retain water, maintain biodiversity, sustain agriculture and resist flooding, erosion and landslides is
58 compromised, unless we can find mechanisms by which soil structure can be restored rapidly. Soil is
59 composed of a solid phase of particles and a pore phase that provides pathways for the transport of
60 water, nutrients and gases, as well as habitats for microorganisms and fauna. The spatial configuration
61 of soil particles and pores is arranged in aggregates and their stability is an important aspect of soil
62 structure. Apart from soil texture (Bissonnais and Arrouays, 1997), diverse biotic and abiotic factors
63 drive the rate of aggregation and its stability, e.g. microorganisms (Tisdall, 1994; Chenu and Sotzky,
64 2002; Lehmann et al., 2017), fauna (Lee and Foster, 1991; Ayuke et al., 2011), litter and root traits
65 (Baumert et al., 2018; Poirier et al., 2018), environmental variables (e.g., tillage, freeze-thaw, wetting
66 and drying cycles) (Lavee et al., 1996; Six et al., 1998; Gispert et al., 2013b) and inorganic binding
67 agents (e.g., clay minerals, oxides and polyvalent cations such as calcium) (Denef and Six, 2005).
68 Although many studies have focused on how individual factors modify soil aggregation and stability,
69 very few have considered combinations of driving factors and their effects (Six et al., 2004; Baumert
70 et al., 2018), particularly complex biotic factors. However, a better understanding of these interactions
71 will improve soil restoration and the production of ecosystem services (Lavelle et al., 2020).

72 The importance of biotic factors for soil aggregation, and in particular soil microorganisms, was first
73 suggested by Tisdall (1994) and has since been recognized worldwide (e.g. Degens, 1997; Chotte,
74 2005; Gupta and Germida, 2015). In a global meta-analysis, Lehmann et al. (2017) found a positive
75 effect of soil biota on aggregation, with bacteria and fungi being generally more important for
76 aggregation than other groups. Good soil structure is usually associated with high microbial biomass
77 (Degens, 1997) and activity (Cui and Holden, 2015), mainly through the chemical binding action of
78 microbial compounds such as bacterial extracellular polymeric substances (EPS) or soil proteins, or
79 through the physical enmeshment of soil particles by fungal hyphae (Oades and Waters, 1991; Rillig

80 and Mummey, 2006). Rillig et al. (2002a,b) attributed the positive effect of plant roots on aggregation
81 to the excretion of a protein by arbuscular mycorrhizal fungi (AMF) called glomalin. This protein is
82 operationally defined by its extraction method and belongs to a group of proteins after called
83 glomalin-related soil proteins (GRSPs) that are no longer believed to be solely of fungal origin.
84 Ergosterol is the most abundant sterol in fungal cell membranes and since fungal hyphae enmeshment
85 is a major factor for the formation of macroaggregates, it should be related to aggregate stability. Yet,
86 the feedback mechanisms between aggregate formation and stability and the influence of microbial
87 communities are not fully understood and identifying the relationships between changes in microbial
88 community structure and GRSP and ergosterol are key for understanding soil aggregation processes.

89 Root system morphology can influence soil aggregation through changes in the distribution of carbon
90 inputs into soil (Carter et al., 1994; Degens, 1997). Fine roots have direct effects on macroaggregate
91 stability through the production of mucilage and exudates and indirect effects through the promotion
92 of hyphal growth and the release of root-derived particulate organic matter (POM) that stimulate
93 microbial activity (Morel et al., 1991; Miller and Jastrow, 1992; Brax et al., 2020). Consequently,
94 plants with fibrous root systems, that have numerous, evenly distributed thin and fine roots, may have
95 more available mucilage/exudates, as well as more easily degradable POM, than plants with taproot
96 systems. Taproot systems generally comprise one large central root and less thin or fine roots (Miller
97 and Jastrow, 1992; Degens, 1997), thus the potential input of total carbon into soil is reduced. We ask
98 therefore, if root system morphology, and individual root traits such as diameter, length and carbon
99 and nitrogen content (Saleem et al., 2018), influence the composition of microbial communities and
100 so can modify macroaggregate formation and stability.

101 Soil aggregates are important for microbial community ecology, evolution and for microbially
102 mediated cycles (Goebel et al., 2009; Rillig et al., 2016; Rillig et al., 2017). The heterogeneity of
103 pores and water films on soil particles leads to a diversity of microhabitats and gradients of abiotic
104 traits (e.g. nutrients and pH) at microscales (Or et al., 2007). Soil carbon type and availability
105 provides strong selective pressures for the different lifestyle strategies among copiotrophs (common
106 in carbon-rich environments) and oligotrophs (present in habitats of low carbon flux) (Fierer et al.,

107 2007; Eilers et al., 2010; Trivedi et al., 2017). Trivedi et al. (2017) showed that aggregate size
108 modulated the effects of management practices on soil carbon, with oligotrophs (e.g. Acidobacteria,
109 Chloroflexi and Verrucomicrobia) inhabiting microaggregates containing higher proportions of
110 recalcitrant carbon and copiotrophs (e.g. α and β Proteobacteria, Bacteroidetes and Actinobacteria)
111 living in macroaggregates with more labile carbon. Hence, soils with high macroaggregate stability
112 should be dominated by copiotrophs. However, aggregate size is not the only factor influencing the
113 abundance of copiotrophs and oligotrophs. Lauber et al. (2009) and Carbonetto et al. (2014) showed
114 that agro-systems had a higher abundance of copiotrophs than in forests, probably because of greater
115 nitrogen fertilization stimulating copiotrophic communities (Fierer et al., 2012). Complex interactions
116 in soil can influence the abundance of different lifestyle strategies, e.g., rhizosphere communities are
117 enriched with rhizodeposits (Dennis et al., 2010) and as a consequence, increase copiotroph
118 abundance compared to bulk soil (Peiffer et al., 2013; Lladó and Baldrian, 2017). However, studies on
119 how root and litter traits affect the abundance of these trophic modes in the different soil niches are
120 notably lacking (Saleem et al., 2018), even though these scaled down studies are still needed to
121 decipher the rules of rhizosphere community assembly (Brunel et al., 2020). The disturbance of soils
122 through management also decreases soil fungal diversity, increasing the prevalence of generalist taxa,
123 reducing the relative abundances of saprotrophic and ectomycorrhizal fungi and increasing the
124 abundance of pathogenic fungi (Mueller et al., 2016; Marín et al., 2017). Managed land use types also
125 possess more Ascomycota and less Basidiomycota compared to undisturbed natural ecosystems (de
126 Castro et al., 2008; Mueller et al., 2016). Here, we ask if differences in the abovementioned bacterial
127 and fungal trophic modes are due mainly to differences in aggregate formation, soil chemical
128 characteristics or plant traits, along a land use gradient.

129 Since the complex feedbacks between aggregate formation and microbial communities are not fully
130 understood, studies that integrate field observation with experimental testing are needed. Here, we
131 performed two studies to determine the combined effects of soil type, root system morphological traits
132 and microbial communities on soil aggregation and stability. We first surveyed these responses in the
133 field across a gradient of different land use types (grassland, pasture, arable land, secondary woodland

134 and ancient woodland) and hypothesized (hypothesis 1) that there are strong relationships between
135 soil aggregate stability and microbial community structure that are driven by land use. Additionally,
136 we hypothesize (hypothesis 2) that soils of higher aggregate stability will have characteristic
137 communities comprising increased bacterial copiotrophs and increased relative abundance of
138 saprotrophic and ectomycorrhizal fungi related to a decrease in the relative abundance of Ascomycota
139 and an increase of Basidiomycota. However, this relationship may be influenced by other edaphic
140 factors (e.g. soil nitrogen content) and root and litter traits, or soil habitat (e.g., with the rhizosphere
141 niche favouring copiotrophic bacterial communities).

142 Then, to experimentally test interactive effects of plants and microbes on aggregate stability, we
143 performed a controlled experiment where we sterilized soil samples from arable land and secondary
144 woodland as well as a control soil; and inoculated into sterilized soil in which was grown two species
145 of herbaceous plants with different root system morphologies. We hypothesize (hypothesis 3) that
146 very fibrous root systems will increase aggregate stability, because more labile carbon will be
147 available for microbial communities from fine root exudates and mucilage. We further hypothesize
148 (hypothesis 4) that the effect of inoculation on aggregate stability will be greater with the inoculum
149 coming from the soil with highest aggregate stability (secondary woodland), and this is related to
150 source microbial community characteristics (measured using sequencing, GRSP, fungal biomass
151 ergosterol).

152 **2. Material and methods**

153 **2.1. Field sampling**

154 The study site is located in Wytham, Oxfordshire, southern England (1° 20 ' W, 51° 47 ' N), where
155 three land uses (grassland, pasture and arable) and two forests (ancient woodland and secondary
156 woodland) were selected (Fig. 1). The forests are located in a 410 ha area of mixed woodlands and
157 grassland, on a hill owned by the University of Oxford since 1943. These forests have been studied
158 for several decades (Watts, 1969; Mihók *et al.*, 2009) and the management history of the site is well
159 documented (Gibson, 1986). In this area, we selected two forests for our study, an ancient woodland

160 and a secondary woodland. The ancient woodland was traditionally managed by coppicing, but this
161 management stopped between 40 and 100 years ago (differing locations were abandoned at different
162 times). *Acer pseudoplatanus* L. and *Quercus robur* L. are abundant and *Corylus avellana* L. was the
163 main coppice species. The dominant tree species of the naturally regenerated secondary woodland are
164 *Fraxinus excelsior* L. and *Acer pseudoplatanus* L. (Savill et al., 2011). The ground flora of these
165 woodlands is characterised by dominant *Mercurialis perennis* (dog's mercury) and changeable
166 combinations of *Endymion non-scripta* (bluebell), *Circaea lutetiana*, *Geum urbanum*, *Arum*
167 *maculatum* and *Viola riviniana/reichenbachiana*. *Rubus fruticosus* and bryophytes are common
168 features of the field layer, in addition to the sedge *Carex sylvatica* and the grasses *Poa trivialis* and
169 *Brachypodium sylvatica* (Butt et al., 2009). The three other land uses that were selected for this study
170 are in an adjacent farmland. The arable site was cultivated with winter oats (*Avena sativa* L.) with no
171 fertilization in 2014 (the sampling year) and with spring wheat (*Triticum aestivum* L.), fertilized at a
172 rate 164 kg N ha⁻¹ and 7 t ha⁻¹ of pig manure in 2013. In this arable site, soil was ploughed prior to our
173 sampling. The other two sites, grassland and pasture, have been permanent pastures since 1992, with
174 the only difference that the grassland had not been actively grazed the year preceding the study (2013-
175 2014), increasing drastically the aboveground biomass of vegetation. Soils in the area span a
176 geological sequence from Oxford Clay to Upper Corallian sand and silt to Upper Corallian limestone
177 and Coral Rag developing soils with characteristic textures (Savill et al., 2011). All sites selected for
178 this study are located on the Oxford Clay formation with Denchworth series soil (Eutric Vertic
179 Stagnosols *sensu* USS Working Group WRB (2014)).

180 We collected soil samples to examine relationships in soil properties and microbial communities
181 between sites and to provide background data for the inoculation experiment (see below). In each land
182 use type, sampling was performed in three plots that were randomly located and samples were
183 collected at a depth of 0-0.1 m except for aggregate stability samples and soil physicochemical
184 properties that were collected at two depths (0-10 cm and 20-30 cm). Depth can influence aggregate
185 stability as it affects soil properties that are related to soil aggregate stability such as total organic
186 carbon, total nitrogen, freeze-thaw cycles, etc. Four different soil samples were collected at each

187 depth using sterilised material: (i) one soil cylinder (1.01 dm³) for fine root analysis, (ii) 10 to 20 g of
188 soil for microbial analysis placed in a plastic bag, (iii) soil for measurements of aggregate stability (at
189 depths of 0-0.1 m and 0.2-0.3 m) placed in 0.75 dm³ containers and (iv) two soil samples for soil
190 physicochemical analyses (at depths of 0-0.1 m and 0.2-0.3 m). Litter from the surface of each soil
191 core was sampled and kept in plastic bags for further analysis. A total of 15 samples of each type were
192 collected: five land uses x three samples at each land use (except for aggregate stability and soil
193 physicochemical properties where a total of 30 samples were collected, since six samples were
194 collected at each plot; three at 0-0.1 m and three at 0.2-0.3 m). Soil samples were collected during
195 July 7th - 10th, 2014, at the peak of the summer season when microbial activity was expected to be at
196 its maximum (Sarithchandra et al., 1988; Bardgett et al., 1997). Soil physicochemical properties of
197 the five sites are given in Table 1.

198 Samples for root, litter and microbial analysis were stored at -20°C prior to analysis. Soil samples for
199 aggregate stability tests were air-dried at ambient temperature and analysed within one month.

200 **2.2. Laboratory inoculation experiment setup and sampling**

201 Intact soil cores with a diameter of 110 mm and a length of 150 mm (1.42 dm³) were collected from
202 two field sites, arable land and secondary woodland in close proximity (300 m) for the laboratory
203 inoculation experiment (Fig. 2). These two land uses were selected based on their large differences in
204 soil aggregate stability and the minimal differences in soil textural properties (Table 1, Fig. 3). Thirty-
205 three intact soil cores in the arable land site were collected on January 7th, 2015 and another set of 33
206 soil cores were collected in Wytham woods on January 8th, 2015 (Figs. S1 and S2a). Thirty-three pots
207 were also filled with a commercial loamy soil (Boughton Kettering Loam, Boughton Loam Ltd, U.K.)
208 to be used as a control treatment because this loamy soil lacks structural stability. Before and after
209 sterilization, three samples from each provenance were collected for the measurement of
210 physicochemical properties (Table S1).

211 After the field collection, the upper 30 mm of each soil core were removed to allow subsequent
212 inoculation with 15 mm of non sterile soil from the arable and woodland soils (0.14 dm³, 10% of total

213 volume) and 15 mm free for water ponding from irrigation (Figs. S2b and c). Thirty soil cores of each
214 treatment (arable, woodland and control) were sterilized by gamma irradiation at Elgin Synergy
215 Health PLC (Swindon, U.K.) with a dose between 50.5 and 52.1 kGy on January 29th, 2015. Three
216 cores of each soil were used to assess aggregate stability after gamma radiation and compared with
217 the three non sterilized cores to verify that there was no effect of gamma irradiation on aggregate
218 stability. Similarly, one soil sample in each of these three cores was collected to assess changes in
219 microbial communities before and after the gamma radiation (three samples assessed before and three
220 samples after).

221 After irradiation, the upper 15 mm of 18 soil cores was inoculated with fresh soil collected on
222 February 9th, 2015 from the arable and woodland fields (nine soil cores from each provenance) and
223 another set of nine soil cores were not inoculated (controls).

224 In three replicates of each combination, seeds from two species with either tap/adventitious (*Urtica*
225 *dioica*) or fibrous (*Brachypodium sylvaticum*) root systems (*sensu* Fitter and Peat, 1994) were sown
226 and a third treatment with no plants was used as a control (see Fig. 2 and Fig. S2 for a detailed
227 illustration of procedure). Seeds of the two selected species, abundant in the Wytham area with
228 contrasting root systems (Savill et al., 2011), were obtained from two different suppliers: Emorsgate
229 Seeds, U.K. (*B. sylvaticum*) and Naturescape British Wildflowers, U.K. (*U. dioica*). Seeds were
230 sterilized by placing them in a mesh bag and left for 30 min in a 2% sodium hypochlorite solution.
231 They were removed from the bag and rinsed three times (for 5 min each) in sterilized deionised water
232 (Commander et al., 2009). Then, seeds were germinated for three weeks in sterilized, deionised water
233 before sowing. Ten seeds of each species were sown in each pot. Plants were grown in a temperature
234 and light controlled facility for 4 months (February 12th to June 15th, 2018, Fig. S2d). Pots were
235 regularly weeded by hand for the first six weeks, but after that, the species that were sown
236 commenced growth and weeding was ceased to avoid disturbance, although some seedlings emerged
237 (particularly in treatments inoculated with arable soil).

238 The first three irrigation events were performed with autoclaved deionised water, but afterwards, pots
239 were watered with tap water using an automatic station (to promote downward movement of
240 inoculum). To identify microbial communities existing in tap water and subtract them from the later
241 microbial analysis on soils, 5 l of tap water were collected at the beginning and the end of the
242 experiment. This water was filtered, and DNA was extracted.

243 **2.3. Sample analysis**

244 **2.3.1. Soil respiration, moisture and physicochemical properties**

245 **2.3.1.1. Field study**

246 Soil collected in the field study was sieved at 2 mm after air drying and the physicochemical
247 properties of the < 2 mm soil fraction was quantified by Natural Resource Management (Berkshire,
248 UK). Soil pH was measured in water as 1:2.5 extract. Soil organic matter content (SOM) was
249 determined via loss-on-ignition at 500 °C (Dean 1974). Total nitrogen (N) and carbon (C) were
250 determined via the DUMAS method (Shea and Watts 1939). Available potassium (K) and magnesium
251 (Mg) were determined through ammonium nitrate extraction and available phosphorus (P) was
252 measured using the Olsen method (extraction with 0.5 M sodium bicarbonate solution, (Olsen et al.
253 1954)). Soil texture was determined by laser-diffraction analysis (McCave et al. 1986). Each soil
254 sample was previously digested in hydrogen peroxide solution to destroy the organic matter and
255 sodium hexametaphosphate to release the bound clay particles.

256 Aggregate stability was determined by the fast wetting standard method, ISO/CD 10930, developed
257 by Le Bissonnais (Le Bissonnais, 1996). This methodology is appropriate to compare the behaviour of
258 a large range of soils during rapid wetting (mimicking heavy rainstorms in summer). A quantity of 5 g
259 of aggregates (3-5 mm) were gently immersed in 50 ml of deionized water for 10 minutes; water was
260 then removed with a pipette and the soil material was transferred to a 50 µm sieve previously
261 immersed in ethanol. The 50 µm sieve immersed in ethanol was gently moved five times to separate
262 fragments smaller and bigger than 50 µm. The >50 µm fraction was collected, oven-dried and gently

263 dry-sieved by hand on a column of six sieves: 2000, 1000, 500, 200, 100 and 50 μm . The mass
264 percentage of each size fraction was calculated, and the aggregate stability was expressed by
265 computation of the mean weight diameter (MWD).

266 **2.3.1.2. Laboratory inoculation experiment**

267 Soil samples collected from the inoculation experiment were sent to the chemical analysis laboratory
268 of Forest Research (Surrey, UK). Soil pH was measured in a suspension of 5 g of soil with 25 ml of
269 water that was shaken on an orbital shaker for 15 min and settled for 45 min prior to pH measurement
270 using a Sentek pH electrode (Reference method: ISO 10390). Total N, C, soil organic carbon (SOC)
271 and inorganic carbon (CaCO_3) were measured with a combustion method using a Carlo Erba CN
272 analyser (Reference method ISO 10694 & 13878). Aggregate stability was measured with the fast
273 wetting standard method described above.

274 We examined if microbial inoculation (through the addition of surface soil) changed soil texture and
275 aggregate stability in deeper soil. The top 15 mm of soil was removed, and aggregate stability
276 measured in the remaining soil. Texture of the aggregates obtained at the end of the experiment was
277 measured by the CIRAD - US Analyses (Montpellier, France). The objective was to test whether the
278 texture of the obtained aggregates was more related to the inoculated core (origin) or to the *inoculum*
279 (inoculation treatment) and so, if there was any possible influence of inoculation with soil on the
280 results of stability tests. Soil texture of aggregates was determined with an automated pipette method
281 (Reference method AFNOR NF X 31-107) using a Texusol granulometer (ISITEC-LAB) on samples
282 pre-treated following the standard method NF ISO 11464 (X31-412).

283 Two additional measurements in soils were made: i) the rate of exchange of CO_2 (assimilation; g
284 $(\text{CO}_2) \text{ m}^{-2} \text{ h}^{-1}$) measured with an EGM-4 Environmental Gas Monitor for CO_2 and a SRC-1 Soil
285 Respiration Chamber (PP Systems, Amesbury, MA, U.S.A.) in June 15th-17th, 2015 and ii) soil water
286 content measured at two dates (June 8th and 12th, 2015) after irrigation with a WET Sensor (Delta-T
287 Devices Ltd, Cambridge, UK).

288 2.3.2. Aboveground vegetation, fine roots and litter

289 In the soil inoculation experiment, all aboveground vegetation biomass was cut at the surface at the
290 end of the experiment and dried at 60°C until constant weight (Fig. S2e). Then, a core (0.295 dm³)
291 was collected in each of the pots for the assessment of root properties (Fig. S2f and Fig. S3).

292 Root samples from both the field study and the inoculation experiment were washed gently with
293 deionised water and divided in two subsamples: 1) a representative subsample that was selected for
294 scanning and later drying in the oven at 40°C (n=3 for each land use in the field experiment and n =
295 27 for each provenance; arable, woodland and control; in the inoculation experiment) and 2)
296 remaining root material of the sample that was dried at 40°C until constant weight (n=3 for each land
297 use in the field experiment and n = 27 for each provenance in the inoculation experiment). Both
298 subsamples were weighed before and after drying. Roots selected for morphological measurement
299 were stained with methylene blue (1 g l⁻¹) to increase the contrast and allow the detection of fine roots.
300 Then, roots were placed in a tray with deionised water and scanned (Epson© V700 Perfection) at a
301 resolution of 1200 dpi. Analysed subsample roots were then recovered, and oven dried at 40 °C and
302 weighed to obtain dry mass. Root images were analysed with the WinRhizo® software (Pro version
303 2007, Regent Instrument, Quebec, Canada) using the automatic thresholding option and Lagarde's
304 mode, with a filter identifying roots when length was five times the width. We measured total root
305 length and the length of roots in seven diameter classes (roots with a width of 0 - 1 mm were
306 separated into 0.2 mm classes and roots with a width of 1 - 2 mm were separated into 0.5 mm classes).
307 Debris was removed with a filter that eliminated fragments that were less than five times longer than
308 their width. Specific root length (SRL) was calculated as the ratio between total root length and root
309 dry mass. The percentages of very fine (VFR, diameter < 0.2 mm) and fine (FR, 0.2 < diameter < 1
310 mm) roots were defined as the ratio of length in the concerned root classes to total root length (Miller
311 and Jastrow, 1990). Total root mass density (RMD) was calculated by the ratio of total root dry mass
312 and the soil volume extracted. Total root length density (RLD) was calculated by the ratio of total root
313 length and the soil volume extracted. Total root dry mass was calculated as the sum of the dry mass of

314 roots selected for morphological analysis and those of the remaining roots. Root dry matter content
315 (RDMC) was calculated as the ratio of the root dry weight and fresh weight.

316 The concentrations of water soluble compounds (cellulose, hemicellulose and lignin; mg g^{-1}) in root
317 and litter samples from the field experiment ($n = 11$ for litter, i.e. 3 replicates x 4 treatments = 12
318 excluding 1 sample with insufficient mass; and $n = 4$ for roots as replicates were combined due to
319 limited sample amount; “arable” was excluded from these measurements), were obtained by the Van
320 Soest Method (Van Soest, 1963) with a Fibersac fibre analyser (Ankom, Macedon, USA). Root C and
321 N concentrations were measured in roots from the field study ($n=12$, i.e. 3 replicates x 4 treatments
322 since arable was excluded from these measurements) and the inoculation experiment ($n = 3$ for each
323 treatment with roots) and litter C and N were measured in samples from the field experiment ($n=15$)
324 using an elemental analyser (Thermo-Finnigan EA1112, Italy).

325 **2.3.3. Soil microbial communities**

326 **2.3.3.1. Obtaining rhizosphere and bulk soil samples**

327 In both the field study and laboratory inoculation experiment, roots were cleaned and rhizosphere and
328 bulk soil samples obtained following the protocols of Bulgarelli et al. (2012, 2015). Briefly, loose soil
329 was manually removed from the root system and stored as the bulk soil sample. Roots were collected
330 in 50 ml falcon tubes containing 10 ml PBS-S buffer (130 mM NaCl, 7 mM Na_2HPO_4 , 3mM
331 NaH_2PO_4 , pH 7.0, 0.02 % Silwet L-77) and washed for 20 minutes at 180 rpm on a shaking platform.
332 These roots were transferred to a new falcon tube and subjected to a second washing treatment (20
333 minutes at 180 rpm in 3 ml PBS-S buffer). The soil suspensions collected in the falcon tubes after the
334 first and second washing treatments were combined, centrifuged at 4000 g for 20 min and the pellet,
335 considered as the rhizosphere sample, was frozen and stored at -20°C until further analysis.

336 **2.3.3.2. DNA extraction, amplicon library construction and sequencing**

337 Total DNA was extracted from soil (0.25 g) and the rhizosphere fractions (0.25 g when possible and
338 the entire material available when quantity was less than 0.25 g). DNA extraction was performed

339 using PowerSoil®-htp96 Well Soil DNA Isolation Kit according to the manufacturer's instructions
340 (MOBIO Laboratories, UK).

341 Bacterial and fungal community biodiversity was assessed using Illumina amplicon sequencing of
342 16S rRNA genes (bacteria) and the Internal transcribed spacer (ITS) region (fungi) to
343 phylogenetically identify responsive taxa. A phylogenetic analysis was also performed for bacterial
344 communities. Amplicon libraries were constructed according to the dual indexing strategy of Kozich
345 et al. (2013), with each primer consisting of the appropriate Illumina adapter, an 8-nt index sequence,
346 a 10-nt pad sequence, a 2-nt linker and the gene specific primer. For 16S, the V3-V4 hypervariable
347 regions of the 16S rRNA gene was targeted using primers based upon the universal primer sequence
348 341F and 806R. For ITS, region 2 (ITS2) was amplified utilising the fITS7 (forward) and ITS4
349 (reverse) primer sequences described in Ihrmark et al. (2012). Additional methodological details of
350 Illumina sequencing are described in Notes S1.

351 **2.3.3.3. Processing high-throughput data for community analysis**

352 Sequenced 16S rRNA paired-end reads were joined using PEAR (Zhang et al., 2014), quality filtered
353 using FASTX tools (hannonlab.cshl.edu), length filtered with the minimum length of 300 bps,
354 presence of PhiX and adapters were checked and removed with BBTools (jgi.doe.gov/data-and-
355 tools/bbtools/), and chimeras were identified and removed with VSEARCH_UCHIME_REF (Rognes
356 et al., 2016) using Greengenes Release 13_5 (at 97%) (DeSantis et al., 2006). Singletons were
357 removed and the resulting sequences were clustered into operational taxonomic units (OTUs) with
358 VSEARCH_CLUSTER (Rognes et al., 2016) at 97% sequence identity (Tindall et al., 2010).
359 Representative sequences for each OTU were taxonomically assigned by RDP Classifier with the
360 bootstrap threshold of 0.8 or greater (Wang et al., 2007) using the Greengenes Release 13_5 (full)
361 (DeSantis et al., 2006) as the reference. Unless stated otherwise, default parameters were used for the
362 steps listed. ITS2 sequences were processed using the PIPITS pipeline (Gweon et al., 2015), where
363 OTUs were taxonomically assigned against the UNITE database (Release 31.01.2016, Koljalg et al.
364 (2013).

365 **2.3.4. Glomalin-Related Soil Proteins (GRSP).**

366 Hurisso et al. (2018) recently proposed the term autoclaved-citrate extractable protein (ACE), to avoid
367 the implied assumption to the fungal origin of these proteins, but here we use the term Glomalin-
368 Related Soil Proteins (GRSP) for ease of comparison with other studies. In the inoculation
369 experiment, soil samples from the middle of each soil core were collected after the incubation period
370 for the assessment of GRSP, thought to be a marker of arbuscular mycorrhizal activity. Soil samples
371 were stored at -20 °C until further processing. Operationally-defined GRSP fractions were obtained
372 using the extraction methods proposed by Wright and Upadhyaya (1996) as follows. Easily
373 extractable soil protein (GRSP_{EE}) was obtained by autoclaving soil in a solution of 20 mM sodium
374 citrate at pH 7 for 30 min. Total GRSP (GRSP_T) was obtained by two successive autoclave extractions
375 of soil in 50 mM sodium citrate at pH 8, each for 60 min. In each case, the soil:solution ratio was 1:8
376 and phases were separated after cooling by centrifugation at 15000 g for 15 min. For GRSP_T, after the
377 first autoclave cycle, solution was removed and replaced by the same volume of new addition of
378 citrate solution, the mixture was vortexed to re-suspend the soil. The solutions were combined to form
379 the total extract. Solutions were frozen until required for analysis. After thawing, samples were
380 centrifuged again at 15000 g for 15 minutes to remove any precipitate that may have formed. Protein
381 in both extracts were assayed using the Bradford technique with Bradford QuickStart kits from
382 BioRad Laboratories (Hercules, CA, USA), calibrated against solutions of bovine serum albumin
383 (BSA) within the working range of 0-200 mg dm⁻³. Following the recommendations proposed by
384 Moragues-Saitua and co-workers (Moragues-Saitua et al., 2019), samples were diluted twofold
385 (absorbance of about 0.1 at 465 nm), to reduce the interference of co-extracted coloured compounds.
386 A sample volume of 20 µl were assayed in microplates with 230 µl Bradford dye reagent. Optical
387 density was measured at 595 nm in ThermoScientific Multiskan GO spectrometer (Waltham, MA,
388 USA) and sample blank colour absorbance at pH=1 corrected.

389 **2.3.5. Ergosterol**

390 Ergosterol, a proxy for active fungal biomass, was measured in soil samples that were obtained from
391 the middle of each core of the inoculation experiment after the incubation period. Ergosterol was
392 assessed following the method of Plassard et al. (2000). Briefly, ergosterol was extracted by mixing
393 very gently 0.5 g of soil in 3 ml of methanol containing polyclar (0.5%, w/v) (CAS n° 9009-39-8) in
394 screw-cap tubes. After firm closure, tubes were incubated at 4°C for 24 h, without shaking as
395 ergosterol is rapidly lost by oxidation. The concentration of ergosterol in filtered (0.45 µm) methanol
396 extracts was determined at 270 nm by high-performance liquid chromatography using a C18 column
397 and eluted with methanol flowing at 1 ml min⁻¹.

398 **2.4. Statistical analysis**

399 **2.4.1. Field study**

400 First, we examined the differences on soil, root and litter physicochemical characteristics and
401 aggregate stability among the different land uses. Then, we performed NMDS data ordinations of
402 microbial data and tested of the effect of land use and sample fraction on their structure
403 (Permutational Multivariate Analysis of Variance, PERMANOVA). Finally, we studied the
404 correlation between microbial community composition and environmental variables (Spearman
405 correlations between Shannon diversity indexes and distance-based redundancy analysis, dbRDA).

406 Soil aggregate stability, soil physicochemical properties and root and litter traits were analysed by
407 analysis of variance (ANOVA) and post hoc Tukey's honestly significant difference (HSD) test. The
408 land use explanatory variable was treated as a factor. Relative abundance of the different phyla and
409 trophic modes were analysed by ANOVA and post hoc HSD with land use and sample fraction as
410 explanatory variables. All variables tested fulfilled ANOVA assumptions except for the Saprotroph
411 trophic mode that was inverse transformed to fulfil ANOVA assumptions. The relationships between
412 the relative abundance of bacterial phyla and the soil physicochemical variables, root and litter traits
413 were tested using Spearman correlations.

414 Similarities/dissimilarities between microbial communities were displayed using non-metric multi-
415 dimensional scaling (NMDS) of Bray–Curtis dissimilarity for bacterial and fungal species matrices.
416 To indicate similarities between treatments on the NMDS configuration, the points on the NMDS
417 ordination were overlapped with spider diagrams and convex hull polygons indicating land use and
418 soil fraction was indicated with different symbols. Surfaces were added to show soil aggregate
419 stability data on the ordinations. PERMANOVA tests (adonis R function) were performed for the
420 bacteria and fungi species matrices as the response and the two different factors (land use and soil
421 fraction) as the explanatory variables.

422 The Shannon diversity index for bacteria and fungi was calculated for each land use and sample
423 fraction (rhizosphere and bulk soil samples). The relationships between the microbial community
424 composition and the soil physicochemical variables, root and litter traits were tested using Spearman
425 correlations between these variables and alpha (Shannon diversity index) and beta diversities (NMDS
426 first and second axis) and a db-RDA. In order to select the environmental parameters to be included in
427 the constrained ordination, an initial db-RDA including all parameters was performed followed by a
428 stepwise model selection using Generalized Akaike Information Criterion (AIC, ordistep function
429 with a backward direction). Finally, the db-RDA analysis was performed only for the variables
430 obtained. ANOVA tests were performed on the final constrained ordination to confirm that the first
431 two axes and the environmental variables and the final constrained ordination were significant.

432 **2.4.2. Laboratory inoculation experiment**

433 First, we explored the differences on soil, vegetation and root traits and aggregate stability among the
434 different combinations of treatments: i) origin (control, arable and woodland), ii) inoculation (not
435 inoculated, arable and woodland) and iii) root treatment (no roots, fibrous and tap). Soil
436 physicochemical properties, aboveground vegetation biomass, root traits, aggregate stability and
437 aggregate fractions were analysed by ANOVA when ANOVA assumptions were fulfilled and with
438 Kruskal Wallis tests when they were not fulfilled. Origin, inoculation, and root treatment were used as
439 explanatory variables and were treated as factors. PCA was performed with a selection of variables

440 used for the description of soil, vegetation, and root properties. Variables included in the PCA were:
441 aboveground biomass, CO₂ assimilation, soil water content, soil pH, SOC, soil C:N, SRL, VFR,
442 MRD, RMD, RLD, root C:N, GRSP_T, ergosterol and MWD. These variables were selected based on
443 their correlation and ecological significance (i.e. variables that were correlated but their source was
444 ecologically different were kept, e.g. soil C:N and root C:N). Apart from the ANOVA performed for
445 MWD with soil origin, inoculation and root treatment as factors, an additional analysis of covariance
446 (ANCOVA) model was tested and included RLD to explore separately the effect of root chemical (i.e.
447 species effect) and root physical properties (quantity of roots) within the effect of root treatment.

448 Non-metric multi-dimensional scaling (NMDS) of Bray–Curtis dissimilarity were performed for
449 bacterial and fungal species matrices obtained for bulk soil fraction (since the rhizosphere sample
450 fraction was only available for treatments where roots were present). To indicate similarities between
451 treatments on the NMDS configuration, the points on the NMDS ordination were overlapped with
452 convex hull polygons indicating the inoculation treatment, origin was indicated with symbols of
453 different colour and root treatment was indicated with different symbol shape. PERMANOVA tests
454 (adonis R function) were performed for the bacteria and fungi species matrices as the response and the
455 three different treatments (origin, inoculation and root treatment) as the explanatory variables using
456 species matrix of bulk soil and rhizosphere fractions. Additional Analysis of similarities (ANOSIM)
457 were performed to test the effects of root treatment, and inoculation in each of the combinations of the
458 experimental design in bulk soil sample fractions.

459 The Shannon diversity index for bacteria and fungi was calculated for each pot. In line with the field
460 study, Spearman correlations between soil, vegetation, root traits and aggregate stability variables and
461 alpha (Shannon diversity index) and beta diversities (NMDS first and second axis) were calculated.
462 Likewise, a db-RDA was performed on species matrices of bacterial and fungal communities obtained
463 from the bulk soil fraction. Environmental parameters included in the db-RDA were selected by a
464 stepwise model selection process using AIC similar to the process performed for the field study.
465 ANOVA tests were performed on the final constrained ordination to confirm that the first two axes
466 and the environmental variables and the final constrained ordination were significant.

467 Finally, we explored the textural class (proportion of clay, fine silt, coarse silt, fine sand, coarse sand)
468 of soil aggregates used for the stability tests, to examine whether there was any possible influence of
469 inoculation on the results of stability tests. We depicted a soil texture triangle to show the texture of
470 the different treatment combinations and overlapped with ellipses discriminating the origin of the
471 sample and symbols of different colours indicating inoculation treatment. Then, we performed a
472 weighted Classical (Metric) Multidimensional Scaling with textural properties to explore the
473 differences among origin and inoculation treatment, the points on this ordination were overlapped
474 with convex hull polygons indicating the origin, and inoculation was indicated with symbols of
475 different colour. Finally, we performed an analysis of similarities (ANOSIM) to test the effects of
476 origin and inoculation treatments on the textural properties of aggregates.

477 All analyses were performed in RStudio Version 3.6.0 (R Core Team, 2019) using the vegan
478 (Oksanen et al., 2019) and ade4 (Dray and Dufour, 2007) packages.

479 **3. Results**

480 **3.1. Field study: Relationships between soil aggregate stability and microbial communities**
481 **among land use types**

482 **3.1.1. Relationships between land use and soil, litter and root properties**

483 Soil physicochemical properties from samples taken at depths of 0-0.1 m and 0.2-0.3 m, differed
484 between land uses (Table 1). The arable field site had a higher pH and nutrient contents (P, K) but
485 lower Mg, SOC, N and C than the other sites. The pasture site had the highest SOC, N and C contents.
486 All soils had a loamy texture, but ancient and secondary woodlands and arable field sites possessed a
487 clay loam texture and both the grassland and pasture had a sandy loam texture. Roots had similar
488 traits between ancient and secondary woodlands (Table S2), with lower SRL (19.68 ± 4.06 and
489 13.16 ± 1.16 m g⁻¹ respectively), lower MRD (0.44 ± 0.01 and 0.56 ± 0.02 mm respectively), lower RLD
490 (5.69 ± 1.26 and 5.12 ± 0.90 km m⁻³), higher N content (1.18 ± 0.06 and 1.48 ± 0.01 mg g⁻¹ respectively)
491 and lower C:N (35.04 ± 2.57 and 27.72 ± 0.69 respectively) than the other land use types. Litter traits
492 had similar trends to that observed in roots: litter from ancient and secondary woodlands had higher
493 hemicellulose (17.02 ± 1.01 and 19.88 ± 0.20 mg g⁻¹ respectively) and N content (1.37 ± 0.07 and
494 1.43 ± 0.02 mg g⁻¹ respectively) and lower C:N (26.62 ± 1.35 and 26.21 ± 1.08 respectively), compared to
495 the other land use types. Soil aggregate stability was significantly different among land uses (Fig. 3),
496 with pasture having the greatest MWD and arable land the lowest, with the other three land uses
497 possessing an intermediate behaviour. A clear effect of soil depth was also identified (Fig. 3),
498 highlighting the role of processes at the surface that promote aggregate stability and justifying the
499 subsequent focus on soil sampled at a depth of 0-0.1 m.

500 **3.1.2. Effects of land use on the structure of bacterial and fungal communities**

501 Land use type affected the relative abundance of Actinobacteria, Proteobacteria and Verrucomicrobia
502 (Fig. 4a) with the arable land having a significantly higher relative abundance of Actinobacteria and
503 lower relative abundance of Proteobacteria and Verrucomicrobia compared to other land uses. In
504 addition, a significant interaction was found between land use and sample fraction for Acidobacteria,
505 with a decreased relative abundance in woodland rhizosphere soils but not in woodland bulk soil

506 samples. Sample fraction also significantly affected the relative abundance of Bacteroidetes and
507 Verrucomicrobia (Fig. 4a), which were more abundant in bulk soils for Bacteroidetes and the opposite
508 for Verrucomicrobia, with only a significant effect of land use type in the rhizosphere for
509 Verrucomicrobia. Correlations between relative abundance of bacterial phyla and soil, root and litter
510 variables showed that oligotrophic Chloroflexi and Verrucomicrobia were significantly correlated to
511 root and litter physicochemical variables (i.e. SRL, VFR, MRD, RDMC, root and litter N and C:N,
512 litter lignin) in the rhizosphere but not in the bulk soil (Table 2). Conversely, copiotrophic phyla
513 (Actinobacteria, Bacteroidetes and Proteobacteria) showed similar correlations in bulk soil but not in
514 the rhizosphere. The relative abundance of these copiotrophic phyla were correlated with some soil
515 physicochemical traits (pH, P, sand content) and for the specific case of Actinobacteria, this phylum
516 showed significant correlations with root and litter traits (SRL, RMD, root N, C:N and hemicellulose
517 and litter fibres, Table 2). For fungi, land use influenced the relative abundance of Ascomycota and
518 Zygomycota, with a significant effect of sample fraction for Ascomycota (Fig. 4b). Arable land had
519 the highest relative abundances of Ascomycota, whilst the secondary woodland possessed the lowest
520 relative abundance of Ascomycota and the highest abundance of Zygomycota. Regarding trophic
521 mode, land use type affected the relative abundance of saprotrophs and symbiotrophs with the arable
522 field having the lowest abundances. The land use type also influenced the relative abundance of fungi
523 belonging to an unspecific category that includes pathotrophs, saprotrophs and symbiotrophs, with the
524 highest relative abundance in secondary woodland (Fig. 4c).

525 NMDS ordinations showed that bacterial and fungal communities were markedly distinct between
526 land use types (Figure 5a). These findings were supported by the PERMANOVA performed for the
527 bacterial and fungal species matrices (Table 3), which showed that land use type was by far the most
528 important factor structuring bacterial and fungal communities ($R^2=0.44$ $p=0.01$ and $R^2=0.53$ $p=0.01$,
529 respectively) irrespective of whether communities were taken from the rhizosphere or bulk soils
530 (Table 3). Significant trends between changes in community structure and soil aggregate stability
531 were also observed, with arable communities having the lowest MWD (Figure 5a).

532 **3.1.3. Relationships between environmental variables and bacterial and fungal communities**
533 **among land use types**

534 The results of the relationships between soil physicochemical properties, root and litter traits (Tables 1
535 and S2, Figure 3) and microbial community composition are shown in the db-RDA (Figure 5b) and
536 Spearman correlations with alpha and beta diversities (Table S4). The final db-RDA analysis (Figure
537 5b) shows only the environmental parameters that were selected through stepwise model selection:
538 MWD, soil pH and P, and MRD for bacterial communities and MWD, RLD, FR and soil pH and P for
539 fungal communities. There was a clear separation between land uses for bacterial communities
540 (Figure 5b), with three main groups: i) woodlands (ancient and secondary), ii) grasslands (actively
541 and not actively grazed) and iii) arable land. Soil from woodlands had significantly greater MWD and
542 higher MRD, whereas arable land had a significantly greater soil pH and P (Figure 5b). For fungal
543 communities, a similar configuration was found among the three land use types, but soil from
544 grasslands had a significantly higher RLD, and woodlands had a greater FR. ANCOVA tests
545 performed on the final constrained ordination confirmed that the first two axes, the environmental
546 variables and the final constrained ordination were significant.

547 Bacterial and fungal alpha diversities (Table S3) were not significantly correlated with any of the
548 variables measured (Table S4). Bacterial beta diversity was significantly and positively correlated
549 with soil pH, clay and magnesium contents, soil C:N, FR, MRD, root and litter N and litter lignin and
550 negatively correlated to MWD, sand content, SRL, VFR and root and litter C:N and hemicellulose.
551 Similarly, fungal beta diversity was significantly and positively correlated ($p < 0.001$ and $R^2 \geq 0.65$)
552 with soil pH, clay and magnesium contents, soil C:N, FR, MRD, root N, and litter lignin, and
553 negatively with sand content, SRL, VFR, RLD, root C:N and root and litter hemicellulose (Table S4).

554 **3.2. Laboratory inoculation experiment: Effect of microbial inoculation on soil aggregate**
555 **stability and interactions with soil origin and root systems.**

556 **3.2.1. Effects of soil origin, inoculation and root systems on soil properties and plant traits**

557 The PCA performed with a selection of variables used for the description of soil, plant and root traits
558 explained 56.7% of the variance (Fig. 6). The first PCA axis (horizontal), accounted for 43.7% of the
559 variation and the second PCA axis accounted for 13% of the variation. This PCA showed a clear
560 differentiation between control, and either arable land or woodland. The main contributors to these
561 differences were soil pH, SOC and MWD. Control soil had significantly higher pH, soil and root C:N
562 ratios and lower SOC, MWD, GRSP_T and ergosterol content compared to soils from the arable land
563 and woodland.

564 All soil properties assessed (e.g. pH, SOC, C, N) were influenced significantly by the origin of the
565 soil (i.e. control, arable and woodland, Table S5, $p < 0.05$). Additionally, a significant effect of
566 inoculation was found for soil water content ($p < 0.001$), which may be because this was the only
567 variable that was measured in the top 0.1 m of the soil profile, therefore including the soil used for
568 inoculation. Aboveground plant biomass was significantly affected by soil origin and root system type
569 ($p < 0.001$) but not by inoculation. Concerning root traits, soil origin affected significantly root biomass
570 ($p = 0.014$), SRL, VFR, FR, MRD, N and C:N ($p < 0.001$) and root system type affected root biomass
571 ($p < 0.001$), SRL ($p < 0.020$), RMD and RLD ($p < 0.001$) but no effect of inoculation was found (Table
572 S5, Fig. 7).

573 **3.2.2. Effects of soil origin, inoculation and root systems on the structure of bacterial and fungal** 574 **communities.**

575 NMDS ordinations showed that bacterial and fungal communities were markedly distinct between
576 inoculation treatments and soil origin but there were weaker effects of root system types (Figure 8a).
577 These findings were supported by the PERMANOVA performed for the bacterial and fungal species
578 matrices (Table 4), which showed that the main factors structuring bacterial communities were
579 inoculation ($R^2 = 0.18$, $p < 0.001$) and origin ($R^2 = 0.05$, $p < 0.001$) in bulk soil, and inoculation ($R^2 = 0.26$,
580 $p < 0.001$), origin ($R^2 = 0.09$, $p < 0.001$) and root system ($R^2 = 0.06$, $p < 0.001$) in the rhizosphere. The
581 results observed for fungal communities were different, with inoculation ($R^2 = 0.16$, $p < 0.001$), origin
582 ($R^2 = 0.03$, $p < 0.001$) and root system ($R^2 = 0.02$, $p < 0.001$) significantly affecting fungal communities in

583 the bulk soil. However, for fungal communities in the rhizosphere, only inoculation treatments
584 ($R^2=0.24$, $p<0.001$) and soil origin ($R^2=0.08$, $p<0.001$) had significant effects. Overall, considering
585 the R^2 values, the influence of root system type was stronger on the structure of bacterial communities
586 than for fungal communities. Furthermore, the interaction between origin and inoculation was
587 significant in all cases ($p<0.001$), indicating that the effect of inoculation was affected by the soil
588 origin. Additionally, for fungal communities in the bulk soil, the effect of root system type was
589 affected by both inoculation (inoculation*root treatment $R^2=0.03$, $p<0.001$) and soil origin
590 (origin*root treatment $R^2=0.03$, $p=0.042$). The overall plant root system effects were minor compared
591 to soil properties, nevertheless significant effects of root systems in certain soils were observed (Table
592 5). The effect of root system type was context-specific: in the non-inoculated control soil both
593 bacterial and fungal communities were significantly affected ($p=0.013$ and 0.012 respectively), while
594 for the field-sampled soils root system significantly influenced fungal communities only in arable soil
595 non-inoculated ($p=0.001$) and inoculated with woodland soil ($p=0.015$). In contrast, inoculation
596 affected both bacterial and fungal communities ($p<0.05$) for all combinations of treatments.

597 The dbRDA constrained ordinations of bacteria and fungi species matrices showed that the main
598 factors affecting the structure of microbial communities were MWD, RMD and soil C:N for bacterial
599 communities and MWD, soil C, RMD, GRSP_T and ergosterol for fungal communities (Fig. 8b).

600 ANCOVA tests performed on the final constrained ordination confirmed that the first two axes, the
601 environmental variables and the final constrained ordination were significant. Bacterial and fungal
602 alpha diversities did not show significant correlations with any of the variables measured (Table S6).

603 Bacterial beta diversity was positively correlated ($p < 0.001$ and $R^2 \geq 0.3$) with some soil properties
604 (SOC, N, C, GRSP_{EE} and MWD) and with root N, and negatively correlated with soil pH, soil C:N,
605 CaCO₃ and root C:N. Fungal beta diversity was significantly and negatively correlated ($p < 0.001$ and
606 $R^2 \geq 0.3$) with soil pH and CaCO₃, but positively correlated with MWD and ergosterol (Table S6). It
607 is interesting to note that GRSP was strongly related to bacterial beta diversity, even more than with
608 fungal beta diversity, lending support to the increasing evidence that GSRP is not solely of AFM
609 fungal origin (Cissé et al., 2020).

610 **3.2.3. Effects of soil origin, inoculation and root treatments on soil aggregate stability.**

611 Soil aggregate stability (MWD) was significantly affected by soil origin ($p < 0.001$) and inoculation
612 ($p = 0.04$) but not by root system type (Table 6, Figure 7a). The inoculation increased aggregate
613 stability in the poorly structured control soil, but not in arable and woodland soils (Figure 7a). The
614 effect of inoculation was mainly on macroaggregates and their dispersion (fractions 1-2 mm, 0.5-1
615 mm and 0.1-0.2 mm; Table S5). The ANCOVA model including RLD showed that the effect of root
616 system type was again not significant, but a significant interaction did occur between inoculation and
617 RLD, revealing that the number of roots modified the effect of inoculation ($p < 0.001$; Table 6). The
618 correlations between aggregate stability and the reputed fungal markers $GRSP_T$, $GRSP_{EE}$ and
619 ergosterol were only significant when all three soils were analysed together (Fig. S4), in agreement
620 with the results of ANOVA tests that showed that only the effect of soil origin was significant (Table
621 S5). The analysis of the textural properties of soil aggregates used for the aggregate stability tests
622 showed that removing the top 15 mm after the experiment for all soil analyses was efficient and there
623 was no possible influence of soil addition on the results of stability tests (Fig. S5a and S5b), since the
624 texture of aggregates used for the tests was related to origin ($p = 0.001$) but not to inoculation
625 ($p = 0.015$, Fig. S5c).

626

627 **4. Discussion**

628 *Hypothesis 1: There are strong relationships between soil aggregate stability and microbial*
629 *community structure that are driven by land use.*

630 In agreement with our first hypothesis, the impact of land use significantly affected soil aggregate
631 stability and the structure of microbial communities. A large body of previous research has found that
632 the conversion of natural forest to cropland results in a decline of soil quality, SOC and aggregate
633 stability and the opposite is true for soils restored after a disturbance (Barto et al., 2010; Duchicela et
634 al., 2013; Delelegn et al., 2017). Positive relationships between SOC and aggregate stability have
635 been found in different soil types, such as black soils (Zhang et al., 2012), Typic Ustochrepts (Saha et
636 al., 2011), and loamy soils (Vermic Haplubrepts) where SOC is thought to increase aggregate stability
637 by lowering the wettability and increasing the cohesion of aggregates (Chenu et al., 2000). Thus,
638 differences in SOC might explain the lowest aggregate stability in the arable field and the highest
639 stability in pasture and grassland. Apart from this influence of SOC on MWD, we found other factors
640 influencing MWD in our land use gradient that impact aggregate stability, as described previously: i)
641 agricultural practices such as the mechanical action of tillage (Balesdent et al., 2000), management
642 inputs (e.g. fertilizer, herbicide (Steenwerth et al., 2002)) and grazing (Barto et al., 2010), ii) the
643 presence of roots (measured through RLD; Hudek et al. (2017)), iii) the higher C:N in roots and litter
644 implying a higher content in recalcitrant matter that influences fungal activity and MWD as a
645 consequence (Bossuyt et al., 2001).

646 We demonstrated a strong impact of land use gradient on the structure of bacterial and fungal
647 communities. A regional study in Great Basin Province (California, USA) identified that agricultural
648 management can have larger effects on soil microbial communities than elevation and precipitation
649 gradients (Drenovsky et al., 2010). However, fungal and bacterial communities differ in their
650 responses to changes in agricultural practices. Previous research has shown that tillage reduces
651 microbial biomass and abundance of AMF (Helgason et al., 2010; Zhang et al., 2013) and that the
652 structure of the microbial communities responds strongly to land use with higher proportions of fungi

653 observed in natural ecosystems compared with agro-systems (Fanin and Bertrand, 2016). In a study
654 across a range of European field sites, land use intensification effects on microbial communities were
655 stronger when land use change affected soil conditions such as pH, carbon, nitrogen and phosphorous
656 contents (Thomson et al., 2015). Here, although we found significant relationships between the
657 changes in microbial community structure, aggregate stability and soil properties along the land use
658 gradient, causality cannot be confirmed.

659 *Hypothesis 2: The arable field site with lower aggregate stability will have a lower relative*
660 *abundance of copiotrophic communities and reduced relative abundances of saprotrophic and*
661 *ectomycorrhizal fungi, but the rhizosphere niche will favour copiotrophic bacterial communities.*

662 Given that microaggregates are considered to hold oligotrophic communities and macroaggregates
663 favour copiotrophic communities due to their differences in contents of labile/recalcitrant carbon
664 (Trivedi et al., 2017), we expected to find an increase in the relative abundance of copiotrophic
665 communities in soils with higher soil aggregate stability (i.e. higher MWD after disruption). In line
666 with this hypothesis, the arable field (with the lowest soil aggregate stability), had a lower relative
667 abundance of copiotrophic Proteobacteria phyla. However, we did not observe similar trends for
668 copiotrophic Actinobacteria and Bacteroidetes, and oligotrophic Acidobacteria, Chloroflexi and
669 Verrucomicrobia. In addition, correlations between relative abundance of bacterial phyla and MWD
670 were not significant. However, other soil physicochemical factors and notably root and litter
671 physicochemical traits, affected the relative abundance of these phyla with a differential effect on bulk
672 and rhizosphere niches. Lauber et al. (2008) also showed that rather than land use itself, changes in
673 soil edaphic properties explained changes in life strategies of bacterial communities. Since
674 Chloroflexi and Verrucomicrobia were correlated to root and litter physicochemical variables in the
675 rhizosphere but not in the bulk soil, we conclude that oligotrophic bacteria are more dependent on
676 rhizosphere conditions where labile carbon originates mostly from rhizodeposits, exudates and
677 mucigel (Dennis et al., 2010). On the contrary, copiotrophic phyla were more dependent on bulk soil
678 conditions where there are more exopolysaccharides released into the soil by the microorganisms
679 themselves. The relative abundance of these copiotrophic phyla was correlated with some soil

680 physicochemical traits (pH, P, sand content) and for the specific case of Actinobacteria, this phylum
681 showed correlations with root and litter traits (SRL, RMD, root N, C:N and hemicellulose and litter
682 fibres, Table 2). This particular behaviour of Actinobacteria could be because Actinobacteria exhibits
683 both copiotrophic and oligotrophic life strategies consistent with their capacity to degrade both labile
684 and complex carbon substrates (Ho et al., 2017). Bergmann et al. (2011) found that Verrucomicrobia
685 are oligotrophic with a highly variable relative abundance across soils, and that members of this
686 phylum were most abundant in soils from grasslands and prairies. However, these authors
687 acknowledged that the ecology of Verrucomicrobia remains poorly understood, and different factors
688 may affect the distribution of this phylum.

689 Studies of bacterial communities inhabiting the rhizosphere in grasslands (Mao et al., 2014; Shi et al.,
690 2015) and croplands (Peiffer et al., 2013; Donn et al., 2015) found that rhizosphere samples are
691 enriched in Proteobacteria, Actinobacteria, and Bacteroidetes. However, here we found an enrichment
692 of Verrucomicrobia (oligotrophic) and diminution of Bacteroidetes (copiotrophic) and Acidobacteria
693 (oligotrophic) in rhizosphere communities. Thus, the hypothesis that the rhizospheric niche favours
694 copiotrophic bacterial taxa (Peiffer et al., 2013; Lladó and Baldrian, 2017) was only confirmed with
695 the reduced relative abundance of the oligotrophic Acidobacteria, highlighting the heterogeneous
696 nature of the rhizosphere and the importance of studying the relationships between soil, root and
697 physicochemical variables and the relative abundance of trophic modes in both rhizosphere and bulk
698 soils that are detailed above.

699 Fungal symbiotrophs and saprotrophs are among the most sensitive to management practices
700 (Hartmann et al., 2012; Orgiazzi et al., 2012; Mueller et al., 2016). For example, Orgiazzi et al. (2012)
701 found that ectomycorrhizal phylotypes were numerous in natural sites with trees but were missing in
702 anthropogenic and grass-covered sites, whereas Lauber et al. (2008) found that coprophilous fungi
703 were common in grazed sites. Management disturbances have been found to increase the relative
704 abundance of Ascomycota and decrease Basidiomycota (Thomson et al., 2015). Similarly, we
705 observed the expected decrease in fungal saprotrophs and symbiotrophs at the arable site and a higher

706 relative abundance of Ascomycota, although a large proportion of unidentified trophic modes may be
707 influencing our results.

708 *Hypothesis 3: Microbial inoculation of soils and dense root systems increase soil aggregate stability*
709 *in unstructured soils.*

710 Inoculation of live soil communities had large impacts on the sterilised soil communities, irrespective
711 of soil origin (sterilised control, arable or woodland). However, the effect of the inoculation on the
712 structure of bacterial communities was dependent on the origin of the soils. This result agrees with
713 previous studies that found that texture and pore size distribution strongly determined the fate of
714 introduced microorganisms (Rutherford and Juma, 1992; Huysman and Verstraete, 1993; van Veen et
715 al., 1997). Additionally, the effect of root system type was only evident on bacterial communities in
716 the rhizosphere, but not on communities inhabiting bulk soil, which is likely due to differences in root
717 length and the increased microbial numbers in the rhizosphere carbon rich habitat (Jones et al., 2009).
718 In a previous study, we also found that the interactions between microbial community structure and
719 root traits were more intense in communities isolated from rhizosphere compared to bulk soil, and that
720 bacterial community composition was better explained by root traits than in fungal communities
721 (Merino-Martín et al., 2020). We showed that root system type only affected the structure of bacterial
722 communities in control soils that were not inoculated, stressing that the role of roots was secondary
723 compared to inoculation and initial soil structure. Correspondingly, root system type affected fungal
724 community structure in non-inoculated control soil, but it also affected significantly the non-
725 inoculated arable soil and arable soil with woodland inoculum, which may be because it had the
726 lowest relative abundance of fungal communities and the least developed mycorrhizal networks.
727 These limited effects of root system type are in agreement with other studies that proposed a hierarchy
728 in the contribution of soils and root systems (traits and exudates), with soil physicochemical
729 properties contributing most strongly to the microbiome and root traits and exudates gradually altering
730 this soil microbiome (Bever et al., 2012; Philippot et al., 2013; van der Putten et al., 2013).

731 As hypothesized, the microbial inoculation with soils increased aggregate stability in the poorly
732 structured control soil, but not in arable (with low aggregate stability) and woodland soils (with high

733 aggregate stability). This finding is in line with Barto et al. (2010), who concluded that, in highly
734 aggregated soils, abiotic factors can be more important for determining soil aggregation than biotic
735 factors. However, besides the lack of initial structure, the control soil also showed very distinct
736 physicochemical properties that can influence the effect of microbial inoculation on aggregate
737 stability, such as lower contents of SOC and soil N, and higher pH and CaCO₃ content (Al-Ani and
738 Dudas, 1988; Chenu et al., 2000; Wu et al., 2017). Even though the focus of our research was on
739 bacterial and fungal communities, we acknowledge that our soil inoculation method may have
740 included soil fauna, such as nematodes, protozoa, or collembolans that were not explored in our study
741 (van de Voorde et al., 2012).

742 *Hypothesis 4: A greater effect of inoculation on aggregate stability is expected in soils inoculated*
743 *with woodland soil.*

744 Surprisingly, we did not observe greater aggregate stability in soil inoculated with woodland
745 compared to arable soil. These two soils, woodland and arable, were intrinsically similar but with
746 significantly different aggregate stability. In a similar experiment, Duchicela et al. (2012) found
747 greater stable aggregates in pots inoculated with soil communities from remnant grasslands, compared
748 to soils inoculated with communities from disturbed post-agricultural sites. These authors suggested
749 that the use of the same suite of native prairie plant species across both remnant and disturbed soils in
750 their experiment could explain this effect, reflecting local coadaptation (Duchicela et al., 2012).
751 Furthermore, the effects of inoculation may also depend on the starting point, the incubation time and
752 conditions after inoculation and not only on the inoculum used.

753 Several laboratory inoculation experiments also revealed that mycorrhizal fungi play an important
754 role in stabilizing aggregates. Harris et al. (1964) found that even if bacteria were more important than
755 fungi in the primary stabilization of microaggregates, fungi were involved in the stabilization of
756 macroaggregates. Bearden and Petersen (2000) found that AM fungi contributed to the stabilization of
757 aggregates in a vertisol, and that the effect was significant after only one growing season. Bossuyt et
758 al. (2001) found that macroaggregate formation was positively influenced by fungal activity but was
759 not significantly influenced by residue quality or bacterial activity. Most of these inoculation studies

760 found that the mechanism by which mycorrhizal fungi increase the stability of aggregates is not only
761 related to hyphae formation, but is also through the stimulation of root growth (Bearden and Petersen,
762 2000; Graf and Frei, 2013; Bast et al., 2016). In agreement with these authors, we found that in the
763 model including root length density (RLD), there was an effect of the interaction between the
764 inoculation treatment and RLD on aggregate stability, which was remarkable, given the absence of the
765 effect of root type.

766 We did not observe any effect of inoculation and root system type on the content of glomalin related
767 soil proteins (GRSP) or ergosterol. A significant correlation between these compounds and aggregate
768 stability was found, but it was driven by the different origins of the soils. The legacy effect of soil
769 origin on these compounds emphasizes their recalcitrant nature that has been widely discussed in
770 previous research (Lorenz et al., 2007; Gispert et al., 2013a; Cissé et al., 2020). In addition, GRSP
771 does not appear to be a good predictor of fungal activity, nor of increased physical stability.

772 **6. Conclusions**

773 Our results, from field and experimental approaches, depict the complex relationships between soil
774 properties, land use and plant traits shaping microbial communities and affecting soil structure. Our
775 field study demonstrated the existence of strong relationships between land use, microbial
776 communities and soil aggregate stability. Even though we did not find consistent relationships
777 between land use gradient, aggregate stability and the relative abundance of specific bacterial trophic
778 modes, the importance of root traits shaping the relative abundance of bacterial phyla was
779 demonstrated. Rhizosphere conditions shaped bacterial communities with oligotrophic bacteria
780 conditioned by the rhizospheric environment, where labile C is found, whilst copiotrophic phyla were
781 more dependent on bulk soil conditions. Fungal communities responded to the land use gradient with
782 a lower relative abundance of saprophytic and symbiotic in disturbed sites. The inoculation
783 experiment showed that microbial inoculation affects aggregate stability in the control soil (that had
784 very low aggregate stability initially), but not in soils sterilized from the arable field and the
785 secondary woodland, possibly because in aggregated soils, abiotic factors can be more important for

786 determining soil aggregation than biotic factors. We did not observe differential effects of the
787 different inoculums (arable *versus* secondary woodland), most likely due to the short period of the
788 inoculation experiment. Again, the importance of roots for bacterial communities in the rhizosphere
789 niche was highlighted in the inoculation experiment, with root system type affecting only the structure
790 of bacterial communities but not fungal communities, and only in the rhizosphere but not in bulk
791 soils. Additionally, root system type only affected the structure of bacterial communities in control
792 soils that were not inoculated, stressing that the role of roots was secondary compared to inoculation
793 and initial soil structure. Finally, even though we did not find a significant effect of root system type
794 on aggregate stability, we have shown that the effect of the microbial inoculation was modulated by
795 root length density, highlighting the role of roots in shaping microbial communities and their effects
796 on soil structure.

797 **Acknowledgements**

798 LMM and this research were funded with a Marie Curie IEF fellowship (FP7 European program, ref.
799 626666/2013). Many thanks to Francois Pailler, Clément Furget-Bretagnon (INRAE France), Jean
800 Marc Souquet and Josiane Abadie (UMR Eco&Sols, Montpellier) for their help with laboratory work.
801 Thanks are due to Oxford University and FAI Farms, U.K. (FAI Field Station) for access to field sites.
802 A major part of this paper was written by the first author in Madrid, Spain, during the COVID-19
803 lockdown. The authors would like to thank health workers in Spain and worldwide, who risked their
804 lives to overcome the pandemic.

805 **Tables:**

806 **Table 1.** Soil physicochemical properties of sites selected for the field study (mean of the six samples
 807 collected at depths of 0-0.1 m and 0.2-0.3 m). Sand, silt, clay, soil organic carbon (SOC), total
 808 nitrogen (N) and total carbon (C) are expressed in % w/w. Available P, K and Mg are expressed in
 809 mg/l. Carbon:nitrogen (C:N) ratio is expressed in :1. F and p values are shown. Post hoc comparisons

	Ancient Woodland	Secondary Woodland	Grassland	Pasture	Arable	F value	Pr(>F)
pH	5.87 ± 0.16 a	6.38 ± 0.15 ab	6.30 ± 0.15 a	6.10 ± 0.06 a	7.00 ± 0.22 b	7.376	<0.001
P	12.73 ± 2.44 a	10.50 ± 2.53 a	14.10 ± 2.03 a	20.07 ± 2.38 a	80.47 ± 2.47 b	157.2	<0.001
K	321.17 ± 28.01 a	253.50 ± 41.88 a	254.67 ± 29.70 a	205.67 ± 15.24 a	595.00 ± 49.26 b	19.99	<0.001
Mg	256.33 ± 10.19 b	247.50 ± 5.91 ab	229.50 ± 7.10 ab	222.50 ± 5.46 ab	209.00 ± 16.65 a	3.661	0.018
Sand	35.00 ± 2.85 ac	25.67 ± 2.63 a	40.17 ± 5.31 bc	52.67 ± 2.76 b	29.17 ± 1.47 ac	10.57	<0.001
Silt	41.50 ± 1.15 a	44.00 ± 0.52 a	38.33 ± 2.62 ab	32.83 ± 1.58 b	40.50 ± 1.02 a	7.416	<0.001
Clay	23.50 ± 1.88 ab	30.33 ± 2.28 a	21.50 ± 2.92 ab	14.50 ± 1.23 b	30.33 ± 2.23 a	9.331	<0.001
Texture	Clay Loam	Clay Loam	Sandy Loam	Sandy Loam	Clay Loam	-	-
SOC	11.28 ± 1.34 bc	13.60 ± 1.38 abc	14.78 ± 1.37 ab	17.43 ± 1.46 a	8.52 ± 0.29 c	7.435	<0.001
Total Nitrogen	0.46 ± 0.06 ab	0.59 ± 0.05 ab	0.66 ± 0.05 ac	0.87 ± 0.06 c	0.42 ± 0.01 b	12.76	<0.001
Total Carbon	5.25 ± 0.73 b	6.52 ± 0.74 ab	6.97 ± 0.68 ab	8.56 ± 0.75 a	4.36 ± 0.12 b	6.202	<0.001
C:N	11.18 ± 0.25 a	11.02 ± 0.40 a	10.42 ± 0.30 a	9.85 ± 0.18 a	10.52 ± 0.46 a	2.51	0.067

810 were performed with Tukey HSD.

811

812 **Table 2.** Spearman's correlation coefficients between relative abundance of the different phyla of
 813 Bacteria and soil properties, root and litter traits for bulk soil and rhizosphere sample fractions in the
 814 field study (****P < 0.0001, ***P < 0.001, **P < 0.01, *P < 0.05). Correlations ≥ 0.65 are
 815 highlighted in bold and correlations present in only one sample fraction are highlighted in red.

	Bulk soil						Rhizosphere					
	Oligotrophic			Copiotrophic			Oligotrophic			Copiotrophic		
	Acidobacteria	Chloroflexi	Verrucomicrobia	Actinobacteria	Bacteroidetes	Proteobacteria	Acidobacteria	Chloroflexi	Verrucomicrobia	Actinobacteria	Bacteroidetes	Proteobacteria
Soil												
MWD	0.01	-0.26	-0.06	-0.13	0.38	0.38	0.41	0.13	-0.18	-0.53	0.38	0.2
pH	-0.3	0.72**	-0.08	0.44	-0.86**	-0.84****	0.2	0.4	-0.08	0.32	0.07	-0.87****
P	-0.08	0.39	-0.48	0.67**	-0.29	-0.44	0.45	0.51	-0.64*	0.2	-0.12	-0.07
Potassium	-0.2	0.49	-0.11	0.390	-0.4	-0.460	-0.09	0.2	-0.06	0.370	-0.35	-0.130
Magnesium	-0.07	-0.38	0.45	-0.71**	0.24	0.38	-0.69**	-0.65*	0.67**	-0.5	0.23	0.38
Sand	0.24	-0.55*	-0.17	-0.12	0.71**	0.70**	0.01	-0.1	-0.11	-0.13	-0.02	0.60*
Silt	-0.39	0.25	0.46	-0.18	-0.61*	-0.43	-0.37	-0.17	0.49	0.04	0.18	-0.46
Clay	-0.09	0.48	0.1	0.04	-0.61*	-0.59*	-0.02	0.04	0.09	0.08	-0.03	-0.52
SOC	0.25	-0.29	-0.07	-0.25	0.47	0.5	0.14	-0.09	-0.02	-0.36	0.15	0.28
N	0.19	-0.17	-0.08	-0.21	0.39	0.4	0.18	-0.03	-0.02	-0.38	0.16	0.21
C	0.3	-0.24	-0.09	-0.25	0.44	0.46	0.14	-0.09	-0.01	-0.36	0.09	0.27
C:N	0.60*	-0.35	0.29	-0.70**	0.24	0.4	-0.52	-0.73**	0.59*	-0.04	-0.4	0.16
Roots												
SRL	-0.24	0.55*	-0.52*	0.82***	-0.18	-0.44	0.43	0.80**	-0.69**	0.47	-0.02	-0.21
VFR	-0.16	0.29	-0.55*	0.54*	0.17	-0.04	0.41	0.75**	-0.68**	0.19	0.15	0.06
FR	0.02	-0.24	0.52*	-0.38	-0.25	-0.08	-0.32	-0.64*	0.64*	-0.15	-0.1	-0.16
MRD	0.2	-0.47	0.56*	-0.73**	0.05	0.28	-0.44	-0.80**	0.72**	-0.42	-0.11	0.14
RDMC	-0.14	-0.60*	0.64*	-0.39	0.16	0.18	-0.42	-0.90**	0.69**	-0.34	-0.15	0.17
RMD	0	-0.35	-0.05	-0.18	0.35	0.52*	0.31	0.14	-0.16	-0.31	0.45	0.22
RLD	-0.07	-0.1	-0.3	-0.03	0.42	0.54*	0.16	0.36	-0.27	-0.2	0.42	0.43
N	0.21	0.08	0.56	-0.77**	-0.38	-0.13	-0.44	-0.59*	0.71**	0.01	-0.19	-0.32
C	0.31	-0.72**	0.04	-0.13	0.46	0.34	0.06	-0.42	-0.05	0.01	-0.09	0.09
C:N	-0.21	-0.08	-0.56	0.77**	0.38	0.13	0.44	0.59*	-0.71**	-0.01	0.19	0.32
lignin	-0.11	0.17	0.35	-0.35	-0.48	-0.32	-0.22	-0.15	0.45	0.22	0.22	-0.65*
celulose	0.06	-0.09	0.24	-0.02	-0.09	-0.26	0.26	-0.24	0.13	-0.3	-0.28	-0.19
hemicelulose	-0.11	0.04	-0.63*	0.76**	0.3	0.13	0.56	0.63*	-0.78**	-0.17	0.15	0.32
Litter												
N	0.11	-0.06	0.5	-0.54*	-0.19	0	-0.57*	-0.53	0.68**	0.13	-0.13	-0.21
C	-0.01	-0.03	-0.36	0.44	0.01	-0.12	0.60*	0.31	-0.45	-0.21	0.14	-0.19
C:N	-0.09	0.05	-0.54*	0.56*	0.17	-0.01	0.61*	0.51	-0.68**	-0.16	0.15	0.09
lignin	0.14	-0.29	0.61*	-0.74**	-0.1	0.24	-0.57	-0.79**	0.76**	0.27	-0.39	0.15
celulose	0.2	0	0.36	-0.65*	0.1	-0.02	-0.2	-0.28	0.35	-0.47	0.09	0.02
hemicelulose	-0.17	0.15	-0.47	0.75**	-0.01	-0.33	0.66*	0.68*	-0.71*	-0.17	0.35	-0.32

816

817 **Table 3.** Effects of land use and sample fraction and their interactions on the structure of bacterial and
 818 fungal communities assessed with PERMANOVA. The degrees of freedom (Df), sum of squares (sum
 819 of sqs), mean of squares (mean of sqs), the F. statistic, the proportion of the variance explained by
 820 each model (R²) and probability (P) are shown.

Factors	Bacteria						Fungi					
	Df	Sums of sqs	Mean of sqs	F.Model	R ²	Pr(>F)	Df	Sums of sqs	Mean of sqs	F.Model	R ²	Pr(>F)
Land use	4	1.5469	0.38673	4.4095	0.438	0.001	4	4.8003	1.20008	6.3216	0.533	0.001
Samplefraction	1	0.1366	0.1366	1.5575	0.039	0.093	1	0.1365	0.13649	0.719	0.015	0.745
Land use*Samplefraction	4	0.1793	0.04483	0.5112	0.051	1	4	0.4629	0.11572	0.6096	0.051	0.993
Residuals	19	1.6663	0.0877		0.472		19	3.6069	0.18984		0.400	
Total	28	3.5292			1		28	9.0067			1	

821 **Table 4.** Effects of inoculation, origin and root treatments and their interactions on the structure of
 822 bacterial and fungal communities assessed with PERMANOVA. The degrees of freedom (Df), sum of
 823 squares (sum of sqs), mean of squares (mean of sqs), the F. statistic, the proportion of the variance
 824 explained by each model (R²) and probability (P) are shown.

Bulk soil												
Factors	Bacteria						Fungi					
	Df	Sums of sqs	Mean of sqs	F.Model	R ²	Pr(>F)	Df	Sums of sqs	Mean of sqs	F.Model	R ²	Pr(>F)
Inoculation	2	9.214	4.6072	18.5425	0.182	0.001	2	10.144	5.0721	15.0325	0.164	0.001
Origin	2	2.603	1.3017	5.239	0.051	0.001	2	1.703	0.8513	2.523	0.028	0.001
Rootsystem	2	0.709	0.3547	1.4277	0.014	0.074	2	1.109	0.5545	1.6434	0.018	0.004
Inoculation*Origin	4	2.185	0.5463	2.1986	0.043	0.001	4	2.622	0.6554	1.9424	0.042	0.001
Inoculation*Rootsystem	4	1.042	0.2604	1.0481	0.02	0.34	4	1.902	0.4756	1.4095	0.03078	0.01
Origin*Rootsystem	4	0.919	0.2297	0.9247	0.018	0.623	4	1.697	0.4243	1.2575	0.027	0.042
Inoculation*Origin*Rootsystem	8	1.745	0.2181	0.8777	0.034	0.818	t 8	3.142	0.3927	1.164	0.051	0.059
Residuals	130	32.301	0.2485		0.64		117	39.477	0.3374		0.63883	
Total	156	50.718			1.000		143	61.795			1.000	

Rhizosphere												
Factors	Bacteria						Fungi					
	Df	Sums of sqs	Mean of sqs	F.Model	R ²	Pr(>F)	Df	Sums of sqs	Mean of sqs	F.Model	R ²	Pr(>F)
Inoculation	2	3.5166	1.7583	12.2209	0.264	0.001	2	4.7153	2.35765	8.9631	0.241	0.001
Origin	2	1.1993	0.59963	4.1677	0.090	0.001	2	1.5272	0.7636	2.903	0.078	0.001
Rootsystem	1	0.8523	0.85235	5.9242	0.064	0.001	1	0.3279	0.32786	1.2464	0.017	0.182
Inoculation*Origin	4	1.4709	0.36771	2.5558	0.110	0.001	4	2.0624	0.51559	1.9601	0.105	0.001
Inoculation*Rootsystem	2	0.394	0.197	1.3692	0.03	0.086	2	0.6301	0.31504	1.1977	0.03218	0.167
Origin*Rootsystem	2	0.3101	0.15505	1.0777	0.023	0.329	2	0.5685	0.28424	1.0806	0.029	0.312
Inoculation*Origin*Rootsystem	4	0.7098	0.17746	1.2334	0.053	0.094	4	1.066	0.2665	1.0132	0.054	0.429
Residuals	34	4.8918	0.14388		0.37		33	8.6803	0.26304		0.44338	
Total	51	13.3448			1.000		50	19.5776			1.000	

825

826

827 **Table 5.** Effects of root system type and inoculation in each of the combinations of the experimental
 828 design. The ANOSIM statistic R and the significance values are shown. The proportion of the
 829 variance explained by each model (R^2) and probability (P) are shown.

Origin	Inoculation	Effect of root system			
		Bacteria		Fungi	
		R^2	Pr(>F)	R^2	Pr(>F)
Arable	Arable	-0.029	0.551	-0.008	0.520
Arable	NotInoculated	0.016	0.381	0.341	0.001
Arable	Woodland	0.005	0.398	0.251	0.015
Control	Arable	-0.080	0.818	-0.075	0.797
Control	NotInoculated	0.172	0.013	0.687	0.012
Control	Woodland	-0.021	0.461	-0.029	0.564
Woodland	Arable	-0.047	0.624	0.019	0.355
Woodland	NotInoculated	0.108	0.121	0.056	0.213
Woodland	Woodland	0.013	0.367	-0.004	0.483
Origin	Root system	Effect of inoculation			
		Bacteria		Fungi	
		R^2	Pr(>F)	R^2	Pr(>F)
Arable	Fibrous	0.460	0.001	0.709	0.001
Arable	Noroot	0.362	0.003	0.577	0.001
Arable	Tap	0.398	0.004	0.753	0.001
Control	Fibrous	0.616	0.001	0.825	0.001
Control	Noroot	0.501	0.002	0.798	0.001
Control	Tap	0.505	0.002	0.497	0.001
Woodland	Fibrous	0.513	0.001	0.789	0.001
Woodland	Noroot	0.590	0.001	0.612	0.001
Woodland	Tap	0.563	0.001	0.820	0.001

830

831

832

833

834

835

836

837

838


839

840 **Table 6.** Effects of inoculation, soil origin and root system type and their interactions on soil
841 aggregate stability assessed with ANOVA. An additional ANCOVA model was tested including root
842 length density (RLD), to separate the effect of species and physical properties (quantity of roots)
843 within the effect of root system treatment. The degrees of freedom (Df), sum of squares (Sum of sqs),
844 mean of squares (Mean of sqs), the F. statistic and p-value (P) associated with the F statistic are
845 shown.

846

Factors	Df	Sums of sqs	Mean of sqs	F.Model	Pr(>F)
Origin	2	45.76	22.881	460.529	<0.001
Inoculation	2	0.33	0.166	3.348	0.043
Root system	2	0.07	0.033	0.656	0.523
Origin*Inoculation	4	1.76	0.439	8.833	<0.001
Origin*Root system	4	0.3	0.076	1.531	0.206
Inoculation*Root system	4	0.06	0.016	0.318	0.865
Origin*Inoculation*Root system	8	0.37	0.046	0.92	0.507
Residuals	54	2.68	0.05		
Model including RLD					
Origin	2	45.76	22.88	871.82	<0.001
Inoculation	2	0.33	0.166	6.343	0.005
Root system	2	0.07	0.033	1.246	0.302
RLD	1	0.04	0.038	1.457	0.237
Origin*Inoculation	4	1.89	0.472	17.994	<0.001
Origin*Root system	4	0.24	0.059	2.264	0.085
Inoculation*Root system	4	0.04	0.01	0.385	0.817
Origin*RLD	2	0.01	0.006	0.22	0.803
Inoculation*RLD	2	0.6	0.301	11.462	0.000
Root system*RLD	2	0.04	0.019	0.707	0.501
Origin*Inoculation*Root system	8	0.38	0.048	1.83	0.109
Origin*Inoculation*RLD	4	0.38	0.096	3.649	0.015
Origin*Root system*RLD	4	0.37	0.092	3.5	0.018
Inoculation*Root system*RLD	3	0.31	0.104	3.955	0.017
Origin*Inoculation*Root system*RLD	5	0.06	0.011	0.43	0.824
Residuals	31	0.81	0.026		

847 **Figures:**



848


849

850 **Figure 1.** Location of sampling plots spanning a range of land uses. Red dots correspond to plots
851 where sampling was performed. (Source: 51°46'46.91''N and 1°19'35.76''. Google Earth
852 v7.3.2.5776, May 20, 2018. May 11, 2020).

853




854




861 **Figure 2.** Schematic illustration of the treatments under study with three soil origins, three root
862 system types (fibrous root system of *Brachypodium sylvaticum*, taproot system of *Urtica dioica* and a
863 control with no roots present) and three inoculation treatments. Different coloured bars on tops of pots
864 refer to the source of soil inoculum (see legend).

865




873 **Figure 3.** Soil aggregate stability (MWD) measured in the field in the different land uses at the
 874 surface and 0.2 – 0.3 m depth. Boxplots represent the minimum, maximum, median, first quartile and
 875 third quartile in the data set. Different letters show post-hoc Tukey honestly significant difference
 876 (HSD) results between land uses for the two different depths.

877




878 **Figure 4.** Boxplots with relative abundance of phyla found in the three plots in each land use in
 879 rhizosphere and bulk soil fractions for bacteria (a) and fungi (b) and proportion of fungal trophic
 880 modes (c). Boxplots represent the minimum, maximum, median, first quartile and third quartile in the
 881 data set. Different letters show post-hoc Tukey honestly significant difference (HSD) results. Letters
 882 out of the boxplot figure show the overall effect including rhizosphere and bulk soils.




892 **Figure 5.** a) NMDS ordinations of bacteria and fungi species matrices with convex hull polygons and
 893 spider diagrams containing plots of the five sites and overlaid gradients of aggregate stability and b)
 894 graphs of dbRDA constrained ordinations of bacteria and fungi species matrices with convex hull
 895 polygons and spider diagrams containing plots of the five sites and significant variables obtained by
 896 automatic backward stepwise model building. Data are shown for rhizosphere (□) and bulk soil (○)
 897 fractions. Acronyms for land use types: AW is ancient woodland, SW is secondary woodland, GR is
 898 grassland, PA is pasture, and AR is arable. Acronyms for factors: S_pH is soil pH, S_P is soil P,
 899 MWD is mean weight diameter, MRD is mean root diameter, FR is % fine roots, RLD is root length
 900 density.

901




909 **Figure 6.** PCA of variables used for the description of soil, vegetation and root properties in the
 910 experiment. Variables showed were selected based on their correlation and ecological significance
 911 (i.e. variables that were correlated but their source was ecologically different were kept, e.g. soil C:N
 912 and root C:N). Acronyms: S_pH is soil pH, S_C:N is soil C:N ratio, S_WC is soil water content,
 913 S_CO2 is rate of exchange of CO₂, S_OC is soil organic carbon, MWD is mean weight diameter,
 914 Ab_biomass is Aboveground biomass, R_C:N is root C:N ratio, VFR is % very fine roots, RMD is
 915 root mean diameter, RLD is root length density, MRD is mean root diameter, SRL is specific root
 916 length, GRSPT is total glomalin related soil proteins.



917

918 **Figure 7.** Selected specific traits of soil samples after incubation period: a) soil aggregate stability
 919 (mean weight diameter, MWD), b) root length density (RLD, km m^{-3}), c) specific root length (SRL, m
 920 gr^{-1}), d) soil pH, e) soil organic carbon (SOC, ww^{-1}), f) soil nitrogen (N, ww^{-1}), g) soil total inorganic
 921 carbon (CaCO_3 , ww^{-1}). Boxplots represent the minimum, maximum, median, first quartile and third
 922 quartile in the data set.



935 **Figure 8.** a) NMDS ordinations of bacteria and fungi species matrices with convex hull polygons
 936 containing plots of the three inoculation treatments (not inoculated, woodland and arable) and b)
 937 graphs of dbRDA constrained ordinations of bacteria and fungi species matrices with convex hull
 938 polygons containing plots of the three inoculation treatments (not inoculated, woodland and arable)
 939 and significant variables obtained by automatic backward stepwise model building. Acronyms are:
 940 S_C is soil total carbon, S_C.N is soil C:N ratio, MWD is mean weight diameter, RMD is root mean
 941 diameter, GRSPT is total glomalin related soil proteins.

942

943 **References:**

- 944 Adhikari, K., Hartemink, A.E., 2016. Linking soils to ecosystem services — A global review. *Geoderma*
945 262, 101-111.
- 946 Al-Ani, A.N., Dudas, M.J., 1988. Influence of calcium carbonate on mean weight diameter of soil. *Soil*
947 *and Tillage Research* 11, 19-26.
- 948 Ayuke, F.O., Brussaard, L., Vanlauwe, B., Six, J., Lelei, D.K., Kibunja, C.N., Pulleman, M.M., 2011. Soil
949 fertility management: Impacts on soil macrofauna, soil aggregation and soil organic matter
950 allocation. *Applied Soil Ecology* 48, 53-62.
- 951 Balesdent, J., Chenu, C., Balabane, M., 2000. Relationship of soil organic matter dynamics to physical
952 protection and tillage. *Soil and Tillage Research* 53, 215-230.
- 953 Bardgett, R.D., Leemans, D.K., Cook, R., Hobbs, P.J., 1997. Seasonality of the soil biota of grazed and
954 ungrazed hill grasslands. *Soil Biology and Biochemistry* 29, 1285-1294.
- 955 Barto, E.K., Alt, F., Oelmann, Y., Wilcke, W., Rillig, M.C., 2010. Contributions of biotic and abiotic
956 factors to soil aggregation across a land use gradient. *Soil Biology and Biochemistry* 42, 2316-2324.
- 957 Bast, A., Wilcke, W., Graf, F., Lüscher, P., Gärtner, H., 2016. Does mycorrhizal inoculation improve
958 plant survival, aggregate stability, and fine root development on a coarse-grained soil in an alpine
959 eco-engineering field experiment? *Journal of Geophysical Research: Biogeosciences* 121, 2158-2171.
- 960 Baumert, V.L., Vasilyeva, N.A., Vladimirov, A.A., Meier, I.C., Kögel-Knabner, I., Mueller, C.W., 2018.
961 Root Exudates Induce Soil Macroaggregation Facilitated by Fungi in Subsoil. *Frontiers in*
962 *Environmental Science* 6.
- 963 Bearden, B., Petersen, L., 2000. Influence of arbuscular mycorrhizal fungi on soil structure and
964 aggregate stability of a vertisol. *Plant and Soil*, 173-183.
- 965 Bergmann, G.T., Bates, S.T., Eilers, K.G., Lauber, C.L., Caporaso, J.G., Walters, W.A., Knight, R., Fierer,
966 N., 2011. The under-recognized dominance of Verrucomicrobia in soil bacterial communities. *Soil*
967 *biology & biochemistry* 43, 1450-1455.
- 968 Bever, J.D., Platt, T.G., Morton, E.R., 2012. Microbial population and community dynamics on plant
969 roots and their feedbacks on plant communities. *Annual Review of Microbiology* 66, 265-283.
- 970 Bossuyt, H., Denef, K., Six, J., Frey, S.D., Merckx, R., Paustian, K., 2001. Influence of microbial
971 populations and residue quality on aggregate stability. *Applied Soil Ecology* 16, 195-208.
- 972 Brax, M., Buchmann, C., Kenngott, K., Schaumann, G.E., Diehl, D., 2020. Influence of the physico-
973 chemical properties of root mucilage and model substances on the microstructural stability of sand.
974 *Biogeochemistry* 147, 35-52.
- 975 Brunel, C., Pouteau, R., Dawson, W., Pester, M., Ramirez, K.S., van Kleunen, M., 2020. Towards
976 Unraveling Macroecological Patterns in Rhizosphere Microbiomes. *Trends in Plant Science*.
- 977 Bulgarelli, D., Garrido-Oter, R., Münch, Philipp C., Weiman, A., Dröge, J., Pan, Y., McHardy, Alice C.,
978 Schulze-Lefert, P., 2015. Structure and Function of the Bacterial Root Microbiota in Wild and
979 Domesticated Barley. *Cell Host & Microbe* 17, 392-403.
- 980 Bulgarelli, D., Rott, M., Schlaeppi, K., Ver Loren van Themaat, E., Ahmadinejad, N., Assenza, F., Rauf,
981 P., Huettel, B., Reinhardt, R., Schmelzer, E., Peplies, J., Gloeckner, F.O., Amann, R., Eickhorst, T.,
982 Schulze-Lefert, P., 2012. Revealing structure and assembly cues for Arabidopsis root-inhabiting
983 bacterial microbiota. *Nature* 488, 91-95.
- 984 Butt, N., Campbell, G., Malhi, Y., Morecroft, M., Fenn, K., Thomas, M., 2009. Initial Results from
985 Establishment of a Long-term Broadleaf Monitoring Plot at Wytham Woods, Oxford, UK.
- 986 Carbonetto, B., Rascovan, N., Álvarez, R., Mentaberry, A., Vázquez, M.P., 2014. Structure,
987 composition and metagenomic profile of soil microbiomes associated to agricultural land use and
988 tillage systems in Argentine Pampas. *PLOS ONE* 9, e99949-e99949.
- 989 Carter, M., Kunelius, H., Angers, D., 1994. Soil Structural Form and Stability, and Organic Matter
990 Under Cool-Season Perennial Grasses. *Soil Science Society of America Journal - SSSAJ* 58.
- 991 Chenu, C., Le Bissonnais, Y., Arrouays, D., 2000. Organic Matter Influence on Clay Wettability and Soil
992 Aggregate Stability. *Soil Science Society of America Journal* 64, 1479-1486.

993 Chenu, C., Sotzky, G., 2002. Interactions Between Microorganisms and Soil Particles, Interactions
994 between soil particles and microorganisms, pp. 3-40.

995 Chotte, J.-L., 2005. Importance of Microorganisms for Soil Aggregation, *Microorganisms in Soils:*
996 *Roles in Genesis and Functions*, pp. 107-119.

997 Cissé, G., van Oort, F., Chenu, C., Essi, M., Staunton, S., 2020. Is the operationally defined fraction of
998 soil organic matter, "GRSP" (glomalin-related soil protein), stable in soils? Evidence from trends in
999 long-term bare fallow soil. *European Journal of Soil Science* 10.1111/ejss.12974.

1000 Commander, L.E., Merritt, D.J., Rokich, D.P., Dixon, K.W., 2009. Seed biology of Australian arid zone
1001 species: Germination of 18 species used for rehabilitation. *Journal of Arid Environments* 73, 617-625.

1002 Cui, J., Holden, N.M., 2015. The relationship between soil microbial activity and microbial biomass,
1003 soil structure and grassland management. *Soil and Tillage Research* 146, 32-38.

1004 de Castro, A.P., Quirino, B.F., Pappas, G., Jr., Kurokawa, A.S., Neto, E.L., Kruger, R.H., 2008. Diversity
1005 of soil fungal communities of Cerrado and its closely surrounding agriculture fields. *Archives of*
1006 *Microbiology* 190, 129-139.

1007 Degens, B.P., 1997. Macro-aggregation of soils by biological bonding and binding mechanisms and
1008 the factors affecting these: a review. *Soil Research* 35, 431-460.

1009 Deleegn, Y.T., Purahong, W., Blazevic, A., Yitafaru, B., Wubet, T., Göransson, H., Godbold, D.L., 2017.
1010 Changes in land use alter soil quality and aggregate stability in the highlands of northern Ethiopia.
1011 *Scientific Reports* 7, 13602.

1012 Denef, K., Six, J., 2005. Clay mineralogy determines the importance of biological versus abiotic
1013 processes for macroaggregate formation and stabilization. *European Journal of Soil Science* 56, 469-
1014 479.

1015 Dennis, P.G., Miller, A.J., Hirsch, P.R., 2010. Are root exudates more important than other sources of
1016 rhizodeposits in structuring rhizosphere bacterial communities? *FEMS Microbiology Ecology* 72, 313-
1017 327.

1018 DeSantis, T.Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E.L., Keller, K., Huber, T., Dalevi, D., Hu,
1019 P., Andersen, G.L., 2006. Greengenes, a Chimera-Checked 16S rRNA Gene Database and Workbench
1020 Compatible with ARB. *Applied and Environmental Microbiology* 72, 5069-5072.

1021 Donn, S., Kirkegaard, J.A., Perera, G., Richardson, A.E., Watt, M., 2015. Evolution of bacterial
1022 communities in the wheat crop rhizosphere. *Environmental Microbiology* 17, 610-621.

1023 Dray, S., Dufour, A.-B., 2007. The ade4 Package: Implementing the Duality Diagram for Ecologists.
1024 *Journal of Statistical Software* 22, 1-20.

1025 Drenovsky, R.E., Steenwerth, K.L., Jackson, L.E., Scow, K.M., 2010. Land use and climatic factors
1026 structure regional patterns in soil microbial communities. *Global Ecology and Biogeography* 19, 27-
1027 39.

1028 Duchicela, J., Sullivan, T.S., Bontti, E., Bever, J.D., 2013. Soil aggregate stability increase is strongly
1029 related to fungal community succession along an abandoned agricultural field chronosequence in
1030 the Bolivian Altiplano. *Journal of Applied Ecology*.

1031 Duchicela, J., Vogelsang, K.M., Schultz, P.a., Kaonongbua, W., Middleton, E.L., Bever, J.D., 2012. Non-
1032 native plants and soil microbes: potential contributors to the consistent reduction in soil aggregate
1033 stability caused by the disturbance of North American grasslands. *New Phytologist* 196, 212-222.

1034 Eilers, K.G., Lauber, C.L., Knight, R., Fierer, N., 2010. Shifts in bacterial community structure
1035 associated with inputs of low molecular weight carbon compounds to soil. *Soil Biology and*
1036 *Biochemistry* 42, 896-903.

1037 Fanin, N., Bertrand, I., 2016. Aboveground litter quality is a better predictor than belowground
1038 microbial communities when estimating carbon mineralization along a land-use gradient. *Soil*
1039 *Biology and Biochemistry* 94, 48-60.

1040 Fierer, N., Bradford, M.A., Jackson, R.B., 2007. Toward an ecological classification of soil bacteria.
1041 *Ecology* 88, 1354-1364.

1042 Fierer, N., Lauber, C.L., Ramirez, K.S., Zaneveld, J., Bradford, M.A., Knight, R., 2012. Comparative
1043 metagenomic, phylogenetic and physiological analyses of soil microbial communities across nitrogen
1044 gradients. *The ISME Journal* 6, 1007-1017.

1045 Fitter, A.H., Peat, H.J., 1994. The Ecological Flora Database. *Journal of Ecology* 82, 415-425.

1046 Gispert, M., Emran, M., Pardini, G., Doni, S., Ceccanti, B., 2013a. The impact of land management
1047 and abandonment on soil enzymatic activity, glomalin content and aggregate stability. *Geoderma*
1048 202-203, 51-61.

1049 Gispert, M., Emran, M., Pardini, G., Doni, S., Ceccanti, B., 2013b. The impact of land management
1050 and abandonment on soil enzymatic activity, glomalin content and aggregate stability. *Geoderma*
1051 202-203, 51-61.

1052 Goebel, M.-O., Woche, S.K., Bachmann, J., 2009. Do soil aggregates really protect encapsulated
1053 organic matter against microbial decomposition? *Biologia* 64, 443-448.

1054 Graf, F., Frei, M., 2013. Soil aggregate stability related to soil density, root length, and mycorrhiza
1055 using site-specific *Alnus incana* and *Melanogaster variegatus* s.l. *Ecological Engineering* 57, 314-323.

1056 Gupta, V.V.S.R., Germida, J.J., 2015. Soil aggregation: Influence on microbial biomass and
1057 implications for biological processes. *Soil Biology and Biochemistry* 80, A3-A9.

1058 Gweon, H.S., Oliver, A., Taylor, J., Booth, T., Gibbs, M., Read, D.S., Griffiths, R.I., Schonrogge, K.,
1059 2015. PIPITS: an automated pipeline for analyses of fungal internal transcribed spacer sequences
1060 from the Illumina sequencing platform. *Methods in Ecology and Evolution* 6, 973-980.

1061 Harris, R.F., Chesters, G., Allen, O.N., Attoe, O.J., 1964. Mechanisms Involved in Soil Aggregate
1062 Stabilization by Fungi and Bacteria. *Soil Science Society of America Journal* 28, 529-532.

1063 Hartmann, M., Howes, C.G., VanInsberghe, D., Yu, H., Bachar, D., Christen, R., Henrik Nilsson, R.,
1064 Hallam, S.J., Mohn, W.W., 2012. Significant and persistent impact of timber harvesting on soil
1065 microbial communities in Northern coniferous forests. *The ISME Journal* 6, 2199-2218.

1066 Helgason, B.L., Walley, F.L., Germida, J.J., 2010. No-till soil management increases microbial biomass
1067 and alters community profiles in soil aggregates. *Applied Soil Ecology* 46, 390-397.

1068 Ho, A., Di Lonardo, D.P., Bodelier, P.L.E., 2017. Revisiting life strategy concepts in environmental
1069 microbial ecology. *FEMS Microbiology Ecology* 93.

1070 Hudek, C., Stanchi, S., D'Amico, M., Freppaz, M., 2017. Quantifying the contribution of the root
1071 system of alpine vegetation in the soil aggregate stability of moraine. *International Soil and Water*
1072 *Conservation Research* 5, 36-42.

1073 Hurisso, T.T., Moebius-Clune, D.J., Culman, S.W., Moebius-Clune, B.N., Thies, J.E., van Es, H.M., 2018.
1074 Soil Protein as a Rapid Soil Health Indicator of Potentially Available Organic Nitrogen. *Agricultural &*
1075 *Environmental Letters* 3, 180006.

1076 Huysman, F., Verstraete, W., 1993. Effect of cell surface characteristics on the adhesion of bacteria
1077 to soil particles. *Biology and Fertility of Soils* 16, 21-26.

1078 Ihrmark, K., Bodeker, I.T., Cruz-Martinez, K., Friberg, H., Kubartova, A., Schenck, J., Strid, Y., Stenlid,
1079 J., Brandstrom-Durling, M., Clemmensen, K.E., Lindahl, B.D., 2012. New primers to amplify the fungal
1080 ITS2 region--evaluation by 454-sequencing of artificial and natural communities. *FEMS Microbiology*
1081 *Ecology* 82, 666-677.

1082 Jie, C., Jing-zhang, C., Man-zhi, T., Zi-tong, G., 2002. Soil degradation: a global problem endangering
1083 sustainable development. *Journal of Geographical Sciences* 12, 243-252.

1084 Jones, D.L., Nguyen, C., Finlay, R.D., 2009. Carbon flow in the rhizosphere: carbon trading at the soil-
1085 root interface. *Plant and Soil* 321, 5-33.

1086 Koljalg, U., Nilsson, R.H., Abarenkov, K., Tedersoo, L., Taylor, A.F., Bahram, M., Bates, S.T., Bruns,
1087 T.D., Bengtsson-Palme, J., Callaghan, T.M., Douglas, B., Drenkhan, T., Eberhardt, U., Duenas, M.,
1088 Grebenc, T., Griffith, G.W., Hartmann, M., Kirk, P.M., Kohout, P., Larsson, E., Lindahl, B.D., Lucking,
1089 R., Martin, M.P., Matheny, P.B., Nguyen, N.H., Niskanen, T., Oja, J., Peay, K.G., Peintner, U., Peterson,
1090 M., Poldmaa, K., Saag, L., Saar, I., Schussler, A., Scott, J.A., Senes, C., Smith, M.E., Suija, A., Taylor,
1091 D.L., Telleria, M.T., Weiss, M., Larsson, K.H., 2013. Towards a unified paradigm for sequence-based
1092 identification of fungi. *Molecular Ecology* 22, 5271-5277.

1093 Kozich, J.J., Westcott, S.L., Baxter, N.T., Highlander, S.K., Schloss, P.D., 2013. Development of a dual-
1094 index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq
1095 Illumina sequencing platform. *Applied and Environmental Microbiology* 79, 5112-5120.

1096 Lauber, C.L., Hamady, M., Knight, R., Fierer, N., 2009. Pyrosequencing-Based Assessment of Soil pH
1097 as a Predictor of Soil Bacterial Community Structure at the Continental Scale. *Applied and*
1098 *Environmental Microbiology* 75, 5111-5120.

1099 Lauber, C.L., Strickland, M.S., Bradford, M.A., Fierer, N., 2008. The influence of soil properties on the
1100 structure of bacterial and fungal communities across land-use types. *Soil Biology and Biochemistry*
1101 40, 2407-2415.

1102 Lavee, H., Sarah, P., Imeson, A.C., 1996. Aggregate Stability Dynamics as Affected by Soil
1103 Temperature and Moisture Regimes. *Geografiska Annaler. Series A, Physical Geography* 78, 73-82.

1104 Lavelle, P., Spain, A., Fonte, S., Bedano, J.C., Blanchart, E., Galindo, V., Grimaldi, M., Jimenez, J.J.,
1105 Velasquez, E., Zangerlé, A., 2020. Soil aggregation, ecosystem engineers and the C cycle. *Acta*
1106 *Oecologica* 105, 103561.

1107 Le Bissonnais, Y., 1996. Aggregate stability and assessment of soil crustability and erodibility: I.
1108 Theory and methodology. *European Journal of Soil Science* 47, 425-437.

1109 Lee, K., Foster, R., 1991. Soil fauna and soil structure. *Soil Research* 29, 745-775.

1110 Lehmann, A., Zheng, W., Rillig, M.C., 2017. Soil biota contributions to soil aggregation. *Nat Ecol Evol*
1111 1, 1828-1835.

1112 Lladó, S., Baldrian, P., 2017. Community-level physiological profiling analyses show potential to
1113 identify the copiotrophic bacteria present in soil environments. *PLOS ONE* 12, e0171638-e0171638.

1114 Lorenz, K., Lal, R., Preston, C.M., Nierop, K.G.J., 2007. Strengthening the soil organic carbon pool by
1115 increasing contributions from recalcitrant aliphatic bio(macro)molecules. *Geoderma* 142, 1-10.

1116 Mao, Y., Li, X., Smyth, E.M., Yannarell, A.C., Mackie, R.I., 2014. Enrichment of specific bacterial and
1117 eukaryotic microbes in the rhizosphere of switchgrass (*Panicum virgatum* L.) through root exudates.
1118 *Environmental Microbiology Reports* 6, 293-306.

1119 Marín, C., Godoy, R., Valenzuela, E., Schloter, M., Wubet, T., Boy, J., Gschwendtner, S., 2017.
1120 Functional land-use change effects on soil fungal communities in Chilean temperate rainforests.
1121 *Journal of Soil Science and Plant Nutrition* 17, 985-1002.

1122 Merino-Martín, L., Griffiths, R.I., Gweon, H.S., Furget-Bretagnon, C., Oliver, A., Mao, Z., Le Bissonnais,
1123 Y., Stokes, A., 2020. Rhizosphere bacteria are more strongly related to plant root traits than fungi in
1124 temperate montane forests: insights from closed and open forest patches along an elevational
1125 gradient. *Plant and Soil*.

1126 Miller, R., Jastrow, J., 1992. The Role of Mycorrhizal Fungi in Soil Conservation, pp. 29-44.

1127 Miller, R.M., Jastrow, J.D., 1990. Hierarchy of root and mycorrhizal fungal interactions with soil
1128 aggregation. *Soil Biology and Biochemistry* 22, 579-584.

1129 Morel, J.L., Habib, L., Plantureux, S., Guckert, A., 1991. Influence of maize root mucilage on soil
1130 aggregate stability. *Plant and Soil* 136, 111-119.

1131 Mueller, R.C., Rodrigues, J.L.M., Nüsslein, K., Bohannan, B.J.M., 2016. Land use change in the
1132 Amazon rain forest favours generalist fungi. *Functional Ecology* 30, 1845-1853.

1133 Oades, J., Waters, A., 1991. Aggregate hierarchy in soils. *Soil Research* 29, 815-828.

1134 Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., Minchin, P.R., O'Hara,
1135 R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., 2019. vegan: Community
1136 Ecology Package, R package version 2.5-6. <https://CRAN.R-project.org/package=vegan>.

1137 Or, D., Smets, B.F., Wraith, J.M., Dechesne, A., Friedman, S.P., 2007. Physical constraints affecting
1138 bacterial habitats and activity in unsaturated porous media – a review. *Advances in Water Resources*
1139 30, 1505-1527.

1140 Orgiazzi, A., Lumini, E., Nilsson, R.H., Girlanda, M., Vizzini, A., Bonfante, P., Bianciotto, V., 2012.
1141 Unravelling Soil Fungal Communities from Different Mediterranean Land-Use Backgrounds. *PLOS*
1142 *ONE* 7, e34847.

1143 Peiffer, J.A., Spor, A., Koren, O., Jin, Z., Tringe, S.G., Dangl, J.L., Buckler, E.S., Ley, R.E., 2013. Diversity
1144 and heritability of the maize rhizosphere microbiome under field conditions. *Proceedings of the*
1145 *National Academy of Sciences* 110, 6548-6553.

1146 Peng, X., Horn, R., Hallett, P., 2015. Soil structure and its functions in ecosystems: Phase matter &
1147 scale matter. *Soil and Tillage Research* 146, 1-3.

1148 Philippot, L., Raaijmakers, J.M., Lemanceau, P., van der Putten, W.H., 2013. Going back to the roots:
1149 the microbial ecology of the rhizosphere. *Nature Reviews. Microbiology* 11, 789-799.

1150 Plassard, C., Bonafos, B., Touraine, B., 2000. Differential effects of mineral and organic N sources,
1151 and of ectomycorrhizal infection by *Hebeloma cylindrosporum*, on growth and N utilization in *Pinus*
1152 *pinaster*. *Plant, Cell & Environment* 23, 1195-1205.

1153 Poirier, V., Roumet, C., Angers, D.A., Munson, A.D., 2018. Species and root traits impact
1154 macroaggregation in the rhizospheric soil of a Mediterranean common garden experiment. *Plant*
1155 *and Soil* 424, 289-302.

1156 R Core Team, 2019. R: A Language and Environment for Statistical Computing. R Foundation for
1157 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

1158 Rillig, M.C., Lehmann, A., Aguilar-Trigueros, C.A., Antonovics, J., Caruso, T., Hempel, S., Lehmann, J.,
1159 Valyi, K., Verbruggen, E., Veresoglou, S.D., Powell, J.R., 2016. Soil microbes and community
1160 coalescence. *Pedobiologia* 59, 37-40.

1161 Rillig, M.C., Muller, L.A.H., Lehmann, A., 2017. Soil aggregates as massively concurrent evolutionary
1162 incubators. *ISME J* 11, 1943-1948.

1163 Rillig, M.C., Mummey, D.L., 2006. Mycorrhizas and soil structure. *New Phytologist* 171, 41-53.

1164 Rillig, M.C., Wright, S.F., Eviner, V.T., 2002a. The role of arbuscular mycorrhizal fungi and glomalin in
1165 soil aggregation: comparing effects of five plant species. *Plant and Soil* 238, 325-333.

1166 Rillig, M.C., Wright, S.F., Shaw, M.R., Field, C.B., 2002b. Artificial climate warming positively affects
1167 arbuscular mycorrhizae but decreases soil aggregate water stability in an annual grassland. *Oikos* 97,
1168 52-58.

1169 Rognes, T., Flouri, T., Nichols, B., Quince, C., Mahé, F., 2016. VSEARCH: a versatile open source tool
1170 for metagenomics. *PeerJ* 4, e2584.

1171 Rutherford, P.M., Juma, N.G., 1992. Influence of texture on habitable pore space and bacterial-
1172 protozoan populations in soil. *Biology and Fertility of Soils* 12, 221-227.

1173 Saha, D., Kukal, S.S., Sharma, S., 2011. Landuse impacts on SOC fractions and aggregate stability in
1174 typic ustochrepts of Northwest India. *Plant and Soil* 339, 457-470.

1175 Saleem, M., Law, A.D., Sahib, M.R., Pervaiz, Z.H., Zhang, Q., 2018. Impact of root system architecture
1176 on rhizosphere and root microbiome. *Rhizosphere* 6, 47-51.

1177 Sarathchandra, S.U., Perrott, K.W., Boase, M.R., Waller, J.E., 1988. Seasonal changes and the effects
1178 of fertiliser on some chemical, biochemical and microbiological characteristics of high-producing
1179 pastoral soil. *Biology and Fertility of Soils* 6, 328-335.

1180 Savill, P.S.S., Perrins, C., Kirby, K., Fisher, N., 2011. Wytham Woods: Oxford's Ecological Laboratory.
1181 *Wytham Woods: Oxford's Ecological Laboratory*, 1-288.

1182 Shi, S., Nuccio, E., Herman, D.J., Rijkers, R., Estera, K., Li, J., da Rocha, U.N., He, Z., Pett-Ridge, J.,
1183 Brodie, E.L., Zhou, J., Firestone, M., 2015. Successional Trajectories of Rhizosphere Bacterial
1184 Communities over Consecutive Seasons. *mBio* 6, e00746-00715.

1185 Six, J., Bossuyt, H., Degryze, S., Denef, K., 2004. A history of research on the link between
1186 (micro)aggregates, soil biota, and soil organic matter dynamics. *Soil and Tillage Research* 79, 7-31.

1187 Six, J., Elliott, E.T., Paustian, K., Doran, J.W., 1998. Aggregation and Soil Organic Matter Accumulation
1188 in Cultivated and Native Grassland Soils. *Soil Science Society of America Journal* 62, 1367-1377.

1189 Steenwerth, K.L., Jackson, L.E., Calderón, F.J., Stromberg, M.R., Scow, K.M., 2002. Soil microbial
1190 community composition and land use history in cultivated and grassland ecosystems of coastal
1191 California. *Soil Biology and Biochemistry* 34, 1599-1611.

1192 Thomson, B.C., Tisserant, E., Plassart, P., Uroz, S., Griffiths, R.I., Hannula, S.E., Buée, M., Mougel, C.,
1193 Ranjard, L., Van Veen, J.A., Martin, F., Bailey, M.J., Lemanceau, P., 2015. Soil conditions and land use

1194 intensification effects on soil microbial communities across a range of European field sites. *Soil*
1195 *Biology and Biochemistry* 88, 403-413.

1196 Tindall, B.J., Rossello-Mora, R., Busse, H.J., Ludwig, W., Kampfer, P., 2010. Notes on the
1197 characterization of prokaryote strains for taxonomic purposes. *Int J Syst Evol Microbiol* 60, 249-266.

1198 Tisdall, J.M., 1994. Possible role of soil microorganisms in aggregation in soils. *Plant and Soil* 159,
1199 115-121.

1200 Trivedi, P., Delgado-Baquerizo, M., Jeffries, T.C., Trivedi, C., Anderson, I.C., Lai, K., McNee, M.,
1201 Flower, K., Pal Singh, B., Minkey, D., Singh, B.K., 2017. Soil aggregation and associated microbial
1202 communities modify the impact of agricultural management on carbon content. *Environmental*
1203 *Microbiology* 19, 3070-3086.

1204 USS Working Group WRB, 2014. World Reference Base for soil resources 2014: international soil
1205 classification system for naming soils and creating legends for soil maps. *World Soil Resources*
1206 *Reports No. 106*. FAO, Rome.

1207 van de Voorde, T.F.J., van der Putten, W.H., Bezemer, T.M., 2012. Soil inoculation method
1208 determines the strength of plant–soil interactions. *Soil Biology and Biochemistry* 55, 1-6.

1209 van der Putten, W.H., Bardgett, R.D., Bever, J.D., Bezemer, T.M., Casper, B.B., Fukami, T., Kardol, P.,
1210 Klironomos, J.N., Kulmatiski, A., Schweitzer, J.A., Suding, K.N., Van de Voorde, T.F.J., Wardle, D.A.,
1211 2013. Plant–soil feedbacks: the past, the present and future challenges. *Journal of Ecology* 101, 265-
1212 276.

1213 Van Soest, P.J., 1963. Use of detergents in the analysis of fibrous feeds. 2. A rapid method for the
1214 determination of fiber and lignin. *Journal of the Association of Official Agricultural Chemists* 46, 829-
1215 835.

1216 van Veen, J.A., van Overbeek, L.S., van Elsas, J.D., 1997. Fate and activity of microorganisms
1217 introduced into soil. *Microbiology and Molecular Biology Reviews* 61, 121-135.

1218 Wang, Q., Garrity, G.M., Tiedje, J.M., Cole, J.R., 2007. Naïve Bayesian Classifier for Rapid Assignment
1219 of rRNA Sequences into the New Bacterial Taxonomy. *Applied and Environmental Microbiology* 73,
1220 5261-5267.

1221 Wright, S.F., Upadhyaya, A., 1996. Extraction of an abundant and unusual protein from soil and
1222 comparison with hyphal protein of arbuscular mycorrhizal fungi. *Soil Science* 161, 575-586.

1223 Wu, X., Wei, Y., Wang, J., Wang, D., She, L., Wang, J., Cai, C., 2017. Effects of soil physicochemical
1224 properties on aggregate stability along a weathering gradient. *CATENA* 156, 205-215.

1225 Zhang, J., Kobert, K., Flouri, T., Stamatakis, A., 2014. PEAR: a fast and accurate Illumina Paired-End
1226 reAd mergeR. *Bioinformatics* 30, 614-620.

1227 Zhang, S., Li, Q., Lü, Y., Zhang, X., Liang, W., 2013. Contributions of soil biota to C sequestration
1228 varied with aggregate fractions under different tillage systems. *Soil Biology and Biochemistry* 62,
1229 147-156.

1230 Zhang, S., Li, Q., Zhang, X., Wei, K., Chen, L., Liang, W., 2012. Effects of conservation tillage on soil
1231 aggregation and aggregate binding agents in black soil of Northeast China. *Soil and Tillage Research*
1232 124, 196-202.

1233

Supplementary Material for online publication only

[Click here to download Supplementary Material for online publication only: Supplementary material.docx](#)

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: