



HAL
open science

Sugar Signaling and Post-transcriptional Regulation in Plants: An Overlooked or an Emerging Topic?

Ming Wang, Lili Zang, Fuchao Jiao, Maria-Dolores Perez-Garcia, Laurent Ogé, Latifa Hamama, José Le Gourrierc, Soulaiman Sakr, Jingtang Chen

► **To cite this version:**

Ming Wang, Lili Zang, Fuchao Jiao, Maria-Dolores Perez-Garcia, Laurent Ogé, et al.. Sugar Signaling and Post-transcriptional Regulation in Plants: An Overlooked or an Emerging Topic?. *Frontiers in Plant Science*, 2020, 11, pp.578096. 10.3389/fpls.2020.578096 . hal-03046616

HAL Id: hal-03046616

<https://hal.inrae.fr/hal-03046616>

Submitted on 11 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Sugar Signaling and Post-transcriptional Regulation in Plants: An Overlooked or an Emerging Topic?

Ming Wang^{1,2†}, Lili Zang^{2†}, Fuchao Jiao^{1†}, Maria-Dolores Perez-Garcia², Laurent Ogé², Latifa Hamama², José Le Gourrierec², Soulayman Sakr^{2*} and Jingtang Chen^{1*}

¹College of Agronomy, Qingdao Agricultural University, Qingdao, China, ²IRHS-UMR1345, INRAE, Institut Agro, SFR 4207 QuaSaV, Université d'Angers, Beaucazoué, France

OPEN ACCESS

Edited by:

Jean Molinier,
UPR2357 Institut de biologie
moléculaire des plantes (IBMP),
France

Reviewed by:

Pablo A. Manavella,
CONICET Santa Fe,
Argentina
Thomas Blein,
UMR9213 Institut des Sciences des
Plantes de Paris Saclay (IP2S),
France

*Correspondence:

Jingtang Chen
chenjingtang@126.com
Soulayman Sakr
soulayman.sakr@agrocampus-ouest.fr

[†]These authors share first authorship

Specialty section:

This article was submitted to
Plant Cell Biology,
a section of the journal
Frontiers in Plant Science

Received: 30 June 2020

Accepted: 02 October 2020

Published: 05 November 2020

Citation:

Wang M, Zang L, Jiao F,
Perez-Garcia M-D, Ogé L,
Hamama L, Le Gourrierec J,
Sakr S and Chen J (2020) Sugar
Signaling and Post-transcriptional
Regulation in Plants: An Overlooked
or an Emerging Topic?
Front. Plant Sci. 11:578096.
doi: 10.3389/fpls.2020.578096

Plants are autotrophic organisms that self-produce sugars through photosynthesis. These sugars serve as an energy source, carbon skeletons, and signaling entities throughout plants' life. Post-transcriptional regulation of gene expression plays an important role in various sugar-related processes. In cells, it is regulated by many factors, such as RNA-binding proteins (RBPs), microRNAs, the spliceosome, etc. To date, most of the investigations into sugar-related gene expression have been focused on the transcriptional level in plants, while only a few studies have been conducted on post-transcriptional mechanisms. The present review provides an overview of the relationships between sugar and post-transcriptional regulation in plants. It addresses the relationships between sugar signaling and RBPs, microRNAs, and mRNA stability. These new items insights will help to reach a comprehensive understanding of the diversity of sugar signaling regulatory networks, and open onto new investigations into the relevance of these regulations for plant growth and development.

Keywords: sugar, RNA-binding protein, post-transcriptional regulation, microRNA, mRNA stability

INTRODUCTION

As living organisms, plants need various compounds to meet the requirements of their global metabolism and to finely adapt to different external stimuli. In this process, one of the most essential compounds is sugar, which has both trophic and signaling activities during plant development – a high-energy-demanding and well-controlled process. Plants synthesize sugar from carbon dioxide and water through photosynthesis and finely tune their sugar status to avoid sugar starvation (Stitt and Zeeman, 2012; Kanwar and Jha, 2019; Signorelli et al., 2019). They have evolved a sophisticated machinery to sense different forms of sugars, including hexoses, sucrose, and various sugar phosphates (e.g., trehalose-6-phosphate), and elicit the appropriate responses. Some responses are sugar-type-specific (Granot et al., 2014; Figueroa and Lunn, 2016; Li and Sheen, 2016; Janse van Rensburg and Van den Ende, 2018; Wingler, 2018). As a signaling entity, sugar can influence a diversity of physiological processes of the plant life cycle and operates on transcriptional, post-transcriptional, translational, and post-translational regulation. Most of the currently available knowledge is focused on sugar-dependent transcriptional

regulation (Lastdrager et al., 2014; Li and Sheen, 2016; Sakr et al., 2018; Wingler, 2018; Jiao et al., 2019; Rodriguez et al., 2019; Sami et al., 2019). Post-transcriptional regulation of gene expression is a pivotal mechanism whereby plants rapidly reprogram their transcriptome and proteome in response to endogenous and environmental cues and involves many factors such as proteins [RNA-binding proteins (RBPs)], microRNAs (miRNAs), and the spliceosome (Guerra et al., 2015; Romanowski and Yanovsky, 2015; Zhang, 2015; Kawa and Testerink, 2017; Samad et al., 2017; Maronedez et al., 2019; Rigo et al., 2019). RBPs are mainly cytosolic and nuclear proteins that govern the processing, cellular localization, and decay of cellular RNA. They contain RNA recognition motifs (RRMs) that allow their binding to a specific sequence in the target transcripts (Chou et al., 2017; Tian et al., 2018a; Lu et al., 2019; Mahalingam and Walling, 2020). miRNAs are small non-coding RNA molecules that function in RNA silencing *via* base-pairing with complementary sequences within mRNA molecules (Bartel, 2004, 2018), leading to mRNA cleavage; they shorten the poly(A) tail of mRNAs, or influence mRNA translation, altogether downregulating gene expression through the target transcript (Zhang, 2015; Meyers and Axtell, 2019). Alternative splicing – also termed alternative RNA splicing – is mediated by the spliceosome, a complex and large molecular machinery mainly located within the nucleus of eukaryotic cells (Wilkinson et al., 2020). Splicing can also have other functions, like the generation of premature stop codons that recruit the nonsense-mediated decay (NMD) machinery (Kesarwani et al., 2019; Wilkinson et al., 2020). Consequently, the proteins translated from alternatively spliced mRNAs are expected to have different amino acid sequences, different protein structures, and even different biological functions.

Sugar signaling-dependent regulation represents an intricate regulatory network that relies on highly diverse mechanisms that coordinate the appropriate use of available energy and sugar to sustain plant development and growth under the ever-changing environment. Sugar-dependent post-transcriptional regulation could be one important mechanistic aspect of this network. Current knowledge in this topic is still fragmented and makes it very difficult to draw a comprehensive scheme and bring out new research questions. The present review addresses the relationship between sugar and post-transcriptional gene regulation in plants and provides first insights into the role of various important mechanisms of post-transcriptional regulation, i.e., RBPs, miRNAs, and mRNA decay/stability in sugar-related pathways. It underlines the physiological relevance of such regulation mechanisms in different biological contexts and raises questions for upcoming studies.

SUGAR AND RNA-BINDING PROTEINS

RNA-binding proteins control nearly all aspects of eukaryotic post-transcriptional gene regulation and consequently determine the fate and expression of the plant transcriptome. Hundreds of RBPs have been identified in *Arabidopsis*. Most of them

are plant specific, and could carry out specific functions in plant physiology (Lorković, 2009; Reichel et al., 2016). They share one or more canonical RNA-binding domains including the RRM, the K-homology (KH) domain, the Pumilio/FBF (PUF) domain, the RRM and KH domains, DEAD/DEAH boxes, zinc-finger structures, the Piwi/Argonaute/Zwille (PAZ) domain, double-stranded RNA-binding domains (DS-RBD), pentatricopeptide-repeat (PPR) domains, etc. (Silverman et al., 2013; Lee and Kang, 2016; Wang et al., 2018a). The link between glucose signaling and RBP-mediated post-transcriptional regulation has been explored in the model plant *Arabidopsis thaliana*. Transgenic *Arabidopsis* plants overexpressing *atRZ-1a*, which encodes a zinc-finger-containing glycine-rich RNA-binding protein (GRP), exhibited delayed germination and seedling growth under abiotic stresses (dehydration or salt stress), and hypersensitivity to glucose and ABA, relatively to the wild type (Kim and Kang, 2006; Kim et al., 2007b). Yet, the molecular function of *atrZ1* in glucose-dependent post-transcriptional regulation of seedling establishment is still unknown. The RBP FLOWERING CONTROL LOCUS A (FCA) contains two RRM domains and one WW domain (Jang et al., 2009; **Table 1**) and operates as an inhibitor of FLOWERING LOCUS C (FLC; Macknight et al., 1997; Liu et al., 2007), one of the repressor integrators, tightly controls flowering signals (Michaels and Amasino, 1999). FLC is positively and transcriptionally regulated by the ABI5 transcription factor (ABA-insensitive 5, **Figure 1**), which is involved in ABA-mediated floral transition (Wang et al., 2013) and in integrating glucose

TABLE 1 | The type and function of sugar related RBPs.

Gene	RNA-binding protein type	Function
AtRZ-1a	Zinc finger-containing glycine-rich RNA-binding protein	Involve in freezing tolerance and cold stress
GLYCINE RICH PROTEIN 2	RRM protein	Response to cold, osmotic stress, water deprivation, and seed germination
FLOWERING CONTROL LOCUS A	RRM protein and contain WW domain	Involved in the promotion of the transition of the vegetative meristem to reproductive development
HYPONASTIC LEAVES 1	No reports	Flowering, leaf, and root development
APUM2, APUM3, and APUM4	Pumilio/FBF protein	No reports
ALDH7B4	No reports	Aldehyde dehydrogenase
RAFFINOSE SYNTHASE 6	No reports	Biosynthesis of the raffinose; response to cold, karrikin, and oxidative stress
SOAR1	PPR protein	ABA responses, probably located upstream of an ABA-responsive transcription factor ABI5
RhPUF4	Pumilio/FBF protein	Involved in bud outgrowth probably

The function of *Arabidopsis* protein is from the annotation of TAIR database (www.arabidopsis.org; Swarbreck et al., 2007; Berardini et al., 2015).

SUGARS AND MICRORNAs

miRNAs are small non-coding RNA molecules that participate in RNA silencing and post-transcriptional regulation of gene expression (Song et al., 2019). miRNA genes are transcribed by RNA polymerase II in the nucleus. This generates long primary transcripts of miRNA (primary miRNAs, pri-miRNA in short), which are converted into a precursor miRNA (pre-miRNA) by endonuclease DICER-like1 (DCL1). After a complex processing involving the C2H2-zinc finger protein SERRATE (SE), DCL1 and the double-stranded RBP HYPONASTIC LEAVES1 (HYL1), the miRNA is loaded onto ARGONAUTE1 (AGO1) to integrate the RNA-induced silencing complex (RISC; Voinnet, 2009; Rogers and Chen, 2013). Several studies support a direct link between sugar signaling and miRNAs in a variety of physiological processes in plants. Duarte et al. (2013) shown that *Arabidopsis* mutants disrupted in miRNA biosynthesis (*hyl1-2* and *dcl1-11*) and miRNA activity (*ago1-25*) exhibited a glucose-hyposensitive phenotype at the early seedling stage, and the expression of several miRNA target genes was deregulated, mainly via hexokinase-independent pathway. miRNA156 is one of the best characterized miRNAs in terms of sugar-dependent regulation. It is conserved in land plants and contributes to diverse physiological processes such as leaf development, heat stress memory, developmental transition, apical dominance, and flowering (Kim et al., 2012; Bhogale et al., 2014; Yu et al., 2015; Zhang, 2015; Gao et al., 2018; Kumar et al., 2020). The biological function of miRNA156 implies the repression of SQUAMOSA-PROMOTER BINDING PROTEIN-LIKEs (SPLs; Wang et al., 2009; Wahl et al., 2013; Xu et al., 2016; Wei et al., 2017; Zheng et al., 2019a,b; Hu et al., 2020, Jiao et al., 2020; Ponnu et al., 2020). A direct link between sugar and miRNA156 abundance is based on the ability of exogenous glucose or sucrose supply to cause the levels of mature miRNA156 to drop and thereby accelerate the vegetative-reproductive phase transition, along with the juvenile-to-adult phase transition. Conversely, defoliation and a reduced photosynthetic rate delay plant developmental transitions (Yang et al., 2013; Yu et al., 2013). The glucose-induced repression of miRNA156 is dependent on the hexokinase 1-signaling pathway (Yang et al., 2013), while trehalose-6-phosphate regulates developmental transition through a distinct mechanism (Wahl et al., 2013; Ponnu et al., 2020). miRNA156 also targets a variety of mRNAs that encode regulatory proteins involved in various physiological processes in plants (Wang and Wang, 2015). Like miRNA156, miRNA399 was determined to be sucrose-responsive through a microRNA array assay and high levels of sucrose inhibited the accumulation of microRNA399 family under phosphate starvation conditions in *Arabidopsis* (Tian et al., 2018b). miRNA398, that is associated with the adaptive plant response to biotic, abiotic, and nutrient stresses and could be involved in sugar-signaling pathway (Sunkar et al., 2006; Dugas and Bartel, 2008; Jia et al., 2009; Feng et al., 2015). miRNA398 accumulation is repressed by carbon depletion (Pant et al., 2009), while sucrose supply induces

its accumulation through the SPL7 transcription factor that directly recognizes the GTAC boxes located in the *miRNA398* promoter (Yamasaki et al., 2009). In line with this, *spl7* knockdown mutants consistently accumulate lower levels of miRNA398 under normal conditions (Ren and Tang, 2012). Targets of miR398a include two ROS-scavenging enzymes (COPPER/ZINC SUPEROXIDE DISMUTASE, CSD1 and CSD2) necessary for detoxification of stress-dependent reactive oxygen species stimulation (Farooq et al., 2019) and this sugar-mediated regulation of *miRNA398* would be an appropriate response to nutrient stress. The *miRNA398* binding site of CSD1 can be eliminated by alternative splicing in peanut and *Arabidopsis*, resulting in different tolerance levels to abiotic stress (Park and Grabau, 2017), indicating how alternative splicing processes influence plant response through interactions with miRNAs. Microarray analyses have also shown responsiveness to sucrose from other mature miRNAs in *Arabidopsis* (Figure 2), including miRNA408 (involved in the response to iron deficiency and in photosynthesis; Pan et al., 2018; Carrió-Seguí et al., 2019), miRNA319 (involved in leaf development; Koyama et al., 2017), and miRNA160 (involved in heat tolerance; Lin et al., 2018). The levels of miRNA319 and miRNA408 are enhanced by sucrose supply, while the levels of miRNA160 are reduced. Moreover, the induction of miRNA408 by sucrose is associated once again with SPL7 (Ren and Tang, 2012), which may play a prominent role in sugar-mediated regulation of miRNA biosynthesis (Figure 2). In sweet sorghum (*Sorghum bicolor*), the expression levels of nine known mature miRNAs and 12 novel mature miRNAs have been found influenced by sugar abundance in the stem (Yu et al., 2015). Although the targets of these mature miRNAs exhibit functions related to shoot apical meristem specification, polar specification of the adaxial/abaxial axis, bilateral symmetry determination, and transcriptional regulation (Yu et al., 2015), the genuine participation of sugar sensing and signaling in this regulatory network remains to be elucidated.

A link between sucrose transporters (SUTs, H⁺/Suc symporters) and miRNAs exists in plants. SUTs are key players in sucrose phloem loading and sugar allocation within plants




FIGURE 2 | The relationship between sugar and reported miRNA, and the function of the related miRNA. Sugar stimulates the transcription of *miRNA398*, *miRNA408*, and *miRNA319* but inhibits that of *miRNA160* and *miRNA156*. SnRK1 inhibits the transcription of *miRNA824A*, *miRNA775A*, and *miRNA159B*. HXK1, hexokinase 1; SPL, SQUAMOSA-PROMOTER BINDING PROTEIN-LIKE.

(Braun et al., 2014; Milne et al., 2017; Doidy et al., 2019). The half-life of SUT mRNAs ranges between 60 and 130 min and is tightly related to miRNA regulation (He et al., 2008; Liesche et al., 2011). Interestingly, the relationship between miRNA biosynthesis and the cellular energy status is also supported by the fact that the transient overexpression of the energy-sensing SnRK1 in protoplasts leads to the repression of a variety of miRNAs (Confraria et al., 2013). These include miRNA159B (involved in leaf senescence, Huo et al., 2015), miRNA161 (induces the expression of PPR genes, Cai et al., 2018), miRNA775A (no function reported to date), and miRNA824A (involved in flowering time regulation, Hu et al., 2014) and might be involved in SnRK1-dependent energy signaling. However, the molecular regulatory network involved in this SnRK1-dependent miRNA biosynthesis remains an open question.

CROSSTALK BETWEEN SUGAR-RELATED RNA-BINDING PROTEINS AND MICRORNAs

Post-translational modifications have been reported as a key regulator of the miRNA-biogenesis machinery. In *Arabidopsis*, HYL1 activity is controlled by its phosphorylation state through complex mechanisms. The Protein Phosphatase 4 (PP4)/Suppressor of MEK 1 (SMEK1) complex and C-TERMINAL DOMAIN PHOSPHATASE-LIKE 1 and 2 (CPL1 and CPL2) dephosphorylate and activate HYL1, while Mitogen-activated Protein Kinase (MPK) phosphorylates and inactivates it (Manavella et al., 2012; Su et al., 2017; Meng et al., 2018; Wang et al., 2019c). MPK might bridge a gap between miRNA biosynthesis and sugar signaling, based on its transducing role the WALL-ASSOCIATED KINASE (WAK)-dependent regulation of vacuolar invertase, a driver of cell expansion and growth and a player of sugar signaling (Kohorn et al., 2006, 2009; **Figure 1**). SnRK2s can also affect the phosphorylation status of HYL and SE (Yan et al., 2017), and it will be very interesting to explore the sensitivity of these two proteins to the kinase activity of SnRK1 and, thereby, its relevance in SnRK1-dependent miRNA biosynthesis regulation (Confraria et al., 2013). Beyond this, the basic helix-loop-helix (bHLH) transcription factor phytochrome-interacting factor 4 (PIF4) interacts directly with DCL1 and HYL1 to promote their destabilization and regulate the processing of primary miRNAs during the dark-to-red-light transition (Sun et al., 2018). PIF4 is also controlled through the trehalose-6-phosphate pathway and SnRK1 to modulate *Arabidopsis* hypocotyl elongation in response to high temperature (Delatte et al., 2011; Hwang et al., 2019), so it might be seen as a hub integrating sugar signaling and environmental cues to modulate the regulation of miRNA biogenesis through the DCL1-SE-HYL complex.

Many previous studies indicate that various RBPs participate in miRNA homeostasis. For instance, the WD-40 protein PLEIOTROPIC REGULATORY LOCUS1 (PRL1) is required for miRNAs and small siRNAs to accumulate, by stabilizing pri-miRNAs through its RNA-binding activity and enhancing DCL1 activity (Zhang et al., 2014). Beyond this function,

PRL1 acts as a global regulator of sugar, stress, and hormone responses, partly through SnRK1 repression (Flores-Pérez et al., 2010). However, additional investigations are required to elucidate the molecular connections between these different PRL1-dependent regulatory mechanisms. miRNA172 is a downstream component of the regulatory cascade involved in the regulation of flowering time by sugar-dependent miRNA156 repression (Wu et al., 2009; Martin et al., 2010), in which miRNA172 acts as an inducer of FLOWERING LOCUS T (FT) expression. FCA, an RBP, binds to the flanking sequences of the stem-loop within primary miRNA172 transcripts (pri-miRNA172) *via* the RRM, and promotes its accumulation in response to ambient temperature (**Figure 1**). FCA also binds to the primary transcripts of other temperature-responsive miRNAs, such as miRNA398 and miRNA399 (Jung et al., 2012). The RBP TOUGH (TGH) contributes to the pri-miRNA-HYL1 interaction (Ren et al., 2012), while MOS2 (MODIFIER OF SNC1, 2) is involved in pri-miRNA processing (Wu et al., 2013). Many other examples exist, including EMBRYO DEFECTIVE 2793 (EMB2793, THO2), MOS4-ASSOCIATED COMPLEX 7 (MAC7), and REGULATOR OF CBF GENE EXPRESSION 3 (RCF3) which participate in the regulation of miRNA biogenesis by interacting with HYL1 (Francisco-Mangilet et al., 2015; Karlsson et al., 2015; Jia et al., 2017). However, whether other core components of miRNA processing are dependent on RBPs and the way sugar signaling could contribute to this regulatory network still remains unclear.

SUGAR AND MRNA DECAY/STABILITY

In plants, mRNA decay/stability is an important control point in the regulation of gene expression and can discard potentially deleterious errors in mRNA synthesis (Nagarajan et al., 2019). The mRNA decay/stability of many sugar-metabolism-related enzymes is controlled through post-transcriptional regulation. This holds true for the maize cell wall invertase gene (*Incw1*) that displays two transcripts – *Incw1-S* (small) and *Incw1-L* (large) – according to the respective lengths of its 3'untranslated regions (UTR; Cheng et al., 1999). Since sucrose and D-glucose appear to be associated with the increased steady-state abundance of *Incw1-S* mRNA and cell wall invertase activity, these authors suggested that the 3'UTR of the *Incw1* gene was a regulatory sensor of carbon starvation and acted as a link between translation activity and the sink metabolism in plants. The 3'UTRs of *OsVIN1* and *AtvacINV2*, encoding vacuolar invertases in rice and *Arabidopsis*, respectively, are involved in this process. Downstream regulatory elements or a motif that participates in the rapid degradation of mRNAs, e.g., small auxin-up RNAs (SAUR; Feldbrügge et al., 2002; van Mourik et al., 2017), may be involved too (Huang et al., 2007). The expression of α -amylase, an endo-amylolytic enzyme that catalyzes starch degradation in plants, is induced by sucrose starvation and suppressed by sucrose availability in rice. Sugar repression of α -amylase 3 (α AMY3) expression in rice suspension cells involves controlling both its transcription rate and mRNA stability (Sheu et al., 1994;

Chan and Yu, 1998). An analysis of reporter mRNA half-lives indicated that two subdomains of the α AMY3 3'UTR contained the UAUUAUGUA motif required for the sugar-dependent destabilization of α AMY3 mRNA (Sheu et al., 1994; Chan and Yu, 1998). The same motif might also be involved in sugar-mediated post-transcriptional downregulation of *RhbRC1* in *R. hybrida* (Wang et al., 2019a), and could be conserved in angiosperms. Such sugar-dependent regulation of mRNA stability is required for the rapid adjustment of gene expression in response to the sugar status of the cell. In *Arabidopsis* cell cultures, the stability of 224 mRNAs was repressed by sucrose limitation, concomitantly with a drop in the cell metabolic activity (Nicolai et al., 2006). The mRNA half-lives of actin (*ACT*), alcohol dehydrogenase 2 (*ADH2*), glyceraldehyde 3-phosphate dehydrogenase (*G3PD*), and sucrose synthase P-2 (*SSP2*) were consistently 1.6- to 2.6-fold longer in sucrose-supplied rice cells (Ho et al., 2001). In line with this, the mRNA stability of the bZIP63 transcription factor, an important mediator of the adaptive response induced by SnRK1 during energy or sugar depletion (Baena-González et al., 2007) decreased following exogenous glucose supply in *Arabidopsis* seedlings (Matioli et al., 2011). The involvement of bZIP63 as a hub integrating the sugar and energy statuses and mRNA stability deserves to be addressed. *Low- β -amylase1* (*lba1*) is a missense mutation of UP-FRAMESHIFT 1 (*UPF1*) RNA helicase, involved in nonsense-mediated mRNA decay (NMD). Its *Arabidopsis* mutant exhibited lower sugar induction of the *At β Amy* transcript, which was restored by complementation of the *lba1* mutation with wild type *UPF1*, further supporting the link between sugar signaling and the fate of the mRNA (Yoine et al., 2006). All these findings clearly indicate a relationship between the sugar status and mRNA stability in a variety of biological contexts, opening the avenue for deciphering the sugar sensing and signaling mechanisms. In line with that, mRNA stability might also be important for the diurnal regulation of mRNA levels of sucrose transporters and in turn in sugar allocation at the whole plant level. For example, a sucrose transporter (*SUT1*) displayed a quick turnover rate in leaves of tomato (*Lycopersicon esculentum*), potato (*Solanum tuberosum*), and tobacco (*Nicotiana tabacum*; Kühn et al., 1997; Kühn and Grof, 2010). The mRNA levels of *StSUT2* and *StSUT4* may be regulated by putative RBPs (He et al., 2008). Two AUUUA motifs exist in the 3'UTR or CDS region of *StSUT2* and *StSUT4* mRNA (He et al., 2008); they have been characterized as the binding sites of proteins involved in mediating mRNA degradation (Chen and Shyu, 1995; He et al., 2008; Liesche et al., 2011). However, the nature of these proteins is still unknown.

SUGAR AND ALTERNATIVE SPLICING

Alternative splicing is a finely regulated process that takes place during gene expression and leads to a single gene coding for multiple proteins. Serine/arginine-rich 45 (SR45) is a serine/arginine-rich splicing factor that participates in 5' and 3' splicing site selection of introns and can bridge the 5' and 3' components of the spliceosome. The SR45 splicing factor regulates glucose

signaling during early seedling development in *Arabidopsis* (Carvalho et al., 2010), more likely through the modulation of SnRK1-stability (Carvalho et al., 2016). The *sr45-1* knockout mutant indeed displays a high level of the energy-limitation-sensing SnRK1 protein under glucose supply, which is in agreement with the upregulation of SnRK1-activated genes such as *SENESCENCE-ASSOCIATED GENE1* (*SEN1*), *GLUTAMINE-DEPENDENT ASPARAGINE SYNTHETASE1* (*ASN1*), and *DORMANCY-ASSOCIATED GENE2* (*DRM2*). Moreover, the glucose hypersensitivity of the *sr45-1* mutant is alleviated when SnRK1 is disrupted (Figure 1). SR45 controls the alternative splicing of *5-phosphatase 13* (*5PTase13*) in *Arabidopsis*, which encodes an inositol polyphosphate 5-phosphatase involved in regulating SnRK1 stability negatively *in vitro* (Carvalho et al., 2016; Figure 1). This link between sugar signaling and RNA splicing has also been reported for the photomorphogenesis-related alternative splicing shifts primarily controlled by a metabolic photosynthesis-derived signal and exogenous sucrose supply, correlated with the expression of dark-induced genes under the control of SnRK1 (Hartmann et al., 2016). *AtTZF1/AtCTH/AtC3H23* (a tandem-arrayed CCCH-type zinc finger motif involved in stress- and hormone-mediated growth), was also identified as a sugar-sensitive gene in *Arabidopsis* (Qu et al., 2014). *AtTZF1* can traffick between the nucleus and cytoplasmic foci and bind both DNA and RNA *in vitro*; it may be involved in RNA regulation and under the control of sugar signaling (Pomeranz et al., 2010). However, the basic molecular mechanisms behind this regulation have not been addressed to date.

CONCLUSION

Post-transcriptional regulation is an essential component of gene expression regulation in plants. Numerous findings have unveiled and characterized various factors involved in post-transcriptional regulation. The present review provides a first comprehensive picture of the relationship between sugar (metabolism and signaling) and post-transcriptional regulation factors in plants, including RBPs, miRNAs, and mRNA stability of sugar-related genes. More work needs to be carried out to figure out the functions and mechanisms related to the involvement of post-transcriptional regulation in sugar-related processes, e.g., whether regulatory mechanisms found in human cells or yeast are also conserved in plants. Considering the frequently observed connection between mRNA abundance and sugar, some recently developed technologies for RNA editing (CRISPR-Cas13), RNA binding (RNA interactome capture, photoactivatable ribonucleoside-enhanced crosslinking), and RNA folding (DMS-seq, SHAPE-seq) will support future studies. Besides, many aspects of RNA decay still need to be studied in depth, such as the spliceosome and the editosome (a large multi-protein complex that catalyzes RNA editing), which play a crucial role in post-transcriptional regulation. Although some reports about the interaction between sugar-related RBPs and miRNAs exist, further investigations are still required to gain a comprehensive understanding of the way sugar signaling operates through each of these post-transcriptional regulation

mechanisms and how they crosstalk to regulate plant growth and development. The hub role of hexokinase, SnRK1, and/or TOR kinase but also the relevance of the trehalose signaling pathway in the different post-transcriptional regulation networks could be two main future lines of research. This further knowledge will also pave the way for discovering a new and complex sugar regulatory network in plants.

AUTHOR CONTRIBUTIONS

All authors listed have made direct contribution to the work, and approved it for publication. MW, LZ, FJ, SS, and JC have

written different part of the manuscript. M-DP-G, LO, and LH have contributed to the section sugar and RNA binding proteins and JL to the section of sugars and miRNA.

ACKNOWLEDGMENTS

This work was supported by the program of China Scholarships Council (No. 201506320203), Talent Introduction Special Funds of Qingdao Agricultural University (663/1120070) and by the ANR (Agence Nationale de la Recherche) project Labcom, called ESTIM (Evaluation de STIMulateurs de vitalité des plantes; Projet ANR- 15-LCV3-000).

REFERENCES

- Arroyo, A., Bossi, F., Finkelstein, R. R., and León, P. (2003). Three genes that affect sugar sensing (abscisic acid insensitive 4, abscisic acid insensitive 5, and constitutive triple response 1) are differentially regulated by glucose in *Arabidopsis*. *Plant Physiol.* 133, 231–242. doi: 10.1104/pp.103.021089
- Baena-González, E., Rolland, R., Thevelein, J. M., and Sheen, J. (2007). A central integrator of transcription networks in plant stress and energy signaling. *Nature* 448, 938–942. doi: 10.1038/nature06069
- Bartel, D. P. (2004). MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 116, 281–297. doi: 10.1016/S0092-8674(04)00045-5
- Bartel, D. P. (2018). Metazoan microRNAs. *Cell* 173, 20–51. doi: 10.1016/j.cell.2018.03.006
- Berardini, T. Z., Reiser, L., Li, D., Mezheritsky, Y., Muller, R., Strait, E., et al. (2015). The *Arabidopsis* information resource: making and mining the “gold standard” annotated reference plant genome. *Genesis* 53, 474–485. doi: 10.1002/dvg.22877
- Bhogale, S., Mahajan, A. S., Natarajan, B., Rajabhoj, M., Thulasiram, H. V., and Banerjee, A. K. (2014). MicroRNA156: a potential graft-transmissible microRNA that modulates plant architecture and tuberization in *Solanum tuberosum* ssp. *andigena*. *Plant Physiol.* 164, 1011–1027. doi: 10.1104/pp.113.230714
- Bi, C., Ma, Y., Jiang, S. C., Mei, C., Wang, X. F., and Zhang, D. P. (2019). *Arabidopsis* translation initiation factors eIF iso4G1/2 link repression of mRNA cap-binding complex eIF iso4F assembly with RNA-binding protein SOAR 1-mediated ABA signaling. *New Phytol.* 223, 1388–1406. doi: 10.1111/nph.15880
- Braun, D. M., Wang, L., and Ruan, Y. L. (2014). Understanding and manipulating sucrose phloem loading, unloading, metabolism, and signalling to enhance crop yield and food security. *J. Exp. Bot.* 65, 1713–1735. doi: 10.1093/jxb/ert416
- Cai, Q., Liang, C., Wang, S., Hou, Y., Gao, L., Liu, L., et al. (2018). The disease resistance protein SNC1 represses the biogenesis of microRNAs and phased siRNAs. *Nat. Commun.* 9:5080. doi: 10.1038/s41467-018-07516-z
- Carrió-Seguí, Á., Ruiz-Rivero, O., Villamayor-Belinchón, L., Puig, S., Perea-García, A., and Peñarrubia, L. (2019). The altered expression of microRNA408 influences the *Arabidopsis* response to iron deficiency. *Front. Plant Sci.* 10:324. doi: 10.3389/fpls.2019.00324
- Carvalho, R. F., Carvalho, S. D., and Duque, P. (2010). The plant-specific SR45 protein negatively regulates glucose and ABA signaling during early seedling development in *Arabidopsis*. *Plant Physiol.* 154, 772–783. doi: 10.1104/pp.110.155523
- Carvalho, R. F., Szakonyi, D., Simpson, C. G., Barbosa, I. C., Brown, J. W., Baena-González, E., et al. (2016). The *Arabidopsis* SR45 splicing factor, a negative regulator of sugar signaling, modulates SNF1-related protein kinase 1 stability. *Plant Cell* 28, 1910–1925. doi: 10.1105/tpc.16.00301
- Chan, M. T., and Yu, S. M. (1998). The 3' untranslated region of a rice α -amylase gene functions as a sugar-dependent mRNA stability determinant. *Proc. Natl. Acad. Sci. U. S. A.* 95, 6543–6547. doi: 10.2307/45417
- Chen, C. Y. A., and Shyu, A. B. (1995). AU-rich elements: characterization and importance in mRNA degradation. *Trends Biochem. Sci.* 20, 465–470. doi: 10.1016/S0968-0004(00)89102-1
- Cheng, W. H., Taliervo, E. W., and Chourey, P. S. (1999). Sugars modulate an unusual mode of control of the cell-wall invertase gene (Incw1) through its 3' untranslated region in a cell suspension culture of maize. *Proc. Natl. Acad. Sci. U. S. A.* 96, 10512–10517. doi: 10.1073/pnas.96.18.10512
- Chou, H. L., Tian, L., Kumamaru, T., Hamada, S., and Okita, T. W. (2017). Multifunctional RNA binding protein OsTudor-SN in storage protein mRNA transport and localization. *Plant Physiol.* 175, 1608–1623. doi: 10.1104/pp.17.01388
- Confraria, A., Martinho, C. S. D. S., Elias, A., Rubio-Somoza, I., and Baena-González, E. (2013). miRNAs mediate SnRK1-dependent energy signaling in *Arabidopsis*. *Front. Plant Sci.* 4:197. doi: 10.3389/fpls.2013.00197
- Corradetti, M. N., and Guan, K. (2006). Upstream of the mammalian target of rapamycin: do all roads pass through mTOR? *Oncogene* 25, 6347–6360. doi: 10.1038/sj.onc.1209885
- Dekkers, B. J., Schuurmans, J. A., and Smeekens, S. C. (2008). Interaction between sugar and abscisic acid signalling during early seedling development in *Arabidopsis*. *Plant Mol. Biol.* 67, 151–167. doi: 10.1007/s11103-008-9308-6
- Delatte, T. L., Sedijani, P., Kondou, Y., Matsui, M., de Jong, G. J., Somsen, G. W., et al. (2011). Growth arrest by trehalose-6-phosphate: an astonishing case of primary metabolite control over growth by way of the SnRK1 signaling pathway. *Plant Physiol.* 157, 160–174. doi: 10.1104/pp.111.180422
- Dobrenel, T., Caldana, C., Hanson, J., Robaglia, C., Vincentz, M., Veit, B., et al. (2016). TOR signaling and nutrient sensing. *Annu. Rev. Plant Biol.* 67, 261–285. doi: 10.1146/annurev-arplant-043014-114648
- Doidy, J., Vidal, U., and Lemoine, R. (2019). Sugar transporters in Fabaceae, featuring SUT MST and SWEET families of the model plant *Medicago truncatula* and the agricultural crop *Pisum sativum*. *PLoS One* 14:e0223173. doi: 10.1371/journal.pone.0223173
- Duarte, G. T., Matioli, C. C., Pant, B. D., Schlereth, A., Scheible, W. R., Stitt, M., et al. (2013). Involvement of microRNA-related regulatory pathways in the glucose-mediated control of *Arabidopsis* early seedling development. *J. Exp. Bot.* 64, 4301–4312. doi: 10.1093/jxb/ert239
- Dugas, D. V., and Bartel, B. (2008). Sucrose induction of *Arabidopsis* miR398 represses two Cu/Zn superoxide dismutases. *Plant Mol. Biol.* 67, 403–417. doi: 10.1007/s11103-008-9329-1
- Farooq, M. A., Niazi, A. K., Akhtar, J., Saifullah, U., Farooq, M., Souri, Z., et al. (2019). Acquiring control: the evolution of ROS-induced oxidative stress and redox signaling pathways in plant stress responses. *Plant Physiol. Biochem.* 141, 353–369. doi: 10.1016/j.plaphy.2019.04.039
- Feldbrügge, M., Arizti, P., Sullivan, M. L., Zamore, P. D., Belasco, J. G., and Green, P. J. (2002). Comparative analysis of the plant mRNA-destabilizing element, DST, in mammalian and tobacco cells. *Plant Mol. Biol.* 49, 215–223. doi: 10.1023/a:1014936824187
- Feng, C., Chen, Y., Wang, C., Kong, Y., Wu, W., and Chen, Y. (2014). *Arabidopsis* RAV1 transcription factor, phosphorylated by SnRK2 kinases, regulates the expression of *ABI3*, *ABI4*, and *ABI5* during seed germination and early seedling development. *Plant J.* 80, 654–668. doi: 10.1111/tpj.12670
- Feng, J. X., Liu, D., Pan, Y., Gong, W., Ma, L. G., Luo, J. C., et al. (2005). An annotation update via cDNA sequence analysis and comprehensive

- profiling of developmental, hormonal or environmental responsiveness of the *Arabidopsis* AP2/EREBP transcription factor gene family. *Plant Mol. Biol.* 59, 853–868. doi: 10.1007/s11103-005-1511-0
- Feng, J., Wang, J., Fan, P., Jia, W., Nie, L., Jiang, P., et al. (2015). High-throughput deep sequencing reveals that microRNAs play important roles in salt tolerance of eukaryote *Salicornia europaea*. *BMC Plant Biol.* 15:63. doi: 10.1186/s12870-015-0451-3
- Figueroa, C. M., and Lunn, J. E. (2016). A tale of two sugars: trehalose 6-phosphate and sucrose. *Plant Physiol.* 172, 7–27. doi: 10.1104/pp.16.00417
- Flores-Pérez, Ú., Pérez-Gil, J., Closa, M., Wright, L. P., Botella-Pavía, P., Phillips, M. A., et al. (2010). Pleiotropic regulatory locus 1 (PRL1) integrates the regulation of sugar responses with isoprenoid metabolism in *Arabidopsis*. *Mol. Plant* 3, 101–112. doi: 10.1093/mp/ssp100
- Francischini, C. W., and Quaggio, R. B. (2009). Molecular characterization of *Arabidopsis thaliana* PUF proteins—binding specificity and target candidates. *FEBS J.* 276, 5456–5470. doi: 10.1111/j.1742-4658.2009.07230.x
- Francisco-Mangilet, A. G., Karlsson, P., Kim, M. H., Eo, H. J., Oh, S. A., Kim, J. H., et al. (2015). THO₂, a core member of the THO/TREX complex, is required for micro RNA production in *Arabidopsis*. *Plant J.* 82, 1018–1029. doi: 10.1111/tj.12874
- Fujiki, Y., Ito, M., Nishida, I., and Watanabe, A. (2000). Multiple signaling pathways in gene expression during sugar starvation. Pharmacological analysis of din gene expression in suspension-cultured cells of *Arabidopsis*. *Plant Physiol.* 124, 1139–1148. doi: 10.1104/pp.124.3.1139
- Fusaro, A. F., Bocca, S. N., Ramos, R. L. B., Barrôco, R. M., Magioli, C., Jorge, V. C., et al. (2007). AtGRP2, a cold-induced nucleocytoplasmic RNA-binding protein, has a role in flower and seed development. *Planta* 225, 1339–1351. doi: 10.1007/s00425-006-0444-4
- Gao, R., Wang, Y., Gruber, M. Y., and Hannoufa, A. (2018). miR156/SPL10 modulates lateral root development, branching and leaf morphology in *Arabidopsis* by silencing AGAMOUS-LIKE 79. *Front. Plant Sci.* 8:2226. doi: 10.3389/fpls.2017.02226
- Garapati, P., Feil, R., Lunn, J. E., Van Dijk, P., Balazadeh, S., and Mueller-Roeber, M. (2015). Transcription factor *Arabidopsis* activating factor1 integrates carbon starvation responses with trehalose metabolism. *Plant Physiol.* 169, 379–390. doi: 10.1104/pp.15.00917
- Gilmonreal, M., Zabalza, A., Missihoun, T. D., Dormann, P., Bartels, D., and Royuela, M. (2017). Induction of the PDH bypass and upregulation of the ALDH7B4 in plants treated with herbicides inhibiting amino acid biosynthesis. *Plant Sci.* 264, 16–28. doi: 10.1016/j.plantsci.2017.08.003
- Granot, D., Kelly, G., Stein, O., and David-Schwartz, R. (2014). Substantial roles of hexokinase and fructokinase in the effects of sugars on plant physiology and development. *J. Exp. Bot.* 65, 809–819. doi: 10.1093/jxb/ert400
- Guerra, D., Crosatti, C., Khoshro, H. H., Mastrangelo, A. M., Mica, E., and Mazzucotelli, E. (2015). Post-transcriptional and post-translational regulations of drought and heat response in plants: a spider's web of mechanisms. *Front. Plant Sci.* 6:57. doi: 10.3389/fpls.2015.00057
- Hartmann, L., Drewe-Boß, P., Wießner, T., Wagner, G., Geue, S., Lee, H. C., et al. (2016). Alternative splicing substantially diversifies the transcriptome during early photomorphogenesis and correlates with the energy availability in *Arabidopsis*. *Plant Cell* 28, 2715–2734. doi: 10.1105/tpc.16.00508
- He, H., Chincinska, I., Hackel, A., Grimm, B., and Kühn, C. (2008). Phloem mobility and stability of sucrose transporter transcripts. *Open Plant Sci. J.* 2, 1–14. doi: 10.2174/1874294700802010001
- Ho, S. L., Chao, Y. C., Tong, W. F., and Yu, S. M. (2001). Sugar coordinately and differentially regulates growth-and stress-related gene expression via a complex signal transduction network and multiple control mechanisms. *Plant Physiol.* 125, 877–890. doi: 10.1104/pp.125.2.877
- Hou, Q., and Bartels, D. (2015). Comparative study of the aldehyde dehydrogenase (ALDH) gene superfamily in the glycophyte *Arabidopsis thaliana* and *Eutrema halophytes*. *Ann. Bot.* 115, 465–479. doi: 10.1093/aob/mcu152
- Hu, J., Liu, Y., Tang, X., Rao, H., and Pei, J. (2020). Transcriptome profiling of the flowering transition in saffron (*Crocus sativus* L.). *Sci. Rep.* 10:9680. doi: 10.1038/s41598-020-66675-6
- Hu, J. Y., Zhou, Y., He, F., Dong, X., Liu, L. Y., Coupland, G., et al. (2014). miR824-regulated AGAMOUS-LIKE16 contributes to flowering time repression in *Arabidopsis*. *Plant Cell* 26, 2024–2037. doi: 10.1105/tpc.114.124685
- Huang, L. F., Bockock, P. N., Davis, J. M., and Koch, K. E. (2007). Regulation of invertase: a 'suite' of transcriptional and post-transcriptional mechanisms. *Funct. Plant Biol.* 34, 499–507. doi: 10.1071/fp06227
- Huo, X., Wang, C., Teng, Y., and Liu, X. (2015). Identification of miRNAs associated with dark-induced senescence in *Arabidopsis*. *BMC Plant Biol.* 15:266. doi: 10.1186/s12870-015-0656-5
- Hwang, G., Kim, S., Cho, J. Y., Paik, I., Kim, J. I., and Oh, E. (2019). Trehalose-6-phosphate signaling regulates thermoresponsive hypocotyl growth in *Arabidopsis thaliana*. *EMBO Rep.* 20:e47828. doi: 10.15252/embr.201947828
- Jang, Y. H., Park, H. Y., Kim, S. K., Lee, J. H., Suh, M. C., Chung, Y. S., et al. (2009). Survey of rice proteins interacting with OsFCA and OsFY proteins which are homologous to the *Arabidopsis* flowering time proteins, FCA and FY. *Plant Cell Physiol.* 50, 1479–1492. doi: 10.1093/pcp/pcp093
- Janse van Rensburg, H. C., and Van den Ende, W. (2018). UDP-glucose: a potential signaling molecule in plants? *Front. Plant Sci.* 8:2230. doi: 10.3389/fpls.2017.02230
- Jia, X., Wang, W., Ren, L., Chen, Q., Mendu, V., Willcutt, B., et al. (2009). Differential and dynamic regulation of miR398 in response to ABA and salt stress in *Populus tremula* and *Arabidopsis thaliana*. *Plant Mol. Biol.* 71, 51–59. doi: 10.1007/s11103-009-9508-8
- Jia, T., Zhang, B., You, C., Zhang, Y., Zeng, L., Li, S., et al. (2017). The *Arabidopsis* MOS4-associated complex promotes microRNA biogenesis and precursor messenger RNA splicing. *Plant Cell* 29, 2626–2643. doi: 10.1105/tpc.17.00370
- Jiang, S. C., Mei, C., Liang, S., Yu, Y. T., Lu, K., Wu, Z., et al. (2015). Crucial roles of the pentatricopeptide repeat protein SOAR1 in *Arabidopsis* response to drought, salt and cold stresses. *Plant Mol. Biol.* 88, 369–385. doi: 10.1007/s11103-015-0327-9
- Jiao, Y., Lei, W., Xu, W., and Chen, W. L. (2019). Glucose signaling, AtRGS1 and plant autophagy. *Plant Signal. Behav.* 14:1607465. doi: 10.1080/15592324.2019.1607465
- Jiao, Y., Wang, Y., Xue, D., Wang, J., Yan, M., Liu, G., et al. (2020). Regulation of OsSPL14 by OsmiR156 defines ideal plant architecture in rice. *Nat. Genet.* 42, 541–544. doi: 10.1038/ng.591
- Jung, J. H., Seo, P. J., Ahn, J. H., and Park, C. M. (2012). *Arabidopsis* RNA-binding protein FCA regulates microRNA172 processing in thermosensory flowering. *J. Biol. Chem.* 287, 16007–16016. doi: 10.1074/jbc.M111.337485
- Kanwar, P., and Jha, G. (2019). Alterations in plant sugar metabolism: signatory of pathogen attack. *Planta* 249, 305–318. doi: 10.1007/s00425-018-3018-3
- Karlsson, P., Christie, M. D., Seymour, D. K., Wang, H., Wang, X., Hagemann, J., et al. (2015). KH domain protein RCF3 is a tissue-biased regulator of the plant miRNA biogenesis cofactor HYL1. *Proc. Natl. Acad. Sci. U. S. A.* 112, 14096–14101. doi: 10.1073/pnas.1512865112
- Kawa, D., and Testerink, C. (2017). Regulation of mRNA decay in plant responses to salt and osmotic stress. *Cell. Mol. Life Sci.* 74, 1165–1176. doi: 10.1007/s00018-016-2376-x
- Kesarwani, A. K., Lee, H. C., Ricca, P. G., Sullivan, G., Faiss, N., Wagner, G., et al. (2019). Multifactorial and species-specific feedback regulation of the RNA surveillance pathway nonsense-mediated decay in plants. *Plant Cell Physiol.* 60, 1986–1999. doi: 10.1093/pcp/pcz141
- Kim, Y. O., and Kang, H. (2006). The role of a zinc finger-containing glycine-rich RNA-binding protein during the cold adaptation process in *Arabidopsis thaliana*. *Plant Cell Physiol.* 47, 793–798. doi: 10.1093/pcp/pcj047
- Kim, J. J., Lee, J. H., Kim, W., Jung, H. S., Huijser, P., and Ahn, J. H. (2012). The microRNA156-SQUAMOSA PROMOTER BINDING PROTEIN-LIKE3 module regulates ambient temperature-responsive flowering via FLOWERING LOCUS T in *Arabidopsis*. *Plant Physiol.* 159, 461–478. doi: 10.1104/pp.111.192369
- Kim, Y. O., Pan, S., Jung, C. H., and Kang, H. (2007b). A zinc finger-containing glycine-rich RNA-binding protein, atRZ-1a, has a negative impact on seed germination and seedling growth of *Arabidopsis thaliana* under salt or drought stress conditions. *Plant Cell Physiol.* 48, 1170–1181. doi: 10.1093/pcp/pcm087
- Kim, J. Y., Park, S. J., Jang, B., Jung, C. H., Ahn, S. J., Goh, C. H., et al. (2007a). Functional characterization of a glycine-rich RNA-binding protein 2 in *Arabidopsis thaliana* under abiotic stress conditions. *Plant J.* 50, 439–451. doi: 10.1111/j.1365-313X.2007.03057.x
- Kirch, H. H., Schlingensiepen, S., Kotchoni, S., Sunkar, R., and Bartels, D. (2005). Detailed expression analysis of selected genes of the aldehyde dehydrogenase (ALDH) gene superfamily in *Arabidopsis thaliana*. *Plant Mol. Biol.* 57, 315–332. doi: 10.1007/s11103-004-7796-6

- Kohorn, B. D., Johansen, S., Shishido, A., Todorova, T., Martinez, R., Defeo, E., et al. (2009). Pectin activation of MAP kinase and gene expression is WAK2 dependent. *Plant J.* 60, 974–982. doi: 10.1111/j.1365-313X.2009.04016.x
- Kohorn, B. D., Kobayashi, M., Johansen, S., Riese, J., Huang, L. F., Koch, K., et al. (2006). An *Arabidopsis* cell wall-associated kinase required for invertase activity and cell growth. *Plant J.* 46, 307–316. doi: 10.1111/j.1365-313X.2006.02695.x
- Koyama, T., Sato, F., and Ohme-Takagi, M. (2017). Roles of miR319 and TCP transcription factors in leaf development. *Plant Physiol.* 175, 874–885. doi: 10.1104/pp.17.00732
- Kühn, C., and Grof, C. P. L. (2010). Sucrose transporters of higher plants. *Curr. Opin. Plant Biol.* 13, 287–297. doi: 10.1016/j.pbi.2010.02.001
- Kühn, C., Franceschi, V. R., Schulz, A., Lemoine, R., and Frommer, W. B. (1997). Macromolecular trafficking indicated by localization and turnover of sucrose transporters in enucleate sieve elements. *Science* 275, 1298–1300. doi: 10.1126/science.275.5304.1298
- Kumar, A., Kondhare, K. R., Vetal, P. V., and Banerjee, A. K. (2020). PcG proteins MSI1 and BMI1 function upstream of miR156 to regulate aerial tuber formation in potato. *Plant Physiol.* 182, 185–203. doi: 10.1104/pp.19.00416
- Lastdrager, J., Hanson, J., and Smeekens, S. (2014). Sugar signals and the control of plant growth and development. *J. Exp. Bot.* 65, 799–807. doi: 10.1093/jxb/ert474
- Lee, K., and Kang, H. (2016). Emerging roles of RNA-binding proteins in plant growth, development, and stress responses. *Mol. Cell* 39, 179–185. doi: 10.14348/molcells.2016.2359
- Li, X., Cai, W., Liu, Y., Li, H., Fu, L., Liu, Z., et al. (2017). Differential TOR activation and cell proliferation in *Arabidopsis* root and shoot apices. *Proc. Natl. Acad. Sci. U. S. A.* 114, 2765–2770. doi: 10.1073/pnas.1618782114
- Li, L., and Sheen, J. (2016). Dynamic and diverse sugar signaling. *Curr. Opin. Plant Biol.* 33, 116–125. doi: 10.1016/j.pbi.2016.06.018
- Liesche, J., Krügel, U., He, H., Chincinska, I., Hackel, A., and Kühn, C. (2011). Sucrose transporter regulation at the transcriptional, post-transcriptional and post-translational level. *J. Plant Physiol.* 168, 1426–1433. doi: 10.1016/j.jplph.2011.02.005
- Lin, J. S., Kuo, C. C., Yang, I. C., Tsai, W. A., Shen, Y. H., Lin, C. C., et al. (2018). MicroRNA160 modulates plant development and heat shock protein gene expression to mediate heat tolerance in *Arabidopsis*. *Front. Plant Sci.* 9:68. doi: 10.3389/fpls.2018.00068
- Liu, Y., and Bassham, D. C. (2010). TOR is a negative regulator of autophagy in *Arabidopsis thaliana*. *PLoS One* 5:e11883. doi: 10.1371/journal.pone.0011883
- Liu, F., Quesada, V., Crevillén, P., Bäurle, I., Swiezewski, S., and Dean, C. (2007). The *Arabidopsis* RNA-binding protein FCA requires a lysine-specific demethylase 1 homolog to downregulate FLC. *Mol. Cell* 28, 398–407. doi: 10.1016/j.molcel.2007.10.018
- Lorković, Z. J. (2009). Role of plant RNA-binding proteins in development, stress response and genome organization. *Trends Plant Sci.* 14, 229–236. doi: 10.1016/j.tplants.2009.01.007
- Lu, Y., Sun, J., Yang, Z., Zhao, C., Zhu, M., Ma, D., et al. (2019). Genome-wide identification and expression analysis of glycine-rich RNA-binding protein family in sweet potato wild relative *Ipomoea trifida*. *Gene* 686, 177–186. doi: 10.1016/j.gene.2018.11.044
- Macknight, R., Bancroft, I., Page, T., Lister, C., Schmidt, R., Love, K., et al. (1997). FCA, a gene controlling flowering time in *Arabidopsis*, encodes a protein containing RNA-binding domains. *Cell* 89, 737–745. doi: 10.1016/S0092-8674(00)80256-1
- Mahalingam, R., and Walling, J. G. (2020). Genomic survey of RNA recognition motif (RRM) containing RNA binding proteins from barley (*Hordeum vulgare* ssp. *vulgare*). *Genomics* 112, 1829–1839. doi: 10.1016/j.ygeno.2019.10.016
- Manavella, P. A., Hagmann, J., Ott, F., Laubinger, S., Franz, M., Macek, B., et al. (2012). Fast-forward genetics identifies plant CPL phosphatases as regulators of miRNA processing factor HYL1. *Cell* 151, 859–870. doi: 10.1016/j.cell.2012.09.039
- Marondedze, C., Thomas, L., Serrano, N. L., Lilley, K. S., and Gehring, C. (2016). The RNA-binding protein repertoire of *Arabidopsis thaliana*. *Sci. Rep.* 6:29766. doi: 10.1038/srep29766
- Marondedze, C., Thomas, L., Gehring, C., and Lilley, K. S. (2019). Changes in the *Arabidopsis* RNA-binding proteome reveal novel stress response mechanisms. *BMC Plant Biol.* 19:139. doi: 10.1186/s12870-019-1750-x
- Martin, R. C., Asahina, M., Liu, P., Kristof, J. R., Coppersmith, J. L., Pluskota, W. E., et al. (2010). The microRNA156 and microRNA172 gene regulation cascades at post-germinative stages in *Arabidopsis*. *Seed Sci. Res.* 20, 79–87. doi: 10.1017/s0960258510000085
- Matiolli, C. C., Tomaz, J. P., Duarte, G. T., Prado, F. M., Del Bem, L. E. V., Silveira, A. B., et al. (2011). The *Arabidopsis* bZIP gene AtbZIP63 is a sensitive integrator of transient abscisic acid and glucose signals. *Plant Physiol.* 157, 692–705. doi: 10.1104/pp.111.181743
- Mei, C., Jiang, S. C., Lu, Y. F., Wu, F. Q., Yu, Y. T., Liang, S., et al. (2014). *Arabidopsis* pentatricopeptide repeat protein SOAR1 plays a critical role in abscisic acid signalling. *J. Exp. Bot.* 65, 5317–5330. doi: 10.1093/jxb/eru293
- Meng, L., Xu, M., Wan, W., Yu, F., Li, C., Wang, J., et al. (2018). Sucrose signaling regulates anthocyanin biosynthesis through a MAPK cascade in *Arabidopsis thaliana*. *Genetics* 210, 607–619. doi: 10.1534/genetics.118.301470
- Meyers, B. C., and Axtell, M. J. (2019). MicroRNAs in plants: key findings from the early years. *Plant Cell* 31:1206. doi: 10.1105/tpc.19.00310
- Michaels, S. D., and Amasino, R. M. (1999). FLOWERING LOCUS C encodes a novel MADS domain protein that acts as a repressor of flowering. *Plant Cell* 11, 949–956. doi: 10.1105/tpc.11.5.949
- Milne, R. J., Perroux, J. M., Rae, A. L., Reinders, A., Ward, J. M., Offler, C. E., et al. (2017). Sucrose transporter localization and function in phloem unloading in developing stems. *Plant Physiol.* 173, 1330–1341. doi: 10.1104/pp.16.01594
- Nagarajan, V. K., Kukulich, P. M., von Hagel, B., and Green, P. J. (2019). RNA degradomes reveal substrates and importance for dark and nitrogen stress responses of *Arabidopsis* XRN4. *Nucleic Acids Res.* 47, 9216–9230. doi: 10.1093/nar/gkz712
- Nicolai, M., Roncato, M. A., Canoy, A. S., Rouquie, D., Sarda, X., Freysson, G., et al. (2006). Large-scale analysis of mRNA translation states during sucrose starvation in *Arabidopsis* cells identifies cell proliferation and chromatin structure as targets of translational control. *Plant Physiol.* 141, 663–673. doi: 10.1104/pp.106.079418
- Pan, J., Huang, D., Guo, Z., Kuang, Z., Zhang, H., Xie, X., et al. (2018). Overexpression of microRNA408 enhances photosynthesis, growth, and seed yield in diverse plants. *J. Integr. Plant Biol.* 60, 323–340. doi: 10.1111/jipb.12634
- Pant, B. D., Musialak-Lange, M., Nuc, P., May, P., Buhtz, A., Kehr, J., et al. (2009). Identification of nutrient-responsive *Arabidopsis* and rapeseed microRNAs by comprehensive real-time polymerase chain reaction profiling and small RNA sequencing. *Plant Physiol.* 150, 1541–1555. doi: 10.1104/pp.109.139139
- Park, S. Y., and Grabau, E. (2017). Bypassing miRNA-mediated gene regulation under drought stress: alternative splicing affects CSD1 gene expression. *Plant Mol. Biol.* 95, 243–252. doi: 10.1007/s11103-017-0642-4
- Pomeranz, M. C., Hah, C., Lin, P. C., Kang, S. G., Finer, J. J., Blackshear, P. J., et al. (2010). The *Arabidopsis* tandem zinc finger protein AtTZF1 traffics between the nucleus and cytoplasmic foci and binds both DNA and RNA. *Plant Physiol.* 152, 151–165. doi: 10.1104/pp.109.145656
- Ponnu, J., Schlereth, A., Zacharaki, V., Działo, M. A., Abel, C., Feil, R., et al. (2020). The trehalose 6-phosphate pathway impacts vegetative phase change in *Arabidopsis thaliana*. *Plant J.* doi: 10.1111/tpj.14965 [Epub ahead of print]
- Qu, J., Kang, S. G., Wang, W., Musier-Forsyth, K., and Jang, J. C. (2014). The *Arabidopsis thaliana* tandem zinc finger 1 (AtTZF1) protein in RNA binding and decay. *Plant J.* 78, 452–467. doi: 10.1111/tpj.12485
- Reichel, M., Liao, Y., Rettel, M., Ragan, C., Evers, M., Alleaume, A. M., et al. (2016). In planta determination of the mRNA-binding proteome of *Arabidopsis* etiolated seedlings. *Plant Cell* 28, 2435–2452. doi: 10.1105/tpc.16.00562
- Ren, L., and Tang, G. (2012). Identification of sucrose-responsive microRNAs reveals sucrose-regulated copper accumulations in an SPL7-dependent and independent manner in *Arabidopsis thaliana*. *Plant Sci.* 187, 59–68. doi: 10.1016/j.plantsci.2012.01.014
- Ren, G., Xie, M., Dou, Y., Zhang, S., Zhang, C., and Yu, B. (2012). Regulation of miRNA abundance by RNA binding protein TOUGH in *Arabidopsis*. *Proc. Natl. Acad. Sci. U. S. A.* 109, 12817–12821. doi: 10.1073/pnas.1204915109
- Riechmann, J. L., Heard, J., Martin, G., Reuber, L., Jiang, C. -Z., Keddie, J., et al. (2000). *Arabidopsis* transcription factors: genome-wide comparative analysis among eukaryotes. *Science* 290, 2105–2110. doi: 10.1126/science.290.5499.2105

- Rigo, R., Bazin, J., Crespi, M., and Charon, C. (2019). Alternative splicing in the regulation of plant–microbe interactions. *Plant Cell Physiol.* 60, 1906–1916. doi: 10.1093/pcp/pcz086
- Rodriguez, M., Parola, R., Andreola, S., Pereyra, C., and Martínez-Noël, G. (2019). TOR and SnRK1 signaling pathways in plant response to abiotic stresses: do they always act according to the “yin-yang” model? *Plant Sci.* 288:110220. doi: 10.1016/j.plantsci.2019.110220
- Rogers, K., and Chen, X. (2013). Biogenesis, turnover, and mode of action of plant microRNAs. *Plant Cell* 25, 2383–2399. doi: 10.1105/tpc.113.113159
- Romanowski, A., and Yanovsky, M. J. (2015). Circadian rhythms and post-transcriptional regulation in higher plants. *Front. Plant Sci.* 6:437. doi: 10.3389/fpls.2015.00437
- Rosenberger, C. L., and Chen, J. (2018). To grow or not to grow: TOR and SnRK2 coordinate growth and stress response in *Arabidopsis*. *Mol. Cell* 69, 3–4. doi: 10.1016/j.molcel.2017.12.013
- Sakr, S., Wang, M., Dédaldéchamp, F., Perez-Garcia, M. D., Ogé, L., Hamama, L., et al. (2018). The sugar-signaling hub: overview of regulators and interaction with the hormonal and metabolic network. *Int. J. Mol. Sci.* 19:2506. doi: 10.3390/ijms19092506
- Samad, A. F. A., Sajad, M., Nazaruddin, N., Fauzi, I. A., Murad, A., Murad, A. M. A., et al. (2017). MicroRNA and transcription factor: key players in plant regulatory network. *Front. Plant Sci.* 8:565. doi: 10.3389/fpls.2017.00565
- Sami, F., Siddiqui, H., and Hayat, S. (2019). Interaction of glucose and phytohormone signaling in plants. *Plant Physiol. Biochem.* 135, 119–126. doi: 10.1016/j.plaphy.2018.11.005
- Sheu, J. J., Jan, S. P., Lee, H. T., and Yu, S. M. (1994). Control of transcription and mRNA turnover as mechanisms of metabolic repression of α -amylase gene expression. *Plant J.* 5, 655–664. doi: 10.1111/j.1365-313x.1994.00655.x
- Shi, L., Wu, Y., and Sheen, J. (2018). TOR signaling in plants: conservation and innovation. *Development* 145:dev160887. doi: 10.1242/dev.160887
- Shinozawa, A., Otake, R., Takezawa, D., Umezawa, T., Komatsu, K., Tanaka, K., et al. (2019). SnRK2 protein kinases represent an ancient system in plants for adaptation to a terrestrial environment. *Commun. Biol.* 2:30. doi: 10.1038/s42003-019-0281-1
- Signorelli, S., Masclaux-Daubresse, C., Moriyasu, Y., Van den Ende, W., and Bassham, D. C. (2019). Sugars and autophagy in plants. *Front. Plant Sci.* 10:1190. doi: 10.3389/fpls.2019.01190
- Silverman, I. M., Li, F., and Gregory, B. D. (2013). Genomic era analyses of RNA secondary structure and RNA-binding proteins reveal their significance to post-transcriptional regulation in plants. *Plant Sci.* 205–206, 55–62. doi: 10.1016/j.plantsci.2013.01.009
- Song, X., Li, Y., Cao, X., and Qi, Y. (2019). MicroRNAs and their regulatory roles in plant–environment interactions. *Annu. Rev. Plant Biol.* 70, 489–525. doi: 10.1146/annurev-arplant-050718-100334
- Stitt, M., and Zeeman, S. C. (2012). Starch turnover: pathways, regulation and role in growth. *Curr. Opin. Plant Biol.* 15, 282–292. doi: 10.1016/j.pbi.2012.03.016
- Su, C., Li, Z., Cheng, J., Li, L., Zhong, S., Liu, L., et al. (2017). The protein phosphatase 4 and SMEK1 complex dephosphorylates HYL1 to promote miRNA biogenesis by antagonizing the MAPK cascade in *Arabidopsis*. *Dev. Cell* 41, 527.e5–539.e5. doi: 10.1016/j.devcel.2017.05.008
- Sun, Z., Li, M., Zhou, Y., Guo, T., Liu, Y., Zhang, H., et al. (2018). Coordinated regulation of *Arabidopsis* microRNA biogenesis and red light signaling through Dicer-like 1 and phytochrome-interacting factor 4. *PLoS Genet.* 14:e1007247. doi: 10.1371/journal.pgen.1007247
- Sunkar, R., Kapoor, A., and Zhu, J. K. (2006). Post-transcriptional induction of two Cu/Zn superoxide dismutase genes in *Arabidopsis* is mediated by downregulation of miR398 and important for oxidative stress tolerance. *Plant Cell* 18, 2051–2065. doi: 10.1105/tpc.106.041673
- Swarbreck, D., Wilks, C., Lamesch, P., Berardini, T. Z., Garcia-Hernandez, M., Foerster, H., et al. (2007). The *Arabidopsis* information resource (TAIR): gene structure and function annotation. *Nucleic Acids Res.* 36(Suppl. 1), D1009–D1014. doi: 10.1093/nar/gkm965
- Tian, L., Chou, H. L., Zhang, L., Hwang, S. K., Starkenburg, S. R., Doroshenko, K. A., et al. (2018a). RNA-binding protein RBP-P is required for glutelin and prolamine mRNA localization in rice endosperm cells. *Plant Cell* 30, 2529–2552. doi: 10.1105/tpc.18.00321
- Tian, L., Liu, H., Ren, L., Ku, L., Wu, L., Li, M., et al. (2018b). MicroRNA 399 as a potential integrator of photo-response, phosphate homeostasis, and sucrose signaling under long day condition. *BMC Plant Biol.* 18:290. doi: 10.1186/s12870-018-1460-9
- Van Leene, J., Han, C., Gadeyne, A., Eeckhout, D., Matthijs, C., Cannoot, B., et al. (2019). Capturing the phosphorylation and protein interaction landscape of the plant TOR kinase. *Nat. Plants* 5:316. doi: 10.1038/s41477-019-0378-z
- van Mourik, H., van Dijk, A. D., Stortenbeker, N., Angenent, G. C., and Bemer, M. (2017). Divergent regulation of *Arabidopsis* SAUR genes: a focus on the SAUR10-clade. *BMC Plant Biol.* 17:245. doi: 10.1186/s12870-017-1210-4
- Voinnet, O. (2009). Origin, biogenesis, and activity of plant microRNAs. *Cell* 136, 669–687. doi: 10.1016/j.cell.2009.01.046
- Wahl, V., Ponnu, J., Schlereth, A., Arrivault, S., Langenecker, T., Franke, A., et al. (2013). Regulation of flowering by trehalose-6-phosphate signaling in *Arabidopsis thaliana*. *Science* 339, 704–707. doi: 10.1126/science.1230406
- Wang, J., Czech, B., and Weigel, D. (2009). miR156-regulated SPL transcription factors define an endogenous flowering pathway in *Arabidopsis thaliana*. *Cell* 138, 738–749. doi: 10.1016/j.cell.2009.06.014
- Wang, Y., Li, L., Ye, T., Lu, Y., Chen, X., and Wu, Y. (2013). The inhibitory effect of ABA on floral transition is mediated by ABI5 in *Arabidopsis*. *J. Exp. Bot.* 64, 675–684. doi: 10.1093/jxb/ers361
- Wang, M., Moigne, M. L., Bertheloot, J., Crespel, L., Perezgarcia, M. D., Oge, L., et al. (2019b). BRANCHED1: a key hub of shoot branching. *Front. Plant Sci.* 10:76. doi: 10.3389/fpls.2019.00076
- Wang, M., Ogé, L., Perez-Garcia, M. D., Hamama, L., and Sakr, S. (2018a). The PUF protein family: overview on PUF RNA targets, biological functions, and post-transcriptional regulation. *Int. J. Mol. Sci.* 19:410. doi: 10.3390/ijms19020410
- Wang, M., Ogé, L., Voisine, L., Perez-Garcia, M. D., Jeuffre, J., Hibrand Saint-Oyant, L., et al. (2019a). Post-transcriptional regulation of *RhBRC1* (*Rosa hybrida* *BRANCHED1*) in response to sugars is mediated via its own 3′ untranslated region, with a potential role of RbPUF4 (Pumilio RNA-binding protein family). *Int. J. Mol. Sci.* 20:3808. doi: 10.3390/ijms20153808
- Wang, S., Quan, L., Li, S., You, C., Zhang, Y., Gao, L., et al. (2019c). The PROTEIN PHOSPHATASE4 complex promotes transcription and processing of primary microRNAs in *Arabidopsis*. *Plant Cell* 31, 486–501. doi: 10.1105/tpc.18.00556
- Wang, H., and Wang, H. (2015). The miR156/SPL module, a regulatory hub and versatile toolbox, gears up crops for enhanced agronomic traits. *Mol. Plant* 8, 677–688. doi: 10.1016/j.molp.2015.01.008
- Wang, P., Zhao, Y., Li, Z., Hsu, C., Liu, X., Fu, L., et al. (2018b). Reciprocal regulation of the TOR kinase and ABA receptor balances plant growth and stress response. *Mol. Cell* 69, 100.e6–112.e6. doi: 10.1016/j.molcel.2017.12.002
- Wei, Q., Ma, C., Xu, Y., Wang, T., Chen, Y., Lü, J., et al. (2017). Control of chrysanthemum flowering through integration with an aging pathway. *Nat. Commun.* 8:829. doi: 10.1038/s41467-017-00812-0
- Wilkinson, M. E., Charenton, C., and Nagai, K. (2020). RNA splicing by the spliceosome. *Annu. Rev. Biochem.* 89, 359–388. doi: 10.1146/annurev-biochem-091719-064225
- Wingler, A. (2018). Transitioning to the next phase: the role of sugar signaling throughout the plant life cycle. *Plant Physiol.* 176, 1075–1084. doi: 10.1104/pp.17.01229
- Wu, G., Park, M. Y., Conway, S. R., Wang, J. W., Weigel, D., and Poethig, R. S. (2009). The sequential action of miR156 and miR172 regulates developmental timing in *Arabidopsis*. *Cell* 138, 750–759. doi: 10.1016/j.cell.2009.06.031
- Wu, X., Shi, Y., Li, J., Xu, L., Fang, Y., Li, X., et al. (2013). A role for the RNA-binding protein MOS2 in microRNA maturation in *Arabidopsis*. *Cell Res.* 23, 645–657. doi: 10.1038/cr.2013.23
- Xu, M., Hu, T., Zhao, J., Park, M. Y., Earley, K. W., Wu, G., et al. (2016). Developmental functions of miR156-regulated SQUAMOSA PROMOTER BINDING PROTEIN-LIKE (SPL) genes in *Arabidopsis thaliana*. *PLoS Genet.* 12:e1006263. doi: 10.1371/journal.pgen.1006263
- Yamasaki, H., Hayashi, M., Fukazawa, M., Kobayashi, Y., and Shikanai, T. (2009). SQUAMOSA promoter binding protein-like7 is a central regulator for copper homeostasis in *Arabidopsis*. *Plant Cell* 21, 347–361. doi: 10.1105/tpc.108.060137
- Yan, J., Wang, P., Wang, B., Hsu, C. C., Tang, K., Zhang, H., et al. (2017). The SnRK2 kinases modulate miRNA accumulation in *Arabidopsis*. *PLoS Genet.* 13:e1006753. doi: 10.1371/journal.pgen.1006753
- Yang, L., Xu, M., Koo, Y., He, J., and Poethig, R. S. (2013). Sugar promotes vegetative phase change in *Arabidopsis thaliana* by repressing the expression of MIR156A and MIR156C. *eLife* 2:e00260. doi: 10.7554/eLife.00260

- Yoine, M., Ohto, M. A., Onai, K., Mita, S., and Nakamura, K. (2006). The lba1 mutation of UPF1 RNA helicase involved in nonsense-mediated mRNA decay causes pleiotropic phenotypic changes and altered sugar signalling in *Arabidopsis*. *Plant J.* 47, 49–62. doi: 10.1111/j.1365-313X.2006.02771.x
- Yu, S., Cao, L., Zhou, C. M., Zhang, T. Q., Lian, H., Sun, Y., et al. (2013). Sugar is an endogenous cue for juvenile-to-adult phase transition in plants. *eLife* 2:e00269. doi: 10.7554/eLife.00269
- Yu, H., Cong, L., Zhu, Z., Wang, C., Zou, J., Tao, C., et al. (2015). Identification of differentially expressed microRNA in the stems and leaves during sugar accumulation in sweet sorghum. *Gene* 571, 221–230. doi: 10.1016/j.gene.2015.06.056
- Zhang, B. (2015). MicroRNA: a new target for improving plant tolerance to abiotic stress. *J. Exp. Bot.* 66, 1749–1761. doi: 10.1093/jxb/erv013
- Zhang, S., Liu, Y., and Yu, B. (2014). PRL1, an RNA-binding protein, positively regulates the accumulation of miRNAs and siRNAs in *Arabidopsis*. *PLoS Genet.* 10:e1004841. doi: 10.1371/journal.pgen.1004841
- Zhang, H., Mao, X., and Jing, R. (2011). SnRK2 acts within an intricate network that links sucrose metabolic and stress signaling in wheat. *Plant Signal. Behav.* 6, 652–654. doi: 10.4161/psb.6.5.14945
- Zhao, J., Missihoun, T. D., and Bartels, D. (2017). The role of *Arabidopsis* aldehyde dehydrogenase genes in response to high temperature and stress combinations. *J. Exp. Bot.* 68, 4295–4308. doi: 10.1093/jxb/erx194
- Zhao, J., Missihoun, T. D., and Bartels, D. (2018). The ATAF1 transcription factor is a key regulator of aldehyde dehydrogenase 7B4 (ALDH7B4) gene expression in *Arabidopsis thaliana*. *Planta* 248, 1017–1027. doi: 10.1007/s00425-018-2955-1
- Zheng, J., Ma, Y., Zhang, M., Lyu, M., Yuan, Y., and Wu, B. (2019a). Expression pattern of FT/TFL1 and miR156-targeted SPL genes associated with developmental stages in *Dendrobium catenatum*. *Int. J. Mol. Sci.* 20:2725. doi: 10.3390/ijms20112725
- Zheng, Z., Xu, X., Crosley, R. A., Greenwalt, S. A., Sun, Y., Blakeslee, B., et al. (2010). The protein kinase SnRK2.6 mediates the regulation of sucrose metabolism and plant growth in *Arabidopsis*. *Plant Physiol.* 153, 99–113. doi: 10.1104/pp.109.150789
- Zheng, C., Ye, M., Sang, M., and Wu, R. (2019b). A regulatory network for miR156-SPL module in *Arabidopsis thaliana*. *Int. J. Mol. Sci.* 20:6166. doi: 10.3390/ijms20246166

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Wang, Zang, Jiao, Perez-Garcia, Ogé, Hamama, Le Gourrierec, Sakr and Chen. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.