A meta-analysis of paleolimnological records reveals the sensitivity of lacustrine carbon burial rates to carbon sources and preservation conditions during the Anthropocene

Jean-Philippe Jenny, C Niemann, P Francus, A Noren, N Carvalhais

A meta-analysis of paleolimnological records reveals the sensitivity of lacustrine carbon burial rates to carbon sources and preservation conditions during the Anthropocene

JP. Jenny[1*], C. Niemann[2], P. Francus[3], A. Noren[4], N. Carvalhais[2,5]

[1]INRA, UMR CARRTEL, Université Savoie Mont Mont Blanc, Thonon les Bains, France,
[2]Max Planck Institute for Biogeochemistry, Jena, Germany,
[3]Centre Eau Terre Environnement, INRS, G1K9A9 Québec (Qc), Canada and GEOTOP Research Center, Montréal, (Qc), Canada, [4]National Lacustrine Core Facility (LacCore), Minneapolis, USA
[5]Departamento de Ciências e Engenharia do Ambiente, DCEA, Faculdade de Ciências e Tecnologia, FCT, Universidade Nova de Lisboa, 2829-516 Caparica, Portugal.
e-mail: jean-philippe.jenny@inrae.fr

Inland waters are sites of active carbon (C) processing and transport along the land to ocean aquatic continuum (LOAC) that need to be accounted for closing the global C budget[1,2]. However, monitoring data are lacking and do not extend back as far as few decades, limiting our comprehension of the LOAC C cycle for the last centuries. Lake sediments provide a key archive for assessing C transport and transformation that occurs in lake catchments. Here, the analysis of large numbers of samples was performed on 420 lakes sediment records of the world to assess C burial rates and sources during the last 300 years. C and N (organic and mineral forms) on discrete samples were analyzed using a Variomax elemental analyzer to assess total C sequestration by lakes, C sources, and long-term changes in the contribution of allochtonous vs autochtonous sources to C transfers in lake-watersheds. Continuous sediment records were generated using core scanners (i.e. micro-XRF) and computed tomography to provide near-annual trends on terrigenous elements, here used as proxies of allochtonous sources (e.g. Al, Ti, K, Fe), and Mn:Fe ratio to infer past oxygen conditions[3]. Our results suggest that establishing a morphology-relevant lake typology that better characterises the types and distribution of oxygen conditions and terrigenous supplies across our sites is therefore the first step in providing a more robust evidence base for explaining the spatial-temporal variation in lake C burial rates.

1. Montgomery, D. R. Soil erosion and agricultural sustainability. *Proc. Natl. Acad. Sci.* **104**, 13268–13272 (2007).
2. Borrelli, P. *et al.* An assessment of the global impact of 21st century land use change on soil erosion. *Nat. Commun.* **8**, 2013 (2017).
3. Jenny, J.-P. *et al.* Contrasted effects of climate change on temperate large lakes oxygen-depletion (Lakes Geneva, Bourget, Annecy). in vol. 15 10075 (2013).