



HAL
open science

Nutritional LCA improves the understanding of the environmental impacts of foods taking into account the diversity of recipes within a same food category: the case of pizzas

Adeline Cortesi, Gwenola Yannou-Le Bris, Isabelle Souchon, Anne Saint-Eve,
Louis Georges Soler, Caroline Pénicaud

► To cite this version:

Adeline Cortesi, Gwenola Yannou-Le Bris, Isabelle Souchon, Anne Saint-Eve, Louis Georges Soler, et al.. Nutritional LCA improves the understanding of the environmental impacts of foods taking into account the diversity of recipes within a same food category: the case of pizzas. 12. International Conference on Life Cycle Assessment of Food 2020 (LCA Food 2020), Oct 2020, Virtual from Berlin, Germany. hal-03086180

HAL Id: hal-03086180

<https://hal.inrae.fr/hal-03086180v1>

Submitted on 5 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Nutritional LCA improves the understanding of the environmental impacts of foods taking into account the diversity of recipes within a same food category: the case of pizzas



Adeline Cortesi^{1,*}, Gwenola Yannou-Le Bris², Isabelle Souchon³, Anne Saint-Eve¹, Louis-Georges Soler⁴, Caroline Pénicaut¹

1: Université Paris-Saclay, INRAE, AgroParisTech, UMR SayFood, 78850, Thiverval-Grignon, France

2: Université Paris-Saclay, INRAE, AgroParisTech, UMR SayFood, 91300, Massy, France

3: Université d'Avignon, INRAE, UMR SQPOV, 84911 Avignon Cedex 9, France

4: Université Paris-Saclay, INRAE, UR ALISS, 94205, Ivry-sur-Seine, France

*Contact: adeline.cortesi@inrae.fr

Introduction

Context:

30% of the European global impact on the environment is directly linked to our food and this number is predicted to increase in the next few years due to the rise of the world population (Tukker *et al.*, 2006).

➔ There is a need to better understand the environmental impact of each food product so food industries can reformulate their products. Also consumers can make more sustainable food choices notably thanks to the environmental labelling.

Problems:

-Most of the studies focus on a single representant of each category of food products (Farahani *et al.*, 2019; Tsarouhas *et al.*, 2015).

➔ Lack of knowledge about the possible variability of environmental impact between several food products belonging to the same category.

-Most of the studies using LCA to evaluate the environmental impact of food products use a mass-based Functional Unit (FU) (Kim *et al.*, 2013; Farahani *et al.*, 2019), which is not representative of food products quality. Furthermore, studies using nutritional FU mainly focus on comparing different food categories (Saarinen *et al.*, 2017). The usefulness of nutritional FU to compare products within a same food category has not been investigated.

Objectives:


-To study the possible differences between a large number of products from the same food category in terms of environmental impact with the example of 80 different commercial pizzas.

-To evaluate how the use of different nutritional FU can affect the conclusions about environmental impact of 80 different pizzas.

Methodology: Pizzas selection

80 pizzas have been selected, using the OQALI database (<https://www.oqali.fr/Donnees-publiques>) to represent the 392 pizzas of the French retail market (2010) in terms of families, sectors, distributors and nutritional properties.

Recipes of the pizzas were calculated using product labelling. Ingredient compositions (such as dough, tomato sauce...) are supposed to be the same for all pizzas and only their amount change between pizzas.



Percentage of pizzas from different families, sectors and distributors among the 392 pizzas of the French retail market and among the selected sample of 80 pizzas

Methodology: LCA

Data collection was obtained from environmental databases (mainly Ecoinvent and Agribalyse), literature (technical informations from suppliers) and experimental measurements of packagings.

The LCA of those pizzas were realized on SimaPro software using the « EF 3.0 Method (adapted) V1.00 / EF 3.0 normalization and weighting set ». All indicators from this methodology have been studied but this poster will present only results related to Climate Change.

Methodology: Different FU

Mass-based FU

The mass-based FU is 1kg of ready-to-eat pizza. The links between impact on climate change of pizzas and their compositions and nutritional characteristics have been studied using a Partial Least Square Regression on XLStat software.

Nutritional FU

• Several nutritional FU have been tested. To do so, the mass of pizza needed to get a certain amount of different nutrients has been determined for each pizza and the associated impact on climate change has been calculated.

- We selected FU based on:
 - Recommended portion (mass of the portion recommended on the packaging)
 - Energy content (mass of pizza needed to obtain 100kcal)
 - Protein content (mass of pizza needed to obtain 1/6 of daily recommendations)
 - Fibre content (mass of pizza needed to obtain 1/6 of daily recommendations)
 - Fibre and protein contents combined (mass of pizza needed to obtain at least 1/6 of daily recommendations in proteins and fibres)


Results with mass-based FU

Impact of each step of pizza production on Climate Change



➔ Ingredients production is the step of the pizza production which has the highest impact on Climate Change.

Variability in the impact of each of the 80 pizzas on Climate Change




➔ The impact of pizzas on Climate Change can be different from a pizza to another (up to more than 4,5 times between the less impacting pizza and the most impacting one).

➔ Families (except from the « bolognese » family), sectors (fresh/frozen) and retailers (data not shown for retailers) don't seem to define homogeneous groups of pizzas in terms of impact on climate change.

➔ Pizzas with the highest impact on Climate Change are those containing beef. Nevertheless this conclusion is very sensitive of the value used for beef in the Life Cycle Inventory and this value can be different between databases.

Results with mass-based FU


Link between impact on Climate Change and compositions and nutritional characteristics of the pizzas




➔ The impact of pizzas on Climate Change is positively impacted by the amount of proteins, fats and energy.

Results with nutritional FU


This chapter presents the evolution of the impact on climate change of 5 different pizzas (from the 80 pizzas) depending on the FU used. These 5 pizzas are shown with an illustrative purpose of the influence of the use of nutritional FUs on pizzas with different nutritional characteristics. The conclusions given here can be applied to the whole set of 80 pizzas.



➔ The use of a FU based on the recommended portion improves the impact on Climate Change of pizzas with the smallest recommended portions which often are the biggest pizzas (which are supposed to be shared) comparing to those with bigger recommended portions.




➔ The use of a FU based on energy improves the impact on Climate Change of the most energetic pizzas compared to those which contain less energy.




➔ The use of a FU based on proteins allows an improvement of the impact on Climate Change of the richest pizzas in proteins compared to those which contain less proteins.

Results with nutritional FU



➔ The use of a FU based on fibres doesn't really have an impact on the pizzas ranking in terms of impact on Climate Change.



➔ The use of a FU based on proteins and fibres combined globally has the same impact as using a FU based only on fibres because fibres are more limiting than proteins in the majority of pizzas.

Conclusion

Ingredients production is the step of pizzas production with the highest impact on Climate Change.

Impact on Climate Change can be very different between products coming from the same food category.

In the case of the 80 pizzas studied, pizzas with the highest impact are those containing beef. Nevertheless, this conclusion is very sensitive of the data chosen for beef.

Surprisingly, frozen pizzas don't seem to have a higher impact on Climate Change than fresh ones. This result may be explained by the fact that we only considered the electricity of the storage and because the use phase doesn't have a strong impact on Climate Change compared to the ingredient production.

Furthermore, the results can vary greatly depending on the FU used. In this case, the use of nutritional FU can help deciding which pizza is the most sustainable depending on what is important for the consumer. Nevertheless, even if the use of nutritional FU might be useful for quite homogeneous food categories in terms of composition, it is more delicate for heterogeneous food products categories such as pizzas.

For this reason, it would be interesting to evaluate the effect of those nutritional FU on a more homogeneous food category. It would also be interesting to evaluate the impact of the use of a more global nutritional FU on different food products belonging to the same category.

Acknowledgements

This work has been funded by INRA metaprogram DID'IT.

References

Farahani S.S., Soheilifard F., Nejad Raini M.G., Kokei D., 2019. Comparison of different tomato puree production phases from an environmental point of view. *The International Journal of Life Cycle Assessment*, 24, 1817–1827.

Kim D., Thoma G., Nutter D., Milani F., Ulrich R., Norris G., 2013. Life cycle assessment of cheese and whey production in the USA. *LCA for energy systems and food products*, 18, 1019-1035.

Tukker A., Huppes G., Geerken T., Nielsen P., 2006. Environmental Impact of Products (EIPRO): Analysis of the life cycle environmental impacts related to the final consumption of the EU-25. Technical Report Series.

Tsarouhas P., Achillas Ch., Aidonis D., Foinas D., Maslis V., 2015. Life Cycle Assessment of olive oil production in Greece. *Journal of Cleaner Production*, 93, 75-83.

Saarinen M., Fogelholm M., Tahoven, R., Kurppa S., 2017. Taking nutrition into account within the life cycle assessment of food products. *Journal of Cleaner Production* 148, 828-844.