



**HAL**  
open science

## **Bartonella gabonensis sp. nov., a new bartonella species from savannah rodent *Lophuromys* sp. in Franceville, Gabon**

J.B. Mangombi, N. N'Dilimabaka, H. Medkour, O.L. Banga, M.L. Tall, M. Ben Khedher, J. Terras, S. Abdi, Mathieu Bourgarel, E. Leroy, et al.

### ► To cite this version:

J.B. Mangombi, N. N'Dilimabaka, H. Medkour, O.L. Banga, M.L. Tall, et al.. Bartonella gabonensis sp. nov., a new bartonella species from savannah rodent Lophuromys sp. in Franceville, Gabon. New Microbes and New Infections, 2020, 38, pp.100796. 10.1016/j.nmni.2020.100796 . hal-03086548

**HAL Id: hal-03086548**

**<https://hal.inrae.fr/hal-03086548>**

Submitted on 22 Dec 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

# Characterization and antibiotic resistance pattern of diffusely adherent *Escherichia coli* (DAEC), isolated from paediatric diarrhoea in Shiraz, southern Iran

K. Javadi<sup>1)</sup>, S. Mohebi<sup>1)</sup>, M. Motamedifar<sup>1),2)</sup> and N. Hadi<sup>1),3)</sup>

1) Department of Bacteriology and Virology, School of Medicine, 2) Shiraz HIV/AIDS Research Centre, Institute of Health and 3) Bioinformatics and Computational Biology Research Centre, Shiraz University of Medical Sciences, Shiraz, Iran

## Abstract

Diarrhoea is a major health concern, especially in developing countries. Research has implicated diffusely adherent *Escherichia coli* (DAEC) strains as a cause of diarrhoea. In this study, we investigated the prevalence, adherence assay, virulence gene profiles and antimicrobial resistance of DAEC at a hospital in southern Iran. In this cross-sectional study, 309 infants and children under the age of 13 years with diarrhoea who had been referred to Shahid Dastgheib Hospital, Shiraz between October 2018 and May 2019 were recruited. Microbiological methods, PCR, HEp-2 adherence assay and antimicrobial susceptibility test were used. Of the 309 stool samples, 207 (66.9%) were found to contain *E. coli* by biochemical tests and culture. Molecular analysis of Afa/Dr and AIDA-I adhesin-encoding genes showed that 14 (6.7%) out of 207 *E. coli* isolates were DAEC. All DAEC isolates in HEp-2 cells showed a diffusely adherent pattern. The virulence genes *sat*, *pet*, *sigA*, *pic*, *astA* and *fimH* were found in 50%, 0%, 14.2%, 14.2%, 21.4% and 100% of DAEC isolates, respectively. The most effective antibiotic against the DAEC isolates was imipenem (92.8%) and the least effective was ampicillin (0%). Our findings expand the knowledge on DAEC prevalence and its characteristics in Iran. It also explains the role of virulence genes in DAEC pathogenesis. The results showed that although the prevalence of DAEC is low, these strains exhibit a high rate of antimicrobial resistance as well as high frequency for carrying virulence genes.

© 2020 Published by Elsevier Ltd.

**Keywords:** Adherence, antibiotic resistance, diarrhoea, diffusely adherent *Escherichia coli*, virulence genes

**Original Submission:** 19 August 2020; **Revised Submission:** 25 September 2020; **Accepted:** 5 October 2020

**Article published online:** 10 October 2020

**Corresponding author:** N. Hadi, Department of Bacteriology and Virology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran.

**E-mails:** [hadina@sums.ac.ir](mailto:hadina@sums.ac.ir), [nahalhadi@gmail.com](mailto:nahalhadi@gmail.com)

## Introduction

Diarrhoea is a major health problem, especially in developing countries, and one of the most important causes of mortality among children under the age of 5 [1]. The aetiological agents of diarrhoea include a wide range of viruses, bacteria and parasites. Among the bacterial pathogens, diarrhoeagenic *Escherichia coli* is an important cause of endemic and epidemic diarrhoea worldwide [2].

Diarrhoeagenic *E. coli* strains are classified into six main pathotypes based on their specific virulence characteristics, association with some serotypes and epidemiological characteristics: enteropathogenic *E. coli*, enterohaemorrhagic *E. coli*, enterotoxigenic *E. coli*, enteroaggregative *E. coli*, enteroinvasive *E. coli* and diffusely adherent *E. coli* (DAEC) [2].

DAEC strains are defined based on the presence of a diffuse adherence pattern (DA) on HeLa and HEp-2 epithelial cells. In the DA pattern, bacteria uniformly cover the cell surface [3].

According to the adhesin expression, two groups of DAEC strains have been identified, Afa/Dr DAEC and AIDA-I DAEC. Furthermore, these adhesins are also responsible for the DA phenotype [4]. Afa/Dr DAEC strains are associated with acute diarrhoea in children, especially in those 6 months and older, with persistent diarrhoea. Therefore, the DAEC pathogroup contains Afa/Dr adhesin-encoding genes and can cause

diarrhoea in patients [5]. The Afa/Dr family includes fimbrial and afimbrial adhesins, afimbrial adhesins Afa-I, Afa-II, Afa-III, Afa-V, Afa-VII, Afa-VIII, plus Dr-2 as well as Dr and F1845 fimbrial adhesins. Many of these adhesins have been identified in *E. coli* strains isolated from human urinary tract infections or diarrhoea, except Afa-VII, which was only found in *E. coli* isolated from bovine faeces [6]. F1845 adhesin was first identified in an *E. coli* strain (C1845) isolated from a child with chronic diarrhoea [6]. Until now, only the genes encoding for Afa-I, Afa-II, Afa-III and Afa-V have been identified in *E. coli* strains isolated from individuals with diarrhoea [7].

Afa-I and Afa-V adhesins only bind to the decay-accelerating factor, whereas AfaE-III, Dr and F1845 can each also bind to carcinoembryonic antigen-related cell adhesin molecules, and the Dr adhesin can also bind to type IV collagen [8–10].

*Enterobacteriaceae* have autotransporters called Serine protease autotransporters of *Enterobacteriaceae* (SPATEs). SPATEs are classified into two classes. Pet (plasmid-encoded toxin), Sat (secreted autotransporter toxin) and SigA are members of Class I SPATEs, which are cytotoxic to epithelial cells; Pic (protease involved in intestinal colonization) are members of Class II SPATEs, and are non-cytotoxic. The distribution of SPATEs among the diarrhoeagenic *E. coli* has been shown in many studies [11].

Type I pili are a type of composite surface fibre present in DAEC and are encoded by the *fim* gene cluster. FimH is the pilus adhesin [12]. Some DAEC strains through activation of Src and the mitogen-activated protein kinase, elicit a secondary interleukin-8 production by polymorphonuclear lymphocytes in a type-I pili-dependent manner [13].

Other virulence factor genes are recorded in DAEC strains, including *astA*, the gene for enteroaggregative heat-stable toxin I (EAST1), which was first identified in the enteroaggregative *E. coli*. EAST1 similarly stimulates the guanylate cyclase receptor to both enterotoxigenic *E. coli* St toxin and guanylin [14].

Enteropathogenic *E. coli* strains might show a DA pattern on HEp-2 and HeLa cells; therefore, we cannot use cell adhesins assay alone to detect Afa/Dr DAEC [15]. PCR is a simpler and faster method to identify Afa/Dr adhesins; consequently, PCR methods are more suitable to describe Afa/Dr DAEC strains [16,17]. Overall, these studies have clearly shown that it is important to determine the prevalence of DAEC selecting Afa or Daa target genes [18].

Although recent epidemiological studies indicated a high prevalence of DAEC strains isolated from diarrhoeal faeces, its pathogenicity has not been identified in adults [19]. It was found that the relative risk of diarrhoea associated with DAEC increases with the child's age from 18 months to 5 years [20]. The reason for such an age-related phenomenon and the mode of DAEC acquisition are yet to be determined [21].

The objectives of this study were to determine the prevalence of DAEC by PCR and HEp-2 cell adhesin assays among bacterial enteropathogens recovered from children with diarrhoea in Shahid Dastgheib Hospital, Shiraz, Iran, as well as to determine the presence of virulence genes in DAEC strains. We also determined the antimicrobial resistance profiles of the DAEC strains.

## Materials and methods

### Sample collection

This cross-sectional study included infants and children under the age of 13 years with diarrhoea who were referred to Shahid Dastgheib Hospital from October 2018 to May 2019. Patient information, such as age, gender, occult blood, pus cells and red blood cells, were obtained. Our exclusion criteria were age >13 years, no diarrhoea, insufficient data, and dry or suspected contaminated sample.

### Laboratory processing

The stool samples were immediately cultured on xylose lysine deoxycholate and MacConkey agar media in the laboratory of Shahid Dastgheib Hospital and then transferred to the laboratory of Bacteriology and Virology at Shiraz University of Medical Sciences for definitive diagnosis and identification. After incubation for 24 h at 37°C, lactose-positive colonies were re-cultured with standard biochemical methods to detect *E. coli*.

The tests included Gram staining, oxidase test, catalase test, motility test, triple-sugar iron fermentation, the citrate test, methyl red staining, the Voges–Proskauer test, indole test, ortho-nitrophenylgalactoside test, and acid production from carbohydrates.

*Escherichia coli* isolates were stored at –70°C in tryptic soy broth containing 20% glycerol (Merck Co., Darmstadt, Germany) for further characterization.

### DNA extraction

DNA was extracted using the boiling method. In this method, a single colony was grown on eosin methylene blue agar plates, mixed with 300 µL distilled water and boiled at 100°C for 10 minutes. After centrifugation at 13 000 g for 15 min, the supernatant containing the extracted DNA was transferred into a new sterile tube and stored at –20°C.

### Identification of DAEC isolates

All *E. coli* isolates were tested by PCR to detect six adherence genes: *afaE-1*, *afaE-2*, *afaE-3*, *afaE-5*, *daaE* and *aidalaah* for identification of DAEC isolates. The primers that were used to amplify these genes are listed in Table 1.

TABLE I. Primers used in PCR analysis

Gene	Primer sequences	Annealing temperature (°C)	Amplicon size (bp)	Reference
<i>afaE-1</i>	5'-CGAAAACGGCACTGACAAG-3' 5'-AGGCTTCCGTGAATACAACC-3'	58°C	230 pb	[26]
<i>afaE-2</i>	5'-TTAGACCGTACTGTTGTGTACC-3' 5'-TTTCCCAGTAGACTGGAATGAAGC-3'	60°C	375 pb	[26]
<i>afaE-3</i>	5'-TTAGACCGTACTGTTGTGTACC-3' 5'-ACCATTGTCGGTCGTCCAGGC-3'	58°C	408 pb	[26]
<i>afaE-5</i>	5'-TTAGACCGTACTGTTGTGTACC-3' 5'-AGCATCGGCGCGGTATACGGT-3'	61°C	429 pb	[26]
<i>daaE</i>	5'-TGACTGTGACCGAAGAGTGC-3' 5'-TTAGTTCGTCCAGTAACCCCC-3'	56°C	380 pb	[26]
<i>aida/aah</i>	5'-TCGATACCGAAACGCATACGCAGA-3' 5'-ACGCCGATCGGTGATGATGAAGAT-3'	52°C	204 pb	[29]
<i>sat</i>	5'-TCAGAAGCTCAGCGAATCATTG-3' 5'-CCATTATCACCAGTAAAACGCACC-3'	58°C	930 pb	[35]
<i>pet</i>	5'-GGCACAGAAT AAAGGGGTGTTT-3' 5'-CCTCTTGTTCACAGACATAC-3'	58°C	302 pb	[35]
<i>sigA</i>	5'-CCGACTTCTCACTTTCTCCCG-3' 5'-CCATCCAGCTGCATAGTGTGG-3'	58°C	430 pb	[35]
<i>pic</i>	5'-ACTGGATCTAAGGCTCAGGAT-3' 5'-GACTTAATGTCACTGTTCAAGC-3'	58°C	572 pb	[35]
<i>astA</i>	5'-ATGCCATCAACACAGTATAT-3' 5'-GCGAGTGACGGCTTTGTAGT-3'	58°C	110 pb	[38]
<i>fimH</i>	5'-TGCAGAACGGATAAGCCGTGG-3' 5'-GCAGTCACCTGCCCTCCGGTA-3'	61°C	508 pb	[39]

The PCR assay was performed in a total volume of 25 mL containing 0.5 mL of each primer (10 pM), 12.5 mL of DNA Polymerase Master Mix RED (Ampliqon Co., Inc., Odense, Denmark), 1 mL of DNA and 10.5 mL of water (DNase- and RNase-free water) in a T100™ thermal cycler (Bio-Rad, Hercules, CA, USA) under the following conditions: initial denaturation at 95°C for 5 min followed by 30 cycles of denaturation at 95°C for 1 min, annealing at 52°C–61°C for 1 min (Table I), and extension at 72°C for 1 min, and a single final extension at 72°C for 5 min. PCR products underwent electrophoresis in 1.5% agarose gels in the 0.5 Tris/EDTA/boric acid buffer and were visualized using ultraviolet light after staining with safe stain load dye (CinnaGen Co., Tehran, Iran).

### HEp-2 adherence assay

DAEC isolates were examined for HEp-2 adherence as described by Scaletsky et al. [22], with slight modifications. Briefly, monolayers of 10<sup>5</sup> HEp-2 cells were grown in Dulbecco's modified Eagle's medium (DMEM) with 10% fetal bovine serum in Leighton tubes containing a cover slip and incubated for 48 h at 37°C. DAEC strains were grown in 3 mL of trypticase soy broth for 16–18 h at 37°C. The tubes were then washed with phosphate-buffered saline and 1 mL of DMEM with 10% fetal bovine serum was added to each tube. The monolayers were infected with 40 µL of bacterial cultures added to 1 mL of DMEM and incubated for 3 h at 37°C. The infected monolayers were washed with sterile phosphate-buffered saline, fixed with methanol, stained with Giemsa stain and examined for DA pattern under a light microscope using a × 100 objective.

Standard strains of *E. coli* were used as the control to detect *daaE* and *aida/aah*, which were C1845 and 2787, respectively [18,23].

### Detection of virulence genes

All DAEC isolates were examined by PCR to detect six virulence genes: *sat*, *pet*, *sigA*, *pic*, *astA* and *fimH*. The primers used to amplify these genes are listed in Table I. The same conditions as for the adherence genes were employed for the detection and amplification of virulence genes expect that the annealing temperature was 56°C–58°C.

### Antimicrobial susceptibility testing

Antibiotic susceptibility testing of the DAEC isolates to ampicillin, ampicillin-sulbactam, cefotaxime, ceftriaxone, ceftazidime, imipenem, gentamicin, nalidixic acid, trimethoprim-sulfamethoxazole, moxifloxacin and chloramphenicol (Mast Group Ltd, Bootle, UK) was carried out on Müller–Hinton agar (Merck Co.) by disc diffusion method according to the Clinical and Laboratory Standards Institute. *Escherichia coli* ATCC25922 was used as the quality control strain.

### DNA sequence analysis

To verify the accuracy of amplified genes of *afaE-1*, *afaE-2*, *afaE-3*, *afaE-5*, the amplicons were submitted for sequencing (Bioneer Co., Munpyeongseoro, Daedeok-gu, Daejeon, South Korea) and the resulting sequences were analysed using online BLAST software (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>).

### Statistical analysis

SPSS software, version 26.0 (IBM Corp., Armonk, NJ, USA) was used for statistical analysis. The chi-square test and Fisher's exact test were performed for analysis of the data and  $p$  values  $< 0.05$  were considered to be statistically significant.

## Results

### Subject

A total of 309 stool samples from children under 13 years old with diarrhoea were included in our study. The samples were divided into three groups, 131 (42.3%) samples from children younger than 3 years; 98 (31.7%) from children aged 3–6 years and 80 (25.8%) from children aged 7–12 years. Male to female ratio was 52.7% to 47.3% (163 males versus 146 female).

### Prevalence of DAEC strains

*Escherichia coli* isolates were identified in 207 (66.9%) samples by biochemical tests and culture. Strains of *E. coli* harbouring the Afa/Dr and AIDA-I adhesin-encoding genes were found in the stool samples of 14 (6.7%) children with diarrhoea (Table 2); positive samples were from six boys and eight girls. Seven (50%) children with DAEC were under 3 years old, five (35.7%) and two (14.3%) were aged 3–6 years and 7–12 years, respectively (Table 3). Most DAEC samples contained occult blood, red blood cells and pus cells (Table 4).

### Adherence pattern analysis of DAEC strains

The analysis of the adherence patterns of the 14 isolates studied in HEp-2 cells showed that 100% (14 isolates) expressed the DA pattern (Fig. 1).

### Presence of virulence genes

All DAEC isolates were tested by PCR to detect the genes of the proposed DAEC virulence factors, including *sat*, *pet*, *sigA*, *pic*, *astA* and *fimH*. Seven DAEC strains were positive for *sat* (50%). The *astA* gene was identified in three (21.4%) DAEC strains and the *pic* and *sigA* genes were found in two (14.2%) of the 14 DAEC strains. All DAEC isolates were negative for the *pet* gene and all DAEC isolates were positive for the *fimH* gene (Table 5). Several different combinations of the virulence markers were found among the DAEC isolates (Table 6).

### Antimicrobial resistance among DAEC isolates

The results of antimicrobial susceptibility testing of the diarrhoeagenic DAEC isolates with 11 antibiotics by disc diffusion method are shown in Table 7. The most effective antibiotic against DAEC was imipenem and the least effective was ampicillin.

**TABLE 2.** Characterization of the adhesin of diffusely adherent *Escherichia coli* (DAEC) isolates from children with diarrhoea

Isolate	<i>afaE-1</i>	<i>afaE-2</i>	<i>afaE-3</i>	<i>afaE-5</i>	<i>daaE</i>	<i>aida/laah</i>	HEp-2 adherence
DAEC1	+	—	—	+	—	—	DA
DAEC2	+	—	+	—	—	—	DA
DAEC3	+	—	—	—	+	—	DA
DAEC4	+	—	—	—	—	—	DA
DAEC5	+	—	—	—	—	—	DA
DAEC6	—	+	+	—	—	—	DA
DAEC7	+	—	—	—	+	—	DA
DAEC8	+	—	—	—	—	—	DA
DAEC9	—	—	+	—	—	—	DA
DAEC10	—	+	—	—	—	—	DA
DAEC11	—	—	—	+	—	—	DA
DAEC12	+	—	—	—	—	—	DA
DAEC13	+	—	—	+	—	—	DA
DAEC14	—	—	+	—	+	—	DA
<i>n</i> (%)	9 (64.3)	2 (14.2)	4 (28.6)	3 (21.5)	3 (21.5)	0 (0)	

DA, diffuse adherence.

**TABLE 3.** Age and gender of 207 *Escherichia coli* isolates and positive diffusely adherent *E. coli* (DAEC) in different age groups




Age (years)	No. of samples	Males, <i>n</i> (%)	Females, <i>n</i> (%)	DAEC, <i>n</i> (%)
<3	113	58 (53.7)	55 (55.5)	7 (6.2)
3–6	56	29 (26.8)	27 (27.3)	5 (8.9)
7–12	38	21 (19.4)	17 (17.2)	2 (5.2)
Total	207	108 (52.7)	99 (47.3)	14 (6.7)

**TABLE 4.** Stool analysis findings of *Escherichia coli* isolates and positive diffusely adherent *E. coli* (DAEC) isolates from children with diarrhoea

Stool characteristic	All <i>E. coli</i> ( <i>n</i> = 207), <i>n</i> (%)	DAEC ( <i>n</i> = 14), <i>n</i> (%)
Occult blood	149 (72)	10 (71.4)
Red blood cells	118 (57)	7 (50)
Pus cells	145 (70)	8 (57.1)

## Discussion

The DAEC strains are a heterogeneous group of *E. coli* strains that generate a diffuse adherence pattern on HeLa and HEp-2 cells [20]. Food or water polluted with human or animal faeces is the main transmission route for the DAEC pathotype [24]. DAEC strains destroy the intestinal epithelium by binding to proteins that accelerate degradation. They account for a large proportion of diarrhoeal cases observed in inpatients who have no other identified enteropathogen [19]. Recent epidemiological studies have implicated DAEC strains as a cause of diarrhoea in children in developing countries [25].



**FIG. 1.** Microscopic appearance of adherence patterns on HEp2 cells of diffusely adherent *Escherichia coli* (DAEC) isolates from children with diarrhoea: (a) DAEC C1845, (b) *E. coli* K12, (c) DAEC isolated from children with diarrhoea.

**TABLE 5.** Incidence of virulence genes among diffusely adherent *Escherichia coli* (DAEC) isolates from children with diarrhoea

	Sat	pet	sigA	pic	astA	fimH
DAEC	7 (50%)	0 (0)	2 (14.2%)	2 (14.2%)	3 (21.4%)	14 (100%)

**TABLE 6.** The prevalence of different virulence genes among diffusely adherent *Escherichia coli* (DAEC) isolates from children with diarrhoea

Genetic profile	No. (%) of DAEC
<i>fimH</i>	6 (42.8)
<i>fimH, sat</i>	2 (14.2)
<i>fimH, sigA</i>	1 (7.1)
<i>fimH, sat, pic</i>	1 (7.1)
<i>fimH, sat, astA</i>	3 (21.5)
<i>fimH, sat, sigA, pic</i>	1 (7.1)

**TABLE 7.** Antimicrobial susceptibility of diffusely adherent *Escherichia coli* isolates from children with diarrhoea

Antimicrobial agent	Resistant, n (%)	Susceptible, n (%)	Intermediate, n (%)
Ampicillin	14 (100)	0 (0)	0 (0)
Ampicillin-sulbactam	10 (71.4)	3 (21.5)	1 (7.1)
Cefotaxime	11 (78.6)	2 (14.3)	1 (7.1)
Ceftriaxone	7 (50)	4 (28.6)	3 (21.5)
Ceftazidime	8 (57.1)	5 (35.7)	1 (7.1)
Imipenem	0 (0)	13 (92.8)	1 (7.1)
Gentamicin	2 (14.3)	12 (85.7)	0 (0)
Nalidixic acid	7 (50)	4 (28.6)	3 (21.5)
Trimethoprim-Sulfamethoxazole	11 (78.6)	2 (14.3)	1 (7.1)
Moxifloxacin	5 (35.7)	6 (42.9)	3 (21.5)
Chloramphenicol	5 (35.7)	7 (50)	2 (14.3)

The proportion of DAEC among *E. coli* in our study was 6.7%, which is comparable to previous reports in Shiraz (7.9%) [25]. DAEC occurrence in our study was higher than in northwest Mexico (6.2%) [26] and in South American countries, such as Peru (4%) and Colombia (1.2%) [27,28]. In contrast, the identified rate of DAEC was 8.18% in China [29], 15.7% in Brazil [30] and 35% in Mexico [14], which is higher than the rate found in this study.

We investigated all sample data to find any association between these data and DAEC positivity. Accordingly, we classified our data based on age, gender and stool analysis findings, such as occult blood, red blood cells and pus cells. There was no significant association between these data and DAEC, which shows that all diarrhoeal stools should be examined for DAEC isolates.

The Afa/Dr adhesin-encoding genes *afaE-1*, *afaE-2*, *afaE-3*, *afaE-5* and *daaE* were identified in 64.3%, 14.2%, 28.6%, 21.5%

and 21.5% of DAEC isolates, respectively (Table 4). Mansan-Almeida et al. detected *afaE-1*, *afaE-2*, *afaE-3*, *afaE-5* and *daaE* in, respectively, 44%, 10%, 2%, 2% and 6% of DAEC isolated from children with diarrhoea [26]. Therefore, the most frequent gene among Afa/Dr DAEC strains was *afaE-1* in our study.

AIDA-1 is the *E. coli* adhesin involved in diffuse adherence [31]. In the current study, *aida/aah* was not detected in any DAEC isolates, which is in line with the Abbasi et al. study [24]. These results indicate that the prevalence of *aida/aah* was low in DAEC isolates.

The PCR results for the Afa/Dr genes showed significant relation with the HEp-2 cell adhesin assay, as 100% of Afa/Dr-positive isolates were confirmed as DAEC. This is the first report on the adhesin assay of DAEC in Iran.

The *sat* gene was detected in seven (50%) DAEC isolates in this study. Guignot et al. showed that *sat* might play a role in Afa/Dr DAEC intestinal pathogenesis by inducing lesions at the junctional barrier in Caco-2/TC7 monolayers [32]. Spano et al. found *sat* in 7.1% of DAEC isolated from children [33], Li et al. reported that 44.44% of DAEC was positive for *sat* [27] and Mansan-Almeida et al. found *sat* in 46% of DAEC isolated from children [26]. The rate of *sat* genes in our study was higher than those reported by Li et al. [27] and Mansan-Almeida et al. [26], and much higher than the percentage reported by Spano et al. [33]. Therefore, it seems that *sat* is common in the pathogenesis of DAEC.

In the present study, *pet* was not identified in any DAEC isolates. *Pet* generates diarrhoeagenic effects by changes in the actin cytoskeleton [34]. Spano et al. reported that 14.2% of DAEC isolated from children was positive for *pet* [33] but other studies did not detect the *pet* gene in any DAEC isolates [14,27,35].

A total of 14.2% of the DAEC isolates harboured *sigA* and *pic* genes in this study. *SigA* is a cytotoxin that was found to have significant cytotoxic effects on HEp-2 cells [36]. *Pic* is the protein involved in intestinal colonization in children with diarrhoea, and unlike *sigA* it does not have a cytotoxic effect on HEp-2 cells [37]. Spano et al. identified *pic* in 2.3% of DAEC isolated from children [33]. Boisen et al. detected *pic* in 10% of DAEC strains, whereas *sigA* was not detected in any DAEC [38]. Our results showed that the existence of *pic* and *sigA* in DAEC isolates was higher than in other studies.

The *astA* gene encodes the EAST-1 toxin with a positive rate of 21.4% in DAEC isolates in this study. EAST-1 toxin is thought to be a supplementary determinant in the pathogenesis of *E. coli* diarrhoea [39]. Spano et al. detected *astA* in 9.5% of DAEC isolated from children [33]. Lopes et al. reported that 15.2% of DAEC was positive for *astA* [35], and Patzi-Vargas et al. identified *astA* in 3.1% of DAEC [14]. These findings are contrary to

our findings, which showed the occurrence of *astA* in DAEC isolates to be much higher.

In this study, the type I fimbriae-encoding gene *fimH* was identified in all DAEC isolates. Li et al. found *fimH* in 100% of DAEC strains [27], and Lopes et al. detected the *fimH* gene in 48.2% of DAEC strains [35]. Johnson and Stell identified this adhesin in nearly all *E. coli* strains [40].

In the present study, we observed high percentages of resistance to ampicillin, cefotaxime and cotrimoxazole; only imipenem and gentamicin were effective against 92.8% and 85.7% of DAEC isolates, respectively. Moreover, resistance to more than one antibiotic was found in 100% of DAEC strains. Different studies have reported that most DAEC strains were resistant to several antibiotics, such as sulfonamide, doxycycline, tetracycline, ampicillin, cefotaxime and cotrimoxazole, whereas the isolates were sensitive to antibiotics such as imipenem, amikacin, gentamicin and nitrofurantoin [25,27,28,33].

---

## Conclusions

Our findings expand our knowledge of DAEC prevalence and characteristics in Iran, and explain the role of virulence genes in DAEC pathogenesis. Although the prevalence of DAEC was low, these strains exhibited high rates of antimicrobial resistance and high frequency for carrying virulence genes. Furthermore, preventing infections caused by this bacterium among children is essential and further researches are warranted, which should include other cities in Iran, using larger number of DAEC isolates.

---

## Conflict of interest

The authors declare that there is no conflict of interest in relation to this article.

---

## Authors' contribution

All authors participated in the research design and contributed to different parts of the research.

---

## Funding

This paper was part of the MSc thesis of K. Javadi supervised by N. Hadi and was supported by Shiraz University of Medical Sciences grant No. 97–18400.

## Ethical approval

The study design was approved by the Ethics Committee of Shiraz University of Medical Sciences IR.SUMS.REC.1398.613 and followed the statements of the Declaration of Helsinki.

## Acknowledgements

The authors wish to thank the staff of the microbiology laboratories of the hospitals for their kind collaboration.

## References

- [1] Bern C, Martinez J, De Zoysa I, Glass R. The magnitude of the global problem of diarrhoeal disease: a ten-year update. *Bull World Health Organ* 1992;70:705.
- [2] Mansour Amin MS, Javaherizadeh H, Motamedifar M, Saki M, Veisi H, Ebrahimi S, et al. Antibiotic resistance pattern and molecular characterization of extended-spectrum  $\beta$ -lactamase producing enteroaggregative *Escherichia coli* isolates in children from southwest Iran. *Infect Drug Resist* 2018;11:1097.
- [3] Scaletsky IC, Fabbriotti SH, Carvalho RL, Nunes CR, Maranhao HS, Morais MB, et al. Diffusely adherent *Escherichia coli* as a cause of acute diarrhea in young children in Northeast Brazil: a case-control study. *J Clin Microbiol* 2002;40:645–8.
- [4] Scaletsky IC, Fabbriotti SH, Silva SO, Morais MB, Fagundes-Neto U. HEp-2–adherent *Escherichia coli* strains associated with acute infantile diarrhea, São Paulo, Brazil. *Emerg Infect Dis* 2002;8:855.
- [5] Lozer DM, Souza TB, Monfardini MV, Vicentini F, Kitagawa SS, Scaletsky IC, et al. Genotypic and phenotypic analysis of diarrheagenic *Escherichia coli* strains isolated from Brazilian children living in low socioeconomic level communities. *BMC Infect Dis* 2013;13:418.
- [6] Lalioui L, Jouve M, Gounon P, Le Bouguenec C. Molecular cloning and characterization of the *afa-7* and *afa-8* gene clusters encoding afimbrial adhesins in *Escherichia coli* strains associated with diarrhea or septicemia in calves. *Infect Immun* 1999;67:5048–59.
- [7] Servin AL. Pathogenesis of human diffusely adhering *Escherichia coli* expressing Afa/Dr adhesins (Afa/Dr DAEC): current insights and future challenges. *Clin Microbiol Rev* 2014;27:823–69.
- [8] Guignot J, Peiffer I, Bernet-Camard M-F, Lublin DM, Carnoy C, Moseley SL, et al. Recruitment of CD55 and CD66e brush border-associated glycosylphosphatidylinositol-anchored proteins by members of the Afa/Dr diffusely adhering family of *Escherichia coli* that infect the human polarized intestinal Caco-2/TC7 cells. *Infect Immun* 2000;68:3554–63.
- [9] Berger CN, Billker O, Meyer TF, Servin AL, Kansau I. Differential recognition of members of the carcinoembryonic antigen family by Afa/Dr adhesins of diffusely adhering *Escherichia coli* (Afa/Dr DAEC). *Mol Microbiol* 2004;52:963–83.
- [10] Nowicki B, Moulds J, Hull R, Hull S. A hemagglutinin of uropathogenic *Escherichia coli* recognizes the Dr blood group antigen. *Infect Immun* 1988;56:1057–60.
- [11] Tapader R, Chatterjee S, Singh A, Dayma P, Haldar S, Pal A, et al. The high prevalence of serine protease autotransporters of Enterobacteriaceae (SPATEs) in *Escherichia coli* causing neonatal septicemia. *Eur J Clin Microbiol Infect Dis* 2014;33:2015–24.
- [12] Pizarro-Cerdá J, Cossart P. Bacterial adhesin and entry into host cells. *Cell* 2006;124:715–27.
- [13] Sémiramoth N, Gleizes A, Turbica I, Sandré C, Gorges R, Kansau I, et al. *Escherichia coli* type I pili trigger late IL-8 production by neutrophil-like differentiated PLB-985 cells through a Src family kinase- and MAPK-dependent mechanism. *J Leukoc Biol* 2009;85:310–21.
- [14] Patzi-Vargas S, Zaidi MB, Perez-Martinez I, León-Cen M, Michel-Ayala A, Chaussabel D, et al. Diarrheagenic *Escherichia coli* carrying supplementary virulence genes are an important cause of moderate to severe diarrhoeal disease in Mexico. *PLoS Negl Trop Dis* 2015;9:e0003510.
- [15] Labigne-Roussel AF, Lark D, Schoolnik G, Falkow S. Cloning and expression of an afimbrial adhesin (AFA-I) responsible for P blood group-independent, mannose-resistant hemagglutination from a pyelonephritic *Escherichia coli* strain. *Infect Immun* 1984;46:251–9.
- [16] Le Bouguenec C, Archambaud M, Labigne A. Rapid and specific detection of the *pap*, *afa*, and *sfa* adhesin-encoding operons in uropathogenic *Escherichia coli* strains by polymerase chain reaction. *J Clin Microbiol* 1992;30:1189–93.
- [17] Le Bouguenec C, Lalioui L, du Merle L, Jouve M, Courcoux P, Bouzari S, et al. Characterization of AfaE adhesins produced by extraintestinal and intestinal human *Escherichia coli* isolates: PCR assays for detection of *afa* adhesins that do or do not recognize Dr blood group Antigens. *J Clin Microbiol* 2001;35:1738–45.
- [18] Bilge S, Clausen C, Lau W, Moseley S. Molecular characterization of a fimbrial adhesin, F1845, mediating diffuse adherence of diarrhea-associated *Escherichia coli* to HEp-2 cells. *J Bacteriol* 1989;171:4281–9.
- [19] Jallat C, Livrelli V, Darfeuille-Michaud A, Rich C, Joly B. *Escherichia coli* strains involved in diarrhea in France: high prevalence and heterogeneity of diffusely adhering strains. *J Clin Microbiol* 1993;31:2031–7.
- [20] Servin AL. Pathogenesis of Afa/Dr diffusely adhering *Escherichia coli*. *Clin Microbiol Rev* 2005;18:264–92.
- [21] Nataro JP, Kaper JB. Diarrheagenic *Escherichia coli*. *Clin Microbiol Rev* 1998;11:142–201.
- [22] Scaletsky I, Silva M, Trabulsi LR. Distinctive patterns of adherence of enteropathogenic *Escherichia coli* to HeLa cells. *Infect Immun* 1984;45:534–6.
- [23] Benz I, Schmidt MA. Cloning and expression of an adhesin (AIDA-I) involved in diffuse adherence of enteropathogenic *Escherichia coli*. *Infect Immun* 1989;57:1506–11.
- [24] Abbasi P, Kargar M, Doosti A, Mardaneh J, Ghorbani-Dalini S, Dehyadegari MA. Molecular detection of diffusely adherent *Escherichia coli* strains associated with diarrhea in Shiraz, Iran. *Arch Pediatr Infect Dis* 2016;5.
- [25] Levine MM, Ferreccio C, Prado V, Cayazzo M, Abrego P, Martinez J, et al. Epidemiologic studies of *Escherichia coli* diarrheal infections in a low socioeconomic level peri-urban community in Santiago, Chile. *Am J Epidemiol* 1993;138:849–69.
- [26] Canizalez-Roman A, Flores-Villaseñor HM, Gonzalez-Nuñez E, Velazquez-Roman J, Vidal JE, Muro-Amador S, et al. Surveillance of diarrheagenic *Escherichia coli* strains isolated from diarrhea cases from children, adults and elderly at Northwest of Mexico. *Front Microbiol* 2016;7:1924.
- [27] Gómez-Duarte OG, Arzuza O, Urbina D, Bai J, Guerra J, Montes O, et al. Detection of *Escherichia coli* enteropathogens by multiplex polymerase chain reaction from children's diarrheal stools in two Caribbean–Colombian cities. *Foodborne Pathog Dis* 2010;7:199–206.
- [28] Ochoa TJ, Ruiz J, Molina M, Del Valle LJ, Vargas M, Gil AI, et al. High frequency of antimicrobial drug resistance of diarrheagenic *Escherichia coli* in infants in Peru. *Am J Trop Med Hyg* 2009;81:296–301.

- [29] Li D, Shen M, Xu Y, Liu C, Wang W, Wu J, et al. Virulence gene profiles and molecular genetic characteristics of diarrheagenic *Escherichia coli* from a hospital in western China. *Gut Pathog* 2018;10:35.
- [30] Mansan-Almeida R, Pereira AL, Giugliano LG. Diffusely adherent *Escherichia coli* strains isolated from children and adults constitute two different populations. *BMC Microbiol* 2013;13:22.
- [31] Charbonneau M-È, Berthiaume F, Mourez M. Proteolytic processing is not essential for multiple functions of the *Escherichia coli* auto-transporter adhesin involved in diffuse adherence (AIDA-I). *J Bacteriol* 2006;188:8504–12.
- [32] Guignot J, Chaplais C, Coconnier-Polter MH, Servin AL. The secreted auto-transporter toxin, Sat, functions as a virulence factor in Afa/Dr diffusely adhering *Escherichia coli* by promoting lesions in tight junction of polarized epithelial cells. *Cell Microbiol* 2007;9:204–21.
- [33] Spano LC, da Cunha KF, Monfardini MV, Fonseca RdCB, Scaletsky ICA. High prevalence of diarrheagenic *Escherichia coli* carrying toxin-encoding genes isolated from children and adults in southeastern Brazil. *BMC Infect Dis* 2017;17:773.
- [34] Navarro-Garcia F, Sonnested M, Teter K. Host-toxin interactions involving EspC and Pet, two serine protease auto-transporters of the Enterobacteriaceae. *Toxins (Basel)* 2010;2:1134–47.
- [35] Lopes LM, Fabbriotti SH, Ferreira AJ, Kato MA, Michalski J, Scaletsky IC. Heterogeneity among strains of diffusely adherent *Escherichia coli* isolated in Brazil. *J Clin Microbiol* 2005;43:1968–72.
- [36] Ruiz-Perez F, Nataro JP. Bacterial serine proteases secreted by the auto-transporter pathway: classification, specificity, and role in virulence. *Cell Mol Life Sci* 2014;71:745–70.
- [37] Navarro-Garcia F, Gutierrez-Jimenez J, Garcia-Tovar C, Castro LA, Salazar-Gonzalez H, Cordova V. Pic, an auto-transporter protein secreted by different pathogens in the Enterobacteriaceae family, is a potent mucus secretagogue. *Infect Immun* 2010;78:4101–9.
- [38] Boisen N, Ruiz-Perez F, Scheutz F, Krogfelt KA, Nataro JP. High prevalence of serine protease auto-transporter cytotoxins among strains of enteroaggregative *Escherichia coli*. *Am J Trop Med Hyg* 2009;80:294–301.
- [39] Yuste M, De La Fuente R, Ruiz-Santa-Quiteria J, Cid D, Orden J. Detection of the astA (EAST1) gene in attaching and effacing *Escherichia coli* from ruminants. *J Vet Med B Infect Dis Vet Public Health* 2006;53:75–7.
- [40] Johnson JR, Stell AL. Extended virulence genotypes of *Escherichia coli* strains from patients with urosepsis in relation to phylogeny and host compromise. *J Infect Dis* 2000;181:261–72.