



# Special features of neuroendocrine interactions between stress and reproduction in teleosts

Karine Rousseau, Patrick Prunet, Sylvie Dufour

## ► To cite this version:

Karine Rousseau, Patrick Prunet, Sylvie Dufour. Special features of neuroendocrine interactions between stress and reproduction in teleosts. *General and Comparative Endocrinology*, 2021, 300, pp.1-23. 10.1016/j.ygcen.2020.113634 . hal-03109982

**HAL Id: hal-03109982**

**<https://hal.inrae.fr/hal-03109982>**

Submitted on 7 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

**Special features of neuroendocrine interactions  
between stress and reproduction in teleosts**

**Rousseau Karine<sup>a</sup>, Prunet Patrick<sup>b</sup> and Dufour Sylvie<sup>a\*</sup>**

<sup>a</sup> Muséum National d'Histoire Naturelle, Research Unit BOREA, Biology of Aquatic  
Organisms and Ecosystems, CNRS, IRD, SU, UCN, UA, Paris, France

<sup>b</sup> INRAE, UR1037, Laboratoire de Physiologie et de Génomique des Poissons (LPGP),  
Rennes, France

\*Corresponding author: [sylvie.dufour@mnhn.fr](mailto:sylvie.dufour@mnhn.fr)

**Keywords:** stress; reproduction; teleosts; gonadotropic axis; corticotropic axis; evolution.

## Introduction

### 1. Various interactions between stress, cortisol and reproduction in teleosts

- 1.1. Negative interactions between stress and reproduction
- 1.2. Positive interactions between cortisol and reproduction related to peculiar life cycles
  - 1.2.1. Cortisol and environmentally-related spawning activity in some teleosts
  - 1.2.2. Cortisol and upstream reproductive migration in salmonids
  - 1.2.3. Cortisol and oceanic reproductive migration in eels
- 1.3. Cortisol and gonadal sex differentiation
- 1.4. Animal welfare, environment, stress and reproduction

### 2. Overview of neuroendocrine stress and reproductive axes in teleosts

- 2.1. Teleost specific traits compared to other vertebrates
  - 2.1.1. Anatomic-functional specificities
    - 2.1.1.1. Pituitary regionalisation and direct innervation
    - 2.1.1.2. Caudal neurosecretory system
  - 2.1.2. Teleost specific whole genome duplication
- 2.2. The corticotrophic axis
  - 2.2.1. Main actors of the corticotrophic axis in vertebrates
  - 2.2.2. Specific features of the corticotrophic axis in teleosts
    - 2.2.2.1. CRH and related peptides and their receptors
    - 2.2.2.2. Dopamine and other neurohormones
    - 2.2.2.3. POMC-derived peptides and their receptors
    - 2.2.2.4. Corticosteroids and their receptors
- 2.3. The gonadotrophic axis
  - 2.3.1. Main actors of the gonadotrophic axis in vertebrates
  - 2.3.2. Specific features of the gonadotrophic axis in teleosts
    - 2.3.2.1. GnRHs and their receptors
    - 2.3.2.2. Dopamine and other neurohormones
    - 2.3.2.3. Gonadotropins and their receptors
    - 2.3.2.4. Sex steroids and their receptors

### 3. Neuroendocrine mechanisms of the interactions between corticotrophic and gonadotrophic axes in teleosts

- 3.1. Regulatory effects of actors of the corticotrophic axis on the gonadotrophic axis
  - 3.1.1. Roles of corticosteroids and their receptors
    - 3.1.1.1. Expression and role along the gonadotrophic axis
    - 3.1.1.2. *In vivo* effects of corticosteroids
    - 3.1.1.3. *In vitro* effects of corticosteroids
  - 3.1.2. Roles POMC-derived peptides
  - 3.1.3. Roles of CRH
- 3.2. Regulatory effects of actors of the gonadotrophic axis on the corticotrophic axis
  - 3.2.1. Roles of sex steroids
  - 3.2.2. Roles of other actors

## Conclusions

## Abstract

Stress and reproduction are both essential functions for vertebrate survival, ensuring on one side adaptative responses to environmental changes and potential life threats, and on the other side production of progeny. With more than 25,000 species, teleosts constitute the largest group of extant vertebrates, and exhibit a large diversity of life cycles, environmental conditions and regulatory processes. Interactions between stress and reproduction are a growing concern both for conservation of fish biodiversity in the frame of global changes and for the development of sustainability of aquaculture including fish welfare. In teleosts, as in other vertebrates, adverse effects of stress on reproduction have been largely documented and will be shortly overviewed. Unexpectedly, stress notably *via* cortisol, may also facilitate reproductive function in some teleost species in relation to their peculiar life cycles and this review will provide some examples. Our review will then mainly address the neuroendocrine axes involved in the control of stress and reproduction, namely the corticotropic and gonadotropic axes, as well as their interactions. After reporting some anatomo-functional specificities of the neuroendocrine systems in teleosts, we will describe the major actors of the corticotropic and gonadotropic axes at the brain-pituitary-peripheral glands (interrenals and gonads) levels, with a special focus on the impact of teleost-specific whole genome duplication (3R) on the number of paralogs and their potential differential functions. We will finally review the current knowledge on the neuroendocrine mechanisms of the various interactions between stress and reproduction at different levels of the two axes in teleosts in a comparative and evolutionary perspective.

## Introduction

The first definition of stress by Selye in 1973 as "... the non-specific response of the body to any demand made upon it", refers to a general adaptation syndrome as a response to a stressor, which is similar irrespective of its nature (Selye, 1973). Since then, various definitions have appeared in the literature but the most complete one proposed by Schreck defines stress as "the physiological cascade of events that occurs when an organism is attempting to resist death or re-establish homeostatic norms in face of an insult" (Schreck, 2000). In response to a stressful situation and the recognition of a threat by the central nervous system, an adaptive compensatory non-specific response takes place with an initial adrenergic response and a subsequent synthesis and secretion of cortisol due to activation of the corticotropic axis, hypothalamus-pituitary-adrenal (HPA) in mammals, birds and reptiles, and hypothalamus-pituitary-interrenal gland/head kidney (HPI) in amphibians and teleosts. Activation of these endocrine pathways constitutes the primary responses, which lead to make energy available for systems involved in the stress responses. Thus, activation of the cardiovascular and respiratory responses accompanied by osmoregulatory disfunctions favour oxygen and energy substrates distribution and constitute the secondary responses. Tertiary responses refer to aspects of whole-animal performance and generally are maladaptive; they include changes in growth, swimming capacity, and modified behavioral patterns (feeding, aggression) [for reviews: (Barton, 2002; Gorissen and Flik, 2016; Schreck and Tort, 2016; Wendelaar Bonga, 1997)].

Stress is, in most cases, reported to affect reproduction in a deleterious way, but may also induce positive outcomes and a substantial body of research has been dedicated to decipher the mechanisms underlying the complex relationships between stress and reproduction [for reviews: (Fuzzen et al., 2011; Leatherland et al., 2010; Milla et al., 2009; Pankhurst, 2016; Schreck, 2010)]. Since the first paper by Selye in 1939 (Selye, 1939) which proposed that

activation of the hypothalamo-pituitary-adrenal (HPA) axis inhibits the hypothalamo-pituitary-gonads (HPG) axis, this paradigm has been established mainly in mammalian species (Chand and Lovejoy, 2011) but also in fish for which numerous reports emphasizing effects of corticosteroids on reproduction have been produced (Fuzzen et al., 2011; Leatherland et al., 2010; Milla et al., 2009; Pankhurst, 2016; Schreck, 2010) . In this paper, we will focus our interest on the relationships between stress and reproductive neuroendocrine axes, trying to decipher direct stress effects and separate them from systemic effects on other biological functions. Thanks to the recent progress brought by genomic studies, we will be able to integrate the most recent informations provided by molecular phylogeny analyses and describe effects of new actors, which appeared *via* different rounds of whole genome duplications. This review will focus on teleost fish but with an evolutionary perspective requiring comparison with other vertebrate species.

## **1. Various interactions between stress, cortisol and reproduction in teleosts**

Many relationships between stress and reproduction, with the influence of environmental conditions, have been documented in natural situation and in aquaculture. In this context, we **first** consider in this **section** the effects of stressors on reproduction, which are frequently associated with high cortisol levels.

### **1.1. Negative interactions between stress and reproduction**

Early works in many teleost species reported the deleterious effects of stress and stress-related increase of cortisol on gonadotropic axis and reproductive performance. Chronically-induced stress such as by confinement or captivity and acutely-induced stress such as by predators or by aquaculture practices (handling, frequent netting, tank draining, crowding, noise) **result** in reduced plasma androgen and oestrogen levels [brown trout *Salmo trutta*: (Pickering et al., 1987); wild spotted seatrout *Cynoscion nebulosus*: (Safford and Thomas, 1987); rainbow trout

143 *Oncorhynchus mykiss*: (Pankhurst and Dedual, 1994); red gurnard *Chelidonichthys kumu*:  
 144 (Clearwater and Pankhurst, 1997); striped trumpeter *Latris lineata*: (Morehead, 1998); roach  
 145 *Rutilus rutilus*: (Pottinger et al., 1999); black bream *Acanthopagrus butcheri*: (Haddy and  
 146 Pankhurst, 1999); sockeye salmon *Oncorhynchus nerka*: (Kubokawa et al., 1999); snapper  
 147 *Pagrus auratus*: (Carragher and Pankhurst, 1991; Cleary et al., 2000); spiny damselfish  
 148 *Acanthochromis polyacanthus*: (Pankhurst, 2001); striped bass *Morone saxatilis*: (Castranova  
 149 et al., 2005); jundia *Rhamdia quelen*: (Soso et al., 2008); tilapia *Oreochromis mossambicus*:  
 150 (Chabbi and Ganesh, 2012)]. Stress is also associated with reduced plasma vitellogenin (Vg)  
 151 levels [brown and rainbow trout: (Campbell et al., 1994)]. It also **decreases** plasma luteinizing  
 152 hormone (LH) levels [white sucker *Catostomus commersoni*: (Van Der Kraak et al., 1992)],  
 153 **suppresses** LH secreting cell activity [tilapia: (Chabbi and Ganesh, 2012)], as well as  
 154 **decreases** hypothalamic gonadotropin-releasing hormone 1 (GnRH1) mRNA levels [jack  
 155 mackerel *Trachurus japonicus*: (Imanaga et al., 2014)]. After applied stress, impacts on final  
 156 stages of reproduction **are** also observed such as delayed ovulation [rainbow trout: (Campbell  
 157 et al., 1992; Contreras-Sanchez et al., 1998)], reduced egg size in females [brown and  
 158 rainbow trout: (Campbell et al., 1994, 1992) ; *Neolamprologus pulcher*: (Mileva et al., 2011)]  
 159 and reduced sperm counts in males [brown and rainbow trout: (Campbell et al., 1994)],  
 160 increased gonadal atresia [(pike *Esox lucius*: (De Montalembert et al., 1978); red gurnard:  
 161 (Clearwater and Pankhurst, 1997); snapper: (Cleary et al., 2000); Atlantic bluefin tuna  
 162 *Thunnus thynnus*: (Corriero et al., 2011)], and reduced fecundity [rainbow trout : (Contreras-  
 163 Sanchez et al., 1998)]. Furthermore, impacts on progeny also **occur** such as reduced length  
 164 [tropical damselfish *Pomacentrus amboinensis*: (McCormick, 2009, 2006, 1999, 1998)],  
 165 decreased survival [brown and rainbow trout: (Campbell et al., 1994, 1992)] or occurrence of  
 166 abnormalities [Atlantic cod *Gadus morhua*: (Morgan et al., 1999)] [for reviews: (Billard,  
 167 1981; Pankhurst and Van der Kraak, 1997; Pankhurst, 2016; Pickering, 1989)].

Socially mediated stress response and downregulated reproductive function are observed in some teleosts, such as in cichlids. For example, subordinate (non-territorial) males of *Astatotilapia (Haplochromis) burtoni* display high expression of hypothalamic and pituitary corticotropin-releasing hormone (CRH) receptor as well as high plasma cortisol levels, but low mRNA levels of hypothalamic GnRH1 and of pituitary GnRH receptor, LH and FSH, as well as low plasma levels of LH and FSH, and androgens, compared to dominant (territorial) males [for review: (Maruska, 2014)].

## **1.2. Positive interactions between cortisol and reproduction related to peculiar life cycles**

### **1.2.1. Cortisol and environmentally-related spawning activity in some teleosts**

The catfish, *Heteropneustes fossilis*, awaits the onset of the monsoon rainfall for spawning in order to get favorable environmental conditions. Interestingly, in this species, plasma cortisol levels exhibit a peak at monsoon (Lamba et al., 1983; Sundararaj and Goswami, 1966a, 1966b), and glucocorticoids are effective both *in vivo* and *in vitro* in inducing ovulation (Sundararaj and Goswami, 1977). *In vivo* administration of an ovulatory dose of LH in this species induces an increase in plasma levels of cortisol followed by an increase in sex steroid (T and E2) levels, while ACTH injection only stimulates cortisol levels. The authors suggest that gonadotropin acts at two loci, the interrenal and the ovary (Goswami et al., 1985). All these data indicate a potential positive role of cortisol in the induction of spawning in the catfish. Another striking example is the killifish *Fundulus heteroclitus* which exhibits synchronized spawning with the semilunar cycle of spring tides and a peak of plasma cortisol level coincides with spawning (Bradford and Taylor, 1987).

### **1.2.2. Cortisol and upstream reproductive migration in salmonids**

Upstream migration and spawning require high energy reserve mobilization; in Atlantic salmon *Salmo salar*, for example, between 60% and 70% of the body reserves are spent during upstream migration and spawning (Jonsson et al., 1997). High plasma cortisol levels and interrenal hyperplasia have long been reported during the later stages of the spawning migration and at the time of spawning in various salmonids [Pacific sockeye salmon: (Donaldson and Fagerlund, 1972; Hane and Robertson, 1959; Idler et al., 1959); migratory (steelhead) rainbow trout: (Robertson et al., 1961); Atlantic salmon: (Schmidt and Idler, 1962); kokanee salmon, landlocked form of sockeye salmon: (Carruth et al., 2000)]. However, one study in sockeye salmon reports that sexual maturation is not necessarily accompanied by elevated plasma cortisol levels (Fagerlund, 1967). Increased cortisol level is also described at spawning in non-migratory rainbow trout (Bry, 1985; Robertson et al., 1961). Altogether these studies indicate that high cortisol levels in salmonids are, at least, not deleterious to sexual maturation and spawning.

### 1.2.3. Cortisol and oceanic reproductive migration in eels

The downstream and oceanic migrations of eels (*Anguilla* species), as well as their whole gonadal development from prepubertal stage to full sexual maturation, require both metabolic energy and metabolites that come exclusively from body stores [for review: (Palstra and van den Thillart, 2010)]. Clevestam and collaborators report that 45% of European eel (*Anguilla Anguilla*) from Baltic sea would be within 90% of complete energy depletion after migration and reproduction and 20% would have completely exhausted their initial fat reserves (Clevestam et al., 2011). Eel, at the silver prepubertal, downstream migratory, stage and during their subsequent reproductive oceanic migration and sexual migration, are fasting and all their metabolic stores, accumulated during the juvenile growth (yellow stage) phase, will be mobilized. In silver eels, an elevation of plasma cortisol levels is observed prior to downstream migration (Van Ginneken et al., 2007). Early studies of hypophysectomy and

adrenalectomy [report](#) the need of [an](#) intact pituitary-interrenal axis for maintenance of liver glycogen levels in eels [*Anguilla anguilla*: (Hatey, 1951); American eel *Anguilla rostrata*: (Butler, 1968)]. Cortisol induces mobilization of lipid and protein stores and stimulates hepatic neoglucogenesis [for review: (Butler, 1973); Japanese eel *Anguilla japonica*: (Chan and Woo, 1978); *Anguilla anguilla*: (Dave et al., 1979); *Anguilla rostrata*: (Butler, 1968; Foster and Moon, 1986)]. Cortisol also induces eel vertebral demineralization, by promoting both osteocytic osteolysis and osteoclastic resorption, allowing mobilization of phospho-calcic stores necessary for vitellogenin synthesis (Sbaihi et al., 2009). Besides these actions on metabolism necessary for the eel migratory and reproductive processes, cortisol may also directly stimulate the gonadotropic axis, as shown by its positive effect on pituitary *lhβ* mRNA and LH protein content *in vivo* and *in vitro* [*Anguilla anguilla*: (Huang et al., 1999)]. Thus, in the eel, cortisol may both coordinate storage mobilization and participate in the induction of sexual maturation during reproductive migration.

### 1.3. Cortisol and gonadal sex differentiation

In some teleost fish species, gonadal sex determination and/or sex change are under the regulation of environmental factors. An increasing number of data [proposes](#) cortisol as a key factor integrating environmental cues (such as temperature or social status) to induce male sex determination/sex change [for reviews: (Fernandino et al., 2013; Fernandino and Hattori, 2019; Goikoetxea et al., 2017; Liu et al., 2017; Perry and Grober, 2003; Solomon-Lane et al., 2013; Todd et al., 2016)]. Cortisol administration or stress-induced cortisol release (including high temperature-induced effects) promotes masculinization in a number of teleosts [rainbow trout: (van den Hurk and van Oordt, 1985); pejerrey *Odontesthes bonariensis*: (Hattori et al., 2009); Japanese flounder *Paralichthys olivaceus*: (Yamaguchi et al., 2010); Southern flounder *Paralichthys lethostigma*: (Mankiewicz et al., 2013); three-spot wrasse *Halichoeres trimaculatus*: (Nozu and Nakamura, 2015); black sea bass *Centropristis striata*: (Miller et al.,

2019); medaka *Oryzias latipes* (Hayashi et al., 2010); orange spotted grouper *Epinephelus coioides* (Chen et al., 2020)]. Several potential pathways are described for cortisol triggering maleness: cross-talk between glucocorticoid and androgen metabolism (11 $\beta$ -hydroxysteroid dehydrogenase, 11 $\beta$ HSD, and 11 $\beta$ -hydroxylase, Cyp11b, enzymes) promoting the synthesis of 11-ketotestosterone (11KT); inhibition of aromatase (*cyp19a1a*) expression, leading to a shift in steroidogenesis from estrogens to androgens; and upregulation of anti-Müllerian hormone (*amh*) expression [for reviews: (Fernandino et al., 2013; Fernandino and Hattori, 2019; Goikoetxea et al., 2017; Liu et al., 2017; Perry and Grober, 2003; Solomon-Lane et al., 2013)]. In the pejerrey, cortisol increases the expression of *hsd11b2*, the gene for 11 $\beta$ HSD, an enzyme which deactivates cortisol to cortisone and catalyzes the final step in 11-KT synthesis (Fernandino et al., 2012). Typical DNA binding sites of glucocorticoid receptor (GR) *i.e.* glucocorticoid response elements (GRE) are identified in the *cyp19a1a* promoter in the goby *Gobiodon histrio* (Gardner et al., 2005) and in the black sea bass (Miller et al., 2019). Interestingly, GR is also shown to directly interact with cAMP-responsive element (CRE) on *cyp19a1a* promoter of Japanese flounder, down-regulating the expression of aromatase (Yamaguchi et al., 2010). Temperature-dependent sex determination (TSD) is shown to involve regulation of DNA methylation of the *cyp19a1a* promoter [European sea bass *Dicentrarchus labrax*: (Navarro-Martín et al., 2011)], and such epigenetic modifications may also be a mechanism by which cortisol regulates aromatase expression.

All these studies highlight the role of cortisol as a mediator between environmental conditions, especially temperature, and male sex determination in various teleost species. This confers to cortisol a key-role in the potential impact of global climatic change on the modification of sex ratio in some teleost species and its consequences on population sustainability.

#### 1.4. Animal welfare, environment, stress and reproduction

Concerns about fish welfare have increased considerably during the recent years owing to the growing public interest for the negative impacts of intensification in aquaculture and for scientific debate about pain in fish. Many authors agree on a definition for which welfare is not only based on physical health but also lack of mental suffering and presence of positive feelings (Huntingford et al., 2006; Segner et al., 2012; Sneddon et al., 2016; Stevens et al., 2017; Toni et al., 2019). In this context, most of the welfare issues are related to stress responses, experience of pains, growth problems, incidence of disease, abnormal behaviors and less frequently to degraded breeding with negative impact on male or female reproduction. These effects observed in aquaculture environment are not only the consequence of exposure of adult fish to stressors (*e.g.* confinement, handling, hypoxia or poor water quality) but also associated with abnormal [behavior](#) or impossibility to develop a normal [behavior](#) created by captive environment (Sneddon et al., 2016). Thus, one potential consequence of confinement is irregular spawners which suffer from a low fertilisation rate and a broken ovulatory rhythm (Kjesbu, 1989; Patterson et al., 2004). Intense swimming activity during chase and capture can also be sufficient to compromise reproduction (Pankhurst and Van der Kraak, 1997). Repeated acute confinement stress during [the](#) spawning season as well as poor body condition or [food](#) deprivation have also negative effects on fecundity in cod, all effects which have been suggested to be related to altered energy allocation (Bogevik et al., 2012; Kjesbu et al., 1991; Kjesbu, 1989; Lambert and Dutil, 2000). Behavioral needs can also be a welfare issue, not only by preventing maltreatment but also by providing adult fish resources to perform natural behavioural repertoire necessary for reproduction, accommodations which are species specific (Sneddon et al., 2016). Sometimes also, search for optimal performance in aquaculture can lead to welfare issue in relation to reproduction. In salmon aquaculture, early maturation can be a significant welfare issue. The maturation process is energetically expensive which is reflected in early maturing salmon by

decreased growth rate and increased mortality through susceptibility to pathogens (Gjerde, 1984; Thorpe, 1994). Selection for both fast growth and late sexual maturation has been considered problematic as it has been suggested that there is a correlation between the phenotypes fast growth and early sexual maturation (Thorpe et al., 1983). The importance of energy cost for reproduction also appears when considering exercised female sockeye salmon compared to non-exercised fish: they displayed lower lipid content, delayed maturity, lower egg deposition rate and higher egg mortality prior to ovulation (Chellappa and Huntingford, 1989; Hansen et al., 2010). Methods used in aquaculture for egg collection from female is also a crucial procedure for fish reproduction in salmonid aquaculture: [comparison](#) between various methods (massage of the fish abdomen versus pressure of gas) in rainbow trout indicated that air stripping led to better quality of eggs and better juvenile survival rate (Kowalski et al., 2018). In recreational fisheries, studies on angling and release of nesting species during the spawning season reported decreased progeny survival and paternal nest abandonment and impaired care (Cooke et al., 2000; Hanson et al., 2007; Philipp et al., 1997). However, specific investigations on gonadal development of final maturation find very few adverse effects (Booth et al., 1995; Hall et al., 2009; Lowerre-Barbieri et al., 2011). Moreover, when testing effects of mild angling and release on golden perch *Macquaria ambigua*, normal gonadal development is observed in angled fish, a result attributed to the flexible reproductive strategy of that species and benignness of the mouth hooking (Hall et al., 2017). These results illustrate the complexity of welfare questions related to reproduction, which certainly need to take [into](#) account the specificity of the fish species and/or the characteristics of the environment and stressors.

## **2. Overview of neuroendocrine stress and reproductive axes in teleosts**

### **2.1. Teleost specific traits compared to other vertebrates**

In mammals, brain-pituitary-peripheral glands axes are a classical schema when describing the endocrine regulation of major physiological functions, such as reproduction with the gonadotropic axis or stress with the corticotropic axis. Similar structures are observed in teleost fish which also present specificities which need to be considered for understanding stress/reproduction relationship in these species.

### **2.1.1. Anatomo-functional specificities**

#### **2.1.1.1. Pituitary regionalisation and direct innervation**

The pituitary gland is a vertebrate innovation [for reviews: (Dufour et al., 2020; Sower, 2018)]. In all vertebrates, the pituitary consists of the adenohypophysis and the neurohypophysis. The adenohypophysis has been subdivided by early studies into the *pars distalis* and the *pars intermedia*, with in tetrapods, an additional *pars tuberalis* [for reviews: (De Beer, 1923; Dore, 2017)]. In contrast to tetrapods in which cells of the *pars distalis* seem widely distributed, in teleosts, the different types of cells of the *pars distalis* are regionalised [*Poecilia formosa* and *latipinna* (Olivereau and Ball, 1964); European eel (Olivereau, 1967); Atlantic halibut *Hippoglossus hippoglossus* (Weltzien et al., 2004); zebrafish *Danio rerio* (Pogoda and Hammerschmidt, 2007); for reviews: (Schreibman et al., 1973; Trudeau and Somoza, 2020; Zohar et al., 2010)]. This allowed an anatomical subdivision of the *pars distalis* (PD), in teleosts, into an anterior region, the rostral *pars distalis* (RPD) composed of lactotropes (prolactin cells) and corticotropes (corticotropin, ACTH cells), and a posterior region, the proximal (or caudal) *pars distalis* (PPD) composed of somatotropes (growth hormone cells), thyrotropes (thyrotropin TSH cells) and gonadotropes (luteinizing hormone LH and follicle stimulating hormone FSH cells). Interestingly, in teleosts, the two gonadotropins, LH and FSH, are expressed in different pituitary cells, in contrast to the situation observed in mammals which produce LH and FSH in the same pituitary cell [for review: (Kanda, 2019)]. This feature allowed Golan and collaborators to investigate in

zebrafish the differential organization of LH and FSH cells and they reported that LH cells are coupled *via* strong gap-junction, while FSH cells show long cytoplasmic extension to maintain contact, possibly explaining their differential release patterns (Golan et al., 2016). In teleosts as in tetrapods, the *pars intermedia* of the pituitary contains **melanotrope** cells which produce MSH (melanocyte-stimulating hormone) from the same precursor as ACTH. In tetrapods, brain hypophysiotropic neurons project to the median eminence at the basis of the hypothalamus and release their neurohormones into the hypophyseal portal vascular system, which carries them to the *pars distalis*. Differently, in teleosts, the axonal endings of the brain hypophysiotropic neurons terminate in close vicinity to the cells of the adenohypophysis providing a direct innervation [for review: (Trudeau and Somoza, 2020; Zohar et al., 2010)].

#### **2.1.1.2. Caudal neurosecretory system**

The caudal neurosecretory system (CNSS) is unique to some non-mammalian vertebrates such as teleosts (Osteichthyes) and elasmobranchs (Chondrichthyes). It is situated in the posterior region of the spinal cord and it comprises large neurosecretory neurons named after their discoverer, the Dahlgren cells (Dahlgren, 1914). These neurons project their axons to a neurohaemal organ, the urophysis, where neuropeptides are released into the renal portal system *via* the caudal vein to the head kidney [for reviews: (Bern and Takasugi, 1962; McCrohan et al., 2007; Winter et al., 2000)]. Two of these neuropeptides were first isolated from CNSS and named urotensins: urotensin I in white sucker *Catostomus commersoni* (Lederis et al., 1982) and common carp *Cyprinus carpio* (Ichikawa et al., 1982) and urotensin II in the goby *Gillichthys mirabilis* (Pearson et al., 1980). Urotensin I is related to CRH [for review: (Lovejoy et al., 2014)] and urotensin II to somatostatin [for review: (Tostivint et al., 2014)]. Apart from these urotensins, **the** CNSS produces two other neuropeptides, CRH and parathyroid **hormone-related** protein, as well as a neurotransmitter, acetylcholine [for

review: (McCrohan et al., 2007)]. In teleosts, together with the preoptic area of the brain, the CNSS is the major source of CRH and urotensins [European flounder *Platichthys flesus*: (Lu et al., 2004); rainbow trout: (Bernier et al., 2008; Craig et al., 2005); zebrafish: (Alderman and Bernier, 2009)], two neuropeptides which are involved in the stress axis (cf § 2.2.2.1).

### **2.1.2. Teleost specific whole genome duplication**

Two whole genome duplications (WGD) likely occurred in ancestral vertebrates and are named 1R and 2R for first and second round of WGD, respectively [for review: (Dehal and Boore, 2005)]. An additional WGD specifically occurred at the basis of the teleost lineage, referred to as teleost specific WGD (TWGD) or as 3R for third round of WGD [for review: (Meyer and Van De Peer, 2005)]. WGD events led to the expansion of gene numbers. This additional WGD in teleosts would have favored the evolutionary success and the remarkable biological diversity of teleosts, the largest vertebrate group. Concerning the neuroendocrine axes, the 3R is at the origin of additional paralogs for a number of actors of both gonadotropic and corticotropic axes in teleosts. Further additional WGD occurred more recently independently in some teleost groups, such as in salmonids (Lien et al., 2016; Robertson et al., 2017) and in carps (Larhammar and Risinger, 1994; Wang et al., 2012) and are referred to as 4R for fourth round of WGD. Conservation of duplicated paralogs may be related to amplification of function, sharing of multiple preexisting functions (subfunctionalization) or acquisition of new function (neofunctionalization) [for review: (Dufour et al., 2020)].

## **2.2. The corticotropic axis**

### **2.2.1. Main actors of the corticotropic axis in vertebrates**

In all vertebrates, stress response is regulated by the corticotropic axis (HPA in mammals and sauropsids and HPI in amphibians and teleosts) [for review: (Gorissen and Flik, 2016)]. The neurohormone, corticotropin-releasing hormone (CRH), as its name indicates, controls the production and release of corticotropin (also named adrenocorticotrophic hormone, ACTH), at

the pituitary level. ACTH is the result of tissue-specific post-translational processing of proopiomelanocortin (POMC), together with melanocyte stimulating hormone (MSH), and  $\beta$ -endorphin ( $\beta$ -END). ACTH and MSH, called melanocortins (MC) act *via* MC receptors (MCR), while  $\beta$ -END acts *via* opioid [receptors](#). ACTH from the corticotrophs of the *pars distalis* of the pituitary controls via type 2 MCR (melanocortin-2 receptor, MC2R), glucocorticoid production and release from adrenal cortex cells in amniotes or interrenal cells in amphibians and teleosts. Major glucocorticoids are cortisol in most mammals and ray finned fish and corticosterone in most birds, amphibians and reptiles [for review: (Aerts, 2018)]. MC2R, in teleosts as in tetrapods, is ligand selective as it can only be activated by ACTH and not by MSH, and requires coexpression with an accessory protein, melanocortin-2 receptor accessory protein (MRAP) for trafficking to the cell surface [for reviews: (Dores, 2016; Dores et al., 2016)]. [Glucocorticoids](#) negatively [feedback](#) on the brain (hypothalamic CRH) / pituitary (ACTH) corticotropic axis [for reviews: (Bernier et al., 2009; Faught et al., 2016; Gorissen and Flik, 2016)]; this negative feedback regulation of [the](#) HPI axis involves glucocorticoid receptor (GR) signaling pathway.

### 2.2.2. Specific features of [the](#) corticotropic axis in teleosts

The hypothalamic regulation of corticotropes and melanotropes in teleosts has already been extensively reviewed by Bernier and collaborators (Bernier et al., 2009).

#### 2.2.2.1. CRH and related peptides and their receptors

CRH, first isolated from sheep hypothalamus, together with urotensin I (UI) isolated from the CNS of the white sucker (Lederis et al., 1982) and the common carp (Ichikawa et al., 1982), and with sauvagine (SVG) isolated from the skin of the amphibian *Phylomedusa sauvagei* (Montecucchi et al., 1980), form a large family of peptides. Later on, urocortins were identified in mammals and also found to be related to CRH: urocortin 1 (Ucn1), ortholog of UI and SVG (Vaughan et al., 1995); urocortin 2 [Ucn2: (Reyes et al., 2001)] and urocortin 3

416 [Ucn3: (Lewis et al., 2001)] [for review: (Lovejoy and Balment, 1999)]. All these peptides,  
 417 CRH and urocortins in mammals and CRH and UI/SVG in teleosts and amphibians, represent  
 418 the CRH/urocortin family.

419 The complex evolutionary scenario of this family was recently clarified by Cardoso and  
 420 collaborators (Cardoso et al., 2016). As previously suggested by Hwang and collaborators,  
 421 two ancestral *crh/ucn1* and *ucn2/ucn3* genes likely arose by specific gene duplication before  
 422 vertebrate WGD events (Hwang et al., 2013). Both ancestral genes were duplicated twice in  
 423 ancestral vertebrates *via* 1R and 2R, followed by some paralog losses, leading to up to 5  
 424 genes (*crh1*, *crh2*, *ucn1* issued from ancestral *crh/ucn1*; *ucn2*, *ucn3*, issued from ancestral  
 425 *ucn2/ucn3*) in extant representative species of some vertebrate lineages such as  
 426 chondrichthyans, holosteans and actinistians (Cardoso et al., 2016). Teleost specific 3R  
 427 resulted in the duplication of *crh1* into two paralogs *crh1a* and *crh1b* conserved in many  
 428 species [for review: (Cardoso et al., 2016)]. *Crh2* may have been lost in recent teleosts  
 429 (Cardoso et al., 2016), while one 3R-*crh2* paralog has been conserved in basal groups of  
 430 teleosts [(Maugars et al., 2016) and Maugars et al. unpublished data].

431 The involvement of CRH in the stress response in teleosts is well-documented [for reviews:  
 432 (Flik et al., 2006; Gorissen and Flik, 2016)]. Teleost *crhb* (*crh1b*) paralog is widely expressed  
 433 in the brain of various teleost species [goldfish *Carassius auratus*: (Bernier et al., 1999);  
 434 flounder *Platichthys flesus*: (Lu et al., 2004); zebrafish: (Alderman and Bernier, 2007);  
 435 Japanese eel: (Amano et al., 2014); *Astatotilapia burtoni*: (Carpenter et al., 2014);  
 436 *Schizothorax prenanti*: (Wang et al., 2014)]. To our knowledge, the expression of the *crha*  
 437 (*crh1a*) paralog was only investigated in *Astatotilapia burtoni* and zebrafish: in the zebrafish,  
 438 *crha* expression is restricted to the lateral tuberal nucleus of the ventral hypothalamus, while  
 439 in *A. burtoni*, no *crha* expression is detected in the brain (Grone and Maruska, 2015). The  
 440 authors also show that, while both forms are expressed in the retina of *Astatotilapia burtoni*,

441 only *crhb* expression is observed in the retina of zebrafish (Grone and Maruska, 2015). These  
 442 first data suggest species-specific variations in the respective roles of 3R-duplicated *crhl*  
 443 paralogs in teleosts.

444 In mammals, CRH binds to G-protein coupled receptors, CRHR1 and CRHR2, which belong  
 445 to the class 2 subfamily B1 of secretin-like receptor superfamily. *Crhr1* was duplicated *via*  
 446 teleost-3R into two paralogs (*crhr1a* and *crhr1b*) which were conserved in many extant  
 447 teleosts, while one of 3R-duplicated *crhr2* paralogs would have been lost [for review:  
 448 (Cardoso et al., 2014)]. CRHR1s are thought to mediate CRH action along the  
 449 corticotropic/stress axis, as, like mammalian CRHR1, teleost CRHR1s have similar affinity  
 450 for CRH and UI/Ucn1 while CRHR2 has higher affinity for UI and urocortins (2 and 3) than  
 451 for CRH [catfish *Ameiurus nebulosus*: (Arai et al., 2001); chum salmon *Oncorhynchus keta*:  
 452 (Pohl et al., 2001); common carp: (Manuel et al., 2014)].

453 The ACTH-releasing action of CRH has been demonstrated *in vitro* in many teleost species  
 454 [goldfish: (Fryer et al., 1984); rainbow trout: (Baker et al., 1996; Pierson et al., 1996);  
 455 gilthead sea bream *Sparus aurata*: (Rotllant et al., 2001, 2000); Mozambique tilapia  
 456 *Oreochromis mossambicus*: (Van Enckevort et al., 2000); common carp: (Metz et al., 2004)],  
 457 as in mammals (Rivier et al., 1983; Rivier and Plotsky, 1986), birds (Carsia et al., 1986) and  
 458 amphibians (Tonon et al., 1986). In the common carp, the release of ACTH is stimulated by  
 459 CRH but only when ACTH cells are submitted to DA inhibition (Metz et al., 2004).

460 CRH stimulates not only ACTH release, but also  $\alpha$ -MSH release, in various teleosts such as  
 461 in Mozambique tilapia (Lamers et al., 1994; Van Enckevort et al., 2000), gilthead sea bream  
 462 (Rotllant et al., 2001), red porgy *Pagrus pagrus* (Van Der Salm et al., 2004) and common  
 463 carp (Van Den Burg et al., 2005). This  $\alpha$ -MSH-releasing effect of CRH is also reported in  
 464 mammals such as rat (Meunier et al., 1982; Proulx-Ferland et al., 1982). In contrast, in an  
 465 amphibian, the frog *Rana ridibunda*, while CRH is a potent stimulator of ACTH release by

466 anterior pituitary cells, it is ineffective on  $\alpha$ -MSH release by neurointermediate lobes *in*  
467 *vitro* (Tonon et al., 1986).

468 Other CRH-related peptides such as UI and sauvagine can induce in goldfish the release *in*  
469 *vitro* of ACTH (Fryer et al., 1984, 1983), as well as of  $\alpha$ -MSH (Tran et al., 1990).

470 Interestingly, UI and SVG are equipotent to CRH in stimulating ACTH release by rat anterior  
471 pituitary cells (Rivier et al., 1983), while in the frog *Rana ridibunda*, they are unable to  
472 stimulate ACTH release (Tonon et al., 1986). Concerning  $\alpha$ -MSH, UI and SVG induce its  
473 release by the neurointermediate lobe *in vitro* in *Xenopus laevis* (Verburg-Van Kemenade et  
474 al., 1987), but not in *Rana ridibunda* (Tonon et al., 1986). In addition to its action on the  
475 pituitary, UI is able to directly stimulate cortisol release, and to potentiate ACTH-stimulatory  
476 effect, on interrenals *in vitro* in rainbow trout (Arnold-Reed and Balment, 1994) and  
477 European flounder (Kelsall and Balment, 1998). Finally, in maturing masou salmon  
478 *Oncorhynchus masou*, UI rather than CRH could be involved in the control of  
479 hypercortisolemia, as hypothalamic *uI* expression rises in correlation with enhanced cortisol  
480 secretion, while *crh* expression does not (Westring et al., 2008).

481 In the rainbow trout, both *crh* and *uI* expressions in the brain (hypothalamus and preoptic  
482 area) increase in response to various stressors: hyperosmotic challenge (Craig et al., 2005),  
483 hyperammonemia, isolation (Bernier et al., 2008) as well as hypoxia (Bernier et al., 2008;  
484 Bernier and Craig, 2005). Notably, *crh* and *uI* expressions increase also in the caudal  
485 neurosecretory system (CNSS) in response to various stressors in the rainbow trout  
486 [hyperosmotic challenge: (Craig et al., 2005); hyperammonemia: (Bernier et al., 2008)] and in  
487 the olive Japanese flounder, *Paralichthys olivaceus* [acute hypothermal stress: (Yuan et al.,  
488 2020)]. In the European flounder, net restraint induces an increase in *crh* expression in the  
489 CNSS, but not in the hypothalamus, as well as an increase in CNSS cortisol receptors in  
490 addition to an increase in plasma cortisol levels (Lu et al., 2004). In this species,

glucocorticoid receptors have been colocalized with UI in the CNSS (Bond et al., 1999). All these data suggest the possible involvement of the CNSS in the stress-specific regulation of cortisol production in teleosts and the existence of an extra-pituitary feedback system on CNSS for the suppression of cortisol secretion. Earlier *in vivo* studies already reported the possible involvement of CNSS in the regulation of the corticotropic axis: injection of urophysis extracts or synthetic *Catostomus commersoni* UI in flounder induces an elevation of cortisol (Arnold-Reed and Balment, 1989); in goldfish, urophysectomy produces marked elevation of hypothalamic UI-like activity, pituitary ACTH and plasma cortisol, which can not be observed in urophyssectomized fish receiving dexamethasone (Woo et al., 1985). More functional investigations are needed to assess the possibility of a major contribution of CNSS CRH-related peptides to the regulation of cortisol secretion and stress responses.

#### 2.2.2.2. Dopamine and other neurohormones

##### \*Dopamine

In mammals, different data have been reported concerning the effects of dopamine (DA) on ACTH release between *in vitro* and *in vivo* studies. Treatment of rat anterior pituitary glands (Van Loon and Kragt, 1970) or human corticotroph adenoma cells (Ishibashi and Yamaji, 1981) with DA lowers ACTH release. In contrast, administration of DA agonists to rats elevates plasma ACTH concentrations and subsequently plasma corticosterone levels, and this elevation is prevented by pretreatment with DA antagonists (Borowsky and Kuhn, 1992; Ježová et al., 1985). An  $\alpha$ -MSH-release inhibitory effect of DA is observed in amphibians [*Rana pipiens*: (Saland et al., 1982); *Rana ridibunda* : (Jenks et al., 1985); *Xenopus laevis* : (Verbug-Van Kemenade et al., 1986)].

Possible involvement of DA in the hypothalamic regulation of teleost corticotropes and melanotropes has been suggested by Metz and colleagues. As basal ACTH release increases slowly and steadily over time when pituitary glands of common carp are incubated *in vitro*,

they **conclude** that ACTH release **is** under a predominant inhibitory control *in vivo* (Metz et al., 2004). They also show that DA **inhibits** ACTH release and **is** necessary for CRH-stimulation of ACTH release (Metz et al., 2004). It was previously reported that DA was able to inhibit *in vitro*  $\alpha$ -MSH release by goldfish (Omeljaniuk et al., 1989) and red porgy (Van Der Salm et al., 2004) pituitaries as well as by Mozambique tilapia neurointermediate lobes (Lamers et al., 1991). *In vitro* treatment with DA D2 receptor agonists **inhibits**  $\alpha$ -MSH release, while DA D1 receptor agonists **have** a stimulatory effect on  $\alpha$ -MSH release, in Mozambique tilapia exposed to water with low pH (acid stress) (Lamers et al., 1997). Overall, these results support an involvement of DA in the regulation of ACTH and  $\alpha$ -MSH release probably interdependently with CRH pathway [for review: (Gorissen and Flik, 2016)].

#### **\*Serotonin**

In mammals, serotonin (5-hydroxytryptamine, 5-HT), a monoamine neurotransmitter, is known to stimulate the HPA axis [for review: (Dinan, 1996)] and to control stress coping [for review: (Puglisi-Allegra and Andolina, 2015)]. 5-HT mainly acts on the brain, but it can also act directly on the adrenal and pituitary glands [for review: (Dinan, 1996)]. The brain network involves CRH, GABA and glutamate neurons [for review: (Puglisi-Allegra and Andolina, 2015)].

Similar results have been reported in teleost fish. Injection of a selective agonist for serotonin receptor 5-HT<sub>1A</sub> (8-OH-DPAT) results in increased plasma cortisol concentrations in some teleosts [rainbow trout: (Winberg et al., 1997); Gulf toadfish *Opsanus beta*: (Medeiros et al., 2010); goldfish: (Lim et al., 2013)]. In the Arctic charr *Salvelinus alpinus*, comparison between stressed and unstressed fish **show** that this selective agonist has a stimulatory effect on the HPI axis in unstressed fish, while having a suppressive effect on the stress-induced activation of the HPI axis in fish stressed by handling and *ip* injections (Höglund et al., 2002). In the Gulf toadfish, injection of **8-OH-DPAT** results in increased hypothalamic *crh* mRNA

541 levels and ACTH release from the pituitary and these effects are inhibited by crowding stress  
542 (Medeiros et al., 2014). However in goldfish, injection of this agonist has no effect at the  
543 brain level but rather acts at interrenal tissue to stimulate cortisol secretion (Lim et al., 2013).  
544 *In vitro*, 5-HT stimulates cortisol release by interrenals in Gulf toadfish [kidney pieces:  
545 (Medeiros and McDonald, 2012)] and in goldfish [superfused head kidney tissue: (Lim et al.,  
546 2013)]. In goldfish, Lim and colleagues demonstrate the involvement of multiple 5-HT  
547 receptor subtypes (5-HT<sub>1A</sub> and 5-HT<sub>4</sub>) in the interrenal paracrine effect of 5-HT (Lim et al.,  
548 2013), while in Gulf toadfish, Medeiros and McDonald show that 5-HT<sub>4</sub> does mediate 5-HT  
549 action but 5-HT<sub>1A</sub> does not (Medeiros and McDonald, 2012).

550 \* *Thyrotropin-releasing hormone*

551 In mammals and in adult amphibians, *thyrotropin-releasing hormone* (TRH) is the main  
552 neurohormone controlling the thyrotropic axis *i.e.* the synthesis and release of thyrotropin  
553 (TSH) at the pituitary level, which stimulates the production of thyroid hormones at the  
554 thyroid level. This TSH-releasing role seems to be taken on by CRH in larval amphibians,  
555 reptiles and birds [for reviews: (De Groef et al., 2006; Galas et al., 2009)]. In teleosts,  
556 investigations on the *in vitro* effect of TRH on pituitary TSH production *show* either no effect  
557 [common carp: synthesis, (Kagabu et al., 1998); mRNA levels, (Geven et al., 2009); coho  
558 salmon *Oncorhynchus kisutch*, release: (Larsen et al., 1998)] or a stimulatory effect [bighead  
559 carp *Aristichthys nobilis*, mRNA levels: (Chatterjee et al., 2001; Chowdhury et al., 2004);  
560 Japanese eel, mRNA levels: (Han et al., 2004)]. CRH acts as a TSH-releasing factor on coho  
561 salmon pituitary cells (Larsen et al., 1998), but has no effect on *tshβ* mRNA levels in  
562 common carp pituitary glands *in vitro* (Geven et al., 2009).

563 Conversely to the effect of CRH on TSH, TRH can exert an ACTH-releasing effect in teleosts  
564 as shown by *in vitro* studies [goldfish pituitary dispersed cells: (Tran et al., 1989); gilthead  
565 sea bream pituitary: (Rotllant et al., 2000)]. TRH is also a potent stimulator of  $\alpha$ -MSH release

by the *pars intermedia* in many teleost species [goldfish: (Tran et al., 1989); tilapia: (Lamers et al., 1991); rainbow trout: (Schwartzentruber et al., 1994); gilthead sea bream: (Rotllant et al., 2000); red porgy: (Van Der Salm et al., 2004)]. This  $\alpha$ -MSH-releasing effect of TRH is also observed in amphibians [*Rana ridibunda*: (Tonon et al., 1980); *Xenopus laevis*: (B. Verburg-Van Kemenade et al., 1987)].

After a stress (prolonged crowding), in the gilthead sea bream, the pituitary presents a differential response of ACTH and  $\alpha$ -MSH to TRH and CRH treatments: the stimulation of ACTH by CRH is attenuated, but not the stimulation of ACTH by TRH, while both CRH and TRH stimulation of  $\alpha$ -MSH are enhanced (Rotllant et al., 2000). In Mozambique tilapia, a dual organization of the stress axis has been demonstrated, as after exposure to low-pH water, the CRH/ACTH axis is replaced by a TRH/di-acetylated  $\alpha$ -MSH axis for the production of cortisol (Lamers et al., 1994). Overall, these results suggest the involvement of TRH in the regulation of corticotropes and melanotropes, notably during exposure to stress.

#### ***\*Arginine vasotocin, isotocin and arginine vasopressin***

Arginine vasotocin (AVT), isotocin (IST) and arginine vasopressin (AVP) are all peptides produced by neurons of the preoptic nucleus and released by the neurohypophysis. AVT and IST are present in non-mammalian vertebrates and closely related to mammalian AVP.

As in mammals, these neuropeptides are able to stimulate the release of ACTH *in vitro* [goldfish: (Fryer et al., 1985); trout: (Baker et al., 1996; Bond et al., 2007; Pierson et al., 1996)] and thus increase plasma cortisol levels in teleosts *in vivo* [goldfish: (Fryer and Leung, 1982); trout: (Baker et al., 1996; Bond et al., 2007; Pierson et al., 1996)]. In addition, in mammals and birds [for review: (Cornett et al., 2012)], they potentiate the stimulatory action of CRH on ACTH *in vivo* and *in vitro* (Gillies et al., 1982; Rivier and Vale, 1983; Turkelson et al., 1982). In teleosts, contradictory data are available concerning a possible synergy with CRH on ACTH release *in vitro*: they do synergize in trout (Baker et al., 1996), while they do

not in goldfish (Fryer et al., 1985)]. In the frog *Rana ridibunda*, no potentiation of CRH-induced ACTH release is observed when anterior pituitary cells are incubated with a combination of AVP and CRH (Tonon et al., 1986).

In rainbow trout, a study shows that AVT mRNA levels in neurons of the preoptic nucleus are elevated during acute stress but not chronic stress (Gilchriest et al., 2000).

### 2.2.2.3. POMC-derived peptides and their receptors

#### *\*POMC-derived peptides*

Pro-opiomelanocortin (POMC), together with proenkephalin (PENK), prodynorphin (PDNY) and proorphanin (PNOC), form the opioid/orphanin gene family [(Sundström et al., 2010); for review: (Dores et al., 2002)]. POMC is post-translationally processed and some of the peptides obtained undergo further modifications such as C-terminal amidation and N-acetylation [for reviews: (Dores and Baron, 2011; Takahashi and Mizusawa, 2013)]. POMC organizational plan varies among vertebrates [for reviews: (Dores and Baron, 2011; Dores and Lecaude, 2005)]. POMC is the precursor for ACTH, melanotropins ( $\alpha$ -,  $\beta$ - and  $\gamma$ - MSH), corticotropin-like intermediate peptide (CLIP), lipotropins ( $\beta$ - and  $\gamma$ - LPH) and  $\beta$ -endorphin, in tetrapods and lungfish (Amemiya et al., 1999a; Nakanishi et al., 1979). Chondrichthyans have an additional melanotropin,  $\delta$ -MSH (Amemiya et al., 1999b). Teleosts lack  $\gamma$ -MSH (Kitahara et al., 1988; Lee et al., 1999), but *pomc*- $\beta$  of cichlids and pomacentrids encodes a novel melanocortin peptide,  $\epsilon$ -MSH, the result of a tandem duplication of the segment encoding ACTH (Harris et al., 2014). In all these vertebrates, the adenohypophysis possesses two cell populations that express POMC: in the *pars distalis* (PD), the corticotropes in which POMC is processed to ACTH (and  $\beta$ -lipotropin), and in the *pars intermedia* (PI), the melanotropes in which POMC is processed to ACTH, which is further cleaved to give  $\alpha$ -MSH, corticotropin-like intermediate peptide and  $\beta$ -endorphin [for review: (Takahashi and Mizusawa, 2013)]. Differently, agnathans possess two ‘POMC’ genes: pro-opiocortin (POC),

which encodes ACTH and a different  $\beta$ -endorphin, is expressed in the *pars distalis*; pro-opiomelanotropin (POM), which encodes  $\alpha$ -, and  $\beta$ - MSH, and  $\beta$ -endorphin, is expressed in the *pars intermedia* (Takahashi et al., 2012, 1995).

Teleost 3R gave rise to *pomc* gene duplicates: *pomc- $\alpha$*  (*a* or *A*) and *pomc- $\beta$*  (*b* or *B*), with *pomc- $\beta$*  having lost a functional  $\beta$ -endorphin (De Souza et al., 2005). Further independent gene duplications during teleost evolution resulted in duplicates of *pomc- $\alpha$*  such as in halibut and carp [*pomc-I* and *-II*: (De Souza et al., 2005)], sea bream [*pomc- $\alpha 1$*  and *- $\alpha 2$* : (Cardoso et al., 2011)] and *Astatotilapia burtoni* [*pomc- $\alpha 1$*  and *- $\alpha 2$* : (Harris et al., 2014)]. In barfin flounder *Verasper moseri*, a third *pomc* gene, named *pomc-C*, was also identified (Takahashi et al., 2006); *pomc-C* belongs to the *pomc- $\beta$*  clade while *pomc-A* and *-B* belong to the *pomc- $\alpha$*  clade and thus should be renamed *- $\beta$* , *- $\alpha 1$*  and *- $\alpha 2$* , respectively, according to Cardoso and collaborators (Cardoso et al., 2011). In rainbow trout, apart from *pomc-B*, *pomc-A1* and *-A2* are likely the result of the salmonid genome duplication (4R) (Leder and Silverstein, 2006).

As in other vertebrates [*e.g.* in rodents: (Kraicer et al., 1973)], ACTH is produced in teleosts from POMC by both the *pars distalis* (PD; corticotropes) and the *pars intermedia* (PI; melanotropes) of the pituitary, but ACTH of the *pars intermedia* is further cleaved to produce smaller peptides and cortisol feedbacks only on the *pars distalis* ACTH [for review: (Fryer and Lederis, 1986)]. In tetraodon *Tetraodon nigroviridis*, *pomc- $\alpha$*  and not *pomc- $\beta$*  is expressed in the PD, while both are expressed in the PI (De Souza et al., 2005). In the sea bream, *pomc- $\alpha 2$*  and not *pomc- $\alpha 1$*  is expressed in the PD, while only *pomc- $\alpha 1$*  is expressed in the PI (Cardoso et al., 2011). In the pituitary of barfin flounder, all three *pomc* genes present in this species are expressed in the PD (Takahashi et al., 2006). In *Astatotilapia burtoni* (Harris et al., 2014) and rainbow trout (Leder and Silverstein, 2006), all three *pomc* are expressed in the pituitary, but no indication of the region is reported. This indicates species-specific variation in the expression of the various *pomc* paralogs by the PD corticotropic cells

through teleost radiation.

Species-specific variations among teleosts may also concern the *pomc* paralog linked to stress response. In the gilthead sea bream, an enhanced pituitary expression of *pomc- $\alpha$ 2* (*pomc-b* in the paper) and a decreased one of *pomc- $\alpha$ 1* (*pomc-a* in the paper) are observed after acute stress due to air exposure (Skrzynska et al., 2018)], leading the authors to link *pomc- $\alpha$ 2* and not *pomc- $\alpha$ 1* to stress response in sea bream. However, in the same species, no change in the expression of both paralogs was previously reported after exposure to other acute stress, chasing and persecution (Toni et al., 2015). In *Rhamdia quelen*, both pituitary *pomc- $\alpha$ 2* (*pomcb* in the paper) and *pomc- $\alpha$ 1* (*pomca* in the paper) expressions are unchanged after stressful situation (transport) (Saccol et al., 2018). In contrast, in the Senegalese sole, *Solea senegalensis*, *pomc- $\alpha$ 1* (*pomc-a* in the paper) expression in the pituitary is down-regulated in juveniles chronically stressed by high stocking density, whereas *pomc- $\alpha$ 2* (*pomc-b* in the paper) expression levels remain unaffected (Wunderink et al., 2012). All these data in three different species do not lead to clear conclusions concerning a preferential involvement of one or another *pomc* paralog in stress response among teleosts.

An *in vitro* corticotropic action (*i.e.* induction of cortisol release from head kidney) of  $\alpha$ -MSH (desacetyl and diacetyl forms) has been reported in rainbow trout (Rance and Baker, 1981), Mozambique tilapia (Lamers et al., 1992) and barfin flounder (Kobayashi et al., 2011), but not in common carp (Metz et al., 2005).

#### **\*Melanocortin receptors**

In tetrapods, five MCRs have been identified. In teleost fish, the number of receptors increases up to six in zebrafish, which has two MC5R paralogs (*mc5ra* and *mc5rb*) (Västermark and Schiöth, 2011), while pufferfish *Fugu* has only four, with no melanocortin *mc3r* and only one copy of melanocortin *mc5r* (Logan et al., 2003). Concerning the ligand selectivity of MCRs, all of the paralogous MCRs can be activated by both ACTH and  $\alpha$ -MSH

in extant cartilaginous fishes, while in extant teleosts and tetrapods, MC2R can be activated only by ACTH. The appearance of MRAP1 paralleled the emergence of this MC2R ligand selectivity [for reviews: (Dores, 2016; Dores et al., 2016)].

In mammals, the MCRs have distinct expression site and functions [for reviews: (Cone, 2006; Dores et al., 2014)]: MC1R, expressed in melanocytes, is involved in skin and hair pigmentation; MC2R, expressed in adrenal cortex, is involved in adrenal steroidogenesis and stress response; MC3R and MC4R, expressed in the brain, are involved in the control of energy homeostasis; MC5R, expressed in a variety of exocrine glands, such as sebaceous, lacrimal and preputial glands, is involved in exocrine gland secretion. These features can be also found in teleosts but some peculiarities can be noted. Of particular interest, *mc5r* is co-expressed with *mc2r* in the interrenal of several teleosts [rainbow trout: (Aluru and Vijayan, 2008; Haitina et al., 2004); common carp *Cyprinus carpio*: (Metz et al., 2005); barfin flounder: (Kobayashi et al., 2011)], as in the chicken adrenal (Takeuchi and Takahashi, 1998) and *Xenopus tropicalis* interrenal/kidney (Dores and Garcia, 2015), suggesting a possible role of MC5R in the regulation of HPI/HPA axis in these non-mammalian vertebrates.

#### 2.2.2.4. Corticosteroids and their receptors

##### **\*Glucocorticoids**

In fish, corticosteroids are synthesized by the interrenal tissue, a tissue embedded inside the anterior part of the kidney and homologous to the adrenal cortex in mammals. Corticosteroids are steroid hormones divided into glucocorticoids and mineralocorticoids. Even if cortisol is the major corticosteroid in teleosts [for review: (Mommensen et al., 1999)], others such as 11-deoxycortisol (17, 21 dihydroxy-4-pregnene-3,20 dione), 11-deoxycorticosterone (DOC) and corticosterone are also detected in plasma of teleosts [for review: (Butler, 1973); winter flounder *Pseudopleuronectes americanus*: (Campbell et al., 1976); rainbow trout *Salmo gairdneri*: (Campbell et al., 1980)]. All teleosts so far studied lack aldosterone, which is the

691 principal mineralocorticoid in mammals (Gilmour, 2005), and it is generally accepted that  
692 cortisol [exerts](#) both glucocorticoid and mineralocorticoid actions in teleosts (McCormick,  
693 2001; McCormick et al., 2008). DOC [is](#) shown to be a potent agonist of mineralocorticoid  
694 receptor (MR) (Sturm et al., 2005). Study in rainbow trout has shown a sustained up-  
695 regulation of plasma DOC levels during a confinement stress time-course. However, the low  
696 DOC levels compared to cortisol measured in the plasma do not favour an activity of DOC  
697 through MR receptors (Kiilerich et al., 2018) and physiological relevance of this hormone  
698 action is still unclear in fish (Prunet et al., 2006).

### 699 ***\*Glucocorticoid and mineralocorticoid receptors***

700 Glucocorticoid and mineralocorticoid receptors (GR and MR respectively) are corticosteroid  
701 receptors which belong to the nuclear receptor superfamily [for review: (Bury, 2017)]. In  
702 teleosts, 3R gave rise to duplicated *gr* (*gr1* and *gr2*) with one receptor (GR1) retaining a 9  
703 aminoacid insert which does not exist in GR2 [for review: (Bury, 2017)]. Zebrafish is an  
704 exception as this species conserved only one of the two *gr* paralogs (Schaaf et al., 2008). In  
705 some teleosts, GR1 has two splice variants, GR1a and GR1b [rainbow trout: (Takeo et al.,  
706 1996); *Haplochromis burtoni*: (Greenwood et al., 2003); Tetraodon and Takifugu species:  
707 (Stolte et al., 2006); marine medaka *Oryzias dancena*: (M. Kim et al., 2011)]. 3R gave also  
708 rise to duplicated *mr* (*mr1* and *mr2*) which have been both conserved in a basal teleost, the  
709 European eel [(Lafont et al., 2014) and Lafont et al. unpublished data], while only the *mr1*  
710 paralog would have been conserved in extant teleosts so far studied [for review: (Baker and  
711 Katsu, 2019)]. As they are issued from the 3R, these two *mr* paralogs in the eel should rather  
712 be named *mra* and *mrb*, according to the commonly used nomenclature for teleost 3R-  
713 [paralogs](#). However, they must be distinguished from the two *mr* forms previously found in [the](#)  
714 rainbow trout, named *rtmra* and *rtmrb* (Sturm et al., 2005), which may represent allelic  
715 variants or paralogs issued from salmonid-4R of *mr1*.

716 GR1 and GR2 present differential affinities (sensitivities) and show distinct functionality with  
717 GR2 having a higher hormonal transcriptional activity at equimolar concentration and higher  
718 sensitivity (Bury et al., 2003). This difference in sensitivity is not restricted to the  
719 salmoniformes and have been also observed with the two GRs in *Pantodon buchholzi* (Li et  
720 al., 2012), carp (Stolte et al., 2008), marine medaka (*Oryzias latipes*) (M. Kim et al., 2011),  
721 and the Japanese medaka (*Oryzias latipes*) (Miyagawa et al., 2015). From such *in vitro*  
722 differences in functionality, one can hypothesize that the two teleost GRs may have different  
723 roles: the hypersensitive GR2 could play a prominent role during basal circulatory cortisol  
724 concentrations (unstressed) whereas the less sensitive GR1 may become prominent during  
725 stressful situations when cortisol levels are high [(Bury et al., 2003); for review: (Bury,  
726 2017)]. So far, such functionally distinct roles have not been established in fish. However, a  
727 recent study of stress effects in salmonids shows differential contribution of the corticosteroid  
728 receptors in the regulation of HPI axis activity (Kiilerich et al., 2018) and suggests a negative  
729 feedback regulation of cortisol release at the pituitary level via MR, while a short loop  
730 regulation occurs at the interrenal level via GR in rainbow trout exposed to 7 day-  
731 confinement. The authors also observe the presence of the 2 GR and the MR in hypophysial  
732 ACTH cells confirming the regulation of the HPI axis at the pituitary level by both GRs and  
733 MR. Similar conclusions have been also suggested when studying in the Atlantic salmon  
734 effects of unpredictable chronic stress or to repeated chasing stress (Madaro et al., 2016,  
735 2015). In the common carp, Stolte and collaborators report that, in fish exposed to prolonged  
736 and strong stressors, mRNA levels of all three corticosteroid receptors (*gr1*, *gr2* and *mr*) are  
737 down-regulated in some brain regions, but not in CRH neurons or pituitary ACTH cells  
738 (Stolte et al., 2008), suggesting a role of all three receptors in stress regulation in this species.  
739 In zebrafish (Schaaf et al., 2008), the single *gr* (*gr2*) gene has two splice variants, *grα* and  
740 *grβ* (Hollenberg et al., 1985), with a variant lacking the C-terminal portion of the GR as

described in human (Hollenberg et al., 1985). In human, *grβ* may act as a dominant-repressor of the wildtype variant (Bamberger et al., 1995). These splicing variants are co-expressed in a number of zebrafish tissues (brain, spleen, liver, intestine, heart, gills and muscle), *grα* mRNA levels being significantly higher than those of *grβ* (Schaaf et al., 2008). Despite initial data suggesting that *grβ* has a key role in the negative regulation of *grα* (Chatzopoulou et al., 2015), a recent study by the same group using transgenic zebrafish with inducible expression of *grβ* suggests that this isoform does not have a functional role in transcription regulation (Chatzopoulou et al., 2017). Functional roles of corticosteroid receptors during stress have been studied in this fish species using mutant fish. Thus, GR knockout zebrafish show an inability to cope with stressor such as placement into an unfamiliar environment, even after repeated exposure (Ziv et al., 2013). These fish also show hypercortisolemia and fail to exhibit a cortisol stress response, while KO zebrafish for MR have a delayed but sustained cortisol response (Faught and Vijayan, 2018). Both KO mutants do not show the hyperactivity in response to light observed in wild type zebrafish (Faught and Vijayan, 2018). These recent data suggest distinct but complementary roles for GR and MR in the development and regulation of the stress axis in zebrafish: MR represses HPI axis during development while GR regulates basal cortisol levels in the context of negative feedback regulation during stress. Both receptors are also involved in the control of stress-related behaviour in zebrafish (Faught and Vijayan, 2018). Such a role in the regulation of stress-related behaviour is suggested in medaka as well (Sakamoto et al., 2016).

## 2.3. The gonadotropic axis

### 2.3.1. Main actors of the gonadotropic axis in vertebrates

The brain-pituitary-gonadal axis is responsible for the control of the reproductive function in all vertebrates. It regulates gamete production, sexual secondary characters and behaviors, as

well as key-steps of the life cycle such as puberty, seasonal reproduction or sex change in some hermaphroditic species including teleost representatives.

The brain integrates various environmental and social cues, such as temperature, photoperiod, habitat conditions, presence of partners, as well as internal cues such as developmental stage, size, nutritional status and energy stores, *via* various neuronal networks which converge into the positive or negative regulation of the activity of gonadotropin-releasing hormone (GnRH) hypophysiotropic neurons considered as the master control of reproduction (Gore, 2002). GnRH binds to its cognate membrane receptor (GnRH-R) expressed by the pituitary gonadotrope cells, and which belongs to the rhodopsin-like, class A, G-protein coupled receptor (GPCR) superfamily [for reviews: (Flanagan and Manilall, 2017; Millar et al., 2004)]. GnRH stimulates the synthesis and release of the gonadotropins, LH and FSH.

The two pituitary gonadotropins, LH and FSH are glycoprotein hormones, composed of two subunits alpha and beta. The alpha subunit, named glycoprotein hormone alpha subunit ( $Gp\alpha$ ), is common to both gonadotropins, LH, FSH and to the thyrotropin, TSH, while the beta subunit is specific to each hormone [for reviews: (Cahoreau et al., 2015; Pierce and Parsons, 1981)]. LH and FSH bind to their respective GPCR membrane receptors LHR and FSHR expressed by the gonads. The gonadotropins activate gonadal functions, gametogenesis and steroidogenesis. The enzymatic pathway of the biosynthesis of sex steroids is largely conserved among vertebrates. Sex steroids, androgens, estrogens, progestagens, bind to their respective nuclear receptors, AR, ER, PR, belonging to the nuclear receptor superfamily [for review: (Mangelsdorf et al., 1995)]. As more recently discovered, steroid may also act *via* binding to GPCR membrane receptors, such as estradiol-membrane receptor and progesterone membrane receptor [for review: (Thomas, 2012)]. Sex steroids act on multiple targets, including the gonads themselves, a variety of peripheral tissues as well as the brain and pituitary, where they exert positive and negative feedback controls on the gonadotropic axis.

## 2.3.2. Specific features of the gonadotropic axis in teleosts

### 2.3.2.1. GnRHs and their receptors

Up to three *gnrh* genes are present in extant teleosts, a larger number than in mammals (up to two genes), which had been initially attributed to the 3R. However, subsequent comparative studies revealed the presence of three *gnrh* genes also in basal vertebrate representatives such as a chondrichthyan, the catshark, *Scyliorhynchus canicula* and a basal sarcopterygian, the coelacanth, *Latimeria chalumnae*. Phylogeny and synteny studies support the 1R/2R origin of the three vertebrate *gnrh* and their inheritance by the teleost lineage. In contrast, 3R had no impact on the *gnrh* gene number in extant teleosts, indicating an early loss of 3R-paralogs in this lineage [for reviews: (D. K. Kim et al., 2011; Roch et al., 2011; Tostivint, 2011)]. A maximum of three *gnrh* (1, 2, 3) genes are thus present in teleosts, such as in medaka, while additional, independent and species-specific losses led to the presence of only two *gnrh* genes in some species, e.g. *gnrh1* and 2 in the eel, or *gnrh2* and 3 in the zebrafish. The situation is more complex concerning GnRH receptors. Recent studies indicate that a large number of GnRH receptors resulted from 1R/2R as well as from local gene duplications in ancestral vertebrates; subsequently, several independent gene loss events led to a variable number of *gnrh* genes throughout vertebrate evolution. In teleosts, 3R-duplicated paralogs as well as additional duplications in some lineages increased the number of GnRH-R [up to 5 in acanthopterygians and 6 in salmonids: (Ciani et al., 2020)] as compared to other actinopterygians, with species-specific variations among teleosts due to independent gene losses [(Ciani et al., 2020); for reviews: (Roch et al., 2014; Sefideh et al., 2014; Williams et al., 2014)].

### 2.3.2.2. Dopamine and other neurohormones

**\*Dopamine**

A crucial control is exerted by dopaminergic neurons which directly inhibit the [gonadotrope](#) cell activity and counteract the effect of GnRH, as first discovered in cyprinids by Peter and collaborators and retrieved in many other teleosts species including basal [representatives](#) such as the eel [for reviews: (Dufour et al., 2010, 2005)]. In relation with environmental and internal cues, and in a species-specific manner, this dopaminergic control of gonadotropic cells plays key roles in various reproductive steps in teleosts such as puberty, final oocyte maturation and ovulation, spermiation, or sex change [for reviews: (Dufour et al., 2010, 2005)]. [A major inhibitory role of DA in regulating reproduction is also reported in amphibians, especially in the control of spawning \[for review: \(Vu and Trudeau, 2016\)\].](#)

Beside GnRH and [DA](#), a variety of other neurohormones can also directly regulate the activity of teleost gonadotrophs, with species-specific variations, such as kisspeptin (Kiss), gonadotropin-inhibitory hormone (GnIH), neurokinin, gamma-aminobutyric acid (GABA), noradrenaline, serotonin *etc* [for reviews: (Dufour et al., 2020; Zohar et al., 2010)]. As proposed by Trudeau, these multiple independent controls of gonadotropic cells may provide compensatory regulatory mechanisms, possibly explaining the low or no impact of KO of major genes such as *kiss* or *gnrh* observed in recent studies (Trudeau, 2018). Species-specific variations in the multiple controls of gonadotrophs highlight the evolutionary plasticity of the brain-pituitary anatomo-functional relationships in the teleost lineage, which may have favored the remarkable diversity of their reproductive cycles.

#### ***\*GnIH***

Gonadotropin-inhibitory hormone (GnIH or RFRP-3 in mammals), a RFamide peptide, was discovered from quail brain by Tsutsui and collaborators and [found to be](#) involved in the direct inhibitory pituitary regulation of LH (Tsutsui et al., 2000). Shortly after, [the](#) GnIH precursor was shown to encode two other peptides (Satake et al., 2001). Since then, *gnih* gene homologs, encoding multiple peptides, have been identified in other vertebrates [for reviews:

(Muñoz-Cueto et al., 2017; Tsutsui et al., 2018)]. These peptides, in mammals, also possess an inhibitory action on gonadotropins, while in amphibians they have GH-releasing effects [for review: (Dufour et al., 2020)]. Recently, it was demonstrated that only a single *gnih* gene was present in representative species of elopomorphs (*Anguilla* species), the most basal group of teleosts, as well as in the other teleosts (Maugars et al., 2020). This result indicates a loss of one of the two 3R-duplicated *gnih* paralogs shortly after the 3R (Maugars et al., 2020). Depending on teleost species, inhibitory [e.g. European eel (Pasquier et al., 2018); common carp (Peng et al., 2016); zebrafish (Spicer et al., 2017)], stimulatory [e.g. sockeye salmon (Amano et al., 2006); grass puffer *Takifugu niphobles* (Shahjahan et al., 2011); tilapia *Oreochromis niloticus* (Biran et al., 2014); catla *Catla catla* (Kumar et al., 2019); half-smooth tongue sole *Cynoglossus semilaevis* (Wang et al., 2019)] or a lack of effect [e.g. goldfish (Qi et al., 2013); *Astyanax altiparanae* (Branco et al., 2019)] of GnIH have been described for gonadotropins *in vitro* [for review: (Muñoz-Cueto et al., 2017; Tsutsui et al., 2018; Ubuka and Parhar, 2018)], suggesting a large variability of GnIH function in reproduction across teleosts.

#### \* *Kisspeptin*

Kisspeptin, another RF-amide peptide, is encoded by the *Kiss-1* gene, a metastasis suppressor gene, isolated in 1996 from melanoma cells (Lee et al., 1996). This 54 amino-acid peptide, also called metastin, is processed into shorter peptides which all bind to kisspeptin receptor, KissR (or GPR54) (Ohtaki et al., 2001). In 2003, three research groups discovered the key role of the kisspeptin system in the activation of gonadotropic axis and the control of reproduction in human and mice (de Roux et al., 2003; Funes et al., 2003; Seminara et al., 2004). *In vivo* and *in vitro* studies further demonstrate the stimulatory role of kisspeptin at different levels of the gonadotropic axis: hypothalamus (GnRH neurons), pituitary (LH cells) and gonads [for reviews: (Pinilla et al., 2012; Putteeraj et al., 2016)]. The few available

studies in amphibians and reptiles also suggest a stimulatory role of kisspeptin on the HPG axis, while this system might have been lost in some birds [for review: (Dufour et al., 2020)]. However, data in teleosts report stimulatory, no or inhibitory effects of kisspeptin peptides on GnRH or gonadotropins, suggesting a large variability of the role of Kiss system in teleost reproduction according to species [for reviews: (Dufour et al., 2020; Tena-Sempere et al., 2012)]. Up to four paralogs of *Kiss* and *KissR* resulted from 1R and 2R in early vertebrates, but subsequent gene losses led to the presence of only a single *Kiss* gene and a single *KissR* gene in human. In teleosts, due to paralog losses, no impact of 3R on the number of both *Kiss* and *KissR* genes has been found, leading to a maximum of two *Kiss* and three *KissR* genes, which is no more than in a non-teleost actinopterygian such as the gar, *Lepisosteus oculatus* [for reviews: (Pasquier et al., 2014, 2012)].

#### 2.3.2.3. Gonadotropins and their receptors

As in mammals, only two gonadotropins, LH and FSH, are present in all extant teleosts, including basal representative species such as the eel (elopomorphs) (Yoshiura et al., 1999), indicating an early loss after 3R of the putative additional 3R-paralogs [for review: (Dufour et al., 2020)]. In tetrapods, LH and FSH bind to their respective receptors, LHR and FSHR. As recently shown, an additional *lhr* paralog arose from a local gene duplication in ancestral actinopterygians, and the two *lhr* (*lhr1* and *lhr2*) were inherited by the teleost lineage (Maugars and Dufour, 2015). Due to gene losses after 3R, the number of gonadotropin receptors did not further increase in teleosts with a single *fshr* and up to two *lhr*. Additional independent and species-specific gene losses led to the presence of a single *lhr* in some teleosts, such as only *lhr1* in medaka and *lhr2* in zebrafish (Maugars and Dufour, 2015).

#### 2.3.2.4. Sex steroids and their receptors

888 Gonadal sex steroids are produced from cholesterol by a succession of enzymatic activities. In  
889 teleosts, 3R duplication of a key-enzyme, the aromatase (*Cyp19a1a* and *b*), has been largely  
890 documented [for review: (Zhang et al., 2014)].

891 Sex steroid nuclear receptors have been duplicated by 3R and the conservation of most of the  
892 paralogs led to the presence of a larger number of nuclear receptors in teleosts as compared to  
893 mammals [for review: (Ogino et al., 2018)]. This is also the case for the steroid membrane  
894 receptors. Concerning estradiol receptors, tetrapods possess two nuclear receptors (ESR1 also  
895 named ER $\alpha$ , and ESR2 also named ER $\beta$ ) and a single membrane receptor (GPER) while  
896 teleosts, as a result of 3R, possess three nuclear receptors (*esr1* and duplicated *esr2a* and  
897 *esr2b*) and two membrane receptors (*gpera* and *gperb*) (Lafont et al., 2016). One of the  
898 duplicated *esr1* paralog would have been lost shortly after 3R. A species-specific gene loss  
899 occurred in zebrafish, which **possesses** a single *gper* paralog, while both *gpera* and *b* paralogs  
900 have been conserved in the other teleosts investigated so far (Lafont et al., 2016). For  
901 androgen receptors, 3R also generated two *ar* paralogs (*ara* and *arb*) in teleosts (Ogino et al.,  
902 2009). Recently, Morini and colleagues revealed the presence of duplicated progesterone  
903 nuclear receptors (*pgra* or *pgr1*, and *pgrb* or *pgr2*) in the eel, which originated from 3R. In  
904 other teleosts however only a single *pgr* (orthologous to eel *pgrb/pgr2*) has been found, and  
905 synteny analysis **indicates** the loss of *pgra* (*pgr1*) paralog in the teleost lineage after the  
906 emergence of the basal teleost group of elopomorphs (Morini et al., 2017). Two *pgr* paralogs  
907 **are** reported in the goldfish, likely resulting from 4R. The evolutionary history of membrane  
908 progestin receptors is even more complex as five isoforms have been characterized in  
909 vertebrates (Morini et al., 2017). The conservation of multiple steroid receptor genes in  
910 teleosts likely reflects subfunctionalization and neofunctionalization processes, which may  
911 have contributed to species-specific diversification of sexual characters, regulatory  
912 processes and reproductive cycles throughout teleost radiation.

### 3. Neuroendocrine mechanisms of the interactions between corticotropic and gonadotropic axes in teleosts

#### 3.1. Regulatory effects of actors of the corticotropic axis on the gonadotropic axis

##### 3.1.1. Roles of corticosteroids and their receptors

##### 3.1.1.1. Expression and role along the gonadotropic axis

In the rainbow trout, GR are expressed in GnRH- and dopaminergic neurons as well as on gonadotrope cells [(Teitsma et al., 1999); for review: (Teitsma et al., 1998)]. In addition, GR and ER colocalize in the same neurons and pituitary cells [for review: (Teitsma et al., 1998)].

In *Astatotilapia burtoni*, both GR1 and GR2 are expressed in GnRH1 neurons in the preoptic area (Korzan et al., 2014). A direct action of cortisol on GnRH neurons and targets is suggested in teleosts by the presence of several GR responsive elements (GRE) in the promoters of zebrafish *gnih* and *gnih* receptor (Ogawa and Parhar, 2014). GRs are also expressed in gonads in various teleosts [midshipman *Porichthys notatus*: (Arterbery et al., 2010) ; *Astatotilapia burtoni*: (Maruska and Fernald, 2011)].

MR is expressed in brain, especially in GnRH1 neurons [*Astatotilapia burtoni*: (Korzan et al., 2014)], and in gonads of various teleosts [*Haplochromis burtoni*: (Greenwood et al., 2003); rainbow trout: (Milla et al., 2008; Sturm et al., 2005); midshipman *Porichthys notatus*: (Arterbery et al., 2010); *Astatotilapia burtoni*: (Maruska and Fernald, 2011); for review: (Takahashi and Sakamoto, 2013)].

These expression patterns of GR and MR along the gonadotropic axis suggest potential major actions of glucocorticoids on reproductive functions at different levels, involving endocrine loops and/or paracrine/autocrine regulations.

Corticosteroids themselves are produced by gonads in teleosts [for review: (Kime, 1993)].

Cortisol is present in sperm and seminal fluid [Pacific herring *Clupea harengus* and North sea

plaice *Pleuronectes platessa*: (Scott et al., 1991b, 1991a)]. 11-deoxycortisol is produced by the ovary [*Gillichthys mirabilis* and *Microgadus proximus*: (Colombo et al., 1973); sea bass: (Colombo et al., 1978); *Jenynsia lineata*: (Tesone and Charreau, 1980); goldfish: (Kime et al., 1992)] and testis [*Jenynsia lineata*: (Tesone and Charreau, 1980)]. Like 11-deoxycortisol, 11-deoxycorticosterone (DOC) is an important product of ovarian [*Leptocottus armatus*, *Gillichthys mirabilis* and *Microgadus proximus*: (Colombo et al., 1973); sea bass: (Colombo et al., 1978)] and/or testicular [*Jenynsia lineata*: (Tesone and Charreau, 1980)] steroidogenesis. Future studies should investigate whether ACTH, gonadotropins or both, associated or not to other factors are regulating such corticosteroid production by gonads in fish.

#### 3.1.1.2. *In vivo effects of corticosteroids*

Pioneer works in loach *Misgurnus fossilis* (Kirshenblatt, 1952), conger *Conger conger* (Mousset, 1957) and European eel (Fontaine et al., 1964) show that injection of glucocorticoids (desoxycorticosterone acetate) has positive effects on reproduction as it is able to induce oocyte maturation and spawning. In the catfish *Heteropneustes fossilis* also, adrenal corticosteroids (cortisol, cortisone and desoxycorticosterone acetate) are shown to be effective in inducing ovulation and spawning in hypophysectomized gravid fish, suggesting a direct positive action on the ovary independently of the pituitary (Sundararaj and Goswami, 1966a). Use of metopiron, an adrenocortical inhibitor which can induce «chemical interrenalectomy», allows to show that LH-induced effect on ovulation and spawning in catfish was interrenal-dependent (Sundararaj and Goswami, 1966b). Treatment with cortisol increases pituitary gonadotropin in juvenile rainbow trout [implant: (Crim et al., 1981)] and prepubertal European eel [injection: (Dufour et al., 1983; Huang et al., 1999)]. Altogether, these data suggest positive interactions of glucocorticoids on the HPG axis at both juvenile and adult stages in some teleost species.

963 In contrast, a lot of studies report negative effects of glucocorticoids on the reproductive axis.  
964 Most of them are conducted in salmonids. In the brown trout, implantation of cortisol to  
965 sexually maturing male and female fishes result in smaller gonads, lower plasma levels of sex  
966 steroids and pituitary gonadotropin content (Carragher et al., 1989). Differently, in the  
967 maturing male rainbow trout, a cortisol implant decreases plasma gonadotropin levels while  
968 having no effects on other parameters (plasma sex steroids, pituitary gonadotropin content  
969 and gonad size) (Carragher et al., 1989). In the rainbow trout, intraperitoneal implant of  
970 cortisol decreases hepatic E2-binding site concentration, but increases plasma E2-binding  
971 capacity in immature females (Pottinger and Pickering, 1990) and decreases ER and Vg  
972 expression in maturing females (Lethimonier et al., 2000). Cortisol injection depresses  
973 plasma E2 and T levels, without affecting plasma gonadotropin levels in vitellogenic rainbow  
974 trout (Pankhurst and Van Der Kraak, 2000). In the male rainbow trout, testicular *mr*  
975 expression and plasma DOC increase around the time of spermiation, and the co-  
976 administration of DOC with 17 $\alpha$ , 20 $\beta$ -dihydroprogesterone decreases the spermatocrit value  
977 and therefore increases the sperm fluidity (Milla et al., 2008). In Atlantic salmon, Eriksen and  
978 collaborators report impact on progeny of maternal exposure to cortisol (implant), including  
979 increased mortality and prevalence of deformities, reduced development and yolk-sac volume  
980 (Eriksen et al., 2007, 2006). During a field study on spawning grounds, use in female pink  
981 Pacific salmon *Oncorhynchus gorbuscha* of cortisol implant which further increases  
982 endogenous cortisol levels, impairs reproductive success (fewer eggs) and survival  
983 (McConnachie et al., 2012). The use of metyrapone, which prevents the synthesis of cortisol  
984 from 11-deoxycortisol by inhibiting 11 $\beta$ -hydroxylase, does not change reproductive success  
985 (McConnachie et al., 2012). Altogether, these studies indicate that endogenous physiological  
986 cortisol levels, which raise at the time of upstream migration and spawning in salmonids (cf  
987 §1.2.2), do not impair reproduction, although higher levels may have a negative effect

988 (McConnachie et al., 2012).

989 Deleterious reproductive *in vivo* effects of corticosteroids have also been shown in other  
990 teleost species. In immature male common carp fed with cortisol-containing food pellets over  
991 pubertal period, brain sGnRH content, pituitary *lh* and *fsh* mRNA levels, plasma LH levels  
992 and testicular androgen (T and 11KT) secretion are decreased (Consten et al., 2001a).  
993 Testicular development in this species **is** retarded by this long-term cortisol treatment  
994 (Consten et al., 2001a, 2001b). In contrast, dietary administered cortisol to channel catfish  
995 *Ictalurus punctatus* does not modify reproduction (Small, 2004). In the Arctic char, co-  
996 exposure **to** cortisol **and** E2 results in reduced plasma Vg levels and no change in *Vg* mRNA  
997 levels, suggesting a post-transcriptional action of cortisol (Berg et al., 2004). Implantation of  
998 cortisol **causes** retardation of the ovarian growth and depression of plasma sex steroid levels  
999 in tilapia *Oreochromis mossambicus* [female: (Foo and Lam, 1993a); male: (Foo and Lam,  
1000 1993b)]. Later, Gennotte and collaborators **show** that cortisol treatment induces both positive  
1001 (reduction of the time before ovulation) and negative (reduction of fecundity) effects on the  
1002 reproduction in other tilapia species, *Oreochromis niloticus* (Gennotte et al., 2012).

1003 In the cinnamon clownfish, *Amphiprion melanopus*, injection of cortisol increases ***gnih***  
1004 **mRNA**, and reduces *sbGnRH* mRNA and plasma levels of LH and FSH (Choi et al., 2017).  
1005 As, in this teleost species, GnIH inhibits gonadotropin expression (Choi et al., 2016), cortisol  
1006 treatment leads to a dual negative effect on gonadotropin *via* **an** increase in GnIH and  
1007 decrease in GnRH. A similar upregulation of hypothalamic *gnih* (*rfrp3*) mRNA levels is  
1008 observed in rats after acute or chronic immobilization/restraint stress (Kirby et al., 2009;  
1009 Yang et al., 2017), upregulation which is abolished by adrenalectomy (Kirby et al., 2009) .

1010 Concerning the effect of stress or corticosteroid treatment on **the** kisspeptin system, all the  
1011 available data have been obtained in rodents [for reviews: (Acevedo-Rodriguez et al., 2018;  
1012 Iwasa et al., 2018)] and show a reduction of hypothalamic *KissI* and *KissRI* mRNA levels

[rat, restraint stress or subcutaneous injection of corticosterone: (Kinsey-Jones et al., 2009); mice, implant containing corticosterone: (Luo et al., 2016); mice, restraint stress: (Yang et al., 2017)]. It would be relevant to investigate the regulation of the kisspeptin system by stress and corticosteroids in teleosts, which show different effects of kisspeptin on the control of gonadotropins according to the species.

### 3.1.1.3. *In vitro* effects of corticosteroids

*In vitro* investigations pin-point the direct actions of cortisol at the different levels of the HPG axis. In the rat hypothalamic rfrp-expressing cell line (rHypoE-23) (Gingerich et al., 2009), treatment with corticosterone increases *gnih(rfrp)* mRNA levels and the application of a GR antagonist blocks this increase (Gojska and Belsham, 2014; Son et al., 2014). *In vitro* investigations should be developed in teleosts, using hypothalamic neurons, to study possible direct effects of corticosteroids on the various neuronal populations (GnRH, DA, GnIH and kisspeptin) involved in the gonadotropic axis.

At the pituitary level, cortisol affects reproduction-related gene expression in Atlantic cod pituitary cultures (von Krogh et al., 2019). In this species, cortisol has dual effects on *fshβ* expression, stimulating expression in cells from mature fish, while inhibiting expression in cells from spent fish (von Krogh et al., 2019). In contrast, cortisol has no direct effect on *lhβ* expression (von Krogh et al., 2019). While *gnrhr2a* transcript levels largely increase following cortisol treatment, *gnrhr1b* expression decreases in cells from spent fish and was unaffected at other maturity stages (von Krogh et al., 2019). In the prepubertal European eel, cortisol increases LH cellular content and *lhβ* mRNA levels by pituitary cells in culture (Huang et al., 1999). Future studies should investigate the effects of corticosteroids on the levels of other pituitary targets such as GnIH and kisspeptin receptors in different teleost species, considering that, in teleosts, GnIH and kisspeptin possess direct inhibitory or stimulatory effects on the pituitary depending on the species.

1038 At the liver level, dexamethasone inhibits both basal and E2-stimulated ER expression and  
1039 subsequent Vg expression in trout hepatocyte aggregates (Lethimonier et al., 2000). This is  
1040 achieved *via* an inhibition by GR of the transcriptional factor CCAAT/enhancer-binding  
1041 protein (C/EPB $\beta$ ) binding to the ER promoter region (Lethimonier et al., 2002).

1042 At the gonadal level, different data are available on the direct effect of cortisol on female  
1043 gonadal steroidogenesis depending on species [(Pankhurst, 1998); for review: (Pankhurst,  
1044 2016)]. A decrease in E2 and T release by cultured ovarian follicles after cortisol treatment  
1045 has been observed in rainbow trout (Carragher and Sumpter, 1990), while no such effect has  
1046 been reported in goldfish, common carp and *Pagrus auratus* (Pankhurst et al., 1995). In the  
1047 rainbow trout, cortisol **is** shown to have suppressive effects on basal and LH- or cAMP-  
1048 stimulated T and E2 production and secretion, but no effect on basal or LH-stimulated  
1049 17,20 $\beta$ -dihydroxy-4-pregnene-3-one (17,20 $\beta$ -P) (Barkataki et al., 2011; Reddy et al., 1999).  
1050 In the same species, cortisol **is** reported to enhance the efficiency of gonadotropin to induce  
1051 intrafollicular oocyte maturation (Jalabert, 1975) and to positively regulate the secretion of  
1052 follicular 17 $\alpha$ -hydroxy-20 $\beta$ -dihydroprogesterone (DHP) induced by gonadotropin *in vitro*  
1053 (Jalabert and Fostier, 1984). All these data suggest that, in the rainbow trout, cortisol has a  
1054 positive (or no) effect on the synthesis of progestagen (Maturation Inducing Steroid, MIS)  
1055 during final oocyte maturation, while being inhibitor at earlier stages of oogenesis, on the  
1056 production of T and E2 during vitellogenesis. In **the** zebrafish, cortisol **does** not inhibit hCG-  
1057 stimulated E2 production, while ACTH did (Alsop et al., 2009). In other teleost species,  
1058 glucocorticoids are also able to act on final oocyte maturation. **Glucocorticoids** have been  
1059 shown to induce *in vitro* oocyte maturation in the goldfish [desoxycorticosterone: (Jalabert et  
1060 al., 1973)] and ovulation in the medaka [cortisol: (Hirose, 1976)]. In **the** amago salmon  
1061 (*Oncorhynchus rhodurus*), DOC **can** enhance oocyte maturation induced by gonadotropin or  
1062 by DHP *in vitro* (Young et al., 1982). Cortisol treatment induces DNA damage on zebrafish

stage I and II follicles, and increases *mc2r* expression in stage II follicles *in vitro* (Sousa et al., 2015).

Data on male gonadal steroidogenesis, even if fewer, seem more consistent than those on female steroidogenesis. In the common carp, treatment of male testes *in vitro* with dexamethasone reduces the induction of androgen secretion by carp pituitary extract (Consten et al., 2002). Consten and collaborators have previously reported a decreased *in vitro* production of testicular androgen during early stages of puberty in the common carp fed with cortisol (Consten et al., 2000). In male rainbow trout, DOC possesses an inhibitory effect on the testicular production of DHP *in vitro* (Milla et al., 2008). In the Japanese eel, *in vitro* cortisol treatment induces DNA replication in spermatogonia and enhances the spermatogonial proliferation stimulated by 11KT, revealing a stimulatory effect of cortisol on spermatogenesis (Ozaki et al., 2006).

### 3.1.2. Roles of POMC-derived peptides

Other stress-related endocrine factors have also been studied for their involvement in the regulation of the HPG axis.

*Pomca*-deficient zebrafish, obtained with a TALENs approach, have enhanced somatic growth, reduced anxiety-related behavioral responses, and exhibit hypocortisolism associated with hyperandrogenism (Shi et al., 2019). The authors suggest that the hypocortisolemia and hyperandrogenism are probably due to the changes in expression of interrenal steroidogenic enzymes involved in the synthesis of cortisol (downregulation of *hsd3* expression) and androgens (upregulation of *cyp17ala* and *cyp19ala* expression), respectively (Shi et al., 2019).

*In vitro* experiments show a direct role of ACTH on zebrafish ovaries: ACTH treatment suppresses hCG-stimulated E2 release from ovarian follicles (Alsop et al., 2009), and induces a high level of oocyte vacuolization, as well as DNA damage on stage I and II follicles (Sousa

et al., 2015). In agreement with the direct action of ACTH on the gonads, the ACTH receptor MC2R is highly expressed in gonads of various teleosts [rainbow trout: (Aluru and Vijayan, 2008); zebrafish: (Agulleiro et al., 2010; Alsop et al., 2009); barfin flounder: (Kobayashi et al., 2011); sea bass: (Agulleiro et al., 2013)].

Another POMC-related peptides, the opioid peptide  $\beta$ -endorphin, is also produced and released in response to stressors in teleosts [gilthead sea bream: (Arends et al., 1999; Mosconi et al., 1998); tilapia: (Chabbi and Ganesh, 2012)]. This peptide may be involved in the reproductive response to stress as a suppression of LH secretion, an inhibition of vitellogenic follicular growth and a reduction of gonado-somatic index (GSI) are observed in tilapia injected with  $\beta$ -endorphin (Chabbi and Ganesh, 2013; Ganesh and Chabbi, 2013). Treatment with naltrexone, an opioid receptor antagonist, attenuates these inhibitory effects of  $\beta$ -endorphin in tilapia (Chabbi and Ganesh, 2013; Ganesh and Chabbi, 2013). In common carp, naltrexone stimulates the *in vitro* LH release from whole pituitary gland as well as from dispersed pituitary cells, revealing direct paracrine pituitary effects of opioids (Socha et al., 2003).

### 3.1.3. Roles of CRH

The direct and indirect effects of CRH on the gonadotropic axis have been well-documented in mammals [for reviews: (Chand and Lovejoy, 2011; Kageyama, 2013)]. CRH, injected centrally [ovariectomized female rat: (Ono et al., 1984; Rivier and Vale, 1984); intact and castrated male rat: (Frias et al., 1999)], but not peripherally [human: (D'Agata et al., 1984); rat: (Ono et al., 1984; Rivier and Vale, 1984)], inhibits LH release. Centrally administered CRH also down-regulates *Kiss1* and *Kiss1r* mRNA levels in ovariectomized E2 replaced female rats (Kinsey-Jones et al., 2009). Central administration of  $\alpha$ -helical CRH, a non-selective CRH-R antagonist, reverses the inhibitory effect of stress on pulsatile release of LH in castrated male rat (Rivier et al., 1986). *In vitro*, CRH down-regulates GnRH release from

rat hypothalamic slices and this inhibition is suppressed by  $\alpha$ -helical CRH (Nikolarakis et al., 1986). In parallel to its central inhibitory effect, CRH is also able to act at the pituitary level through CRH-R2 to inhibit LH release in mice (Raftogianni et al., 2018). Previous data in rat failed to show any effect of CRH on LH release *in vitro* (Ono et al., 1984). In teleosts, there are very few investigations regarding CRH action on the gonadotropic axis. Injection of CRH to stripped tilapia *Oreochromis mossambicus* resulted in diminished immunoreactive LH content in the pituitary, lower GSI and absence of vitellogenic follicles in the ovary (Chabbi and Ganesh, 2014). The glucocorticoid synthesis inhibitor, metyrapone, abolishes the inhibitory effects of CRH treatment on LH secretion and follicle growth, demonstrating an action *via* cortisol (Chabbi and Ganesh, 2014). Direct pituitary or gonadal effect of CRH needs to be investigated through *in vitro* studies. CRH-R mRNAs have already been detected in the pituitary [catfish *Ameiurus nebulosus*: (Arai et al., 2001); common carp: (Huising et al., 2004); *Astatotilapia burtoni*: (Chen and Fernald, 2008); tilapia *Oreochromis mossambicus*: (Aruna et al., 2015, 2012)], but future investigations should aim at determining whether these receptors are expressed in other cell types than corticotrophs, notably in gonadotrophs. CRH-R are also expressed in the ovary [*Astatotilapia burtoni*: (Chen and Fernald, 2008); olive flounder *Paralichthys olivaceus*: (Zhou et al., 2019)], which suggests a direct action of CRH on the gonads.

### 3.2. Regulatory effects of actors of the gonadotropic axis on the corticotropic axis

Conversely, some studies reveal a role of actors of the gonadotropic axis on the corticotropic axis in teleosts.

#### 3.2.1. Roles of sex steroids

In the sockeye salmon, gonadectomy blocks the cortisol prespawning rise (Donaldson and Fagerlund, 1970; Robertson, 1961) and sex steroids, estrogens and androgens, can induce

interrenal hyperplasia as well as increase in plasma cortisol levels in both intact and gonadectomized fish [(Donaldson and Fagerlund, 1969; Fagerlund and Donaldson, 1969); for review: (Dickhoff, 1989)]. The situation is different in the rainbow and brown trout in which implantation of T and 11-KT attenuates ACTH and cortisol-elevation after confinement stress, while E2 implantation increases stress-induced ACTH and cortisol levels (Pottinger et al., 1996). In the rainbow trout, *in vivo* treatment with 11KT suppresses interrenal responsiveness to ACTH *in vitro* (Young et al., 1996), while DHP stimulates cortisol production by interrenal tissue cultured *in vitro* (Barry et al., 1997). McQuillan and collaborators compared the *in vitro* effect of E2 on cortisol synthesis by interrenals of the chinook salmon, *Oncorhynchus tshawytscha* and rainbow trout: they show that E2 suppresses the ability of juvenile and mature chinook salmon interrenals to produce cortisol, whereas it has no effect in the rainbow trout (McQuillan et al., 2003). These data reveal potential differences among species and reproductive stage concerning the role of sexual steroids in the induction of hypercortisolemia in salmonids.

Interestingly, the interrenal in fish has been reported to produce androgen [rainbow trout: (Arai et al., 1969); coho salmon: (Schreck et al., 1989); tilapia *Oreochromis mossambicus*: (Balm et al., 1989); African catfish, *Clarias gariepinus*: (Vermeulen et al., 1995)], as in mammals [for review: (Rainey and Nakamura, 2008)]. In human, the production and secretion of adrenal androgens (dehydroepiandrosterone, DHEA and its sulfate ester, DHEAS) in the midchildhood, termed adrenarche, has been involved in the timing of puberty (Remer et al., 2010). Such a role of interrenal androgens could be also envisioned in teleosts.

A potential remarkable interaction between reproductive and stress actors is illustrated by the recent finding of the ability of progesterone to bind to the mineralocorticoid receptor MR. Recent *in vitro* studies indicate that progesterone can activate MR in the elephant shark *Callorhynchus milii*, in ray-finned fishes including trout (Sturm et al., 2005) and zebrafish

(Fuller et al., 2019; Katsu and Baker, 2018) and in chickens, with an EC50 of less than 1 nM for these MRs, which makes this steroid a potential physiological activator of these MRs [for review: (Baker and Katsu, 2020)]. Moreover, in the elephant shark and ray-finned fish, MR are expressed in several tissues, including ovaries and testis (Katsu et al., 2019; Milla et al., 2008; Sturm et al., 2005). This led Baker and Katsu to suggest that progesterone may have multifaceted physiological roles through MR in these species [for review: (Baker and Katsu, 2020)].

### 3.2.2. Roles of other actors

Injection of salmon gonadotropin to sockeye salmon has no impact on plasma cortisol or cortisone levels nor on interrenal activity (Donaldson and McBride, 1974). However, in catfish *Heteropneustes fossilis*, *in vitro* studies suggest that gonadotropin (LH) stimulates the interrenal to produce corticosteroids which in turn act on the oocytes to induce maturation (Sundararaj and Goswami, 1977). Such studies in *Heteropneustes fossilis* highlight the complex synergy between gonadotropic and corticotropic axes in the induction of oocyte final maturation, ovulation and spawning in response to monsoon-related environmental cues (cf §1.2.1).

Other actors of the gonadotropic axis, such as GnRH, might act on the corticotropic axis in teleosts as evidenced in mammals. GnRH is shown to stimulate ACTH release from the rat pituitary (Gambacciani et al., 1988) and also to regulate the expression of pituitary corticotropin-releasing hormone binding protein (CRH-BP) by gonadotrope-like cell line (Westphal and Seasholtz, 2005), highlighting the potential importance of GnRH/CRH interactions at the pituitary level in the interface between stress and reproductive axes. To our knowledge such investigations have not yet been performed in teleosts.

Recent cyto-anatomical studies suggest a potential role of GnIH not only in the gonadotropic axis but also in the corticotropic axis in teleosts. In tilapia, GnIH (LPXRFa) axons are closely

associated with various pituitary cell types including gonadotrophs, corticotrophs and melanotrophs, and GnIH receptors (LPXRFa-Receptor) are expressed in LH, ACTH and alpha MSH cells [(Ogawa et al., 2016); for review: (Muñoz-Cueto et al., 2017)]. It would be very interesting now to clarify whether GnIH is involved in the regulation of these cell types and in the production of ACTH and MSH.

## **Conclusion and perspectives**

Stress impact on reproduction has long been studied in teleosts, especially in the frame of aquaculture. However, most of the accumulated data come from studies on the possible action of peripheral hormone, cortisol, on sexual maturation. Emerging data in mammals show that actors of higher levels (especially brain) should catch researcher attention. The teleost specific whole genome duplication (3R) had made such studies perhaps more difficult, multiplying the number of genes, but thanks to the recent publications of an increasing number of teleost genomes and the development of molecular and genome editing technology tools, it is now possible to investigate their respective functions across teleost species diversity. Thus future directions should aim, for example, at dissecting mechanisms of action of the different paralogs of genes involved in the corticotropic axis and in the caudal neurosecretory system in the regulation of various levels of the gonadotropic axis. Such studies are all the more important that the present review gives multiple examples of species-specific regulatory mechanisms which probably reflect the wide species-specific plasticity of fish reproductive responses when fish are exposed to stressful situations.

## **References**

Acevedo-Rodriguez, A., Kauffman, A.S., Cherrington, B.D., Borges, C.S., Roepke, T.A.,

1213 Laconi, M., 2018. Emerging insights into hypothalamic-pituitary-gonadal axis regulation  
 1214 and interaction with stress signalling. *J. Neuroendocrinol.* 30, 0–3.  
 1215 <https://doi.org/10.1111/jne.12590>

1216 Aerts, J., 2018. Quantification of a glucocorticoid profile in non-pooled samples is pivotal in  
 1217 stress research across vertebrates. *Front. Endocrinol. (Lausanne).* 9, 1–9.  
 1218 <https://doi.org/10.3389/fendo.2018.00635>

1219 Agulleiro, M.J., Roy, S., Sánchez, E., Puchol, S., Gallo-Payet, N., Cerdá-Reverter, J.M.,  
 1220 2010. Role of melanocortin receptor accessory proteins in the function of zebrafish  
 1221 melanocortin receptor type 2. *Mol. Cell. Endocrinol.* 320, 145–152.  
 1222 <https://doi.org/10.1016/j.mce.2010.01.032>

1223 Agulleiro, M.J., Sánchez, E., Leal, E., Cortés, R., Fernández-Durán, B., Guillot, R., Davis, P.,  
 1224 Does, R.M., Gallo-Payet, N., Cerdá-Reverter, J.M., 2013. Molecular Characterization  
 1225 and Functional Regulation of Melanocortin 2 Receptor (MC2R) in the Sea Bass. A  
 1226 Putative Role in the Adaptation to Stress. *PLoS One* 8.  
 1227 <https://doi.org/10.1371/journal.pone.0065450>

1228 Alderman, S., Bernier, N., 2007. Localization of corticotropin-releasing factor, urotensin I,  
 1229 and CRF-binding protein gene expression in the brain of the zebrafish, *Danio rerio*. *J.*  
 1230 *Comp. Neurol.* 502, 783–793. <https://doi.org/10.1002/cne>

1231 Alderman, S.L., Bernier, N.J., 2009. Ontogeny of the corticotropin-releasing factor system in  
 1232 zebrafish. *Gen. Comp. Endocrinol.* 164, 61–69.  
 1233 <https://doi.org/10.1016/j.ygcen.2009.04.007>

1234 Alsop, D., Ings, J.S., Vijayan, M.M., 2009. Adrenocorticotrophic hormone suppresses  
 1235 gonadotropin-stimulated estradiol release from zebrafish ovarian follicles. *PLoS One* 4.  
 1236 <https://doi.org/10.1371/journal.pone.0006463>

1237 Aluru, N., Vijayan, M.M., 2008. Molecular characterization, tissue-specific expression, and

1238 regulation of melanocortin 2 receptor in rainbow trout. *Endocrinology* 149, 4577–4588.  
 1239 <https://doi.org/10.1210/en.2008-0435>

1240 Amano, M., Mizusawa, N., Okubo, K., Amiya, N., Mizusawa, K., Chiba, H., Yamamoto, N.,  
 1241 Takahashi, A., 2014. Cloning of corticotropin-releasing hormone (CRH) precursor  
 1242 cDNA and immunohistochemical detection of CRH peptide in the brain of the Japanese  
 1243 eel, paying special attention to gonadotropin-releasing hormone. *Cell Tissue Res.* 356,  
 1244 243–251. <https://doi.org/10.1007/s00441-013-1784-6>

1245 Amano, M., Moriyama, S., Ligo, M., Kitamura, S., Amiya, N., Yamamori, K., Ukena, K.,  
 1246 Tsutsui, K., 2006. Novel fish hypothalamic neuropeptides stimulate the release of  
 1247 gonadotrophins and growth hormone from the pituitary of sockeye salmon. *J.*  
 1248 *Endocrinol.* 188, 417–423. <https://doi.org/10.1677/joe.1.06494>

1249 Amemiya, Y., Takahashi, A., Meguro, H., Kawauchi, H., 1999a. Molecular cloning of  
 1250 lungfish proopiomelanocortin cDNA. *Gen. Comp. Endocrinol.* 115, 415–421.  
 1251 <https://doi.org/10.1006/gcen.1999.7327>

1252 Amemiya, Y., Takahashi, A., Suzuki, N., Sasayama, Y., Kawauchi, H., 1999b. A newly  
 1253 characterized melanotropin in proopiomelanocortin in pituitaries of an elasmobranch,  
 1254 *Squalus acanthias*. *Gen. Comp. Endocrinol.* 114, 387–395.  
 1255 <https://doi.org/10.1006/gcen.1999.7256>

1256 Arai, M., Assil, I., Abou-Samra, A., 2001. Characterization of Three Corticotropin-Releasing  
 1257 Factor Receptors in Catfish: A Novel Third Receptor Is Predominantly Expressed in  
 1258 Pituitary and Urophysis. *Endocrinology* 142, 446–454.  
 1259 <https://doi.org/10.1210/en.142.1.446>

1260 Arai, R., Tajima, H., Tamaoki, B., 1969. In Vitro Transformation of Steroids by the Head the  
 1261 Body Kidney , and the Corpuscles of Stannius of the Rainbow Trout (*Salmo gairdneri*).  
 1262 *Gen. Comp. Endocrinol.* 12, 99–109.

1263 Arends, R.J., Mancera, J.M., Muñoz, J.L., Wendelaar Bonga, S.E., Flik, G., 1999. The stress  
 1264 response of the gilthead sea bream (*Sparus aurata* L.) to air exposure and confinement. *J.*  
 1265 *Endocrinol.* 163, 149–157. <https://doi.org/10.1677/joe.0.1630149>

1266 Arnold-Reed, D.E., Balment, R.J., 1994. Peptide hormones influence in vitro interrenal  
 1267 secretion of cortisol in the trout, *Oncorhynchus mykiss*. *Gen. Comp. Endocrinol.* 96, 85–  
 1268 91. <https://doi.org/10.1006/gcen.1994.1161>

1269 Arnold-Reed, D.E., Balment, R.J., 1989. Steroidogenic role of the caudal neurosecretory  
 1270 system in the flounder, *Platichthys flesus*. *Gen. Comp. Endocrinol.* 76, 267–273.  
 1271 [https://doi.org/10.1016/0016-6480\(89\)90158-5](https://doi.org/10.1016/0016-6480(89)90158-5)

1272 Arterbery, A.S., Deitcher, D.L., Bass, A.H., 2010. Corticosteroid receptor expression in a  
 1273 teleost fish that displays alternative male reproductive tactics. *Gen. Comp. Endocrinol.*  
 1274 165, 83–90. <https://doi.org/10.1016/j.ygcen.2009.06.004>

1275 Aruna, A., Nagarajan, G., Chang, C.F., 2015. The acute salinity changes activate the dual  
 1276 pathways of endocrine responses in the brain and pituitary of tilapia. *Gen. Comp.*  
 1277 *Endocrinol.* 211, 154–164. <https://doi.org/10.1016/j.ygcen.2014.12.005>

1278 Aruna, A., Nagarajan, G., Chang, C.F., 2012. Involvement of Corticotrophin-Releasing  
 1279 Hormone and Corticosteroid Receptors in the Brain-Pituitary-Gill of Tilapia During the  
 1280 Course of Seawater Acclimation. *J. Neuroendocrinol.* 24, 818–830.  
 1281 <https://doi.org/10.1111/j.1365-2826.2012.02282.x>

1282 Baker, B.I., Bird, D.J., Buckingham, J.C., 1996. In the trout, CRH and AVT synergize to  
 1283 stimulate ACTH release. *Regul. Pept.* 67, 207–210. [https://doi.org/10.1016/S0167-](https://doi.org/10.1016/S0167-0115(96)00130-9)  
 1284 [0115\(96\)00130-9](https://doi.org/10.1016/S0167-0115(96)00130-9)

1285 Baker, M.E., Katsu, Y., 2020. Progesterone: An enigmatic ligand for the mineralocorticoid  
 1286 receptor. *Biochem. Pharmacol.* 177. <https://doi.org/10.1016/j.bcp.2020.113976>

1287 Baker, M.E., Katsu, Y., 2019. Evolution of the Mineralocorticoid Receptor. *Vitam. Horm.*

1288 109, 17–36. <https://doi.org/10.1016/bs.vh.2018.10.009>

1289 Balm, P.H.M., Lambert, J.D.G., Wendelaar Bonga, S.E., 1989. Corticosteroid biosynthesis in  
 1290 the interrenal cells of the teleost fish, *Oreochromis mossambicus*. *Gen. Comp.*  
 1291 *Endocrinol.* 76, 53–62. [https://doi.org/10.1016/0016-6480\(89\)90032-4](https://doi.org/10.1016/0016-6480(89)90032-4)

1292 Bamberger, C.M., Bamberger, A.M., De Castro, M., Chrousos, G.P., 1995. Glucocorticoid  
 1293 receptor  $\beta$ , a potential endogenous inhibitor of glucocorticoid action in humans. *J. Clin.*  
 1294 *Invest.* 95, 2435–2441. <https://doi.org/10.1172/JCI117943>

1295 Barkataki, S., Aluru, N., Li, M., Lin, L., Christie, H., Vijayan, M.M., Leatherland, J.F., 2011.  
 1296 Cortisol inhibition of 17 $\beta$ -estradiol secretion by rainbow trout ovarian follicles involves  
 1297 modulation of star and P450scc gene expression. *J. Aquac. Res. Dev.* 1–8.  
 1298 <https://doi.org/10.4172/2155-9546.S2-001>

1299 Barry, T.P., Riebe, J.D., Parrish, J.J., Malison, J.A., 1997. Effects of 17 $\alpha$ ,20 $\beta$ -dihydroxy-4-  
 1300 pregnen-3-one on cortisol production by rainbow trout interrenal tissue in vitro. *Gen.*  
 1301 *Comp. Endocrinol.* 107, 172–181. <https://doi.org/10.1006/gcen.1997.6910>

1302 Barton, B.A., 2002. Stress in fishes: A diversity of responses with particular reference to  
 1303 changes in circulating corticosteroids. *Integr. Comp. Biol.* 42, 517–525.  
 1304 <https://doi.org/10.1093/icb/42.3.517>

1305 Berg, H., Modig, C., Olsson, P.E., 2004. 17 $\beta$ -estradiol induced vitellogenesis is inhibited  
 1306 by cortisol at the post-transcriptional level in Arctic char (*Salvelinus alpinus*). *Reprod.*  
 1307 *Biol. Endocrinol.* 2, 1–10. <https://doi.org/10.1186/1477-7827-2-62>

1308 Bern, H.A., Takasugi, N., 1962. The caudal neurosecretory of fishes. *Gen. Comp. Endocrinol.*  
 1309 2, 96–110.

1310 Bernier, N.J., Alderman, S.L., Bristow, E.N., 2008. Heads or tails? Stressor-specific  
 1311 expression of corticotropin-releasing factor and urotensin I in the preoptic area and  
 1312 caudal neurosecretory system of rainbow trout. *J. Endocrinol.* 196, 637–648.

1313 <https://doi.org/10.1677/JOE-07-0568>

1314 Bernier, N.J., Craig, P.M., 2005. CRF-related peptides contribute to stress response and  
 1315 regulation of appetite in hypoxic rainbow trout. *Am. J. Physiol. - Regul. Integr. Comp.*  
 1316 *Physiol.* 289, 982–990. <https://doi.org/10.1152/ajpregu.00668.2004>

1317 Bernier, N.J., Flik, G., Klaren, P.H.M., 2009. Regulation And Contribution Of Corticotropic,  
 1318 Melanotropic And Thyrotropic Axes To The Stress Response In Fishes, in: *Fish*  
 1319 *Physiology*. Elsevier Inc., pp. 235–311. [https://doi.org/10.1016/S1546-5098\(09\)28006-X](https://doi.org/10.1016/S1546-5098(09)28006-X)

1320 Bernier, N.J., Lin, X., Peter, R.E., 1999. Differential expression of corticotropin-releasing  
 1321 factor (CRF) and urotensin I precursor genes, and evidence of CRF gene expression  
 1322 regulated by cortisol in goldfish brain. *Gen. Comp. Endocrinol.* 116, 461–477.  
 1323 <https://doi.org/10.1006/gcen.1999.7386>

1324 Billard, R., 1981. Stress, environment and reproduction in teleost fish, in: *Stress and Fish*.  
 1325 Academic Press, pp. 185–208.

1326 Biran, J., Golan, M., Mizrahi, N., Ogawa, S., Parhar, I.S., Levavi-Sivan, B., 2014. LPXRFa,  
 1327 the piscine ortholog of GnIH, and LPXRF receptor positively regulate gonadotropin  
 1328 secretion in tilapia (*Oreochromis niloticus*). *Endocrinology* 155, 4391–4401.  
 1329 <https://doi.org/10.1210/en.2013-2047>

1330 Bøgevik, A.S., Natário, S., Karlsen, O., Thorsen, A., Hamre, K., Rosenlund, G., Norberg, B.,  
 1331 2012. The effect of dietary lipid content and stress on egg quality in farmed Atlantic cod  
 1332 *Gadus morhua*. *J. Fish Biol.* 81, 1391–1405. [https://doi.org/10.1111/j.1095-](https://doi.org/10.1111/j.1095-8649.2012.03425.x)  
 1333 [8649.2012.03425.x](https://doi.org/10.1111/j.1095-8649.2012.03425.x)

1334 Bond, H., Kelsall, C., Teitsma, C., Balment, R., 1999. In teleost fish the caudal  
 1335 neurosecretory system (CNSS) affords pituitary-independent control of cortisol  
 1336 secretion. *Comp Biochem Physiol* 124A, S89.

1337 Bond, H., Warne, J.M., Balment, R.J., 2007. Effect of acute restraint on hypothalamic pro-

1338 vasotocin mRNA expression in flounder, *Platichthys flesus*. *Gen. Comp. Endocrinol.*  
 1339 153, 221–227. <https://doi.org/10.1016/j.ygcen.2007.03.014>  
 1340 Booth, R., Kieffer, J., Tufts, B., Davidson, K., Bielak, A., 1995. Effects of late-season catch  
 1341 and release angling on anaerobic metabolism, acid-base status, survival, and gamete  
 1342 viability in wild Atlantic salmon (*Salmo salar*). *J. Can. des Sci. halieutiques Aquat.* 52,  
 1343 283–290. <https://doi.org/10.1083/jcb.200508014>  
 1344 Borowsky, B., Kuhn, C.M., 1992. D1 and D2 dopamine receptors stimulate hypothalamo-  
 1345 pituitary-adrenal activity in rats. *Neuropharmacology* 31, 671–678.  
 1346 [https://doi.org/10.1016/0028-3908\(92\)90145-F](https://doi.org/10.1016/0028-3908(92)90145-F)  
 1347 Bradford, C., Taylor, M., 1987. Semilunar Changes in Estradiol and Cortisol Coincident with  
 1348 Gonadal Maturation and Spawning in the Killifish *Fundulus heteroclitus*. *Gen. Comp.*  
 1349 *Endocrinol.* 66, 71–78.  
 1350 Branco, G.S., Melo, A.G., Ricci, J.M.B., Digmayer, M., de Jesus, L.W.O., Habibi, H.R.,  
 1351 Nóbrega, R.H., 2019. Effects of GnRH and the dual regulatory actions of GnIH in the  
 1352 pituitary explants and brain slices of *Astyanax altiparanae* males. *Gen. Comp.*  
 1353 *Endocrinol.* 273, 209–217. <https://doi.org/10.1016/j.ygcen.2018.08.006>  
 1354 Bry, C., 1985. Plasma cortisol levels of female rainbow trout (*Salmo gairdneri*) at the end of  
 1355 the reproductive cycle: Relationship with oocyte stages. *Gen. Comp. Endocrinol.* 57, 47–  
 1356 52. [https://doi.org/10.1016/0016-6480\(85\)90199-6](https://doi.org/10.1016/0016-6480(85)90199-6)  
 1357 Bury, N.R., 2017. The evolution, structure and function of the ray finned fish (Actinopterygii)  
 1358 glucocorticoid receptors. *Gen. Comp. Endocrinol.* 251, 4–11.  
 1359 <https://doi.org/10.1016/j.ygcen.2016.06.030>  
 1360 Bury, N.R., Sturm, A., Le Rouzic, P., Lethimonier, C., Ducouret, B., Guiguen, Y., Robinson-  
 1361 Rechavi, M., Laudet, V., Rafestin-Oblin, M.E., Prunet, P., 2003. Evidence for two  
 1362 distinct functional glucocorticoid receptors in teleost fish. *J. Mol. Endocrinol.* 31, 141–

1363 156. <https://doi.org/10.1677/jme.0.0310141>

1364 Butler, D.G., 1973. Structure and function of the adrenal gland of fishes. *Integr. Comp. Biol.*

1365 13, 839–879. <https://doi.org/10.1093/icb/13.3.839>

1366 Butler, D.G., 1968. Hormonal control of gluconeogenesis in the North American eel

1367 (*Anguilla rostrata*). *Gen. Comp. Endocrinol.* 10, 85–91. [https://doi.org/10.1016/0016-](https://doi.org/10.1016/0016-6480(68)90012-9)

1368 6480(68)90012-9

1369 Cahoreau, C., Klett, D., Combarnous, Y., 2015. Structure-function relationships of

1370 glycoprotein hormones and their subunits' ancestors. *Front. Endocrinol. (Lausanne)*. 6,

1371 1–14. <https://doi.org/10.3389/fendo.2015.00026>

1372 Campbell, C., Fostier, A., Jalabert, B., Truscott, B., 1980. Identification and quantification of

1373 steroids in the serum of rainbow trout during spermiation and oocyte maturation. *J*

1374 *Endocrinol* 85, 371–378.

1375 Campbell, C.M., Walsh, J.M., Idler, D.R., 1976. Steroids in the plasma of the winter flounder

1376 (*Pseudopleuronectes americanus walbaum*). A seasonal study and investigation of

1377 steroid involvement in oocyte maturation. *Gen. Comp. Endocrinol.* 29, 14–20.

1378 [https://doi.org/10.1016/0016-6480\(76\)90002-2](https://doi.org/10.1016/0016-6480(76)90002-2)

1379 Campbell, P.M., Pottinger, T.G., Sumpter, J.P., 1994. Preliminary evidence that chronic

1380 confinement stress reduces the quality of gametes produced by brown and rainbow trout.

1381 *Aquaculture* 120, 151–169. [https://doi.org/10.1016/0044-8486\(94\)90230-5](https://doi.org/10.1016/0044-8486(94)90230-5)

1382 Campbell, P.M., Pottinger, T.G., Sumpter, J.P., 1992. Stress Reduces the Quality of Gametes

1383 Produced by Rainbow Trout. *Biol. Reprod.* 47, 1140–1150.

1384 <https://doi.org/10.1095/biolreprod47.6.1140>

1385 Cardoso, J.C.R., Bergqvist, C.A., Félix, R.C., Larhammar, D., 2016. Corticotropin-releasing

1386 hormone family evolution: Five ancestral genes remain in some lineages. *J. Mol.*

1387 *Endocrinol.* 57, 73–86. <https://doi.org/10.1530/JME-16-0051>

1388 Cardoso, J.C.R., Félix, R.C., Bergqvist, C.A., Larhammar, D., 2014. New insights into the  
 1389 evolution of vertebrate CRH (corticotropin-releasing hormone) and invertebrate DH44  
 1390 (diuretic hormone 44) receptors in metazoans. *Gen. Comp. Endocrinol.* 209, 162–170.  
 1391 <https://doi.org/10.1016/j.ygcen.2014.09.004>  
 1392 Cardoso, J.C.R., Laiz-Carrión, R., Louro, B., Silva, N., Canario, A.V.M., Mancera, J.M.,  
 1393 Power, D.M., 2011. Divergence of duplicate POMC genes in gilthead sea bream *Sparus*  
 1394 *auratus*. *Gen. Comp. Endocrinol.* 173, 396–404.  
 1395 <https://doi.org/10.1016/j.ygcen.2010.12.001>  
 1396 Carpenter, R.E., Maruska, K.P., Becker, L., Fernald, R.D., 2014. Social opportunity rapidly  
 1397 regulates expression of CRF and CRF receptors in the brain during social ascent of a  
 1398 teleost fish, *Astatotilapia burtoni*. *PLoS One* 9.  
 1399 <https://doi.org/10.1371/journal.pone.0096632>  
 1400 Carragher, J., Pankhurst, N., 1991. Stress and reproduction in a commercially important  
 1401 marine fish, *Pagrus auratus* (Sparidae), in: Scott, A., Sumpter, J., Kime, D., Rolfe, M.  
 1402 (Eds.), *Proceedings of the Fourth International Symposium on Reproductive Physiology*  
 1403 *of Fish*. pp. 253–255.  
 1404 Carragher, J.F., Sumpter, J.P., 1990. The effect of cortisol on the secretion of sex steroids  
 1405 from cultured ovarian follicles of rainbow trout. *Gen. Comp. Endocrinol.* 77, 403–407.  
 1406 [https://doi.org/10.1016/0016-6480\(90\)90230-J](https://doi.org/10.1016/0016-6480(90)90230-J)  
 1407 Carragher, J.F., Sumpter, J.P., Pottinger, T.G., Pickering, A.D., 1989. The deleterious effects  
 1408 of cortisol implantation on reproductive function in two species of trout, *Salmo trutta* L.  
 1409 and *Salmo gairdneri* Richardson. *Gen. Comp. Endocrinol.* 76, 310–321.  
 1410 [https://doi.org/10.1016/0016-6480\(89\)90163-9](https://doi.org/10.1016/0016-6480(89)90163-9)  
 1411 Carruth, L.L., Dores, R.M., Maldonado, T.A., Norris, D.O., Ruth, T., Jones, R.E., 2000.  
 1412 Elevation of plasma cortisol during the spawning migration of landlocked kokanee

1413 salmon (*Oncorhynchus nerka kennerlyi*). *Comp. Biochem. Physiol. - C Pharmacol.*  
 1414 *Toxicol. Endocrinol.* 127, 123–131. [https://doi.org/10.1016/S0742-8413\(00\)00140-7](https://doi.org/10.1016/S0742-8413(00)00140-7)  
 1415 Carsia, R. V., Weber, H., Perez, F.M., 1986. Corticotropin-releasing factor stimulates the  
 1416 release of adrenocorticotropin from domestic fowl pituitary cells. *Endocrinology* 118,  
 1417 143–148. <https://doi.org/10.1210/endo-118-1-143>  
 1418 Castranova, D.A., King, W., Woods, L.C., 2005. The effects of stress on androgen  
 1419 production, spermiation response and sperm quality in high and low cortisol responsive  
 1420 domesticated male striped bass. *Aquaculture* 246, 413–422.  
 1421 <https://doi.org/10.1016/j.aquaculture.2004.12.019>  
 1422 Chabbi, A., Ganesh, C.B., 2014. Glucocorticoid synthesis inhibitor metyrapone blocks stress-  
 1423 induced suppression along luteinizing hormone secreting cells-ovary axis in the fish  
 1424 *Oreochromis mossambicus*. *J. Exp. Zool. Part A Ecol. Genet. Physiol.* 321, 125–134.  
 1425 <https://doi.org/10.1002/jez.1842>  
 1426 Chabbi, A., Ganesh, C.B., 2013.  $\beta$ -Endorphin-induced Inhibition of Vitellogenic Follicular  
 1427 Growth in the Fish *Oreochromis mossambicus*: Evidence for Opioidergic Mediation of  
 1428 Ovarian Stress Response. *J. Exp. Zool. Part A Ecol. Genet. Physiol.* 319, 156–165.  
 1429 <https://doi.org/10.1002/jez.1781>  
 1430 Chabbi, A., Ganesh, C.B., 2012. Stress-induced inhibition of recruitment of ovarian follicles  
 1431 for vitellogenic growth and interruption of spawning cycle in the fish *Oreochromis*  
 1432 *mossambicus*. *Fish Physiol. Biochem.* 38, 1521–1532. [https://doi.org/10.1007/s10695-](https://doi.org/10.1007/s10695-012-9643-z)  
 1433 [012-9643-z](https://doi.org/10.1007/s10695-012-9643-z)  
 1434 Chan, D.K.O., Woo, N.Y.S., 1978. Effect of cortisol on the metabolism of the eel, *Anguilla*  
 1435 *japonica*. *Gen. Comp. Endocrinol.* 35, 205–215. [https://doi.org/10.1016/0016-](https://doi.org/10.1016/0016-6480(78)90064-3)  
 1436 [6480\(78\)90064-3](https://doi.org/10.1016/0016-6480(78)90064-3)  
 1437 Chand, D., Lovejoy, D.A., 2011. Stress and reproduction: Controversies and challenges. *Gen.*

1438 Comp. Endocrinol. 171, 253–257. <https://doi.org/10.1016/j.ygcen.2011.02.022>

1439 Chatterjee, A., Hsieh, Y.L., Yu, J.Y.L., 2001. Molecular cloning of cDNA encoding thyroid  
 1440 stimulating hormone  $\beta$  subunit of bighead carp *Aristichthys nobilis* and regulation of its  
 1441 gene expression. Mol. Cell. Endocrinol. 174, 1–9. [https://doi.org/10.1016/S0303-](https://doi.org/10.1016/S0303-7207(01)00392-6)  
 1442 7207(01)00392-6

1443 Chatzopoulou, A., Roy, U., Meijer, A.H., Alia, A., Spaink, H.P., Schaaf, M.J.M., 2015.  
 1444 Transcriptional and metabolic effects of glucocorticoid receptor  $\alpha$  and  $\beta$  signaling in  
 1445 zebrafish. Endocrinology 156, 1757–1769. <https://doi.org/10.1210/en.2014-1941>

1446 Chatzopoulou, A., Schoonheim, P.J., Torraca, V., Meijer, A.H., Spaink, H.P., Schaaf, M.J.M.,  
 1447 2017. Functional analysis reveals no transcriptional role for the glucocorticoid receptor  
 1448  $\beta$ -isoform in zebrafish. Mol. Cell. Endocrinol. 447, 61–70.  
 1449 <https://doi.org/10.1016/j.mce.2017.02.036>

1450 Chellappa, S., Huntingford, F., 1989. Depletion of energy reserves during reproductive  
 1451 aggression in male three-spined stickleback, *Gasterosteus aculeatus* L. J. Fish Biol. 35,  
 1452 315–316.

1453 Chen, C.C., Fernald, R.D., 2008. Sequences, expression patterns and regulation of the  
 1454 corticotropin-releasing factor system in a teleost. Gen. Comp. Endocrinol. 157, 148–155.  
 1455 <https://doi.org/10.1016/j.ygcen.2008.04.003>

1456 Chen, J., Peng, C., Yu, Z., Xiao, L., Yu, Q., Li, S., Zhang, H., Lin, H., Zhang, Y., 2020. The  
 1457 Administration of Cortisol Induces Female-to-Male Sex Change in the Protogynous  
 1458 Orange-Spotted Grouper, *Epinephelus coioides*. Front. Endocrinol. (Lausanne). 11, 1–  
 1459 15. <https://doi.org/10.3389/fendo.2020.00012>

1460 Choi, Y.J., Habibi, H.R., Kil, G.S., Jung, M.M., Choi, C.Y., 2017. Effect of cortisol on  
 1461 gonadotropin inhibitory hormone (GnIH) in the cinnamon clownfish, *Amphiprion*  
 1462 *melanopus*. Biochem. Biophys. Res. Commun. 485, 342–348.

1463 <https://doi.org/10.1016/j.bbrc.2017.02.078>

1464 Choi, Y.J., Kim, N.N., Habibi, H.R., Choi, C.Y., 2016. Effects of gonadotropin inhibitory  
 1465 hormone or gonadotropin-releasing hormone on reproduction-related genes in the  
 1466 protandrous cinnamon clownfish, *Amphiprion melanopus*. *Gen. Comp. Endocrinol.* 235,  
 1467 89–99. <https://doi.org/10.1016/j.ygcen.2016.06.010>

1468 Chowdhury, I., Chien, J.T., Chatterjee, A., Yu, J.Y.L., 2004. In vitro effects of mammalian  
 1469 leptin, neuropeptide-Y,  $\beta$ -endorphin and galanin on transcript levels of thyrotropin  $\beta$  and  
 1470 common  $\alpha$  subunit mRNAs in the pituitary of bighead carp (*aristichthys nobilis*). *Comp.*  
 1471 *Biochem. Physiol. - B Biochem. Mol. Biol.* 139, 87–98.  
 1472 <https://doi.org/10.1016/j.cbpc.2004.06.007>

1473 Ciani, E., Fontaine, R., Maugars, G., Nourizadeh-Lillabadi, R., Andersson, E., Bogerd, J., von  
 1474 Krogh, K., Weltzien, F.A., 2020. GnRH receptor *gnhr2bb $\alpha$*  is expressed exclusively in  
 1475 *lhb*-expressing cells in Atlantic salmon male parr. *Gen. Comp. Endocrinol.* 285.  
 1476 <https://doi.org/10.1016/j.ygcen.2019.113293>

1477 Clearwater, S.J., Pankhurst, N.W., 1997. The response to capture and confinement stress of  
 1478 plasma cortisol, plasma sex steroids and vitellogenic oocytes in the marine teleost, red  
 1479 gurnard. *J. Fish Biol.* 50, 429–441. <https://doi.org/10.1006/jfbi.1996.0311>

1480 Cleary, J.J., Pankhurst, N.W., Battaglione, S.C., 2000. The Effect of Capture and Handling  
 1481 Stress on Plasma Steroid Levels and Gonadal Condition in Wild and Farmed Snapper  
 1482 *Pagrus auratus* (Sparidae). *J. World Aquac. Soc.* 31, 558–569.  
 1483 <https://doi.org/10.1111/j.1749-7345.2000.tb00905.x>

1484 Clevestam, P.D., Ogonowski, M., Sjöberg, N.B., Wickström, H., 2011. Too short to spawn?  
 1485 Implications of small body size and swimming distance on successful migration and  
 1486 maturation of the European eel *Anguilla anguilla*. *J. Fish Biol.* 78, 1073–1089.  
 1487 <https://doi.org/10.1111/j.1095-8649.2011.02920.x>

1488 Colombo, L., Bern, H.A., Pieprzyk, J., Johnson, D.W., 1973. Biosynthesis of 11-  
1489 deoxycorticosteroids by teleost ovaries and discussion of their possible role in oocyte  
1490 maturation and ovulation. *Gen. Comp. Endocrinol.* 21, 168–178.  
1491 [https://doi.org/10.1016/0016-6480\(73\)90168-8](https://doi.org/10.1016/0016-6480(73)90168-8)

1492 Colombo, L., Colombo Belvedere, P., Arcarese, G., 1978. Emergence of Ovarian 11-  
1493 Deoxycorticosteroid Biosynthesis at ovulation time in the sea bass, *Dicentrarchus labrax*  
1494 L. *Ann Biol anim Bioch Biophys* 18, 937–941.

1495 Cone, R.D., 2006. Studies on the physiological functions of the melanocortin system. *Endocr.*  
1496 *Rev.* 27, 736–749. <https://doi.org/10.1210/er.2006-0034>

1497 Consten, D., Bogerd, J., Komen, J., Lambert, J.G.D., Goos, H.J.T., 2001a. Long-Term  
1498 Cortisol Treatment Inhibits Pubertal Development in Male Common Carp, *Cyprinus*  
1499 *carpio* L. *Biol. Reprod.* 64, 1063–1071. <https://doi.org/10.1095/biolreprod64.4.1063>

1500 Consten, D., Lambert, J., Goos, H.J.T., 2000. Inhibitory effects of cortisol on in vivo and in  
1501 vitro androgen secretion in male common carp, *Cyprinus carpio* L., in: Norberg, B.,  
1502 Kjesbu, O., Taranger, G., Andersson, E., Stefansson, S. (Eds.), *Proceedings of the 6th*  
1503 *International Symposium on the Reproductive Physiology of Fish.* p. 192.

1504 Consten, D., Lambert, J.G.D., Goos, H.J.T., 2001b. Cortisol affects testicular development in  
1505 male common carp, *Cyprinus carpio* L., but not via an effect on LH secretion. *Comp.*  
1506 *Biochem. Physiol. - B Biochem. Mol. Biol.* 129, 671–677.  
1507 [https://doi.org/10.1016/S1096-4959\(01\)00368-2](https://doi.org/10.1016/S1096-4959(01)00368-2)

1508 Consten, D., Lambert, J.G.D., Komen, H., Goos, H.J.T., 2002. Corticosteroids Affect the  
1509 Testicular Androgen Production in Male Common Carp (*Cyprinus carpio* L.). *Biol.*  
1510 *Reprod.* 66, 106–111. <https://doi.org/10.1095/biolreprod66.1.106>

1511 Contreras-Sanchez, W., Schreck, C., Fitzpatrick, M., Pereira, C., 1998. Effects of stress on the  
1512 reproductive performance of rainbow trout (*Oncorhynchus mykiss*). *Biol. Reprod.* 58,

1513 439–447.

1514 Cooke, S., Philipp, D., Schreer, J., McFinley, R., 2000. Locomotory impairment of nesting  
 1515 male largemouth bass following catch-and-release angling. *North Am. J. Fish. Manag.*  
 1516 20, 968–977.

1517 Cornett, L.E., Kang, S.W., Kuenzel, W.J., 2012. A possible mechanism contributing to the  
 1518 synergistic action of vasotocin (VT) and corticotropin-releasing hormone (CRH)  
 1519 receptors on corticosterone release in birds. *Gen. Comp. Endocrinol.* 188, 46–53.  
 1520 <https://doi.org/10.1016/j.ygcen.2013.02.032>

1521 Corriero, A., Zupa, R., Bello, G., Mylonas, C.C., Deflorio, M., Genovese, S., Basilone, G.,  
 1522 Buscaino, G., Buffa, G., Pousis, C., De Metrio, G., Santamaria, N., 2011. Evidence that  
 1523 severe acute stress and starvation induce rapid atresia of ovarian vitellogenic follicles in  
 1524 Atlantic bluefin tuna, *Thunnus thynnus* (L.) (Osteichthyes: Scombridae). *J. Fish Dis.* 34,  
 1525 853–860. <https://doi.org/10.1111/j.1365-2761.2011.01303.x>

1526 Craig, P.M., Al-Timimi, H., Bernier, N.J., 2005. Differential increase in forebrain and caudal  
 1527 neurosecretory system CRF and urotensin I gene expression associated with seawater  
 1528 transfer in rainbow trout. *Endocrinology* 146, 3851–3860.  
 1529 <https://doi.org/10.1210/en.2005-0004>

1530 Crim, L.W., Peter, R.E., Billard, R., 1981. Onset of gonadotropic hormone accumulation in  
 1531 the immature trout pituitary gland in response to estrogen or aromatizable androgen  
 1532 steroid hormones. *Gen. Comp. Endocrinol.* 44, 374–381. [https://doi.org/10.1016/0016-](https://doi.org/10.1016/0016-6480(81)90015-0)  
 1533 [6480\(81\)90015-0](https://doi.org/10.1016/0016-6480(81)90015-0)

1534 D’Agata, R., Cavagnini, F., Invitti, C., Mongioi, A., Fossati, R., Scapagnini, U., Müller, E.,  
 1535 1984. Effect of CRF on the release of anterior pituitary hormones in normal subjects and  
 1536 patients with Cushing’s disease. *Pharmacol. Res. Commun.* 16, 303–311.

1537 Dahlgren, U., 1914. The electric motor nerve centers in the skate (Rajidae). *Science* (80- ).

1538 40, 862–863.

1539 Dave, G., Johansson-Sjöbeck, M.L., Larsson, Å., Lewander, K., Lidman, U., 1979. Effects of  
 1540 cortisol on the fatty acid composition of the total blood plasma lipids in the European  
 1541 eel, *Anguilla anguilla* L. *Comp. Biochem. Physiol. -- Part A Physiol.* 64, 37–40.  
 1542 [https://doi.org/10.1016/0300-9629\(79\)90427-4](https://doi.org/10.1016/0300-9629(79)90427-4)

1543 De Beer, G., 1923. The Evolution of the Pituitary. *Ref. Modul. Biomed. Sci.* 271–291.  
 1544 <https://doi.org/10.1016/b978-0-12-801238-3.95790-8>

1545 De Groef, B., Van Der Geyten, S., Darras, V.M., Kühn, E.R., 2006. Role of corticotropin-  
 1546 releasing hormone as a thyrotropin-releasing factor in non-mammalian vertebrates. *Gen.*  
 1547 *Comp. Endocrinol.* 146, 62–68. <https://doi.org/10.1016/j.ygcen.2005.10.014>

1548 De Montalembert, G., Jalabert, B., Bry, C., 1978. Precocious induction of maturation and  
 1549 ovulation in northern pike (*Esox lucius*). *Ann. Biol. Anim. Biochim. Biophys.* 18, 969–  
 1550 975. <https://doi.org/10.1051/rnd:19780533>

1551 de Roux, N., Genin, E., Carel, J.-C., Matsuda, F., Chaussain, J.-L., Milgrom, E., 2003.  
 1552 Hypogonadotropic hypogonadism due to loss of function of the KiSS1-derived peptide  
 1553 receptor GPR54. *Proc. Natl. Acad. Sci.* 100, 10972–10976.  
 1554 <https://doi.org/10.1073/pnas.1834399100>

1555 De Souza, F.S.J., Bumashny, V.F., Low, M.J., Rubinstein, M., 2005. Subfunctionalization of  
 1556 expression and peptide domains following the ancient duplication of the  
 1557 proopiomelanocortin gene in teleost fishes. *Mol. Biol. Evol.* 22, 2417–2427.  
 1558 <https://doi.org/10.1093/molbev/msi236>

1559 Dehal, P., Boore, J.L., 2005. Two rounds of whole genome duplication in the ancestral  
 1560 vertebrate. *PLoS Biol.* 3. <https://doi.org/10.1371/journal.pbio.0030314>

1561 Dickhoff, W., 1989. Salmonids and annual fishes: death after sex., in: *Development,*  
 1562 *Maturation and Senescence of Neuroendocrine Systems: A Comparative Approach.*

1563 Academic Press, New York, pp. 253–266.

1564 Dinan, T.G., 1996. Serotonin and the regulation of hypothalamic-pituitary-adrenal axis  
 1565 function. *Life Sci.* 58, 1683–1694. [https://doi.org/10.1016/0024-3205\(96\)00066-5](https://doi.org/10.1016/0024-3205(96)00066-5)

1566 Donaldson, E., Fagerlund, U.H., 1969. Cortisol secretion rate in gonadectomized female  
 1567 sockeye (*Oncorhynchus nerka*): effects of estrogen and cortisol treatment. *J. Fish Res.*  
 1568 Board Canada 26, 1789–1799.

1569 Donaldson, E., Fagerlund, U., 1970. Effect of sexual maturation and gonadectomy at sexual  
 1570 maturity on cortisol secretion rate in sockeye salmon (*Oncorhynchus nerka*). *J. l'Office*  
 1571 *des Rech. sur les pêcheries du Canada* 27, 2287–2296.

1572 Donaldson, E., McBride, J., 1974. Effect of ACTH and salmon gonadotropin on interrenal  
 1573 and thyroid activity of gonadectomized adult sockeye salmon (*Oncorhynchus nerka*). *J.*  
 1574 *l'Office sur les pêcheries du Canada* 31, 1211–1214.

1575 Donaldson, E.M., Fagerlund, U.H.M., 1972. Corticosteroid dynamics in Pacific salmon. *Gen.*  
 1576 *Comp. Endocrinol.* 3, 254–265. [https://doi.org/10.1016/0016-6480\(72\)90155-4](https://doi.org/10.1016/0016-6480(72)90155-4)

1577 Does, R., 2017. *The evolution of the pituitary*. Elsevier.

1578 Does, R., Lecaude, S., Bauer, D., Danielson, P., 2002. Analyzing the evolution of the  
 1579 opioid/orphanin gene family. *Mass Spectrom. Rev.* 21, 220–243.  
 1580 <https://doi.org/10.1002/mas.10029>

1581 Does, R.M., 2016. Hypothesis and Theory: Revisiting Views on the Co-evolution of the  
 1582 Melanocortin Receptors and the Accessory Proteins, MRAP1 and MRAP2. *Front.*  
 1583 *Endocrinol. (Lausanne)*. 7, 1–12. <https://doi.org/10.3389/fendo.2016.00079>

1584 Does, R.M., Baron, A.J., 2011. Evolution of POMC: Origin, phylogeny, posttranslational  
 1585 processing, and the melanocortins. *Ann. N. Y. Acad. Sci.* 1220, 34–48.  
 1586 <https://doi.org/10.1111/j.1749-6632.2010.05928.x>

1587 Does, R.M., Garcia, Y., 2015. Views on the co-evolution of the melanocortin-2 receptor,

1588 MRAPs, and the hypothalamus/pituitary/adrenal-interrenal axis. *Mol. Cell. Endocrinol.*  
 1589 408, 12–22. <https://doi.org/10.1016/j.mce.2014.12.022>  
 1590 Does, R.M., Lecaude, S., 2005. Trends in the evolution of the proopiomelanocortin gene.  
 1591 *Gen. Comp. Endocrinol.* 142, 81–93. <https://doi.org/10.1016/j.ygcen.2005.02.003>  
 1592 Does, R.M., Liang, L., Davis, P., Thomas, A.L., Petko, B., 2016. Melanocortin receptors:  
 1593 Evolution of ligand selectivity for melanocortin peptides. *J. Mol. Endocrinol.* 56, T119–  
 1594 T133. <https://doi.org/10.1530/JME-15-0292>  
 1595 Does, R.M., Londraville, R.L., Prokop, J., Davis, P., Dewey, N., Lesinski, N., 2014.  
 1596 Molecular evolution of GPCRs: Melanocortin/melanocortin receptors. *J. Mol.*  
 1597 *Endocrinol.* 52. <https://doi.org/10.1530/JME-14-0050>  
 1598 Dufour, S., Delerue-Le Belle, N., Fontaine, Y.A., 1983. Effects of steroid hormones on  
 1599 pituitary immunoreactive gonadotropin in European freshwater eel, *Anguilla anguilla* L.  
 1600 *Gen. Comp. Endocrinol.* 52, 190–197. [https://doi.org/10.1016/0016-6480\(83\)90112-0](https://doi.org/10.1016/0016-6480(83)90112-0)  
 1601 Dufour, S., Quérat, B., Tostivint, H., Pasqualini, C., Vaudry, C., Rousseau, K., 2020. Origin  
 1602 and evolution of the neuroendocrine control of reproduction in vertebrates, with special  
 1603 focus on genome and gene duplications. *Physiol. Rev.* 100, 869–943.  
 1604 <https://doi.org/10.1037/0033-2909.126.1.78>  
 1605 Dufour, S., Sebert, M.-E., Weltzien, F.-A., Rousseau, K., Pasqualini, C., 2010.  
 1606 Neuroendocrine control by dopamine of teleost reproduction. *J. Fish Biol.* 76.  
 1607 <https://doi.org/10.1111/j.1095-8649.2009.02499.x>  
 1608 Dufour, S., Weltzien, F.A., Sebert, M.E., Le Belle, N., Vidal, B., Vernier, P., Pasqualini, C.,  
 1609 2005. Dopaminergic inhibition of reproduction in teleost fishes: Ecophysiological and  
 1610 evolutionary implications. *Ann. N. Y. Acad. Sci.* 1040, 9–21.  
 1611 <https://doi.org/10.1196/annals.1327.002>  
 1612 Eriksen, M.S., Bakken, M., Espmark, Å., Braastad, B.O., Salte, R., 2006. Prespawning stress

1613 in farmed Atlantic salmon *Salmo salar*: Maternal cortisol exposure and hyperthermia  
 1614 during embryonic development affect offspring survival, growth and incidence of  
 1615 malformations. *J. Fish Biol.* 69, 114–129. [https://doi.org/10.1111/j.1095-](https://doi.org/10.1111/j.1095-8649.2006.01071.x)  
 1616 8649.2006.01071.x  
 1617 Eriksen, M.S., Espmark, Å., Braastad, B.O., Salte, R., Bakken, M., 2007. Long-term effects  
 1618 of maternal cortisol exposure and mild hyperthermia during embryogeny on survival,  
 1619 growth and morphological anomalies in farmed Atlantic salmon *Salmo salar* offspring. *J.*  
 1620 *Fish Biol.* 70, 462–473. <https://doi.org/10.1111/j.1095-8649.2007.01317.x>  
 1621 Fagerlund, U.H.M., 1967. Plasma cortisol concentration in relation to stress in adult sockeye  
 1622 salmon during the freshwater stage of their life cycle. *Gen. Comp. Endocrinol.* 8, 197–  
 1623 207. [https://doi.org/10.1016/0016-6480\(67\)90066-4](https://doi.org/10.1016/0016-6480(67)90066-4)  
 1624 Fagerlund, U.H.M., Donaldson, E.M., 1969. The effect of androgens on the distribution and  
 1625 secretion of cortisol in gonadectomized male sockeye salmon (*Oncorhynchus nerka*).  
 1626 *Gen. Comp. Endocrinol.* 12, 438–448. [https://doi.org/10.1016/0016-6480\(69\)90160-9](https://doi.org/10.1016/0016-6480(69)90160-9)  
 1627 Faught, E., Aluru, N., Vijayan, M.M., 2016. The Molecular Stress Response, in: *Fish*  
 1628 *Physiology*. Elsevier Inc., pp. 113–166. [https://doi.org/10.1016/B978-0-12-802728-](https://doi.org/10.1016/B978-0-12-802728-8.00004-7)  
 1629 8.00004-7  
 1630 Faught, E., Vijayan, M.M., 2018. The mineralocorticoid receptor is essential for stress axis  
 1631 regulation in zebrafish larvae. *Sci. Rep.* 8, 1–11. [https://doi.org/10.1038/s41598-018-](https://doi.org/10.1038/s41598-018-36681-w)  
 1632 36681-w  
 1633 Fernandino, J.I., Hattori, R.S., 2019. Sex determination in Neotropical fish: Implications  
 1634 ranging from aquaculture technology to ecological assessment. *Gen. Comp. Endocrinol.*  
 1635 273, 172–183. <https://doi.org/10.1016/j.ygcen.2018.07.002>  
 1636 Fernandino, J.I., Hattori, R.S., Kishii, A., Strüssmann, C.A., Somoza, G.M., 2012. The  
 1637 cortisol and androgen pathways cross talk in high temperature-induced masculinization:

1638 The  $11\beta$ -hydroxysteroid dehydrogenase as a key enzyme. *Endocrinology* 153, 6003–  
 1639 6011. <https://doi.org/10.1210/en.2012-1517>  
 1640 Fernandino, J.I., Hattori, R.S., Moreno Acosta, O.D., Strüssmann, C.A., Somoza, G.M., 2013.  
 1641 Environmental stress-induced testis differentiation: Androgen as a by-product of cortisol  
 1642 inactivation. *Gen. Comp. Endocrinol.* 192, 36–44.  
 1643 <https://doi.org/10.1016/j.ygcen.2013.05.024>  
 1644 Flanagan, C.A., Manilall, A., 2017. Gonadotropin-releasing hormone (GnRH) receptor  
 1645 structure and GnRH binding. *Front. Endocrinol. (Lausanne)*. 8, 1–14.  
 1646 <https://doi.org/10.3389/fendo.2017.00274>  
 1647 Flik, G., Klaren, P.H.M., Van Den Burg, E.H., Metz, J.R., Huising, M.O., 2006. CRF and  
 1648 stress in fish. *Gen. Comp. Endocrinol.* 146, 36–44.  
 1649 <https://doi.org/10.1016/j.ygcen.2005.11.005>  
 1650 Fontaine, M., Bertrand, E., Lopez, E., Callamand, O., 1964. Sur la maturation des organes  
 1651 génitaux de l'Anguille femelle (*Anguilla anguilla* L.) et l'émission spontanée des oeufs  
 1652 en aquarium. *C R Acad Sc Paris* 259, 2907–2910.  
 1653 Foo, J.T.W., Lam, T.J., 1993a. Retardation of ovarian growth and depression of serum steroid  
 1654 levels in the tilapia, *Oreochromis mossambicus*, by cortisol implantation. *Aquaculture*  
 1655 115, 133–143. [https://doi.org/10.1016/0044-8486\(93\)90364-5](https://doi.org/10.1016/0044-8486(93)90364-5)  
 1656 Foo, J.T.W., Lam, T.J., 1993b. Serum cortisol response to handling stress and the effect of  
 1657 cortisol implantation on testosterone level in the tilapia, *Oreochromis mossambicus*.  
 1658 *Aquaculture* 115, 145–158. [https://doi.org/10.1016/0044-8486\(93\)90365-6](https://doi.org/10.1016/0044-8486(93)90365-6)  
 1659 Foster, G.D., Moon, T.W., 1986. Cortisol and liver metabolism of immature American eels,  
 1660 *Anguilla rostrata* (LeSueur). *Fish Physiol. Biochem.* 1, 113–124.  
 1661 <https://doi.org/10.1007/BF02290211>  
 1662 Frias, J., Ruiz, E., Ortega, E., 1999. Effect of corticotropin releasing factor injected into the

1663 Median Eminence on growth hormone secretion in male rats. *Neurochem. Res.* 24, 715–  
 1664 718. <https://doi.org/10.1023/A:1020719227235>  
 1665 Fryer, J., K, L., Rivier, J., 1983. Urotensin I, a CRF-like neuropeptide stimulates ACTH  
 1666 release from the teleost pituitary. *Endocrinology* 113, 2308–2310.  
 1667 Fryer, J., Lederis, K., Rivier, J., 1985. ACTH-releasing activity of urotensin I and ovine CRF:  
 1668 Interactions with arginine vasotocin, isotocin and arginine vasopressin. *Regul. Pept.* 11,  
 1669 11–15. [https://doi.org/10.1016/0167-0115\(85\)90026-6](https://doi.org/10.1016/0167-0115(85)90026-6)  
 1670 Fryer, J., Lederis, K., Rivier, J., 1984. Cortisol inhibits the ACTH-releasing activity of  
 1671 urotensin I, CRF and sauvagine observed with superfused goldfish pituitary cells.  
 1672 *Peptides* 5, 925–930. [https://doi.org/10.1016/0196-9781\(84\)90118-9](https://doi.org/10.1016/0196-9781(84)90118-9)  
 1673 Fryer, J.N., Lederis, K., 1986. Control of Corticotropin Secretion in Teleost Fishes. *Amer*  
 1674 *Zool* 26, 1017–1026.  
 1675 Fryer, J.N., Leung, E., 1982. Neurohypophysial hormonal control of cortisol secretion in the  
 1676 teleost *Carassius auratus*. *Gen. Comp. Endocrinol.* 48, 425–431.  
 1677 [https://doi.org/10.1016/0016-6480\(82\)90177-0](https://doi.org/10.1016/0016-6480(82)90177-0)  
 1678 Fuller, P.J., Yao, Y.Z., Jin, R., He, S., Martín-Fernández, B., Young, M.J., Smith, B.J., 2019.  
 1679 Molecular evolution of the switch for progesterone and spironolactone from  
 1680 mineralocorticoid receptor agonist to antagonist. *Proc. Natl. Acad. Sci. U. S. A.* 116,  
 1681 18578–18583. <https://doi.org/10.1073/pnas.1903172116>  
 1682 Funes, S., Hedrick, J.A., Vassileva, G., Markowitz, L., Abbondanzo, S., Golovko, A., Yang,  
 1683 S., Monsma, F.J., Gustafson, E.L., 2003. The KiSS-1 receptor GPR54 is essential for the  
 1684 development of the murine reproductive system. *Biochem. Biophys. Res. Commun.* 312,  
 1685 1357–1363. <https://doi.org/10.1016/j.bbrc.2003.11.066>  
 1686 Fuzzen, M.L.M., Bernier, N.J., Glen Van Der, K., 2011. Stress and Reproduction, Hormones  
 1687 and Reproduction of Vertebrates - Volume 1. Elsevier. <https://doi.org/10.1016/B978-0->

12-375009-9.10006-2

Galas, L., Raoult, E., Tonon, M.C., Okada, R., Jenks, B.G., Castaño, J.P., Kikuyama, S., Malagon, M., Roubos, E.W., Vaudry, H., 2009. TRH acts as a multifunctional hypophysiotropic factor in vertebrates. *Gen. Comp. Endocrinol.* 164, 40–50. <https://doi.org/10.1016/j.ygcen.2009.05.003>

Gambacciani, M., Yen, S., Rasmussen, D., 1988. GnRH stimulates ACTH and immunoreactive beta-endorphin release from the rat pituitary in vitro. *Life Sci.* 43, 755–760.

Ganesh, C.B., Chabbi, A., 2013. Naltrexone attenuates stress-induced suppression of LH secretion in the pituitary gland in the Cichlid fish *Oreochromis mossambicus*: Evidence for the opioidergic mediation of reproductive stress response. *Fish Physiol. Biochem.* 39, 627–636. <https://doi.org/10.1007/s10695-012-9725-y>

Gardner, L., Anderson, T., Place, A.R., Dixon, B., Elizur, A., 2005. Sex change strategy and the aromatase genes. *J. Steroid Biochem. Mol. Biol.* 94, 395–404. <https://doi.org/10.1016/j.jsbmb.2004.12.045>

Gennotte, V., Sawadogo, P., Milla, S., Kestemont, P., Mélard, C., Rougeot, C., 2012. Cortisol is responsible for positive and negative effects in the ovarian maturation induced by the exposure to acute stressors in Nile tilapia, *Oreochromis niloticus*. *Fish Physiol. Biochem.* 38, 1619–1626. <https://doi.org/10.1007/s10695-012-9656-7>

Geven, E.J.W., Flik, G., Klaren, P.H.M., 2009. Central and peripheral integration of interrenal and thyroid axes signals in common carp (*Cyprinus carpio* L.). *J. Endocrinol.* 200, 117–123. <https://doi.org/10.1677/JOE-08-0410>

Gilchrist, B.J., Tipping, D.R., Hake, L., Levy, A., Baker, B.I., 2000. The effects of acute and chronic stresses on vasotocin gene transcripts in the brain of the rainbow trout (*Oncorhynchus mykiss*). *J. Neuroendocrinol.* 12, 795–801.

1713 <https://doi.org/10.1046/j.1365-2826.2000.00522.x>

1714 Gillies, G., Linton, E., Lowry, P., 1982. Corticotropin-releasing activity of the new CRF is  
 1715 potentiated several times by vasopressin. *Nature* 299, 355–357.

1716 Gilmour, K.M., 2005. Mineralocorticoid receptors and hormones: Fishing for answers.  
 1717 *Endocrinology* 146, 44–46. <https://doi.org/10.1210/en.2004-1390>

1718 Gingerich, S., Wang, X., Lee, P.K.P., Dhillon, S.S., Chalmers, J.A., Koletar, M.M., Belsham,  
 1719 D.D., 2009. The generation of an array of clonal, immortalized cell models from the rat  
 1720 hypothalamus: analysis of melatonin effects on kisspeptin and gonadotropin-inhibitory  
 1721 hormone neurons. *Neuroscience* 162, 1134–1140.  
 1722 <https://doi.org/10.1016/j.neuroscience.2009.05.026>

1723 Gjerde, B., 1984. Response to individual selection for age at sexual maturity in Atlantic  
 1724 salmon. *Aquaculture* 38, 229–240. [https://doi.org/10.1016/0044-8486\(84\)90147-9](https://doi.org/10.1016/0044-8486(84)90147-9)

1725 Goikoetxea, A., Todd, E. V., Gemmell, N.J., 2017. Stress and sex: Does cortisol mediate sex  
 1726 change in fish? *Reproduction* 154, R149–R160. <https://doi.org/10.1530/REP-17-0408>

1727 Gojska, N.M., Belsham, D.D., 2014. Glucocorticoid receptor-mediated regulation of Rfrp  
 1728 (GnIH) and Gpr147 (GnIH-R) synthesis in immortalized hypothalamic neurons. *Mol.*  
 1729 *Cell. Endocrinol.* 384, 23–31. <https://doi.org/10.1016/j.mce.2013.12.015>

1730 Golan, M., Martin, A.O., Mollard, P., Levavi-Sivan, B., 2016. Anatomical and functional  
 1731 gonadotrope networks in the teleost pituitary. *Sci. Rep.* 6, 1–8.  
 1732 <https://doi.org/10.1038/srep23777>

1733 Gore, A., 2002. GnRH: The Master Molecule of Reproduction. Kluwer Academic Publishers.

1734 Gorissen, M., Flik, G., 2016. The endocrinology of the stress response in fish : an adaptation-  
 1735 physiological view., in: Schreck, C., Tort, L., Farrell, A., Brauner, C. (Eds.), *Biology of*  
 1736 *Stress in Fish. Fish Physiology Vol 35.* pp. 75–111.

1737 Goswami, S. V., Lamba, V.J., Sundararaj, B.I., 1985. Gonadotrophin-induced oocyte

1738 maturation in the catfish, *Heteropneustes fossilis* (Bloch), requires steroidogenesis in  
 1739 both interrenal and ovary. *Gen. Comp. Endocrinol.* 57, 53–63.  
 1740 [https://doi.org/10.1016/0016-6480\(85\)90200-X](https://doi.org/10.1016/0016-6480(85)90200-X)

1741 Greenwood, A.K., Butler, P.C., White, R.B., Demarco, U., Pearce, D., Fernald, R.D., 2003.  
 1742 Multiple corticosteroid receptors in a teleost fish: Distinct sequences, expression  
 1743 patterns, and transcriptional activities. *Endocrinology* 144, 4226–4236.  
 1744 <https://doi.org/10.1210/en.2003-0566>

1745 Grone, B.P., Maruska, K.P., 2015. Divergent evolution of two corticotropin-releasing  
 1746 hormone (CRH) genes in teleost fishes. *Front. Neurosci.* 9, 1–13.  
 1747 <https://doi.org/10.3389/fnins.2015.00365>

1748 Haddy, J.A., Pankhurst, N.W., 1999. Stress-induced changes in concentrations of plasma sex  
 1749 steroids in black bream. *J. Fish Biol.* 55, 1304–1316.  
 1750 <https://doi.org/10.1006/jfbi.1999.1128>

1751 Haitina, T., Klovins, J., Andersson, J., Fredriksson, R., Lagerström, M.C., Larhammar, D.,  
 1752 Larson, E.T., Schiöth, H.B., 2004. Cloning, tissue distribution, pharmacology and three-  
 1753 dimensional modelling of melanocortin receptors 4 and 5 in rainbow trout suggest close  
 1754 evolutionary relationship of these subtypes. *Biochem. J.* 380, 475–486.  
 1755 <https://doi.org/10.1042/BJ20031934>

1756 Hall, K.C., Broadhurst, M.K., Butcher, P.A., Cameron, L., Rowland, S.J., Millar, R.B., 2017.  
 1757 Sublethal effects of angling and release on golden perch *Macquaria ambigua*:  
 1758 implications for reproduction and fish health. *J. Fish Biol.* 90, 1980–1998.  
 1759 <https://doi.org/10.1111/jfb.13282>

1760 Hall, K.C., Broadhurst, M.K., Butcher, P.A., Rowland, S.J., 2009. Effects of angling on post-  
 1761 release mortality, gonadal development and somatic condition of Australian bass  
 1762 *Macquaria novemaculeata*. *J. Fish Biol.* 75, 2737–2755. <https://doi.org/10.1111/j.1095->

1763 8649.2009.02474.x

1764 Han, Y.S., Liao, I.C., Tzeng, W.N., Yu, J.Y.L., 2004. Cloning of the cDNA for thyroid  
 1765 stimulating hormone  $\beta$  subunit and changes in activity of the pituitary - Thyroid axis  
 1766 during silvering of the Japanese eel, *Anguilla japonica*. J. Mol. Endocrinol. 32, 179–194.  
 1767 <https://doi.org/10.1677/jme.0.0320179>

1768 Hane, S., Robertson, O.H., 1959. Changes in Plasma 17-Hydroxycorticosteroids  
 1769 Accompanying Sexual Maturation and Spawning of the Pacific Salmon (*Oncorhynchus*  
 1770 *Tschawytscha*) and Rainbow Trout (*Salmo Gairdnerii*). Proc. Natl. Acad. Sci. 45, 886–  
 1771 893. <https://doi.org/10.1073/pnas.45.6.886>

1772 Hansen, K.K., Stafflinger, E., Schneider, M., Hauser, F., Cazzamali, G., Williamson, M.,  
 1773 Kollmann, M., Schachtner, J., Grimmelikhuijzen, C.J.P., 2010. Discovery of a novel  
 1774 insect neuropeptide signaling system closely related to the insect adipokinetic hormone  
 1775 and corazonin hormonal systems. J. Biol. Chem. 285, 10736–10747.  
 1776 <https://doi.org/10.1074/jbc.M109.045369>

1777 Hanson, K.C., Cooke, S.J., Suski, C.D., Philipp, D.P., 2007. Effects of different angling  
 1778 practices on post-release behaviour of nest-guarding male black bass, *Micropterus* spp.  
 1779 Fish. Manag. Ecol. 14, 141–148. <https://doi.org/10.1111/j.1365-2400.2007.00534.x>

1780 Harris, R.M., Dijkstra, P.D., Hofmann, H.A., 2014. Complex structural and regulatory  
 1781 evolution of the pro-opiomelanocortin gene family. Gen. Comp. Endocrinol. 195, 107–  
 1782 115. <https://doi.org/10.1016/j.ygcen.2013.10.007>

1783 Hatey, J., 1951. La fonction glycogénique du foie de l'anguille (*Anguilla anguilla* L.) après  
 1784 hypophysectomie. Compt Rend Soc Biol Paris 145, 315–318.

1785 Hattori, R.S., Fernandino, J.I., Kishil, A., Kimura, H., Kinno, T., Oura, M., Somoza, G.M.,  
 1786 Yokota, M., Strüssmann, C.A., Watanabe, S., 2009. Cortisol-induced masculinization:  
 1787 Does thermal stress affect gonadal fate in pejerrey, a teleost fish with temperature-

1788 dependent sex determination? PLoS One 4.  
1789 <https://doi.org/10.1371/journal.pone.0006548>  
1790 Hayashi, Y., Kobira, H., Yamaguchi, T., Shiraishi, E., Yazawa, T., Hirai, T., Kamei, Y.,  
1791 Kitano, T., 2010. High temperature causes masculinization of genetically female medaka  
1792 by elevation of cortisol. *Mol. Reprod. Dev.* 77, 679–686.  
1793 <https://doi.org/10.1002/mrd.21203>  
1794 Hirose, K., 1976. Endocrine control of ovulation in medaka (*Oryzias latipes*) and Ayu  
1795 (*Plecoglossus altivelis*). *J. l'Office des Rech. sur les pêcheries du Canada* 33, 989–994.  
1796 Höglund, E., Balm, P.H.M., Winberg, S., 2002. Stimulatory and inhibitory effects of 5-HT1A  
1797 receptors on adrenocorticotrophic hormone and cortisol secretion in a teleost fish, the  
1798 Arctic charr (*Salvelinus alpinus*). *Neurosci. Lett.* 324, 193–196.  
1799 [https://doi.org/10.1016/S0304-3940\(02\)00200-8](https://doi.org/10.1016/S0304-3940(02)00200-8)  
1800 Hollenberg, S., Weinberger, C., Ong, E., Cerelli, G., Oro, A., Lebo, R., Thompson, E.,  
1801 Rosenfeld, M., Evans, R., 1985. Primary structure and expression of a functionakl  
1802 human glucocorticoid receptor cDNA. *Nature* 318, 635–641.  
1803 Huang, Y.-S., Rousseau, K., Sbahi, M., Le Belle, N., Schmitz, M., Dufour, S., 1999. Cortisol  
1804 selectively stimulates pituitary gonadotropin  $\beta$ -subunit in a primitive teleost, *Anguilla*  
1805 *anguilla*. *Endocrinology* 140, 1228–1235. <https://doi.org/10.1210/endo.140.3.6598>  
1806 Huising, M.O., Metz, J.R., van Schooten, C., Taverne-Thiele, A.J., Hermesen, T., Verburg-van  
1807 Kemenade, B.M.L., Flik, G., 2004. Structural characterisation of a cyprinid (*Cyprinus*  
1808 *carpio* L.) CRH, CRH-BP and CRH-R1, and the role of these proteins in the acute stress  
1809 response. *J. Mol. Endocrinol.* 32, 627–648. <https://doi.org/10.1677/jme.0.0320627>  
1810 Huntingford, F., Adams, C., Braithwaite, V., Kadri, S., Pottinger, T., Sandoe, P., Turnbull, J.,  
1811 2006. Current issues in fish welfare. *J. Fish Biol.* 68, 332–372.  
1812 <https://doi.org/10.1111/j.1095-8649.2005.01046.x>

1813 Hwang, J.I., Moon, M.J., Park, S., Kim, D.K., Cho, E.B., Ha, N., Son, G.H., Kim, K., Vaudry,  
1814 H., Seong, J.Y., 2013. Expansion of secretin-like G protein-coupled receptors and their  
1815 peptide ligands via local duplications before and after two rounds of whole-genome  
1816 duplication. *Mol. Biol. Evol.* 30, 1119–1130. <https://doi.org/10.1093/molbev/mst031>

1817 Ichikawa, T., McMaster, D., Lederis, K., Kobayashi, H., 1982. Isolation and amino acid  
1818 sequence of urotensin I, a vasoactive and ACTH-releasing neuropeptide, from the carp  
1819 (*Cyprinus carpio*) urophysis. *Peptides* 3, 859–867. [https://doi.org/10.1016/0196-](https://doi.org/10.1016/0196-9781(82)90028-6)  
1820 9781(82)90028-6

1821 Idler, D., Ronald, A., Schmidt, P., 1959. Biochemical studies on sockeye salmon during  
1822 spawning migration. VII. Steroid hormones in plasma. *Can. J. Biochem. Physiol.* 37,  
1823 1227–1238. <https://doi.org/10.1139/y60-067>

1824 Imanaga, Y., Nyuji, M., Amano, M., Takahashi, A., Kitano, H., Yamaguchi, A., Matsuyama,  
1825 M., 2014. Characterization of gonadotropin-releasing hormone and gonadotropin in jack  
1826 mackerel (*Trachurus japonicus*): Comparative gene expression analysis with respect to  
1827 reproductive dysfunction in captive and wild fish. *Aquaculture* 428–429, 226–235.  
1828 <https://doi.org/10.1016/j.aquaculture.2014.03.003>

1829 Ishibashi, M., Yamaji, T., 1981. Direct effects of thyrotropin-releasing hormone,  
1830 cyproheptadine, and dopamine on adrenocorticotropin secretion from human  
1831 corticotroph adenoma cells in vitro. *J. Clin. Invest.* 68, 1018–1027.  
1832 <https://doi.org/10.1172/JCI110324>

1833 Iwasa, T., Matsuzaki, T., Yano, K., Mayila, Y., Irahara, M., 2018. The roles of kisspeptin and  
1834 gonadotropin inhibitory hormone in stress-induced reproductive disorders. *Endocr. J.* 65,  
1835 133–140. <https://doi.org/10.1507/endocrj.EJ18-0026>

1836 Jalabert, B., 1975. Modulation par différents stéroïdes non maturants de l'efficacité de la 17β-  
1837 hydroxy-20 β-dihydroprogestérone ou d'un extrait gonadotrope sur la maturation

1838 intrafolliculaire in vitro des ovocytes de la Truite arc-en-ciel *Salmo gairdnerii*. C. R.  
 1839 Acad. Sci. Paris, Sér. D 281, 811–814.

1840 Jalabert, B., Bry, C., Szöllösi, D., Fostier, A., 1973. Étude Comparée de l'action des  
 1841 hormones hypophysaires et stéroïdes sur la maturation in vitro des ovocytes de la truite  
 1842 et du carassin (poissons téléostéens). 59–73.

1843 Jalabert, B., Fostier, A., 1984. The modulatory effect in vitro of oestradiol 17 beta,  
 1844 testosterone or cortisol on the output of 17alpha;-hydroxy-20beta;-dihydroprogesterone  
 1845 by rainbow trout (*Salmo gairdneri*) ovarian follicles stimulated by the maturational  
 1846 gonadotropin s-GtH. *Reprod Nutr Dévelop* 24, 127–136.

1847 [Jenks, B.G., Verburg van Kemenade, B.M.L., Tonon, M.C., Vaudry, H., 1985. Regulation of](#)  
 1848 [biosynthesis and release of pars intermedia peptides in \*Rana ridibunda\*: Dopamine](#)  
 1849 [affects both acetylation and release of  \$\alpha\$ -MSH. \*Peptides\* 6, 913–921.](#)  
 1850 [https://doi.org/10.1016/0196-9781\(85\)90322-5](https://doi.org/10.1016/0196-9781(85)90322-5)

1851 Ježová, D., Jurčovičová, J., Vigaš, M., Murgaš, K., Labrie, F., 1985. Increase in plasma  
 1852 ACTH after dopaminergic stimulation in rats. *Psychopharmacology (Berl)*. 85, 201–203.  
 1853 <https://doi.org/10.1007/BF00428414>

1854 Jonsson, N., Jonsson, B., Hansen, L.P., 1997. Changes in Proximate Composition and  
 1855 Estimates of Energetic Costs During Upstream Migration and Spawning in Atlantic  
 1856 Salmon *Salmo salar*. *J. Anim. Ecol.* 66, 425–436. <https://doi.org/10.2307/5987>

1857 Kagabu, Y., Mishiba, T., Okino, T., Yanagisawa, T., 1998. Effects of thyrotropin-releasing  
 1858 hormone and its metabolites, cyclo(His- Pro) and TRH-OH, on growth hormone and  
 1859 prolactin synthesis in primary cultured pituitary cells of the common carp, *Cyprinus*  
 1860 *carpio*. *Gen. Comp. Endocrinol.* 111, 395–403. <https://doi.org/10.1006/gcen.1998.7124>

1861 Kageyama, K., 2013. Regulation of gonadotropins by corticotropin-releasing factor and  
 1862 urocortin. *Front. Endocrinol. (Lausanne)*. 4, 1–7.

1863 <https://doi.org/10.3389/fendo.2013.00012>

1864 Kanda, S., 2019. Evolution of the regulatory mechanisms for the hypothalamic-pituitary-

1865 gonadal axis in vertebrates—hypothesis from a comparative view. *Gen. Comp.*

1866 *Endocrinol.* 284, 0–1. <https://doi.org/10.1016/j.ygcen.2018.11.014>

1867 Katsu, Y., Baker, M.E., 2018. Progesterone activation of zebrafish mineralocorticoid receptor

1868 may influence growth of some transplanted tumors. *Proc. Natl. Acad. Sci. U. S. A.* 115,

1869 E2908–E2909. <https://doi.org/10.1073/pnas.1802441115>

1870 Katsu, Y., Kohno, S., Oka, K., Lin, X., Otake, S., Pillai, N., Takagi, W., Hyodo, S.,

1871 Venkatesh, B., Baker, M., 2019. Transcriptional activation of elephant shark

1872 mineralocorticoid receptor by corticosteroids, progesterone, and spironolactone. *Sci.*

1873 *Signal.* 12, eaar2668.

1874 Kelsall, C.J., Balment, R.J., 1998. Native urotensins influence cortisol secretion and plasma

1875 cortisol concentration in the euryhaline flounder, *Platichthys flesus*. *Gen. Comp.*

1876 *Endocrinol.* 112, 210–219. <https://doi.org/10.1006/gcen.1998.7166>

1877 Kiilerich, P., Servili, A., Péron, S., Valotaire, C., Goardon, L., Leguen, I., Prunet, P., 2018.

1878 Regulation of the corticosteroid signalling system in rainbow trout HPI axis during

1879 confinement stress. *Gen. Comp. Endocrinol.* 258, 184–193.

1880 <https://doi.org/10.1016/j.ygcen.2017.08.013>

1881 Kim, D.K., Cho, E.B., Moon, M.J., Park, S., Hwang, J.I., Kah, O., Sower, S.A., Vaudry, H.,

1882 Seong, J.Y., 2011. Revisiting the evolution of gonadotropin-releasing hormones and

1883 their receptors in vertebrates: Secrets hidden in genomes. *Gen. Comp. Endocrinol.* 170,

1884 68–78. <https://doi.org/10.1016/j.ygcen.2010.10.018>

1885 Kim, M., Kim, D., Sohn, Y., 2011. Characterization of two functional glucocorticoid

1886 receptors in the marine medaka *Oryzias dancena*. *Gen. Comp. Endocrinol.* 171, 341–349.

1887 <https://doi.org/10.1016/j.ygcen.2011.02.015>

- 1888 Kime, D.E., 1993. Classical and non-classical reproductive steroids in fish. *Rev. Fish Biol.*  
 1889 *Fish.* 3, 160–180.
- 1890 Kime, D.E., Scott, A.P., Canario, A.V.M., 1992. In vitro biosynthesis of steroids, including  
 1891 11-deoxycortisol and 5 $\alpha$ -pregnane-3 $\beta$ ,7 $\alpha$ ,17,20 $\beta$ -tetrol, by ovaries of the goldfish  
 1892 *Carassius auratus* during the stage of oocyte final maturation. *Gen. Comp. Endocrinol.*  
 1893 87, 375–384. [https://doi.org/10.1016/0016-6480\(92\)90044-K](https://doi.org/10.1016/0016-6480(92)90044-K)
- 1894 Kinsey-Jones, J.S., Li, X.F., Knox, A.M.I., Wilkinson, E.S., Zhu, X.L., Chaudhary, A.A.,  
 1895 Milligan, S.R., Lightman, S.L., O’Byrne, K.T., 2009. Down-regulation of hypothalamic  
 1896 kisspeptin and its receptor, Kiss1r, mRNA expression is associated with stress-induced  
 1897 suppression of luteinising hormone secretion in the female rat. *J. Neuroendocrinol.* 21,  
 1898 20–29. <https://doi.org/10.1111/j.1365-2826.2008.01807.x>
- 1899 Kirby, E.D., Geraghty, A.C., Ubuka, T., Bentley, G.E., Kaufer, D., 2009. Stress increases  
 1900 putative gonadotropin inhibitory hormone and decreases luteinizing hormone in male  
 1901 rats. *Proc. Natl. Acad. Sci. U. S. A.* 106, 11324–11329.  
 1902 <https://doi.org/10.1073/pnas.0901176106>
- 1903 Kirshenblatt, I., 1952. The action of steroid hormones in female vy’un. *Dokl Akad Nauk*  
 1904 *USSR* 83, 629–632.
- 1905 Kitahara, N., Nishizawa, T., Iida, K., Okazaki, H., Andoh, T., Soma, G., 1988. Absence of a  
 1906 gamma-melanocyte-stimulating hormone sequence in proopiomelanocortin mRNA of  
 1907 chum salmon *Oncorhynchus keta*. *Comp. Biochem. Physiol. - B Biochem. Mol. Biol.* 91,  
 1908 365–370.
- 1909 Kjesbu, O., Klungsoyr, J., Kryvi, H., Witthames, P., Greer Walker, M., 1991. Fecundity,  
 1910 atresia, and egg size of captive Atlantic cod (*Gadus morhua*) in relation to proximate body  
 1911 composition. *J. Can. des Sci. halieutiques Aquat.* 48, 2333–2343.
- 1912 Kjesbu, O.S., 1989. The spawning activity of cod, *Gadus morhua* L. *J. Fish Biol.* 34, 195–

1913 206. <https://doi.org/10.1111/j.1095-8649.1989.tb03302.x>

1914 Kobayashi, Y., Chiba, H., Yamanome, T., Schiöth, H.B., Takahashi, A., 2011. Melanocortin  
 1915 receptor subtypes in interrenal cells and corticotropic activity of  $\alpha$ -melanocyte-  
 1916 stimulating hormones in barfin flounder, *Verasper moseri*. *Gen. Comp. Endocrinol.* 170,  
 1917 558–568. <https://doi.org/10.1016/j.ygcen.2010.11.019>

1918 Korzan, W.J., Grone, B.P., Fernald, R.D., 2014. Social regulation of cortisol receptor gene  
 1919 expression. *J. Exp. Biol.* 217, 3221–3228. <https://doi.org/10.1242/jeb.104430>

1920 Kowalski, R., Sarosiek, B., JUdycka, S., Dryl, K., Grudniewska, J., Dobosz, S., Cejko, B.,  
 1921 2018. Effectiveness of the air stripping in two salmonid fish, rainbow trout  
 1922 (*Oncorhynchus mykiss*) and brown trout (*Salmo trutta morpha fario*). *J Vis Exp* 16.

1923 Kraicer, J., Gosbee, J., Bencosme, S., 1973. Pars intermedia and pars distalis: two sites of  
 1924 ACTH production in the rat hypophysis. *Neuroendocrinology* 11, 156–176.

1925 Kubokawa, K., Watanabe, T., Yoshioka, M., Iwata, M., 1999. Effects of acute stress on  
 1926 plasma cortisol, sex steroid hormone and glucose levels in male and female sockeye  
 1927 salmon during the breeding season. *Aquaculture* 172, 335–349.  
 1928 [https://doi.org/10.1016/S0044-8486\(98\)00504-3](https://doi.org/10.1016/S0044-8486(98)00504-3)

1929 Kumar, P., Wisdom, K.S., Bhat, I.A., Pathakota, G.B., Nayak, S.K., Reang, D., Nagpure,  
 1930 N.S., Sharma, R., 2019. Molecular characterization of gonadotropin-inhibitory hormone  
 1931 (GnIH) gene and effect of intramuscular injection of GnIH peptide on the reproductive  
 1932 axis in *Catla catla*. *Anim. Biotechnol.* 0, 1–15.  
 1933 <https://doi.org/10.1080/10495398.2019.1597730>

1934 Lafont, A., Hardman, L., Dirks, R., von den Thillart, G., Tomkiewicz, J., Dufour, S., 2014.  
 1935 Characterization and regulation during reproduction of four nuclear corticosteroid  
 1936 receptors in the European eel, *Anguilla anguilla*., in: 10th International Symposium on  
 1937 Reproductive Physiology of Fish.

- 1938 Lafont, A.G., Rousseau, K., Tomkiewicz, J., Dufour, S., 2016. Three nuclear and two  
1939 membrane estrogen receptors in basal teleosts, *Anguilla* sp.: Identification, evolutionary  
1940 history and differential expression regulation. *Gen. Comp. Endocrinol.* 235, 177–191.  
1941 <https://doi.org/10.1016/j.ygcen.2015.11.021>
- 1942 Lamba, V.J., Goswami, S. V., Sundararaj, B.I., 1983. Circannual and circadian variations in  
1943 plasma levels of steroids (cortisol, estradiol-17 $\beta$  estrone, and testosterone) correlated  
1944 with the annual gonadal cycle in the catfish, *Heteropneustes fossilis* (Bloch). *Gen.*  
1945 *Comp. Endocrinol.* 50, 205–225. [https://doi.org/10.1016/0016-6480\(83\)90221-6](https://doi.org/10.1016/0016-6480(83)90221-6)
- 1946 Lambert, Y., Dutil, J.-D., 2000. Energetic consequences of reproduction in Atlantic cod  
1947 (*Gadus morhua*) in relation to spawning level of somatic energy reserves. *J. Can. des Sci.*  
1948 *halieutiques Aquat.* 57, 815–825.
- 1949 Lamers, A., Balm, P., Haenen, H., Jenks, B., Wendelaar-Bonga, S.E., 1991. Regulation of  
1950 differential release of alpha-melanocyte-stimulating hormone forms from the pituitary pf  
1951 a teleost fish, *Oreochromis mossambicus*. *J Endocrinol* 129, 179–187.
- 1952 Lamers, A., Flik, G., Atsma, W., SE, W.-B., 1992. A role for di-acetyl-alpha-melanocyte-  
1953 stimulating hormone in the control of cortisol release in the teleost *Oreochromis*  
1954 *mossambicus*. *J Endocrinol* 135, 285–292.
- 1955 Lamers, A., Flik, G., Wendelaar-Bonga, S., 1994. A specific role for TRH in release of  
1956 diacetyl alpha-MSH in tilapia stressed by acid water. *Am J Physiol* 267, R1302-1308.
- 1957 Lamers, A., Ter Brugge, P., Flik, G., Wendelaar-Bonga, S., 1997. Acid stress induces a D1-  
1958 like dopamine receptor in pituitary MSH cells of *Oreochromis mossambicus*. *Am J*  
1959 *Physiol* 273, R387-392.
- 1960 Larhammar, D., Risinger, C., 1994. Molecular genetic aspects of tetraploidy in the common  
1961 carp *cyprinus carpio*. *Mol. Phylogenet. Evol.* <https://doi.org/10.1006/mpev.1994.1007>
- 1962 Larsen, D.A., Swanson, P., Dickey, J.T., Rivier, J., Dickhoff, W.W., 1998. In vitro

1963 thyrotropin-releasing activity of corticotropin-releasing hormone-family peptides in coho  
 1964 salmon, *Oncorhynchus kisutch*. *Gen. Comp. Endocrinol.* 109, 276–285.  
 1965 <https://doi.org/10.1006/gcen.1997.7031>

1966 Leatherland, J.F., Li, M., Barkataki, S., 2010. Stressors, glucocorticoids and ovarian function  
 1967 in teleosts. *J. Fish Biol.* 76, 86–111. <https://doi.org/10.1111/j.1095-8649.2009.02514.x>

1968 Leder, E.H., Silverstein, J.T., 2006. The pro-opiomelanocortin genes in rainbow trout  
 1969 (*Oncorhynchus mykiss*): Duplications, splice variants, and differential expression. *J.*  
 1970 *Endocrinol.* 188, 355–363. <https://doi.org/10.1677/joe.1.06283>

1971 Lederis, K., Letter, A., McMaster, D., Moore, G., Schlesinger, D., 1982. Complete amino acid  
 1972 sequence of urotensin I, a hypotensive and corticotropin-releasing neuropeptide from  
 1973 *Catostomus*. *Science* (80-. ). 218, 162–165.

1974 Lee, J., Danielson, P., Sollars, C., Alrubaiian, J., Balm, P., Dorés, R.M., 1999. Cloning of a  
 1975 neoteleost (*Oreochromis mossambicus*) pro-opiomelanocortin (POMC) cDNA reveals a  
 1976 deletion of the  $\gamma$ -melanotropin region and most of the joining peptide region:  
 1977 Implications for POMC processing. *Peptides* 20, 1391–1399.  
 1978 [https://doi.org/10.1016/S0196-9781\(99\)00148-5](https://doi.org/10.1016/S0196-9781(99)00148-5)

1979 Lee, J., Miele, M.E., Hicks, D.J., Karen, K., Trent, J., Weissman, B., Welch, D.R., 1996.  
 1980 KiSS-1 , a Novel Human Malignant Melanoma. *J. Natl. Cancer Inst.* 88, 1731–1737.

1981 Lethimonier, C., Flouriot, G., Kah, O., Ducouret, B., 2002. The glucocorticoid receptor  
 1982 represses the positive autoregulation of the trout estrogen receptor gene by preventing  
 1983 the enhancer effect of a C/EBP $\beta$ -like protein. *Endocrinology* 143, 2961–2974.  
 1984 <https://doi.org/10.1210/endo.143.8.8958>

1985 Lethimonier, C., Flouriot, G., Valotaire, Y., Kah, O., Ducouret, B., 2000. Transcriptional  
 1986 Interference Between Glucocorticoid Receptor and Estradiol Receptor Mediates the  
 1987 Inhibitory Effect of Cortisol on Fish Vitellogenesis<sup>1</sup>. *Biol. Reprod.* 62, 1763–1771.

1988 <https://doi.org/10.1095/biolreprod62.6.1763>

1989 Lewis, K., Li, C., Perrin, M.H., Blount, A., Kunitake, K., Donaldson, C., Vaughan, J., Reyes,  
1990 T.M., Gulyas, J., Fischer, W., Bilezikjian, L., Rivier, J., Sawchenko, P.E., Vale, W.W.,  
1991 2001. Identification of urocortin III, an additional member of the corticotropin-releasing  
1992 factor (CRF) family with high affinity for the CRF2 receptor. *Proc. Natl. Acad. Sci. U.*  
1993 *S. A.* 98, 7570–7575. <https://doi.org/10.1073/pnas.121165198>

1994 Li, Y., Sturm, A., Cunningham, P., Bury, N.R., 2012. Evidence for a divergence in function  
1995 between two glucocorticoid receptors from a basal teleost. *BMC Evol. Biol.* 12.  
1996 <https://doi.org/10.1186/1471-2148-12-137>

1997 Lien, S., Koop, B.F., Sandve, S.R., Miller, J.R., Kent, M.P., Nome, T., Hvidsten, T.R., Leong,  
1998 J.S., Minkley, D.R., Zimin, A., Grammes, F., Grove, H., Gjuvsland, A., Walenz, B.,  
1999 Hermansen, R.A., Von Schalburg, K., Rondeau, E.B., Di Genova, A., Samy, J.K.A.,  
2000 Olav Vik, J., Vigeland, M.D., Caler, L., Grimholt, U., Jentoft, S., Inge Våge, D., De  
2001 Jong, P., Moen, T., Baranski, M., Palti, Y., Smith, D.R., Yorke, J.A., Nederbragt, A.J.,  
2002 Tooming-Klunderud, A., Jakobsen, K.S., Jiang, X., Fan, D., Hu, Y., Liberles, D.A.,  
2003 Vidal, R., Iturra, P., Jones, S.J.M., Jonassen, I., Maass, A., Omholt, S.W., Davidson,  
2004 W.S., 2016. The Atlantic salmon genome provides insights into rediploidization. *Nature*  
2005 533, 200–205. <https://doi.org/10.1038/nature17164>

2006 Lim, J.E., Porteus, C.S., Bernier, N.J., 2013. Serotonin directly stimulates cortisol secretion  
2007 from the interrenals in goldfish. *Gen. Comp. Endocrinol.* 192, 246–255.  
2008 <https://doi.org/10.1016/j.ygcen.2013.08.008>

2009 Liu, H., Todd, E. V., Lokman, P.M., Lamm, M.S., Godwin, J.R., Gemmell, N.J., 2017. Sexual  
2010 plasticity: A fishy tale. *Mol. Reprod. Dev.* 84, 171–194.  
2011 <https://doi.org/10.1002/mrd.22691>

2012 Logan, D.W., Bryson-Richardson, R.J., Pagán, K.E., Taylor, M.S., Currie, P.D., Jackson, I.J.,

2013 2003. The structure and evolution of the melanocortin and MCH receptors in fish and  
 2014 mammals. *Genomics* 81, 184–191. [https://doi.org/10.1016/S0888-7543\(02\)00037-X](https://doi.org/10.1016/S0888-7543(02)00037-X)  
 2015 Lovejoy, D.A., Balment, R.J., 1999. Evolution and physiology of the corticotropin-releasing  
 2016 factor (CRF) family of neuropeptides in vertebrates. *Gen. Comp. Endocrinol.* 115, 1–22.  
 2017 <https://doi.org/10.1006/gcen.1999.7298>  
 2018 Lovejoy, D.A., Chang, B.S.W., Lovejoy, N.R., del Castillo, J., 2014. Molecular evolution of  
 2019 GPCRs: CRH/CRH receptors. *J. Mol. Endocrinol.* 52. [https://doi.org/10.1530/JME-13-](https://doi.org/10.1530/JME-13-0238)  
 2020 0238  
 2021 Lowerre-Barbieri, S.K., Brown-Peterson, N.J., Murua, H., Tomkiewicz, J., Wyanski, D.M.,  
 2022 Saborido-Rey, F., 2011. Emerging issues and methodological advances in fisheries  
 2023 reproductive biology. *Mar. Coast. Fish.* 3, 32–51.  
 2024 <https://doi.org/10.1080/19425120.2011.555725>  
 2025 Lu, E., Dow, L., Gumusgoz, S., Brierley, M.J., Warne, J.M., McCrohan, C.R., Balment, R.J.,  
 2026 Riccardi, D., 2004. Coexpression of corticotropin-releasing hormone and urotensin I  
 2027 precursor genes in the caudal neurosecretory system of the euryhaline flounder  
 2028 (*Platichthys flesus*): A possible shared role in peripheral regulation. *Endocrinology* 145,  
 2029 5786–5797. <https://doi.org/10.1210/en.2004-0144>  
 2030 Luo, E., Stephens, S., Chaing, S., Munaganuru, N., Kauffman, A., Breen, K., 2016.  
 2031 Corticosterone blocks ovarian cyclicity and the LH surge via decreased kisspeptin  
 2032 neurons activation in female mice. *Endocrinology* 157, 1187–1199.  
 2033 Madaro, A., Olsen, R.E., Kristiansen, T.S., Ebbesson, L.O.E., Flik, G., Gorissen, M., 2016. A  
 2034 comparative study of the response to repeated chasing stress in Atlantic salmon (*Salmo*  
 2035 *salar* L.) parr and post-smolts. *Comp. Biochem. Physiol. -Part A Mol. Integr. Physiol.*  
 2036 192, 7–16. <https://doi.org/10.1016/j.cbpa.2015.11.005>  
 2037 Madaro, A., Olsen, R.E., Kristiansen, T.S., Ebbesson, L.O.E., Nilsen, T.O., Flik, G.,

2038 Gorissen, M., 2015. Stress in Atlantic salmon: Response to unpredictable chronic stress.  
 2039 J. Exp. Biol. 218, 2538–2550. <https://doi.org/10.1242/jeb.120535>

2040 Mangelsdorf, D.J., Thummel, C., Beato, M., Herrlich, P., Schütz, G., Umesono, K.,  
 2041 Blumberg, B., Kastner, P., Mark, M., Chambon, P., Evans, R.M., 1995. The nuclear  
 2042 receptor superfamily: The second decade. Cell 83, 835–839.  
 2043 [https://doi.org/10.1016/0092-8674\(95\)90199-X](https://doi.org/10.1016/0092-8674(95)90199-X)

2044 Mankiewicz, J.L., Godwin, J., Holler, B.L., Turner, P.M., Murashige, R., Shamey, R.,  
 2045 Daniels, H. V., Borski, R.J., 2013. Masculinizing effect of background color and cortisol  
 2046 in a flatfish with environmental sex-determination. Integr. Comp. Biol. 53, 755–765.  
 2047 <https://doi.org/10.1093/icb/ict093>

2048 Manuel, R., Metz, J.R., Flik, G., Vale, W.W., Huising, M.O., 2014. Corticotropin-releasing  
 2049 factor-binding protein (CRF-BP) inhibits CRF- and urotensin-I-mediated activation of  
 2050 CRF receptor-1 and -2 in common carp. Gen. Comp. Endocrinol. 202, 69–75.  
 2051 <https://doi.org/10.1016/j.ygcen.2014.04.010>

2052 Maruska, K.P., 2014. Social regulation of reproduction in male cichlid fishes. Gen. Comp.  
 2053 Endocrinol. 207, 2–12. <https://doi.org/10.1016/j.ygcen.2014.04.038>

2054 Maruska, K.P., Fernald, R.D., 2011. Plasticity of the reproductive axis caused by social status  
 2055 change in an african cichlid fish: II. Testicular gene expression and spermatogenesis.  
 2056 Endocrinology 152, 291–302. <https://doi.org/10.1210/en.2010-0876>

2057 Maugars, G., Dufour, S., 2015. Demonstration of the coexistence of duplicated LH receptors  
 2058 in teleosts, and their origin in ancestral actinopterygians. PLoS One 10, 1–29.  
 2059 <https://doi.org/10.1371/journal.pone.0135184>

2060 Maugars, G., Mauvois, X., Rousseau, K., Dufour, S., 2016. Evolution of the corticotropin-  
 2061 releasing hormone paralogs in teleosts., in: 8th International Symposium on Fish  
 2062 Endocrinology.

2063 Maugars, G., Pasquier, J., Atkinson, C., Lafont, A., Campo, A., Kamech, N., Lefranc, B.,  
 2064 2020. Gonadotropin-inhibitory hormone in teleosts : New insights from a basal  
 2065 representative , the eel. *Gen. Comp. Endocrinol.* 287, 113350.  
 2066 <https://doi.org/10.1016/j.ygcen.2019.113350>  
 2067 McConnachie, S.H., Cook, K. V., Patterson, D.A., Gilmour, K.M., Hinch, S.G., Farrell, A.P.,  
 2068 Cooke, S.J., 2012. Consequences of acute stress and cortisol manipulation on the  
 2069 physiology, behavior, and reproductive outcome of female Pacific salmon on spawning  
 2070 grounds. *Horm. Behav.* 62, 67–76. <https://doi.org/10.1016/j.yhbeh.2012.05.001>  
 2071 McCormick, M.I., 2009. Indirect effects of heterospecific interactions on progeny size  
 2072 through maternal stress. *Oikos* 118, 744–752. [https://doi.org/10.1111/j.1600-](https://doi.org/10.1111/j.1600-0706.2008.17410.x)  
 2073 [0706.2008.17410.x](https://doi.org/10.1111/j.1600-0706.2008.17410.x)  
 2074 McCormick, M.I., 2006. Mothers matter: Crowding leads to stressed mothers and smaller  
 2075 offspring in marine fish. *Ecology* 87, 1104–1109. [https://doi.org/10.1890/0012-](https://doi.org/10.1890/0012-9658(2006)87[1104:MMCLTS]2.0.CO;2)  
 2076 [9658\(2006\)87\[1104:MMCLTS\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[1104:MMCLTS]2.0.CO;2)  
 2077 McCormick, M.I., 1999. Experimental test of the effect of maternal hormones on larval  
 2078 quality of a coral reef fish. *Oecologia* 118, 412–422.  
 2079 <https://doi.org/10.1007/s004420050743>  
 2080 McCormick, M.I., 1998. Behaviorally induced maternal stress in a fish influences progeny  
 2081 quality by a hormonal mechanism. *Ecology* 79, 1873–1883.  
 2082 [https://doi.org/10.1890/0012-9658\(1998\)079\[1873:BIMSIA\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1998)079[1873:BIMSIA]2.0.CO;2)  
 2083 McCormick, S.D., 2001. Endocrine control of osmoregulation in teleost fish. *Am. Zool.* 41,  
 2084 781–794. <https://doi.org/10.1093/icb/41.4.781>  
 2085 McCormick, S.D., Regish, A., O’Dea, M.F., Shrimpton, J.M., 2008. Are we missing a  
 2086 mineralocorticoid in teleost fish? Effects of cortisol, deoxycorticosterone and  
 2087 aldosterone on osmoregulation, gill Na<sup>+</sup>,K<sup>+</sup>-ATPase activity and isoform mRNA levels

2088 in Atlantic salmon. *Gen. Comp. Endocrinol.* 157, 35–40.

2089 <https://doi.org/10.1016/j.ygcen.2008.03.024>

2090 McCrohan, C.R., Lu, W., Brierley, M.J., Dow, L., Balment, R.J., 2007. Fish caudal

2091 neurosecretory system: A model for the study of neuroendocrine secretion. *Gen. Comp.*

2092 *Endocrinol.* 153, 243–250. <https://doi.org/10.1016/j.ygcen.2006.12.027>

2093 McQuillan, H.J., Lokman, P.M., Young, G., 2003. Effects of sex steroids, sex, and sexual

2094 maturity on cortisol production: An in vitro comparison of chinook salmon and rainbow

2095 trout interrenals. *Gen. Comp. Endocrinol.* 133, 154–163. [https://doi.org/10.1016/S0016-](https://doi.org/10.1016/S0016-6480(03)00163-1)

2096 [6480\(03\)00163-1](https://doi.org/10.1016/S0016-6480(03)00163-1)

2097 Medeiros, L.R., Cartolano, M.C., McDonald, M.D., 2014. Crowding stress inhibits serotonin

2098 1A receptor-mediated increases in corticotropin-releasing factor mRNA expression and

2099 adrenocorticotropin hormone secretion in the Gulf toadfish. *J. Comp. Physiol. B*

2100 *Biochem. Syst. Environ. Physiol.* 184, 259–271. [https://doi.org/10.1007/s00360-013-](https://doi.org/10.1007/s00360-013-0793-9)

2101 [0793-9](https://doi.org/10.1007/s00360-013-0793-9)

2102 Medeiros, L.R., Mager, E.M., Grosell, M., McDonald, M.D., 2010. The serotonin subtype 1A

2103 receptor regulates cortisol secretion in the Gulf toadfish, *Opsanus beta*. *Gen. Comp.*

2104 *Endocrinol.* 168, 377–387. <https://doi.org/10.1016/j.ygcen.2010.05.004>

2105 Medeiros, L.R., McDonald, M.D., 2012. Elevated cortisol inhibits adrenocorticotropic

2106 hormone- and serotonin-stimulated cortisol secretion from the interrenal cells of the gulf

2107 toadfish (*Opsanus beta*). *Gen. Comp. Endocrinol.* 179, 414–420.

2108 <https://doi.org/10.1016/j.ygcen.2012.09.011>

2109 Metz, J.R., Geven, E.J.W., Van Den Burg, E.H., Flik, G., 2005. ACTH,  $\alpha$ -MSH, and control

2110 of cortisol release: Cloning, sequencing, and functional expression of the melanocortin-2

2111 and melanocortin-5 receptor in *Cyprinus carpio*. *Am. J. Physiol. - Regul. Integr. Comp.*

2112 *Physiol.* 289, 814–826. <https://doi.org/10.1152/ajpregu.00826.2004>

2113 Metz, J.R., Huising, M.O., Meek, J., Taverne-Thiele, A.J., Bonga, S.E.W., Flik, G., 2004.  
 2114       Localization, expression and control of adrenocorticotrophic hormone in the nucleus  
 2115       preopticus and pituitary gland of common carp (*Cyprinus carpio* L.). *J. Endocrinol.* 182,  
 2116       23–31. <https://doi.org/10.1677/joe.0.1820023>

2117 Meunier, H., Lefèvre, G., Dumont, D., Labrie, F., 1982. CRF stimulates alpha-MSH secretion  
 2118       and cyclic AMP accumulation in rat pars intermedia cells. *Life Sci.* 31, 2129–2135.

2119 Meyer, A., Van De Peer, Y., 2005. From 2R to 3R: Evidence for a fish-specific genome  
 2120       duplication (FSGD). *BioEssays* 27, 937–945. <https://doi.org/10.1002/bies.20293>

2121 Mileva, V.R., Gilmour, K.M., Balshine, S., 2011. Effects of maternal stress on egg  
 2122       characteristics in a cooperatively breeding fish. *Comp. Biochem. Physiol. - A Mol.*  
 2123       *Integr. Physiol.* 158, 22–29. <https://doi.org/10.1016/j.cbpa.2010.08.017>

2124 Milla, S., Terrien, X., Sturm, A., Ibrahim, F., Giton, F., Fiet, J., Prunet, P., Le Gac, F., 2008.  
 2125       Plasma 11-deoxycorticosterone (DOC) and mineralocorticoid receptor testicular  
 2126       expression during rainbow trout *Oncorhynchus mykiss* spermiation: Implication with  
 2127       17alpha, 20beta-dihydroxyprogesterone on the milt fluidity? *Reprod. Biol. Endocrinol.*  
 2128       6, 1–13. <https://doi.org/10.1186/1477-7827-6-19>

2129 Milla, S., Wang, N., Mandiki, S.N.M., Kestemont, P., 2009. Corticosteroids: Friends or foes  
 2130       of teleost fish reproduction? *Comp. Biochem. Physiol. - A Mol. Integr. Physiol.* 153,  
 2131       242–251. <https://doi.org/10.1016/j.cbpa.2009.02.027>

2132 Millar, R.P., Lu, Z., Pawson, A.J., Flanagan, C.A., Morgan, K., Maudsley, S.R., Human, C.,  
 2133       Sciences, R., 2004. Gonadotropin-Releasing Hormone Receptors. *Endocr. Rev.* 25, 235–  
 2134       275. <https://doi.org/10.1210/er.2003-0002>

2135 Miller, K.A., Kenter, L.W., Breton, T.S., Berlinsky, D.L., 2019. The effects of stress, cortisol  
 2136       administration and cortisol inhibition on black sea bass (*Centropristis striata*) sex  
 2137       differentiation. *Comp. Biochem. Physiol. -Part A Mol. Integr. Physiol.* 227, 154–160.

2138 <https://doi.org/10.1016/j.cbpa.2018.10.009>

2139 Miyagawa, S., Lange, A., Tohyama, S., Ogino, Y., Mizutani, T., Kobayashi, T., Tatarazako,  
 2140 N., Tyler, C.R., Iguchi, T., 2015. Characterization of *Oryzias latipes* glucocorticoid  
 2141 receptors and their unique response to progestins. *J. Appl. Toxicol.* 35, 302–309.  
 2142 <https://doi.org/10.1002/jat.3020>

2143 Mommsen, T.P., Vijayan, M.M., Moon, T.W., 1999. Cortisol in teleosts: Dynamics,  
 2144 mechanisms of action, and metabolic regulation. *Rev. Fish Biol. Fish.* 9, 211–268.  
 2145 <https://doi.org/10.1023/A:1008924418720>

2146 Montecucchi, P., Anastasi, A., de Castaglione, R., Erspamer, V., 1980. Isolation and amino  
 2147 acid composition of sauvagine. *Int J Pept. Protein Res* 16, 191–199.

2148 Morehead, D., 1998. Effect of capture, confinement and repeated sampling on plasma steroid  
 2149 concentrations and oocyte size in female striped trumpeter *Latris lineata* (Latrididae).  
 2150 *Mar. Freshw. Res.* 49, 373–377.

2151 Morgan, M.J., Wilson, C.E., Crim, L.W., 1999. The effect of stress on reproduction in  
 2152 Atlantic cod. *J. Fish Biol.* 54, 477–488. <https://doi.org/10.1006/jfbi.1998.0884>

2153 Morini, M., Peñaranda, D.S., Vilchez, M.C., Nourizadeh-Lillabadi, R., Lafont, A.G., Dufour,  
 2154 S., Asturiano, J.F., Weltzien, F.A., Pérez, L., 2017. Nuclear and membrane progestin  
 2155 receptors in the European eel: Characterization and expression in vivo through  
 2156 spermatogenesis. *Comp. Biochem. Physiol. -Part A Mol. Integr. Physiol.* 207, 79–92.  
 2157 <https://doi.org/10.1016/j.cbpa.2017.02.009>

2158 Mosconi, G., Gallinelli, A., Polzonetti-Magni, A., Facchinetti, F., 1998. Acetyl salmon  
 2159 endorphin-like and interrenal stress response in male gilthead sea bream, *Sparus aurata*.  
 2160 *Neuroendocrinology* 68, 129–134.

2161 Mousset, G., 1957. Sur la maturation sexuelle du congre. *Bull. Cent. Rech. Sc. Biarritz* 1, 559–  
 2162 561.

2163 Muñoz-Cueto, J.A., Paullada-Salmerón, J.A., Aliaga-Guerrero, M., Cowan, M.E., Parhar, I.S.,  
 2164 Ubuka, T., 2017. A journey through the gonadotropin-inhibitory hormone system of fish.  
 2165 Front. Endocrinol. (Lausanne). 8, 1–18. <https://doi.org/10.3389/fendo.2017.00285>  
 2166 Nakanishi, S., Inoue, A., Kita, T., Inoue, A., Nakamura, M., Chang, A.C.Y., Cohen, S.N.,  
 2167 Numa, S., 1979. Nucleotide sequence of cloned cDNA for bovine corticotropin- $\beta$ -  
 2168 lipotropin precursor. Nature 278, 423–427. <https://doi.org/10.1038/278423a0>  
 2169 Navarro-Martín, L., Viñas, J., Ribas, L., Díaz, N., Gutiérrez, A., Di Croce, L., Piferrer, F.,  
 2170 2011. DNA methylation of the gonadal aromatase (cyp19a) promoter is involved in  
 2171 temperature-dependent sex ratio shifts in the European sea bass. PLoS Genet. 7.  
 2172 <https://doi.org/10.1371/journal.pgen.1002447>  
 2173 Nikolarakis, K.E., Almeida, O.F.X., Herz, A., 1986. Corticotropin-releasing factor (CRF)  
 2174 inhibits gonadotropin-releasing hormone (GnRH) release from superfused rat  
 2175 hypothalami in vitro. Brain Res. 377, 388–390. [https://doi.org/10.1016/0006-](https://doi.org/10.1016/0006-8993(86)90887-5)  
 2176 8993(86)90887-5  
 2177 Nozu, R., Nakamura, M., 2015. Cortisol administration induces sex change from ovary to  
 2178 testis in the protogynous wrasse, Halichoeres trimaculatus. Sex. Dev. 9, 118–124.  
 2179 <https://doi.org/10.1159/000373902>  
 2180 Ogawa, S., Parhar, I.S., 2014. Structural and functional divergence of gonadotropin-inhibitory  
 2181 hormone from jawless fish to mammals. Front. Endocrinol. (Lausanne). 5, 1–17.  
 2182 <https://doi.org/10.3389/fendo.2014.00177>  
 2183 Ogawa, S., Sivalingam, M., Biran, J., Golan, M., Anthonysamy, R.S., Levavi-Sivan, B.,  
 2184 Parhar, I.S., 2016. Distribution of LPXRFa, a gonadotropin-inhibitory hormone ortholog  
 2185 peptide, and LPXRFa receptor in the brain and pituitary of the tilapia. J. Comp. Neurol.  
 2186 524, 2753–2775. <https://doi.org/10.1002/cne.23990>  
 2187 Ogino, Y., Katoh, H., Kuraku, S., Yamada, G., 2009. Evolutionary history and functional

2188 characterization of androgen receptor genes in jawed vertebrates. *Endocrinology* 150,  
 2189 5415–5427. <https://doi.org/10.1210/en.2009-0523>  
 2190 Ogino, Y., Tohyama, S., Kohno, S., Toyota, K., Yamada, G., Yatsu, R., Kobayashi, T.,  
 2191 Tatarazako, N., Sato, T., Matsubara, H., Lange, A., Tyler, C.R., Katsu, Y., Iguchi, T.,  
 2192 Miyagawa, S., 2018. Functional distinctions associated with the diversity of sex steroid  
 2193 hormone receptors ESR and AR. *J. Steroid Biochem. Mol. Biol.* 184, 38–46.  
 2194 <https://doi.org/10.1016/j.jsbmb.2018.06.002>  
 2195 Ohtaki, T., Shintani, Y., Honda, S., Matsumoto, H., Hori, A., Kanehashi, K., Terao, Y.,  
 2196 Kumano, S., Takatsu, Y., Masuda, Y., Ishibashi, Y., Watanabe, T., Asada, M., Yamada,  
 2197 T., Suenaga, M., Kitada, C., Usuki, S., Kurokawa, T., Onda, H., Nishimura, O., Fujino,  
 2198 M., 2001. Metastasis suppressor gene KiSS-1 encodes peptide ligand of a G-protein-  
 2199 coupled receptor. *Nature* 411, 613–617. <https://doi.org/10.1038/35079135>  
 2200 Olivereau, M., 1967. Observations sur l’hypophyse de l’Anguille femelle, en particulier lors  
 2201 de la maturation sexuelle. *Zeitschrift für Zellforsch. und Mikroskopische Anat.* 80, 286–  
 2202 306. <https://doi.org/10.1007/BF00337462>  
 2203 Olivereau, M., Ball, J., 1964. Contribution à l’histophysiologie de l’hypophyse des  
 2204 téléostéens, en particulier de celle de *Poecilia species*. *Gen. Comp. Endocrinol.* 4, 523–  
 2205 532.  
 2206 Omeljaniuk, R.J., Tonon, M.C., Peter, R.E., 1989. Dopamine inhibition of gonadotropin and  
 2207  $\alpha$ -melanocyte-stimulating hormone release in vitro from the pituitary of the goldfish  
 2208 (*Carassius auratus*). *Gen. Comp. Endocrinol.* 74, 451–467.  
 2209 [https://doi.org/10.1016/S0016-6480\(89\)80043-7](https://doi.org/10.1016/S0016-6480(89)80043-7)  
 2210 Ono, N., Lumpkin, M., Samson, W., McDonald, J., McCann, S., 1984. Intrahypothalamic  
 2211 action of corticotrophin-releasing factor (CRF) to inhibit growth hormone and LH  
 2212 release in the rat. *Life Sci.* 35, 1117–1123.

2213 <https://doi.org/10.1017/CBO9781107415324.004>

2214 Ozaki, Y., Higuchi, M., Miura, C., Yamaguchi, S., Tozawa, Y., Miura, T., 2006. Roles of  
 2215  $11\beta$ -hydroxysteroid dehydrogenase in fish spermatogenesis. *Endocrinology* 147, 5139–  
 2216 5146. <https://doi.org/10.1210/en.2006-0391>

2217 Palstra, A.P., van den Thillart, G.E.E.J.M., 2010. Swimming physiology of European silver  
 2218 eels (*Anguilla anguilla* L.): Energetic costs and effects on sexual maturation and  
 2219 reproduction. *Fish Physiol. Biochem.* 36, 297–322. [https://doi.org/10.1007/s10695-010-](https://doi.org/10.1007/s10695-010-9397-4)  
 2220 9397-4

2221 Pankhurst, N., Van der Kraak, G., 1997. Effects of stress on reproduction and growth of fish.  
 2222 In *Fish stress and health in aquaculture.*, in: Iwama, G., Pickering, A., Sumpter, J.,  
 2223 Schreck, C. (Eds.), *Fish Stress and Health in Aquaculture*. Society for Experimental  
 2224 Biology Seminar Series 62. University Press, Cambridge, pp. 73–93.

2225 Pankhurst, N., Van der Kraak, G., Peter, R., 1995. Evidence that the inhibitory effects of  
 2226 stress on reproduction in teleost fish are not mediated by the action of cortisol on ovarian  
 2227 steroidogenesis. *Gen. Comp. Endocrinol.* 99, 249–257.

2228 Pankhurst, N.W., 2016. Reproduction and development., in: Schreck, C., Tort, L., Farrell, A.,  
 2229 Brauner, C. (Eds.), *Biology of Stress in Fish*. *Fish Physiology Vol 35*. pp. 295–331.

2230 Pankhurst, N.W., 2001. Stress inhibition of reproductive endocrine processes in a natural  
 2231 population of the spiny damselfish *Acanthochromis polyacanthus*. *Mar Freshwater Res*  
 2232 52, 753–761.

2233 Pankhurst, N.W., 1998. Further evidence of the equivocal effects of cortisol on in vitro  
 2234 steroidogenesis by ovarian follicles of rainbow trout *Oncorhynchus mykiss*. *Fish*  
 2235 *Physiol. Biochem.* 19, 315–323. <https://doi.org/10.1023/A:1007751403189>

2236 Pankhurst, N.W., Dedual, M., 1994. Effects of capture and recovery on plasma levels of  
 2237 cortisol, lactate and gonadal steroids in a natural population of rainbow trout. *J. Fish*

2238 Biol. 45, 1013–1025.

2239 Pankhurst, N.W., Van Der Kraak, G., 2000. Evidence that acute stress inhibits ovarian  
 2240 steroidogenesis in rainbow trout in vivo, through the action of cortisol. *Gen. Comp.*  
 2241 *Endocrinol.* 117, 225–237. <https://doi.org/10.1006/gcen.1999.7401>

2242 Pasquier, J., Kamech, N., Lafont, A.G., Vaudry, H., Rousseau, K., Dufour, S., 2014.  
 2243 Molecular evolution of GPCRs: Kisspeptin/kisspeptin receptors. *J. Mol. Endocrinol.* 52.  
 2244 <https://doi.org/10.1530/JME-13-0224>

2245 Pasquier, J., Lafont, A.-G., Denis, F., Lefranc, B., Dubessy, C., Moreno-Herrera, A., Vaudry,  
 2246 H., Leprince, J., Dufour, S., Rousseau, K., 2018. Eel Kisspeptins: Identification,  
 2247 functional activity, and inhibition on both pituitary LH and GnRH receptor expression.  
 2248 *Front. Endocrinol. (Lausanne)*. 8. <https://doi.org/10.3389/fendo.2017.00353>

2249 Pasquier, J., Lafont, A.G., Tostivint, H., Vaudry, H., Rousseau, K., Dufour, S., 2012.  
 2250 Comparative evolutionary histories of kisspeptins and kisspeptin receptors in vertebrates  
 2251 reveal both parallel and divergent features. *Front. Endocrinol. (Lausanne)*. 3, 1–15.  
 2252 <https://doi.org/10.3389/fendo.2012.00173>

2253 Patterson, D.A., Macdonald, J.S., Hinch, S.G., Healey, M.C., Farrell, A.P., 2004. The effect  
 2254 of exercise and captivity on energy partitioning, reproductive maturation and fertilization  
 2255 success in adult sockeye salmon. *J. Fish Biol.* 64, 1039–1059.  
 2256 <https://doi.org/10.1111/j.1095-8649.2004.0370.x>

2257 Pearson, D., Shively, J., Clark, B., Geschwind, I., Barkley, M., Nishioka, R., Bern, H., 1980.  
 2258 Urotensin II: a somatostatin-like peptide in the caudal neurosecretory system of fishes.  
 2259 *Proc Natl Acad Sci USA* 77, 5021–5024. [https://doi.org/10.1111/j.1399-](https://doi.org/10.1111/j.1399-3011.1982.tb02629.x)  
 2260 [3011.1982.tb02629.x](https://doi.org/10.1111/j.1399-3011.1982.tb02629.x)

2261 Peng, W., Cao, M., Chen, J., Li, Y., Wang, Y., Zhu, Z., Hu, W., 2016. GnIH plays a negative  
 2262 role in regulating GtH expression in the common carp, *Cyprinus carpio* L. *Gen. Comp.*

2263       Endocrinol. 235, 18–28. <https://doi.org/10.1016/j.ygcen.2016.06.001>

2264   Perry, A.N., Grober, M.S., 2003. A model for social control of sex change: Interactions of  
2265       behavior, neuropeptides, glucocorticoids, and sex steroids. *Horm. Behav.* 43, 31–38.  
2266       [https://doi.org/10.1016/S0018-506X\(02\)00036-3](https://doi.org/10.1016/S0018-506X(02)00036-3)

2267   Philipp, D.P., Toline, C.A., Kubacki, M.F., Philipp, D.B.F., Phelan, F.J.S., 1997. The Impact  
2268       of Catch-and-Release Angling on the Reproductive Success of Smallmouth Bass and  
2269       Largemouth Bass. *North Am. J. Fish. Manag.* 17, 557–567.  
2270       [https://doi.org/10.1577/1548-8675\(1997\)017<0557:tiocar>2.3.co;2](https://doi.org/10.1577/1548-8675(1997)017<0557:tiocar>2.3.co;2)

2271   Pickering, A.D., 1989. Environmental stress and the survival of brown trout, *Salmo trutta*.  
2272       *Freshw. Biol.* 21, 47–55. <https://doi.org/10.1111/j.1365-2427.1989.tb01347.x>

2273   Pickering, A.D., Pottinger, T.G., Carragher, J., Sumpter, J.P., 1987. The effects of acute and  
2274       chronic stress on the levels of reproductive hormones in the plasma of mature male  
2275       brown trout, *Salmo trutta* L. *Gen. Comp. Endocrinol.* 68, 249–259.  
2276       [https://doi.org/10.1016/0016-6480\(87\)90036-0](https://doi.org/10.1016/0016-6480(87)90036-0)

2277   Pierce, J.G., Parsons, T.F., 1981. Glycoprotein Hormones: Structure and Function. *Annu.*  
2278       *Rev. Biochem.* 50, 465–495. <https://doi.org/10.1146/annurev.bi.50.070181.002341>

2279   Pierson, P., Guibbolini, M., Lahlou, B., 1996. A V1-type receptor for mediating the  
2280       neurohypophysial hormone-induced ACTH release in trout pituitary. *J Endocrinol* 149,  
2281       109–115.

2282   Pinilla, L., Aguilar, E., Dieguez, C., Millar, R.P., Tena-Sempere, M., 2012. Kisspeptins and  
2283       reproduction: Physiological roles and regulatory mechanisms. *Physiol. Rev.* 92, 1235–  
2284       1316. <https://doi.org/10.1152/physrev.00037.2010>

2285   Pogoda, H.M., Hammerschmidt, M., 2007. Molecular genetics of pituitary development in  
2286       zebrafish. *Semin. Cell Dev. Biol.* 18, 543–558.  
2287       <https://doi.org/10.1016/j.semcdb.2007.04.004>

2288 Pohl, S., Darlison, M.G., Clarke, W.C., Lederis, K., Richter, D., 2001. Cloning and functional  
 2289 pharmacology of two corticotropin-releasing factor receptors from a teleost fish. *Eur. J.*  
 2290 *Pharmacol.* 430, 193–202. [https://doi.org/10.1016/S0014-2999\(01\)01391-7](https://doi.org/10.1016/S0014-2999(01)01391-7)

2291 Pottinger, T., Yeomans, W., Carrick, T., 1999. Plasma cortisol and 17 -oestradiol levels in  
 2292 roach exposed to acute and chronic stress. *J. Fish Biol.* 54, 525–532.

2293 Pottinger, T.G., Carrick, T.R., Hughes, S.E., Balm, P.H.M., 1996. Testosterone, 11-  
 2294 ketotestosterone, and estradiol-17 $\beta$  modify baseline and stress-induced interrenal and  
 2295 corticotropic activity in trout. *Gen. Comp. Endocrinol.* 104, 284–295.  
 2296 <https://doi.org/10.1006/gcen.1996.0173>

2297 Pottinger, T.G., Pickering, A.D., 1990. The effect of cortisol administration on hepatic and  
 2298 plasma estradiol-binding capacity in immature female rainbow trout (*Oncorhynchus*  
 2299 *mykiss*). *Gen. Comp. Endocrinol.* 80, 264–273. [https://doi.org/10.1016/0016-](https://doi.org/10.1016/0016-6480(90)90171-H)  
 2300 [6480\(90\)90171-H](https://doi.org/10.1016/0016-6480(90)90171-H)

2301 Proulx-Ferland, L., Labrie, F., Dumont, D., Côté, J., 1982. Corticotropin-Releasing Factor.  
 2302 *Science* (80-. ). 217, 62–63.

2303 Prunet, P., Sturm, A., Milla, S., 2006. Multiple corticosteroid receptors in fish: From old ideas  
 2304 to new concepts. *Gen. Comp. Endocrinol.* 147, 17–23.  
 2305 <https://doi.org/10.1016/j.ygcen.2006.01.015>

2306 Puglisi-Allegra, S., Andolina, D., 2015. Serotonin and stress coping. *Behav. Brain Res.* 277,  
 2307 58–67. <https://doi.org/10.1016/j.bbr.2014.07.052>

2308 Putteeraj, M., Soga, T., Ubuka, T., Parhar, I.S., 2016. A “timed” kiss is essential for  
 2309 reproduction: Lessons from mammalian studies. *Front. Endocrinol. (Lausanne).* 7, 1–10.  
 2310 <https://doi.org/10.3389/fendo.2016.00121>

2311 Qi, X., Zhou, W., Li, S., Lu, D., Yi, S., Xie, R., Liu, X., Zhang, Y., Lin, H., 2013. Evidences  
 2312 for the regulation of GnRH and GTH expression by GnIH in the goldfish, *Carassius*

2313        auratus. *Mol. Cell. Endocrinol.* 366, 9–20. <https://doi.org/10.1016/j.mce.2012.11.001>  
 2314    Raftogianni, A., Roth, L.C., García-González, D., Bus, T., Kühne, C., Monyer, H., Spergel,  
 2315        D.J., Deussing, J.M., Grinevich, V., 2018. Deciphering the Contributions of CRH  
 2316        Receptors in the Brain and Pituitary to Stress-Induced Inhibition of the Reproductive  
 2317        Axis. *Front. Mol. Neurosci.* 11, 1–20. <https://doi.org/10.3389/fnmol.2018.00305>  
 2318    Rainey, W.E., Nakamura, Y., 2008. Regulation of the adrenal androgen biosynthesis. *J.*  
 2319        *Steroid Biochem. Mol. Biol.* 108, 281–286. <https://doi.org/10.1016/j.jsbmb.2007.09.015>  
 2320    Rance, T.A., Baker, B.I., 1981. The in vitro response of the trout interrenal to various  
 2321        fragments of ACTH. *Gen. Comp. Endocrinol.* 45, 497–503.  
 2322        [https://doi.org/10.1016/0016-6480\(81\)90054-X](https://doi.org/10.1016/0016-6480(81)90054-X)  
 2323    Reddy, P.K., Renaud, R., Leatherland, J.F., 1999. Effects of cortisol and triiodo-L-thyronine  
 2324        on the steroidogenic capacity of rainbow trout ovarian follicles at two stages of oocyte  
 2325        maturation. *Fish Physiol. Biochem.* 21, 129–140.  
 2326        <https://doi.org/10.1023/A:1007800707094>  
 2327    Remer, T., Shi, L., Buyken, A.E., Maser-Gluth, C., Hartmann, M.F., Wudy, S.A., 2010.  
 2328        Prepubertal adrenarchal androgens and animal protein intake independently and  
 2329        differentially influence pubertal timing. *J. Clin. Endocrinol. Metab.* 95, 3002–3009.  
 2330        <https://doi.org/10.1210/jc.2009-2583>  
 2331    Reyes, T.M., Lewis, K., Perrin, M.H., Kunitake, K.S., Vaughan, J., Arias, C.A., Hogenesch,  
 2332        J.B., Gulyas, J., Rivier, J., Vale, W.W., Sawchenko, P.E., 2001. Urocortin II: A member  
 2333        of the corticotropin-releasing factor (CRF) neuropeptide family that is selectively bound  
 2334        by type 2 CRF receptors. *Proc. Natl. Acad. Sci. U. S. A.* 98, 2843–2848.  
 2335        <https://doi.org/10.1073/pnas.051626398>  
 2336    Rivier, C., Plotsky, P., 1986. Mediation by Corticotropin Releasing Factor (CRF) of  
 2337        Adenohypophysial Hormone Secretion. *Annu. Rev. Physiol.* 48, 475–494.

2338 <https://doi.org/10.1146/annurev.physiol.48.1.475>

2339 Rivier, C., Rivier, J., Lederis, K., Vale, W., 1983. In vitro and in vivo ACTH-releasing  
 2340 activity of ovine CRF, sauvagine and urotensin I. Regul. Pept. 5, 139–143.  
 2341 [https://doi.org/10.1016/0167-0115\(83\)90121-0](https://doi.org/10.1016/0167-0115(83)90121-0)

2342 Rivier, C., Rivier, J., Vale, W., 1986. Stress-induced inhibition of reproductive functions: role  
 2343 of endogenous corticotropin-releasing factor. Science (80-. ). 231, 607–609.

2344 Rivier, C., Vale, W., 1984. Influence of corticotropin-releasing factor on reproductive  
 2345 functions in the rat. Endocrinology 114, 914–921. [https://doi.org/10.1210/endo-114-3-](https://doi.org/10.1210/endo-114-3-914)  
 2346 914

2347 Rivier, C., Vale, W., 1983. Interaction of corticotropin-releasing factor and arginine  
 2348 vasopressin on corticotropin secretion in vivo. Endocrinology 113, 939–942.

2349 Robertson, F.M., Gundappa, M.K., Grammes, F., Hvidsten, T.R., Redmond, A.K., Lien, S.,  
 2350 Martin, S.A.M., Holland, P.W.H., Sandve, S.R., Macqueen, D.J., 2017. Lineage-specific  
 2351 rediploidization is a mechanism to explain time-lags between genome duplication and  
 2352 evolutionary diversification. Genome Biol. 18, 1–14. [https://doi.org/10.1186/s13059-](https://doi.org/10.1186/s13059-017-1241-z)  
 2353 017-1241-z

2354 Robertson, O., Krupp, M., Thomas, S.F., Favourjz, C., Hane, S., Wexler, B.C., 1961.  
 2355 Hyperadrenocorticism in Spawning migratory and Nonmigratory Rainbow Trout (*Salmo*  
 2356 *gairdnerii*) ; Comparison with Pacific Salmon (Genus *Oncorhynchus*). Gen. Comp.  
 2357 Endocrinol. 1, 473–484.

2358 Robertson, O.H., 1961. Prolongation of the Life Span of Kokanee Salmon (*Oncorhynchus*  
 2359 *Nerka Kennerlyi*) By Castration Before Beginning of Gonad Development. Proc. Natl.  
 2360 Acad. Sci. 47, 609–621. <https://doi.org/10.1073/pnas.47.4.609>

2361 Roch, G.J., Busby, E.R., Sherwood, N.M., 2014. GnRH receptors and peptides: Skating  
 2362 backward. Gen. Comp. Endocrinol. 209, 118–134.

2363        <https://doi.org/10.1016/j.ygcen.2014.07.025>

2364    Roch, G.J., Busby, E.R., Sherwood, N.M., 2011. Evolution of GnRH: Diving deeper. *Gen.*

2365        *Comp. Endocrinol.* 171, 1–16. <https://doi.org/10.1016/j.ygcen.2010.12.014>

2366    Rotllant, J., Balm, P.H.M., Pérez-Sánchez, J., Wendelaar-Bonga, S.E., Tort, L., 2001.

2367        Pituitary and interrenal function in gilthead sea bream (*Sparus aurata* L., Teleostei) after

2368        handling and confinement stress. *Gen. Comp. Endocrinol.* 121, 333–342.

2369        <https://doi.org/10.1006/gcen.2001.7604>

2370    Rotllant, J., Balm, P.H.M., Ruane, N.M., Pérez-Sánchez, J., Wendelaar-Bonga, S.E., Tort, L.,

2371        2000. Pituitary proopiomelanocortin-derived peptides and hypothalamus-pituitary-

2372        interrenal axis activity in gilthead sea bream (*sparus aurata*) during prolonged crowding

2373        stress: Differential regulation of adrenocorticotropin hormone and  $\alpha$ -melanocyte-

2374        stimulating . *Gen. Comp. Endocrinol.* 119, 152–163.

2375        <https://doi.org/10.1006/gcen.2000.7508>

2376    Saccol, E.M.H., Jerez-Cepa, I., Ourique, G.M., Pês, T.S., Gressler, L.T., Mourão, R.H.V.,

2377        Martínez-Rodríguez, G., Mancera, J.M., Baldisserotto, B., Pavanato, M.A., Martos-

2378        Sitcha, J.A., 2018. *Myrcia sylvatica* essential oil mitigates molecular, biochemical and

2379        physiological alterations in *Rhamdia quelen* under different stress events associated to

2380        transport. *Res. Vet. Sci.* 117, 150–160. <https://doi.org/10.1016/j.rvsc.2017.12.009>

2381    Safford, S., Thomas, P., 1987. Effects of capture and handling on circulating levels of gonadal

2382        steroids and cortisol in the spotted seatrout, *Cynoscion nebulosus*., in: Idler, D., Crim, L.,

2383        Walsh, J. (Eds.), *Reproductive Physiology of Fish*. St John's: Memorial University of

2384        Newfoundland., p. 312.

2385    Sakamoto, T., Yoshiki, M., Takahashi, H., Yoshida, M., Ogino, Y., Ikeuchi, T., Nakamachi,

2386        T., Konno, N., Matsuda, K., Sakamoto, H., 2016. Principal function of mineralocorticoid

2387        signaling suggested by constitutive knockout of the mineralocorticoid receptor in

2388 medaka fish. *Sci. Rep.* 6, 1–10. <https://doi.org/10.1038/srep37991>

2389 Saland, L.C., Mennin, S.P., Selinfreund, R., Rasmussen, P., 1982. Interaction of  $\beta$ -endorphin,  
 2390 naloxone and dopamine: effects on melanocyte-stimulating hormone secretion of  
 2391 amphibian pituitaries in vitro. *Regul. Pept.* 3, 371–381. [https://doi.org/10.1016/0167-](https://doi.org/10.1016/0167-0115(82)90060-X)  
 2392 [0115\(82\)90060-X](https://doi.org/10.1016/0167-0115(82)90060-X)

2393 Satake, H., Hisada, M., Kawada, T., Minakata, H., Ukena, K., Tsutsui, K., 2001.  
 2394 Characterization of a cDNA encoding a novel avian hypothalamic neuropeptide exerting  
 2395 an inhibitory effect on gonadotropin release. *Biochem. J.* 354, 379–385.

2396 Sbaihi, M., Rousseau, K., Baloché, S., Meunier, F., Fouchereau-Peron, M., Dufour, S., 2009.  
 2397 Cortisol mobilizes mineral stores from vertebral skeleton in the European eel: An  
 2398 ancestral origin for glucocorticoid-induced osteoporosis? *J. Endocrinol.* 201, 241–252.  
 2399 <https://doi.org/10.1677/JOE-08-0492>

2400 Schaaf, M.J.M., Champagne, D., Van Laanen, I.H.C., Van Wijk, D.C.W.A., Meijer, A.H.,  
 2401 Meijer, O.C., Spaijk, H.P., Richardson, M.K., 2008. Discovery of a functional  
 2402 glucocorticoid receptor  $\beta$ -isoform in zebrafish. *Endocrinology* 149, 1591–1598.  
 2403 <https://doi.org/10.1210/en.2007-1364>

2404 Schmidt, P.J., Idler, D.R., 1962. Steroid hormones in the plasma of salmon at various states of  
 2405 maturation. *Gen. Comp. Endocrinol.* 2, 204–214. [https://doi.org/10.1016/0016-](https://doi.org/10.1016/0016-6480(62)90005-9)  
 2406 [6480\(62\)90005-9](https://doi.org/10.1016/0016-6480(62)90005-9)

2407 Schreck, C., 2000. Accumulation and long-term effects of stress in fish., in: Moberg, G.,  
 2408 Mench, J. (Eds.), *The Biology of Animal Stress*. CAB International, Wallingford, pp.  
 2409 147–158.

2410 Schreck, C., Tort, L., 2016. The concept of stress in fish., in: Schreck, C., Tort, L., Farrell, A.,  
 2411 Brauner, C. (Eds.), *Biology of Stress in Fish*. *Fish Physiology Vol 35*. pp. 1–34.

2412 Schreck, C.B., 2010. Stress and fish reproduction: The roles of allostasis and hormesis. *Gen.*

2413 Comp. Endocrinol. 165, 549–556. <https://doi.org/10.1016/j.ygcen.2009.07.004>

2414 Schreck, C.B., Bradford, C.S., Fitzpatrick, M.S., Patiño, R., 1989. Regulation of the interrenal  
2415 of fishes: non-classical control mechanisms. *Fish Physiol. Biochem.* 7, 259–265.  
2416 <https://doi.org/10.1007/BF00004715>

2417 Schreibman, M., Leatherland, J., McKeown, B., 1973. Functional morphology of the teleost  
2418 pituitary gland. *Amer Zool* 13, 719–742.

2419 Schwartzentruber, R., Vaudry, H., Omeljaniuk, R., 1994. Structural requirements for  
2420 stimulation by thyrotropin-releasing hormone of alpha-MSH release from rainbow trout  
2421 (*Oncorhynchus mykiss*) pituitary fragments in vitro. *Rev. Can. Zool.* 72, 1872–1879.

2422 Scott, P., Canario, A., Sherwood, N., Warby, C., 1991a. Levels of steroids, including cortisol  
2423 and 17 $\alpha$ ,20 $\beta$ -dihydroxy-4-pregnen-3-one, in plasma, seminal fluid, and urine of Pacific  
2424 herring (*Clupea harengus pallasi*) and North Sea plaice (*Pleuronectes platessa* L.). *Can. J.*  
2425 *Zool.* 69, 111–116.

2426 Scott, P., Sherwood, N., Canario, A., Warby, C., 1991b. Identification of free and conjugated  
2427 steroids, including cortisol and 17 $\alpha$ ,20 $\beta$ -dihydroxy-4-pregnen-3-one, in the milt of  
2428 Pacific herring, *Clupea harengus*. *Can. J. Zool.* 69, 104–110.

2429 Sefideh, F.A., Moon, M.J., Yun, S., Hong, S.I., Hwang, J.I., Seong, J.Y., 2014. Local  
2430 duplication of gonadotropin-releasing hormone (GnRH) receptor before two rounds of  
2431 whole genome duplication and origin of the mammalian GnRH receptor. *PLoS One* 9.  
2432 <https://doi.org/10.1371/journal.pone.0087901>

2433 Segner, H., Sundh, H., Buchmann, K., Dourfils, J., Sundell, K.S., Mathieu, C., Ruane, N.,  
2434 Jutfelt, F., Toften, H., Vaughan, L., 2012. Health of farmed fish: Its relation to fish  
2435 welfare and its utility as welfare indicator. *Fish Physiol. Biochem.* 38, 85–105.  
2436 <https://doi.org/10.1007/s10695-011-9517-9>

2437 Selye, H., 1973. The evolution of the stress concept. *J.Int.Acad.Prev.Med.* 2, 1–21.

2438 [https://doi.org/10.1016/0002-9149\(70\)90796-4](https://doi.org/10.1016/0002-9149(70)90796-4)

2439 Selye, H., 1939. Effect of adaptation to various damaging agents on the female sex organs in  
 2440 the rat. *Endocrinology* 25, 615–624.

2441 Seminara, S.B., Messenger, S., Chatzidaki, E.E., Thresher, R.R., Acierno, J.S., Shagoury, J.K.,  
 2442 Bo-Abbas, Y., Kuohung, W., Schwinof, K.M., Hendrick, A.G., Zahn, D., Dixon, J.,  
 2443 Kaiser, U.B., Slaugenhaupt, S.A., Gusella, J.F., O’Rahilly, S., Carlton, M.B.L., Crowley,  
 2444 W.F., Aparicio, S.A.J.R., Colledge, W.H., 2004. The GPR54 Gene as a Regulator of  
 2445 Puberty. *N. Engl. J. Med.* 59, 351–353. [https://doi.org/10.1097/00006254-200405000-](https://doi.org/10.1097/00006254-200405000-00020)  
 2446 00020

2447 Shahjahan, M., Ikegami, T., Osugi, T., Ukena, K., Doi, H., Hattori, A., Tsutsui, K., Ando, H.,  
 2448 2011. Synchronised expressions of LPXRFamide peptide and its receptor genes:  
 2449 Seasonal, diurnal and circadian changes during spawning period in grass puffer. *J.*  
 2450 *Neuroendocrinol.* 23, 39–51. <https://doi.org/10.1111/j.1365-2826.2010.02081.x>

2451 Shi, C., Lu, Y., Zhai, G., Huang, J., Shang, G., Lou, Q., Li, D., Jin, X., He, J., Du, Z., Gui, J.,  
 2452 Yin, Z., 2019. Hyperandrogenism in POMCa-deficient zebrafish enhances somatic  
 2453 growth without increasing adiposity. *J. Mol. Cell Biol.* 1–31.  
 2454 <https://doi.org/10.1093/jmcb/mjz053>

2455 Skrzynska, A.K., Maiorano, E., Bastaroli, M., Naderi, F., Míguez, J.M., Martínez-Rodríguez,  
 2456 G., Mancera, J.M., Martos-Sitcha, J.A., 2018. Impact of air exposure on vasotocinergic  
 2457 and isotocinergic systems in gilthead sea bream (*Sparus aurata*): New insights on fish  
 2458 stress response. *Front. Physiol.* 9. <https://doi.org/10.3389/fphys.2018.00096>

2459 Small, B.C., 2004. Effect of dietary cortisol administration on growth and reproductive  
 2460 success of channel catfish. *J. Fish Biol.* 64, 589–596. [https://doi.org/10.1111/j.1095-](https://doi.org/10.1111/j.1095-8649.2004.00322.x)  
 2461 8649.2004.00322.x

2462 Sneddon, L., Wolfenden, D., Thomson, J., 2016. Stress management and welfare., in:

2463 Schreck, C., Tort, L., Farell, A., Brauner, C. (Eds.), *Fish Physiology: Biology of Stress*  
 2464 *in Fish Vol 35*. Academic Press, San Diego, pp. 463–539.  
 2465 Socha, M., Sokolowska-Mikolajczyk, M., Mikoajczyk, T., Chyb, J., Epler, P., 2003.  
 2466 Naltrexone Stimulates LH Secretion in Vitro From the common carpio (*Cyprinus carpio*  
 2467 L.) pituitary gland. *Arch. Polish Fish.* 11, 5–16.  
 2468 Solomon-Lane, T.K., Crespi, E.J., Grober, M.S., 2013. Stress and serial adult metamorphosis:  
 2469 Multiple roles for the stress axis in socially regulated sex change. *Front. Neurosci.* 7, 1–  
 2470 12. <https://doi.org/10.3389/fnins.2013.00210>  
 2471 Son, Y.L., Ubuka, T., Narihiro, M., Fukuda, Y., Hasunuma, I., Yamamoto, K., Belsham,  
 2472 D.D., Tsutsui, K., 2014. Molecular basis for the activation of gonadotropin-inhibitory  
 2473 hormone gene transcription by corticosterone. *Endocrinology* 155, 1817–1826.  
 2474 <https://doi.org/10.1210/en.2013-2076>  
 2475 Soso, A.B., Gil Barcellos, L.J., Ranzani-paiva, M.J., Kreutz, L.C., Quevedo, R.M., Lima, M.,  
 2476 Bolognesi da Silva, L., Ritter, F., Bedin, A.C., Finco, J.A., 2008. The effects of stressful  
 2477 broodstock handling on hormonal profiles and reproductive performance of rhamdia  
 2478 quelen (*Quoy & Gaimard*) females. *J. World Aquac. Soc.* 39, 835–841.  
 2479 <https://doi.org/10.1111/j.1749-7345.2008.00221.x>  
 2480 Sousa, M.L., Figueiredo, F., Pinheiro, C., Silva, A., Malhão, F., Rocha, M.J., Rocha, E.,  
 2481 Urbatzka, R., 2015. Morphological and molecular effects of cortisol and ACTH on  
 2482 zebrafish stage I and II Follicles. *Reproduction* 150, 429–436.  
 2483 <https://doi.org/10.1530/REP-15-0271>  
 2484 Sower, S.A., 2018. Landmark discoveries in elucidating the origins of the hypothalamic-  
 2485 pituitary system from the perspective of a basal vertebrate, sea lamprey. *Gen. Comp.*  
 2486 *Endocrinol.* 264, 3–15. <https://doi.org/10.1016/j.ygcen.2017.10.016>  
 2487 Spicer, O.S., Zmora, N., Wong, T.T., Golan, M., Levavi-Sivan, B., Gothilf, Y., Zohar, Y.,

2488 2017. The gonadotropin-inhibitory hormone (Lpxrfa) system's regulation of  
 2489 reproduction in the brain-pituitary axis of the zebrafish (*Danio rerio*). *Biol. Reprod.* 96,  
 2490 1031–1042. <https://doi.org/10.1093/biolre/iox032>

2491 Stevens, C.H., Croft, D.P., Paull, G.C., Tyler, C.R., 2017. Stress and welfare in ornamental  
 2492 fishes: what can be learned from aquaculture? *J. Fish Biol.* 91, 409–428.  
 2493 <https://doi.org/10.1111/jfb.13377>

2494 Stolte, E.H., Mazon, A.F. De, Leon-koosterziel, K.M., Jesiak, M., Bury, N.R., Sturm, A.,  
 2495 Savelkoul, H.F.J., Kemenade, B.M.L.V. Van, Flik, G., 2008. Corticosteroid receptors  
 2496 involved in stress regulation in common carp, *Cyprinus carpio*. *J. Endocrinol.* 198, 403–  
 2497 417. <https://doi.org/10.1677/J0E-08-0100>

2498 Stolte, E.H., Verburg van Kemenade, B.M.L., Savelkoul, H.F.J., Flik, G., 2006. Evolution of  
 2499 glucocorticoid receptors with different glucocorticoid sensitivity. *J. Endocrinol.* 190, 17–  
 2500 28. <https://doi.org/10.1677/joe.1.06703>

2501 Sturm, A., Bury, N., Dengreville, L., Fagart, J., Flouriot, G., Rafestin-Oblin, M.E., Prunet, P.,  
 2502 2005. 11-Deoxycorticosterone is a potent agonist of the rainbow trout (*Oncorhynchus*  
 2503 *mykiss*) mineralocorticoid receptor. *Endocrinology* 146, 47–55.  
 2504 <https://doi.org/10.1210/en.2004-0128>

2505 Sundararaj, B., Goswami, S., 1966a. Effects of mammalian hypophysial hormones, placental  
 2506 gonadotropins, gonadal hormones, and adrenal corticosteroids on ovulation and  
 2507 spawning in hypophysectomized catfish, *Heteropneustes fossilis* (Bloch). *J. Exp. Zool.* 161,  
 2508 287–296.

2509 Sundararaj, B., Goswami, S., 1966b. Effect of metopiron (SU-4885) on luteinizing hormone  
 2510 and corticosteroid-induced ovulation and spawning in hypophysectomized catfish,  
 2511 *Heteropneustes fossilis* (Bloch). *J. Exp. Zool.* 163, 49–54.  
 2512 <https://doi.org/10.1002/jez.1401630105>

2513 Sundararaj, B.I., Goswami, S. V., 1977. Hormonal regulation of in vivo and in vitro oocyte  
 2514 maturation in the catfish, *Heteropneustes fossilis* (bloch). *Gen. Comp. Endocrinol.* 32,  
 2515 17–28. [https://doi.org/10.1016/0016-6480\(77\)90079-X](https://doi.org/10.1016/0016-6480(77)90079-X)  
 2516 Sundström, G., Dreborg, S., Larhammar, D., 2010. Concomitant duplications of opioid  
 2517 peptide and receptor genes before the origin of jawed vertebrates. *PLoS One* 5.  
 2518 <https://doi.org/10.1371/journal.pone.0010512>  
 2519 Takahashi, A., Amano, M., Amiya, N., Yamanome, T., Yamamori, K., Kawauchi, H., 2006.  
 2520 Expression of three proopiomelanocortin subtype genes and mass spectrometric  
 2521 identification of POMC-derived peptides in pars distalis and pars intermedia of barfin  
 2522 flounder pituitary. *Gen. Comp. Endocrinol.* 145, 280–286.  
 2523 <https://doi.org/10.1016/j.ygcen.2005.09.005>  
 2524 Takahashi, A., Amemiya, Y., Sarashi, M., Sower, S., Kawauchi, H., 1995. Melanotropin and  
 2525 corticotropin are encoded on two distinct genes in the lamprey, the earliest evolved  
 2526 extant vertebrate. *Biochem. Biophys. Res. Commun.* 213, 490–498.  
 2527 Takahashi, A., Kobayashi, Y., Mizusawa, K., 2012. The pituitary-interrenal axis of fish: A  
 2528 review focusing on the lamprey and flounder. *Gen. Comp. Endocrinol.* 188, 54–59.  
 2529 <https://doi.org/10.1016/j.ygcen.2013.03.005>  
 2530 Takahashi, A., Mizusawa, K., 2013. Posttranslational Modifications of Proopiomelanocortin  
 2531 in Vertebrates and Their Biological Significance. *Front. Endocrinol. (Lausanne)*. 4, 1–9.  
 2532 <https://doi.org/10.3389/fendo.2013.00143>  
 2533 Takahashi, H., Sakamoto, T., 2013. The role of “mineralocorticoids” in teleost fish: Relative  
 2534 importance of glucocorticoid signaling in the osmoregulation and “central” actions of  
 2535 mineralocorticoid receptor. *Gen. Comp. Endocrinol.* 181, 223–228.  
 2536 <https://doi.org/10.1016/j.ygcen.2012.11.016>  
 2537 Takeo, J., Hata, J. ichiro, Segawa, C., Toyohara, H., Yamashita, S., 1996. Fish glucocorticoid

2538 receptor with splicing variants in the DNA binding domain. *FEBS Lett.* 389, 244–248.  
 2539 [https://doi.org/10.1016/0014-5793\(96\)00596-0](https://doi.org/10.1016/0014-5793(96)00596-0)  
 2540 Takeuchi, S., Takahashi, S., 1998. Melanocortin receptor genes in the chicken - Tissue  
 2541 distributions. *Gen. Comp. Endocrinol.* 112, 220–231.  
 2542 <https://doi.org/10.1006/gcen.1998.7167>  
 2543 Teitsma, C., Lethimonier, C., Tujague, M., Anglade, I., Saligaut, D., Bailhache, T., Pakdel,  
 2544 F., Kah, O., Ducouret, B., 1998. Identification of potential sites of cortisol actions on the  
 2545 reproductive axis in rainbow trout. *Comp. Biochem. Physiol. - C Pharmacol. Toxicol.*  
 2546 *Endocrinol.* 119, 243–249. [https://doi.org/10.1016/S0742-8413\(98\)00013-9](https://doi.org/10.1016/S0742-8413(98)00013-9)  
 2547 Teitsma, C.A., Anglade, I., Lethimonier, C., Le Dréan, G., Saligaut, D., Ducouret, B., Kah,  
 2548 O., 1999. Glucocorticoid Receptor Immunoreactivity in Neurons and Pituitary Cells  
 2549 Implicated in Reproductive Functions in Rainbow Trout: A Double  
 2550 Immunohistochemical Study. *Biol. Reprod.* 60, 642–650.  
 2551 <https://doi.org/10.1095/biolreprod60.3.642>  
 2552 Tena-Sempere, M., Felip, A., Gómez, A., Zanuy, S., Carrillo, M., 2012. Comparative insights  
 2553 of the kisspeptin/kisspeptin receptor system: Lessons from non-mammalian vertebrates.  
 2554 *Gen. Comp. Endocrinol.* 175, 234–243. <https://doi.org/10.1016/j.ygcen.2011.11.015>  
 2555 Tesone, M., Charreau, E., 1980. Steroid biosynthesis in the gonads of the teleost fish *Jenynsia*  
 2556 *lineata*. *Comp Biochem Physiol B* 65, 631–637.  
 2557 Thomas, P., 2012. Rapid steroid hormone actions initiated at the cell surface and the receptors  
 2558 that mediate them with an emphasis on recent progress in fish models. *Gen. Comp.*  
 2559 *Endocrinol.* 175, 367–383. <https://doi.org/10.1016/j.ygcen.2011.11.032>  
 2560 Thorpe, J.E., 1994. Reproductive strategies in Atlantic salmon, *Salmo salar* L. *Aquac. Res.*  
 2561 25, 77–87. <https://doi.org/10.1111/j.1365-2109.1994.tb00668.x>  
 2562 Thorpe, J.E., Morgan, R.I.G., Talbot, C., Miles, M.S., 1983. Inheritance of developmental

2563 rates in Atlantic salmon, *Salmo salar* L. *Aquaculture* 33, 119–128.  
 2564 [https://doi.org/10.1016/0044-8486\(83\)90392-7](https://doi.org/10.1016/0044-8486(83)90392-7)  
 2565 Todd, E. V., Liu, H., Muncaster, S., Gemmell, N.J., 2016. Bending Genders: The Biology of  
 2566 Natural Sex Change in Fish. *Sex. Dev.* 10, 223–241. <https://doi.org/10.1159/000449297>  
 2567 Toni, C., Martos-Sitcha, J.A., Baldisserotto, B., Heinzmann, B.M., de Lima Silva, L.,  
 2568 Martínez-Rodríguez, G., Mancera, J.M., 2015. Sedative effect of 2-phenoxyethanol and  
 2569 essential oil of *Lippia alba* on stress response in gilthead sea bream (*Sparus aurata*). *Res.*  
 2570 *Vet. Sci.* 103, 20–27. <https://doi.org/10.1016/j.rvsc.2015.09.006>  
 2571 Toni, M., Manciocco, A., Angiulli, E., Alleva, E., Cioni, C., Malavasi, S., 2019. Review:  
 2572 Assessing fish welfare in research and aquaculture, with a focus on European directives.  
 2573 *Animal* 13, 161–170. <https://doi.org/10.1017/S1751731118000940>  
 2574 Tonon, M.C., Cuet, P., Lamacz, M., Jégou, S., Côté, J., Gouteux, L., Ling, N., Pelletier, G.,  
 2575 Vaudry, H., 1986. Comparative effects of corticotropin-releasing factor, arginine  
 2576 vasopressin, and related neuropeptides on the secretion of ACTH and  $\alpha$ -MSH by frog  
 2577 anterior pituitary cells and neurointermediate lobes in vitro. *Gen. Comp. Endocrinol.* 61,  
 2578 438–445. [https://doi.org/10.1016/0016-6480\(86\)90231-5](https://doi.org/10.1016/0016-6480(86)90231-5)  
 2579 [Tonon, M.C., Leroux, P., Leboulenger, F., Delarue, C., Jégou, S., Vaudry, H., 1980.](#)  
 2580 [Thyrotropin-releasing hormone stimulates the release of melanotropin from frog](#)  
 2581 [neurointermediate lobes in vitro. \*Life Sci.\* 26, 869–875.](#)  
 2582 <https://doi.org/10.1017/CBO9781107415324.004>  
 2583 Tostivint, H., 2011. Evolution of the gonadotropin-releasing hormone (GnRH) gene family in  
 2584 relation to vertebrate tetraploidizations. *Gen. Comp. Endocrinol.* 170, 575–581.  
 2585 <https://doi.org/10.1016/j.ygcen.2010.11.017>  
 2586 Tostivint, H., Ocampo Daza, D., Bergqvist, C.A., Quan, F.B., Bougerol, M., Lihmann, I.,  
 2587 Larhammar, D., 2014. Molecular evolution of GPCRs: Somatostatin/urotensin II

2588 receptors. *J. Mol. Endocrinol.* 52. <https://doi.org/10.1530/JME-13-0274>

2589 Tran, T.N., Fryer, J.N., Bennett, H.P.J., Tonon, M.C., Vaudry, H., 1989. TRH stimulates the  
 2590 release of POMC-derived peptides from goldfish melanotropes. *Peptides* 10, 835–841.  
 2591 [https://doi.org/10.1016/0196-9781\(89\)90122-8](https://doi.org/10.1016/0196-9781(89)90122-8)

2592 Tran, T.N., Fryer, J.N., Lederis, K., Vaudry, H., 1990. CRF, Urotensin I, and sauvagine  
 2593 stimulate the release of POMC-derived peptides from goldfish neurointermediate lobe  
 2594 cells. *Gen. Comp. Endocrinol.* 78, 351–360. [https://doi.org/10.1016/0016-](https://doi.org/10.1016/0016-6480(90)90025-H)  
 2595 [6480\(90\)90025-H](https://doi.org/10.1016/0016-6480(90)90025-H)

2596 Trudeau, V.L., 2018. Facing the challenges of neuropeptide gene knockouts: Why do they not  
 2597 inhibit reproduction in adult teleost fish? *Front. Neurosci.* 12, 1–8.  
 2598 <https://doi.org/10.3389/fnins.2018.00302>

2599 Trudeau, V.L., Somoza, G.M., 2020. Multimodal hypothalamo-hypophysial communication  
 2600 in the vertebrates. *Gen. Comp. Endocrinol.* 293, 113475.  
 2601 <https://doi.org/10.1016/j.ygcen.2020.113475>

2602 Tsutsui, K., Osugi, T., Son, Y.L., Ubuka, T., 2018. Review: Structure, function and evolution  
 2603 of GnIH. *Gen. Comp. Endocrinol.* 264, 48–57.  
 2604 <https://doi.org/10.1016/j.ygcen.2017.07.024>

2605 Tsutsui, K., Saigoh, E., Ukena, K., Teranishi, H., Fujisawa, Y., Kikuchi, M., Ishii, S., Sharp,  
 2606 P.J., 2000. A novel avian hypothalamic peptide inhibiting gonadotropin release.  
 2607 *Biochem. Biophys. Res. Commun.* 275, 661–667.  
 2608 <https://doi.org/10.1006/bbrc.2000.3350>

2609 Turkelson, C., Thomas, C., Zrimura, A., Chang, D., Chang, J., Shimizu, M., 1982. In vitro  
 2610 potentiation of the activity of synthetic ovine corticotropin-releasing factor by arginine  
 2611 vasopressin. *Peptides* 1, 111–113.

2612 Ubuka, T., Parhar, I., 2018. Dual actions of mammalian and piscine gonadotropin-inhibitory

2613 hormones, RFamide-related peptides and LPXRFamide peptides, in the hypothalamic-  
 2614 pituitary-gonadal axis. *Front. Endocrinol. (Lausanne)*. 8.  
 2615 <https://doi.org/10.3389/fendo.2017.00377>  
 2616 Van Den Burg, E.H., Metz, J.R., Spanings, F.A.T., Wendelaar Bonga, S.E., Flik, G., 2005.  
 2617 Plasma  $\alpha$ -MSH and acetylated  $\beta$ -endorphin levels following stress vary according to  
 2618 CRH sensitivity of the pituitary melanotropes in common carp, *Cyprinus carpio*. *Gen.*  
 2619 *Comp. Endocrinol.* 140, 210–221. <https://doi.org/10.1016/j.ygcen.2004.11.010>  
 2620 van den Hurk, R., van Oordt, P.G.W.J., 1985. Effects of natural androgens and corticosteroids  
 2621 on gonad differentiation in the rainbow trout, *Salmo gairdneri*. *Gen. Comp. Endocrinol.*  
 2622 57, 216–222. [https://doi.org/10.1016/0016-6480\(85\)90266-7](https://doi.org/10.1016/0016-6480(85)90266-7)  
 2623 Van Der Kraak, G.J., Munkittrick, K.R., McMaster, M.E., Portt, C.B., Chang, J.P., 1992.  
 2624 Exposure to bleached kraft pulp mill effluent disrupts the pituitary-gonadal axis of white  
 2625 sucker at multiple sites. *Toxicol. Appl. Pharmacol.* 115, 224–233.  
 2626 [https://doi.org/10.1016/0041-008X\(92\)90327-O](https://doi.org/10.1016/0041-008X(92)90327-O)  
 2627 Van Der Salm, A.L., Pavlidis, M., Flik, G., Wendelaar Bonga, S.E., 2004. Differential release  
 2628 of  $\alpha$ -melanophore stimulating hormone isoforms by the pituitary gland of red porgy,  
 2629 *Pagrus pagrus*. *Gen. Comp. Endocrinol.* 135, 126–133.  
 2630 <https://doi.org/10.1016/j.ygcen.2003.09.002>  
 2631 Van Enckevort, F.H.J., Pepels, P.P.L.M., Leunissen, J.A.M., Martens, G.J.M., Wendelaar  
 2632 Bonga, S.E., Balm, P.H.M., 2000. *Oreochromis mossambicus* (tilapia) corticotropin-  
 2633 releasing hormone: cDNA sequence and bioactivity. *J. Neuroendocrinol.* 12, 177–186.  
 2634 <https://doi.org/10.1046/j.1365-2826.2000.00434.x>  
 2635 Van Ginneken, V., Durif, C., Balm, S.P., Boot, R., Verstegen, M.W.A., Antonissen, E., Van  
 2636 Den Thillart, G., 2007. Silvering of European eel (*Anguilla anguilla* L.): Seasonal  
 2637 changes of morphological and metabolic parameters. *Anim. Biol.* 57, 63–77.

2638 <https://doi.org/10.1163/157075607780002014>

2639 Van Loon, G., Kragt, C.L., 1970. Effect of dopamine on the biological activity and in vitro  
2640 release of ACTH and FSH. *Proc. Soc. Exp. Biol. Med.* 133, 1137–1141.

2641 Västermark, Å., Schiöth, H.B., 2011. The early origin of melanocortin receptors, agouti-  
2642 related peptide, agouti signalling peptide, and melanocortin receptor-accessory proteins,  
2643 with emphasis on pufferfishes, elephant shark, lampreys, and amphioxus. *Eur. J.*  
2644 *Pharmacol.* 660, 61–69. <https://doi.org/10.1016/j.ejphar.2010.10.106>

2645 Vaughan, J., Donaldson, C., Bittencourt, J., Perrin, M.H., Lewis, K., Sutton, S., Chan, R.,  
2646 Turnbull, A. V., Lovejoy, D., Rivier, C., Rivier, J., Sawchenko, P.E., Vale, W., 1995.  
2647 Urocortin, a mammalian neuropeptide related to fish urotensin I and to corticotropin-  
2648 releasing factor. *Nature* 378, 287–292. <https://doi.org/10.1038/378287a0>

2649 Verbug-Van Kemenade, B.M.L., Jenks, B.G., Driessen, A.G.J., 1986. GABA and Dopamine  
2650 Act Directly on Melanotropes of *Xenopus* to inhibit MSH secretion. *Brain Res. Bull.* 17,  
2651 697–704. [https://doi.org/10.1016/0361-9230\(86\)90203-0](https://doi.org/10.1016/0361-9230(86)90203-0)

2652 Verburg-Van Kemenade, B., Jenks, B.G., Visser, T.J., Tonon, M.C., Vaudry, H., 1987.  
2653 Assessment of TRH as a potential MSH release stimulating factor in *Xenopus laevis*.  
2654 *Peptides* 8, 69–76. [https://doi.org/10.1016/0196-9781\(87\)90167-7](https://doi.org/10.1016/0196-9781(87)90167-7)

2655 Verburg-Van Kemenade, B.M.L., Jenks, B.G., Cruijssen, P.M.J.M., Dings, A., Tonon, M.C.,  
2656 Vaudry, H., 1987. Regulation of MSH release from the neurointermediate lobe of  
2657 *Xenopus laevis* by CRF-like peptides. *Peptides* 8, 1093–1100.  
2658 [https://doi.org/10.1016/0196-9781\(87\)90142-2](https://doi.org/10.1016/0196-9781(87)90142-2)

2659 Vermeulen, G.J., Lambert, J.G.D., Teitsma, C.A., Zandbergen, M.A., Goos, H.J.T., 1995.  
2660 Adrenal tissue in the male African catfish, *Clarias gariepinus*: localization and steroid  
2661 hormone secretion. *Cell Tissue Res.* 280, 653–657. <https://doi.org/10.1007/BF00318367>

2662 von Krogh, K., Bjørndal, G.T., Nourizadeh-Lillabadi, R., Ropstad, E., Haug, T.M., Weltzien,

2663 F.A., 2019. Cortisol differentially affects cell viability and reproduction-related gene  
 2664 expression in Atlantic cod pituitary cultures dependent on stage of sexual maturation.  
 2665 *Comp. Biochem. Physiol. -Part A Mol. Integr. Physiol.* 236.  
 2666 <https://doi.org/10.1016/j.cbpa.2019.06.017>

2667 Vu, M., Trudeau, V., 2016. Neuroendocrine control of spawning in amphibians and its  
 2668 practical applications. *Gen. Comp. Endocrinol.* 234, 28–39.  
 2669 <https://doi.org/10.1016/j.ygcen.2016.03.024>

2670 Wang, B., Yang, G., Xu, Y., Zhang, Y., Liu, X., 2019. In vitro effects of tongue sole LPXRFa  
 2671 and kisspeptin on relative abundance of pituitary hormone mRNA and inhibitory action  
 2672 of LPXRFa on kisspeptin activation in the PKC pathway. *Anim. Reprod. Sci.* 203, 1–9.  
 2673 <https://doi.org/10.1016/j.anireprosci.2019.01.009>

2674 Wang, J., Li, J., Zhang, X., Sun, X., 2012. Transcriptome analysis reveals the time of the  
 2675 fourth round of genome duplication in common carp (*Cyprinus carpio*). *BMC Genomics*  
 2676 13, 96. <https://doi.org/10.1186/1471-2164-13-96>

2677 Wang, T., Zhou, C., Yuan, D., Lin, F., Chen, H., Wu, H., Wei, R., Xin, Z., Liu, J., Gao, Y.,  
 2678 Li, Z., 2014. Schizothorax prenanti corticotropin-releasing hormone (CRH): Molecular  
 2679 cloning, tissue expression, and the function of feeding regulation. *Fish Physiol.*  
 2680 *Biochem.* 40, 1407–1415. <https://doi.org/10.1007/s10695-014-9935-6>

2681 Weltzien, F.A., Andersson, E., Andersen, Ø., Shalchian-Tabrizi, K., Norberg, B., 2004. The  
 2682 brain-pituitary-gonad axis in male teleosts, with special emphasis on flatfish  
 2683 (*Pleuronectiformes*). *Comp. Biochem. Physiol. - A Mol. Integr. Physiol.* 137, 447–477.  
 2684 <https://doi.org/10.1016/j.cbpb.2003.11.007>

2685 Wendelaar Bonga, S.E., 1997. The stress response in fish. *Physiol. Rev.* 77, 591–625.

2686 Westphal, N.J., Seasholtz, A.F., 2005. Gonadotropin-releasing hormone (GnRH) positively  
 2687 regulates corticotropin-releasing hormone-binding protein expression via multiple

intracellular signaling pathways and a multipartite GnRH response element in  $\alpha$ T3-1 cells. *Mol. Endocrinol.* 19, 2780–2797. <https://doi.org/10.1210/me.2004-0519>

Westring, C.G., Ando, H., Kitahashi, T., Bhandari, R.K., Ueda, H., Urano, A., Dores, R.M., Sher, A.A., Danielson, P.B., 2008. Seasonal changes in CRF-I and urotensin I transcript levels in masu salmon: Correlation with cortisol secretion during spawning. *Gen. Comp. Endocrinol.* 155, 126–140. <https://doi.org/10.1016/j.ygcen.2007.03.013>

Williams, B.L., Akazome, Y., Oka, Y., Eisthen, H.L., 2014. Dynamic evolution of the GnRH receptor gene family in vertebrates. *BMC Evol. Biol.* 14, 1–22. <https://doi.org/10.1186/s12862-014-0215-y>

Winberg, S., Nilsson, A., Hylland, P., Söderstöm, V., Nilsson, G.E., 1997. Serotonin as a regulator of hypothalamic-pituitary-interrenal activity in teleost fish. *Neurosci. Lett.* 230, 113–116. [https://doi.org/10.1016/S0304-3940\(97\)00488-6](https://doi.org/10.1016/S0304-3940(97)00488-6)

Winter, M.J., Ashworth, A., Bond, H., Brierley, M.J., McCrohan, C.R., Balment, R.J., 2000. The caudal neurosecretory system: Control and function of a novel neuroendocrine system in fish. *Biochem. Cell Biol.* 78, 193–203. <https://doi.org/10.1139/o00-059>

Woo, N., Hontela, A., Fryer, J., Kobayashi, Y., Lederis, K., 1985. Activation of hypothalamo-hypophysial-interrenal system by urophysectomy in goldfish. *Am J Physiol* 248, R197-201.

Wunderink, Y.S., de Vrieze, E., Metz, J.R., Halm, S., Martínez-Rodríguez, G., Flik, G., Klaren, P.H.M., Mancera, J.M., 2012. Subfunctionalization of pomc paralogues in senegalese sole (*solea senegalensis*). *Gen. Comp. Endocrinol.* 175, 407–415. <https://doi.org/10.1016/j.ygcen.2011.11.026>

Yamaguchi, T., Yoshinaga, N., Yazawa, T., Gen, K., Kitano, T., 2010. Cortisol is involved in temperature-dependent sex determination in the Japanese flounder. *Endocrinology* 151, 3900–3908. <https://doi.org/10.1210/en.2010-0228>

2713 Yang, J.A., Song, C.I., Hughes, J.K., Kreisman, M.J., Parra, R.A., Haisenleder, D.J.,  
 2714 Kauffman, A.S., Breen, K.M., 2017. Acute psychosocial stress inhibits lh pulsatility and  
 2715 kiss1 neuronal activation in female mice. *Endocrinology* 158, 3716–3723.  
 2716 <https://doi.org/10.1210/en.2017-00301>  
 2717 Yoshiura, Y., Suetake, H., Aida, K., 1999. Duality of gonadotropin in a primitive teleost,  
 2718 Japanese eel (*Anguilla japonica*). *Gen. Comp. Endocrinol.* 114, 121–131.  
 2719 <https://doi.org/10.1006/gcen.1998.7242>  
 2720 Young, G., Kagawa, H., Nagahama, Y., 1982. Oocyte maturation in the Amago salmon  
 2721 (*Oncorhynchus rhodurus*): In Vitro Effects of Salmon Gonadotropin, Steroids, and  
 2722 cyanoketone (an inhibitor of 3 $\beta$ -Hydroxy-A5-Steroid Dehydrogenase). *J. Exp. Zool.* 224,  
 2723 265–275.  
 2724 Young, G., Thorarensen, H., Davie, P.S., 1996. 11-Ketotestosterone suppresses interrenal  
 2725 activity in rainbow trout (*Oncorhynchus mykiss*). *Gen. Comp. Endocrinol.* 103, 301–  
 2726 307. <https://doi.org/10.1006/gcen.1996.0125>  
 2727 Yuan, M., Li, X., Long, T., Chen, Y., Lu, W., 2020. Dynamic Responses of the Caudal  
 2728 Neurosecretory System (CNSS) Under Thermal Stress in Olive Flounder (*Paralichthys*  
 2729 *olivaceus*). *Front. Physiol.* 10, 1–9. <https://doi.org/10.3389/fphys.2019.01560>  
 2730 Zhang, Y., Zhang, S., Lu, H., Zhang, L., Zhang, W., 2014. Genes encoding aromatases in  
 2731 teleosts: Evolution and expression regulation. *Gen. Comp. Endocrinol.* 205, 151–158.  
 2732 <https://doi.org/10.1016/j.ygcen.2014.05.008>  
 2733 Zhou, H., Ge, C., Chen, A., Lu, W., 2019. Dynamic Expression and Regulation of Urotensin I  
 2734 and Corticotropin-Releasing Hormone Receptors in Ovary of Olive Flounder  
 2735 *Paralichthys olivaceus*. *Front. Physiol.* 10, 1–10.  
 2736 <https://doi.org/10.3389/fphys.2019.01045>  
 2737 Ziv, L., Muto, A., Schoonheim, P.J., Meijsing, S.H., Strasser, D., Ingraham, H.A., Schaaf,

2738 M.J.M., Yamamoto, K.R., Baier, H., 2013. An affective disorder in zebrafish with  
2739 mutation of the glucocorticoid receptor. *Mol. Psychiatry* 18, 681–691.  
2740 <https://doi.org/10.1038/mp.2012.64>  
2741 Zohar, Y., Muñoz-Cueto, J.A., Elizur, A., Kah, O., 2010. Neuroendocrinology of  
2742 reproduction in teleost fish. *Gen. Comp. Endocrinol.* 165, 438–455.  
2743 <https://doi.org/10.1016/j.ygcen.2009.04.017>  
2744  
2745

2745  
2746

## 2747 **Figure legends**

2748

### 2749 **Figure 1: Stress and reproduction neuroendocrine axes in teleosts.**

2750 The figure displays the main actors and regulatory pathways of the stress neuroendocrine axis  
2751 (corticotropic axis: brain-pituitary-interrenals; green arrows) and of the reproduction  
2752 neuroendocrine axis (gonadotropic axis: brain-pituitary-gonads; blue arrows). The Caudal  
2753 Neurosecretory System (CNSS) which may be involved in the stress axis in teleost is  
2754 indicated in relation to the corticotropic axis. The liver, which plays a key role in reproduction  
2755 in teleosts as in other oviparous vertebrates by producing vitellogenin, is indicated in relation  
2756 to the gonadotropic axis. Abbreviations: AVP, arginine vasopressin; AR, androgen receptor;  
2757 CNSS, caudal neurosecretory system; CRH, corticotropin-releasing hormone; CRH-R,  
2758 corticotropin-releasing hormone receptor; DA, dopamine; DHP, 17-hydroxy-20-  
2759 dihydroprogesterone; DOC, 11-deoxycorticosterone; E2, estradiol; ER, estrogen receptor;  
2760 FSH, follicle stimulating hormone; FSH-R, follicle stimulating hormone receptor; GnIH,  
2761 gonadotropin inhibitory hormone; GnRH, gonadotropin-releasing hormone; GnRH-R,  
2762 gonadotropin-releasing hormone receptor; GR, glucocorticoid receptor; kiss, kisspeptin; kiss-  
2763 R, kisspeptin receptor; LH, luteinising hormone; LH-R, luteinising hormone receptor; MC2R,  
2764 melanocortin receptor 2; other P, other progestins; POMC, proopiomelanocortin; PR,  
2765 progestin receptor; T, testosterone; TRH, thyrotropin releasing hormone; UI, urotensin I; Vg,  
2766 vitellogenin; 5-HT, 5 hydroxytryptamine, or serotonin; 11-KT, 11-ketotestosterone.

2767

### 2768 **Figure 2: Effects of corticotropic axis on gonadotropic axis in vertebrates.**

2769 The figure illustrates current knowledge on the regulatory actions (green arrows) exerted by  
2770 typical actors of the corticotropic axis (brain-pituitary-interrenal) on the gonadotropic and

2771 liver axis (brain-pituitary-gonad-liver). Abbreviations: ACTH, adrenocorticotropin;  $\beta$ -end,  $\beta$ -  
2772 endorphin; CRH, corticotropin-releasing hormone; E2, estradiol; FSH, follicle stimulating  
2773 hormone; GnIH, gonadotropin inhibitory hormone; GnRH, gonadotropin-releasing hormone;  
2774 GR, glucocorticoid receptor; kiss, kisspeptin; LH, luteinising hormone; MC2R, melanocortin  
2775 receptor 2; MR, mineralocorticoid receptor; T, testosterone; 11-KT, 11-ketotestosterone;.

2776

2777 **Figure 3: Effects of gonadotropic axis on corticotropic axis in vertebrates.**

2778 The figure illustrates current knowledge on the regulatory actions (blue arrows) exerted by  
2779 typical actors of the gonadotropic axis (brain-pituitary-gonad) on the corticotropic axis (brain-  
2780 pituitary-interrenal). Abbreviations: ACTH, adrenocorticotropin; AR, androgen receptor;  
2781 CRH, corticotropin-releasing hormone; CRH-BP, corticotropin-releasing hormone binding  
2782 protein; DHP, 17-hydroxy-20-dihydroprogesterone; E2, estradiol; ER, estrogen receptor;  
2783 FSH, follicle stimulating hormone; GnIH, gonadotropin inhibitory hormone; GnRH,  
2784 gonadotropin-releasing hormone; LH, luteinising hormone; LH-R, luteinising hormone  
2785 receptor; other P, other progestins; PR, progestin receptor; T, testosterone; Vg, vitellogenin;  
2786 11-KT, 11-ketotestosterone.

2787

2788 **Table Legends**

2789

2790 **Table 1: Major actors of the corticotropic axis in human and teleosts.**

2791 This table underlines the expansion of gene numbers in teleosts due to teleost-specific whole  
2792 genome duplication (3R). In teleosts, some of the paralogs have been lost in some species.  
2793 Supplemental paralogs issued from additional whole genome duplication (4R, which occurred  
2794 in salmonids and carps) are not indicated. Abbreviations: CRH, corticotropin-releasing  
2795 hormone; CRH-R, corticotropin-releasing hormone receptor; POMC, proopiomelanocortin;

2796 MCR, melanocortin receptor; GR, glucocorticoid receptor; MR, mineralocorticoid receptor;  
2797 UI, urotensin I.

2798

2799 **Table 2: Major actors of the gonadotropic axis in human and teleosts.**


2800 This table underlines the larger gene numbers in teleosts due to teleost-specific whole genome  
2801 duplication (3R) but also to the conservation of paralogs resulting from early vertebrate whole  
2802 genome duplications (1R and 2R), or from local gene duplication (see the text for details). In  
2803 teleosts, some of the paralogs have been lost in some species. Supplemental paralogs issued  
2804 from additional whole genome duplication (4R, which occurred in salmonids and carps) are  
2805 not indicated. Abbreviations: GnRH, gonadotropin-releasing hormone; GnRH-R,  
2806 gonadotropin-releasing hormone receptor; GnIH, gonadotropin inhibitory hormone; kiss,  
2807 kisspeptin; kiss-R, kisspeptin receptor; LH $\beta$ , luteinising hormone; FSH $\beta$ , follicle stimulating  
2808 hormone; LH-R, luteinising hormone receptor; FSH-R, follicle stimulating hormone receptor;  
2809 ER, estrogen receptor; esr, nuclear estrogen receptor; gper, membrane estrogen receptor; AR,  
2810 androgen receptor; PR, progestin receptor; pgr, nuclear progestin receptor; mpgr, membrane  
2811 progestin receptor.

2812

2813


## Corticotropic axis

## Gonadotropic axis




## Corticotropic axis

## Gonadotropic axis



## Corticotropic axis

## Gonadotropic axis



**Table 1: Major actors of the corticotropic axis in human and teleosts.**

Protein	Gene	Human	Teleosts	References / Review articles
CRH	<i>crh</i>	<i>crhl</i>  <i>crh2</i>	<i>crh1a</i>  <i>crh1b</i>  <i>crh2</i>	Lovejoy et al 2014; Cardoso et al 2016  Maugars et al 2016
CRH-R	<i>crhr</i>	<i>crhr1</i>  <i>crhr2</i>	<i>crhr1a</i>  <i>crhr1b</i>  <i>crhr2</i>	Cardoso et al 2014; Lovejoy et al 2014
POMC	<i>pomc</i>	<i>pomc</i>	<i>pomc-<math>\alpha 1</math></i>  <i>pomc-<math>\alpha 2</math></i>  <i>pomc-<math>\beta</math></i>	Dores and Lecaude 2005; Cardoso et al 2011; Dores and Baron 2011
MC1R MC2R MC3R MC4R MC5R	<i>mclr</i> <i>mc2r</i> <i>mc3r</i> <i>mc4r</i> <i>mc5r</i>	<i>mclr</i> <i>mc2r</i> <i>mc3r</i> <i>mc4r</i> <i>mc5r</i>	<i>mclr</i> <i>mc2r</i> <i>mc3r</i> <i>mc4r</i> <i>mc5ra</i>  <i>mc5rb</i>	Vastermark and Schioth 2011; Dores et al 2014, 2016
GR	<i>gr</i>	<i>gr</i>	<i>gr1</i>  <i>gr2</i>	Bury 2017
MR	<i>mr</i>	<i>mr</i>	<i>mr1</i>  <i>mr2</i>	Lafont et al 2014; Baker and Katsu 2019
UI	<i>uI</i>		<i>uI</i>	Lovejoy et al 2014

**Table 2: Major actors of the gonadotropic axis in human and teleosts.**

Protein	Gene	Human	Teleosts	References / Review articles
GnRH	<i>gnrh</i>	<i>gnrh1</i> <i>gnrh2</i>	<i>gnrh1</i> <i>gnrh2</i> <i>gnrh3</i>	Kim et al 2011; Roch et al 2011; Tostivint 2011
GnRH-R	<i>gnrhr</i>	<i>gnrhr-1</i>	Up to 5 <i>gnrhr</i>	Roch et al 2014; Sefideh et al 2014; William et al 2014; Ciani et al 2020
GnIH	<i>gnih</i>	<i>gnih/rfrp</i>	<i>gnih</i>	Tsutsui et al 2018; Maugars et al 2020
Kiss	<i>kiss</i>	<i>kiss1</i>	<i>kiss1</i> <i>kiss2</i>	Pasquier et al 2012, 2014
Kiss-R	<i>kissR</i>	<i>kiss1R</i>	<i>kiss1R</i> <i>kiss2R</i> <i>kiss3R</i>	Pasquier et al 2012, 2014
LH $\beta$	<i>lh<math>\beta</math></i>	<i>lh<math>\beta</math></i>	<i>lh<math>\beta</math></i>	Dufour et al 2020
FSH $\beta$	<i>fsh<math>\beta</math></i>	<i>fsh<math>\beta</math></i>	<i>fsh<math>\beta</math></i>	Dufour et al 2020
LH-R	<i>lhr</i>	<i>lhcgr</i>	<i>lhr1</i> <i>lhr2</i>	Maugars and Dufour 2015
FSH-R	<i>fshr</i>	<i>fshr</i>	<i>fshr</i>	Maugars and Dufour 2015
ER	<i>esr</i>  <i>gper</i>	<i>esr1</i> <i>esr2</i>  <i>gper</i>	<i>esr1</i> <i>esr2a</i> <i>esr2b</i>  <i>gpera</i> <i>gperb</i>	Lafont et al 2016; Ogino et al 2018  Lafont et al 2016
AR	<i>ar</i>	<i>ar</i>	<i>ara</i>  <i>arb</i>	Ogino et al 2009
PR	<i>pgr</i>  <i>mpgr</i>	<i>pgr</i>  multiple	<i>pgra</i>  <i>pgrb</i>  multiple	Morini et al 2019  Morini et al 2019