



**HAL**  
open science

# Mitigating the impact of bad rainy seasons in poor agricultural regions to tackle deforestation

Antoine Leblois

► **To cite this version:**

Antoine Leblois. Mitigating the impact of bad rainy seasons in poor agricultural regions to tackle deforestation. *Environmental Research Letters*, In press, 16 (5), pp.054003. 10.1088/1748-9326/abdbf0 . hal-03111007

**HAL Id: hal-03111007**

**<https://hal.inrae.fr/hal-03111007>**

Submitted on 15 Jan 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

ACCEPTED MANUSCRIPT • OPEN ACCESS

## Mitigating the impact of bad rainy seasons in poor agricultural regions to tackle deforestation

To cite this article before publication: Antoine Leblois 2021 *Environ. Res. Lett.* in press <https://doi.org/10.1088/1748-9326/abdbf0>

### Manuscript version: Accepted Manuscript

Accepted Manuscript is “the version of the article accepted for publication including all changes made as a result of the peer review process, and which may also include the addition to the article by IOP Publishing of a header, an article ID, a cover sheet and/or an ‘Accepted Manuscript’ watermark, but excluding any other editing, typesetting or other changes made by IOP Publishing and/or its licensors”

This Accepted Manuscript is © 2020 The Author(s). Published by IOP Publishing Ltd.

As the Version of Record of this article is going to be / has been published on a gold open access basis under a CC BY 3.0 licence, this Accepted Manuscript is available for reuse under a CC BY 3.0 licence immediately.

Everyone is permitted to use all or part of the original content in this article, provided that they adhere to all the terms of the licence <https://creativecommons.org/licenses/by/3.0>

Although reasonable endeavours have been taken to obtain all necessary permissions from third parties to include their copyrighted content within this article, their full citation and copyright line may not be present in this Accepted Manuscript version. Before using any content from this article, please refer to the Version of Record on IOPscience once published for full citation and copyright details, as permissions may be required. All third party content is fully copyright protected and is not published on a gold open access basis under a CC BY licence, unless that is specifically stated in the figure caption in the Version of Record.

View the [article online](#) for updates and enhancements.

# Mitigating the impact of bad rainy seasons in poor agricultural regions to tackle deforestation

Antoine Leblois\*

January 13, 2021

## Abstract

Land use changes are known to account for over 20% of human greenhouse gas emissions and tree cover losses can significantly influence land-climate dynamics. Land-climate feedbacks have been identified and evaluated for a long time. However, in addition to the direct effect of climate change on forest biomes, recent sparse evidence has shown that land use changes may increase as a result of weather shocks. In Western and Central Africa, agriculture is the main source of income and employment for rural populations. Economies rely on agricultural production, which is largely rainfed, and therefore dependent predominantly upon seasonal rainfall. In this article, I explore the impact of seasonal rainfall quality on deforestation, by combining high-resolution remotely-sensed annual tree cover loss, land cover, human activity and daily rainfall data. I show that in poor regions that are mainly reliant on rainfed agriculture, a bad rainy season leads to large deforestation shocks. These shocks notably depend on the proportion of agricultural land and on the remoteness of the areas in question, as remoteness determines the ability to import food and the existence of alternative income sources. In areas with significant forest cover, a short rainfall season leads to a 15% increase in deforestation. In unconnected areas with small proportions of crop area, the increase in deforestation reaches 20%. Findings suggest that a refined understanding of the land use changes caused by rainfall shocks might be used to improve the design and effectiveness of development, adaptation and conservation policies.

Keywords: Deforestation; Rainfall shocks; West Africa.

---

\*CEE-M, Univ Montpellier, CNRS, INRAE, Institut Agro, 2 place Viala, 34000, Montpellier, France. I thank two anonymous referees for their constructive comments, A. Lobianco for providing a useful remote computation server, P. Delacote, S. Desbureaux, P. Quirion, A. Leblois and L. Pellegrin for careful reading. All remaining errors are mine.

## Contents

<b>1 Introduction: the influence of the poverty-environment nexus on land-climate dynamics</b>	<b>3</b>
<b>2 Study area</b>	<b>4</b>
2.1 Tree cover losses in a region with a strong pressure on land use . . . . .	5
2.2 Quality of the rainfall season and small scale rainfed agriculture . . . . .	5
<b>3 Materials and methods</b>	<b>7</b>
3.1 Data . . . . .	7
3.2 Methodology . . . . .	7
3.2.1 Rainfall season quality . . . . .	8
3.2.2 Land cover . . . . .	8
3.2.3 Remoteness/connectivity: population, energy and markets . . . . .	10
3.3 Statistical identification . . . . .	10
<b>4 Main results: heterogeneous but significant impacts of a bad rainy season on deforestation</b>	<b>11</b>
4.1 Weather shocks and deforestation in the 4 eco-climatic zones . . . . .	11
4.2 Robustness checks . . . . .	12
4.3 Deforestation and remoteness . . . . .	13
<b>5 Discussion</b>	<b>15</b>
<b>6 Data and code availability</b>	<b>16</b>
<b>References</b>	<b>16</b>
<b>7 Appendix</b>	<b>23</b>
<b>A Data</b>	<b>23</b>
A.1 Tree cover . . . . .	23
A.2 Deforestation . . . . .	23
A.3 Daily rainfall estimates . . . . .	24
A.3.1 Annual rainfall season quality indices . . . . .	24
A.3.2 Definition of the rainfall season . . . . .	25
A.4 Administrative boundaries . . . . .	25
A.5 Land cover classification . . . . .	26
A.6 Relative remoteness . . . . .	28
A.6.1 Night-time lights: distance to powered settlements . . . . .	28
A.6.2 Accessibility: travel time to cities . . . . .	28
<b>B Descriptive statistics</b>	<b>29</b>
B.1 Whole sample . . . . .	29
B.2 Eco-climatic zones . . . . .	31
<b>C Statistical analysis</b>	<b>31</b>
C.1 Transmission channels: impacts mainly driven by human activities . . . . .	31
C.2 Whole study region . . . . .	31
C.3 Results by eco-climatic zones . . . . .	31
C.4 Deforestation and remoteness . . . . .	34

<b>D Robustness checks</b>	<b>35</b>
D.1 Considering USGS bioclimatic regions instead of FAO eco-climatic zones . . . . .	35
D.2 Forest definition: thresholds of canopy cover . . . . .	35
D.3 Rainfall shocks severity: SPI thresholds . . . . .	38
D.4 Deforestation and remoteness: Guinean and Guineo-congolian eco-climatic zones . . . . .	39
D.5 Standard error adjustment for spatial correlation and serial correlation in panel data . . . . .	39

## 1 Introduction: the influence of the poverty-environment nexus on land-climate dynamics

There is a consensus regarding the impact of deforestation on local climate, namely warmer and drier conditions (Lawrence and Vandecar, 2015; Wolff et al., 2018; Leite-Filho et al., 2019). However, the potential reverse relationship – *i.e.* of weather shocks on deforestation – has not been largely investigated. The direct effect of rainfall on carbon storage (Brandt et al., 2018) and on forest health (Phillips et al., 2009; Anderegg et al., 2013; Zemp et al., 2017; Verbesselt et al., 2016; Jiang et al., 2019; Aleixo et al., 2019; Brodrigg et al., 2020) could create a vicious circle. Another vicious circle could be created by the indirect effects of drought on deforestation, via human activities, with weather variability creating agricultural income shocks that, in turn, affect land use decisions. Indeed, Zaveri et al. (2020) found that, over the past two decades, dry anomalies have accounted for 7.4% of the global cropland expansion rate and 9% of developing world’s cropland expansion rate. Desbureaux and Damania (2018) showed that, in Madagascar, deforestation increases during drought years and protected areas were partially effective at buffering against these upsurges in deforestation. Staal et al. (2020) found that for every mm of water deficit, deforestation tends to increase by 0.13% in the Amazon.

Yet, short-term effects of income shocks and poverty reduction on the environment constitutes a great empirical puzzle. Recent evidence suggests there is a positive relation between income and environmental degradation in poor countries, that is inconsistent with above-mentioned results. Although for poor populations of many regions, forest biomes act as a buffer against external shocks (Agarwal, 1991; Pattanayak and Sills, 2001; Wunder, 2001; Angelsen and Wunder, 2002; Baland and Francois, 2005; Tschardt et al., 2012; Somorin et al., 2012; Noack et al., 2019), short-term economic opportunities may create incentives for forest degradation. Indeed, Baland et al. (2010) showed that poorer households in rural Nepal collect significantly less firewood than wealthier households in the same village, and likewise, in the context of poverty alleviation policies in Mexico and Gambia, recent robust evidence confirmed that positive income shocks lead to more environmental degradation and deforestation (Alix-Garcia et al., 2013; Heß et al., 2019). Furthermore, Assunção et al. (2019) showed that, in the Brazilian Amazon, enforcement of stricter requirements for rural credit concession led to lower deforestation, especially in the municipalities in which cattle ranching is a primary activity.

1  
2  
3  
4  
5  
6 Alternatively, Ferraro and Simorangkir (2020) observed that poverty alleviation cash  
7 transfer programme in Indonesia was associated with 10 to 50% decrease in deforestation,  
8 suggesting that targeting the very poor may help achieve environmental goals. Defor-  
9 estation can also possibly be curbed by using conditional cash transfers programmes  
10 contingent upon conservation, as shown by Jayachandran et al. (2017)'s study of a pay-  
11 ment for ecosystem-services programme in Uganda.

12 However, these two conflicting strands of evidence can be reconciled in the case of a  
13 weather shock. While a production loss could push smallholders to increase the size of  
14 their cultivated area to meet a subsistence constraint in remote, isolated villages, it could  
15 also reduce deforestation in the presence of alternative income sources by lowering the  
16 relative attractiveness of farming activities.<sup>1</sup> In a nutshell, the outcome of an agricultural  
17 income shock depends upon whether producers are faced with a price inelastic demand  
18 for food, a situation which may occur in remote locations unconnected to markets.  
19 Therefore, the impact should then depend upon the remoteness of agricultural activities,  
20 presence of alternative activities to farming, accessibility of a subsistence food market  
21 and, potentially, the presence of a poverty trap. The relationship between weather shocks  
22 and land use changes are therefore *a priori* ambiguous.

23 This study attempts to unravel this ambiguity and understand how forest cover losses  
24 may react to an income shock stemming – in the very short run – from a weather shock.  
25 I used high resolution tree cover loss, land cover and daily precipitation data in West  
26 Africa to look for a land use response to bad rainy seasons at the very local level. By  
27 considering high-resolution, daily precipitations, it is possible to assess rainfall season  
28 quality, known to be subject to high spatial variations and largely dependent upon rain-  
29 fall timing, and notably on the onset of the season. I found that both the timing and  
30 level of water availability for crops have significant impacts on tree cover losses, and  
31 that these relationships depends upon remoteness, which I estimated using population,  
32 proximity-to-powered-settlements and travel-time-to-the-nearest-city data. The results  
33 also demonstrate that freely available data can help shed light on the mechanisms be-  
34 hind the poverty-environment nexus, and give credence to socio-economically focused  
35 deforestation models that could be of service to the designing of deforestation reduction  
36 policies.

## 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

## 2 Study area

The study area (Figure 1(a)), delimited by the equator and the 20th parallel north and  
the 20th East and West Meridians, includes 22 countries of Western and Central Africa.

---

<sup>1</sup>Alternative income sources has indeed been found to lower deforestation (Bluffstone, 1995) in Burkina  
Faso, the Philippines and the Amazon (Shively and Pagiola, 2004; Etongo et al., 2015; Araujo et al.,  
2019).

## 2.1 Tree cover losses in a region with a strong pressure on land use

Agriculture is the main sector in terms of GDP and labour force in Western and Central Africa (51% of labour force on average in the region, ranging from 25% in Burkina Faso to 76% in Chad, The World Bank (2020)). It is mainly rainfed and largely consists in smallholder agriculture. Curtis et al. (2018) showed that, in Sub-Saharan Africa, 92% of deforested land is attributable to a shift to agriculture. Tyukavina et al. (2018) and Molinario et al. (2020) showed this is particularly true in Congo basin countries, in which forest loss is mainly driven by smallholder clearing.

During the last few decades, Western and Central Africa has thus seen a large increase in agricultural land, notably due to considerable pressure from smallholder agriculture. Between 1975 and 2013, the surface area covered by crops doubled in Western Africa, reaching a total of 1.1 million square km, or 22.4% of total land surface. During this same period, forest cover was reduced by 37%, and is now highly fragmented (Cotillon and Tappan, 2016).

Over 86 thousand square km were deforested during the study period (2001-2019), an area the size of Austria. The spatial distribution of tree cover loss across the study region is shown in Figure 1, both as a share of 2000 tree cover (Figure 1(c)) and as its level in ha (Figure 1(d), in logarithmic scale) for each 0.05 degree pixel (of approximately 30 square km). The average deforestation per pixel over the whole 19-year period is 60 ha (as shown in Supplementary Table 2). A large proportion of the area was already impacted by cropping in 2000 (Figure 2(b)), demonstrating intense pressure on land-use.

## 2.2 Quality of the rainfall season and small scale rainfed agriculture

Rural income in Western and Central Africa largely relies upon the quality of the rainfall season for agriculture, and this quality depends upon rainfall distribution throughout the cropping season. Roudier et al. (2011) showed in a meta analysis that, despite a large dispersion of predicted impact, climate change is expected to have a net negative impact on crop yields in West Africa.

Recently, cumulative annual precipitation in the Sahel has shown a positive trend. However, the observed increase in extreme weather events is suspected to be causing additional damage to agriculture and societies (Panthou et al., 2018; Taylor et al., 2017). Indeed, the recent recovery in cumulative annual rainfall is associated with an increase in rainfall intensity and with fewer rainy days during the rainfall season (Zhang et al., 2017; Bichet and Diedhiou, 2018; Biasutti, 2019). The increase in annual rainfall quantity hides higher intra-seasonal variations, as well as the potential cooccurrence of damaging events (dry spells long enough to harm annual crop yield and/or simply a decrease in number of rainy days) and heavy rains that can also harm agricultural yields and infrastructures. Rainy season onset controls the best planting dates, while rainy season cessation and length determine the type of seed to plant (Marteau et al., 2011). Agricultural activities and production therefore become extremely vulnerable to variability in rainy season onset, cessation date and length (Dodd and Jolliffe, 2001).




Figure 1: Political borders in Africa and region of the study (a), Tree cover in 2000 (b), Tree cover loss as a share of tree cover (c) and Log of tree cover loss in ha (d) during the study period (2001-2019), according to Hansen et al. (2013) and averaged by 0.05 degree pixels (see section 3.2.2 and Appendices A.1 & A.2).



## 3 Materials and methods

### 3.1 Data

I constructed an original database of over 2.5 million observations (see Supplementary Table 3), tracking remotely-sensed tree cover losses throughout approximately 140 thousand pixels (see Supplementary Table 2) over 19 years. By combining these observations with daily rainfall estimates, I was able to measure the impact of a bad rainy season on estimated tree cover loss in the current year.

Tree cover, tree cover loss, daily rainfall, land use, night-time lights and time-distance to cities (respectively described in Appendices A.1, A.2, A.3, A.5, A.6.2 and A.6.1 of Supplementary material) data were aggregated all to the lowest resolution, i.e. the resolution of the rainfall estimates (0.05 degree). In order to consider a panel of pixels potentially concerned by both tree cover losses and a bad rainy season, the final sample is composed of every pixel with at least 1 ha of forest cover in 2000<sup>2</sup> and with over 100 mm of average cumulative annual rainfall.

### 3.2 Methodology

In order to assess the quality of the rainfall season for agricultural production, I computed and tested different indices<sup>3</sup>, out of which two were retained: the rainy season length (measured in days) and the cumulative daily water availability for crops over this rainy season (measured in mm). Each index is described in Section 3.1. Both indices are standardised, using a pixel-specific Standardised Precipitation Index (SPI, Svoboda et al. (2012), see section 3.2.1 below and A.3.1 of Supplementary material). Hereafter, significant negative shocks<sup>4</sup>, with SPI annual values inferior to -1, are referred to as “short rainfall season” when tied to rainy season length and “drought” when tied to cumulative rainfall over the rainy season.

While cumulative rainfall is one of the simplest indicators of drought, it is widely accepted that precipitation is the dominant controlling effect on vegetation phenology in the tropics, and that intra-seasonal rainfall shocks are key parameter for understanding the impact of seasonal cycle changes on African agriculture. It is notably the case of rainy season onset, cessation and length (Dunning et al., 2016; Boyard-Micheau et al., 2013). Furthermore, rainy season onset, which largely determines rainy season length, is perceived by farmers in the Sahel as a good indicator of change in seasonal rainfall quality (Kosmowski et al., 2016).

---

<sup>2</sup>Dropping approximately 0.033% of the pixel covered with forest.

<sup>3</sup>These other indices included the number of rainy days, occurrence of heavy rainfall, onset and offset dates, simpler and non standardised cumulative rainfall, other rainfall season definitions and the number of dry spells (7, 10 and 14 days) during the rainfall season.

<sup>4</sup>*i.e.* negative variations superior to one standard deviation. Robustness checks with other SPI values thresholds are displayed in section D.3 of Supplementary material.

### 3.2.1 Rainfall season quality

**Rainy season length** (offset minus onset), in days, is based on the definition of Liebmann and Marengo (2001) (see Section A.3.2 of Supplementary material). This definition has the advantage of being compatible with the occurrence of two consecutive rainy seasons in one year, which is the case in the Gulf of Guinea.

**Cumulative rainfall** only considers daily rainfall that may actually be used by crops, i.e.: daily rainfall superior to 1 mm (thereby excluding rain that may be entirely evaporated, according to Odekunle (2004)) and inferior to 30 mm (as rains exceeding 30 mm may be subject to direct runoff, according to Baron et al. (2005)). By only considering seasonal rainfall it is possible to focus on the agricultural impact transmission channel (as discussed in Supplementary Section C.1) and lower the noise of wild fires, which mainly occur during the dry season (Desbureaux and Damania, 2018).

Rainy season length and cumulative rainfall are computed using daily rainfall data (CHIRPS estimates, described in Supplementary material, Section A.3) and standardised at the pixel level (19 years of observations).

Because of the high heterogeneity in cumulative annual rainfall (see Figure 5 of Supplementary material, I divided the study region into 4 eco-climatic zones (Figure 2(a))<sup>5</sup>. The Sahelian (and Sudano-Sahelian) climate, corresponds to pixels with over 100 mm and under 900 mm of annual rainfall. The Sudanian climate corresponds to pixels with between 900 and 1100 mm of annual rainfall. The Guinean climate corresponds to pixels with annual rainfall between 1100 and 1700 mm and the Guineo-Congolian climate corresponds to pixels with over 1700 mm. Descriptive statistics for all the variables considered in this paper are provided for each eco-climatic zones in the Section B.2 and robustness of the results with an alternative definition of eco-climatic zones in Section D.1 of Supplementary material.

### 3.2.2 Land cover

I consider Hansen et al. (2013)'s estimates of annual **tree cover loss** for 2001-2019 (see Appendix A.2), available at a 1 arc-second resolution and averaged at 0.05 degree. Most definitions of forests using satellite measurements rely on a minimal tree cover percentage criterion. Given that 40% of dryland forests and 25% of all forests of Central and Western Africa are open forests with a canopy cover inferior to 40%<sup>6</sup> (Patriarca et al., 2019; Mayaux et al., 1998) it appeared necessary to consider a relatively low canopy cover threshold in the study region. Moreover, Sexton et al. (2016) show that when one considers a tree cover threshold superior to 30%, a significant proportion of West Africa's forest biomes are overlooked. Therefore, forest (and deforestation), are

<sup>5</sup>Classification used by the FAO, notably in its global information and early warning system (FAO, 2007) used by the FAO, notably in its global information and early warning system (FAO, 2007). It is very close to the USGS classification of West Africa's bioclimatic regions.

<sup>6</sup>The 40% canopy cover threshold is widely used to distinguish closed forests from open fragmented forests, while the 10% canopy cover threshold is often used to distinguish forest biomes from trees (UNEP, 2001; FAO, 2020).




Figure 2: Definition of the 4 eco-climatic zones (a), Percent of crop area in each 0.05 degree pixel (b) and Log of population density (c) in the study region.

Crop area is derived from ESA land cover data (see Supplementary section A.5), the eco-climatic zones are defined by isohyets of daily rainfall data over the 2000-2019 period (see Supplementary section A.3) and the population data is taken from WorldPop (see Section 3.2.3)

hereafter, restricted to tree cover (and tree cover loss) in 30m sided pixels with over 20% of tree cover in 2000.

**Percent of crop area** (Figure 2(b)) is derived from ESA land cover data in 2000, available at a 300m resolution (described in Section A.5 of Supplementary material).

### 3.2.3 Remoteness/connectivity: population, energy and markets

**Population density** and **remoteness** from energy sources and economic activities are considered in order to look for heterogeneities in the impact of a bad rainy season on deforestation. The log of population density (ind./km<sup>2</sup>) for the year 2000 is taken from *WorldPop* population data (1 km resolution, Tatem (2017), see Figure 2(c)). Remoteness is assessed based upon information relating to existence of powered human settlements in the vicinity of a given pixel (distance to night-time lights, see Supplementary section A.6.1) and the time-distance to the nearest city (travel time to the nearest location with 1500 inhabitants per km<sup>2</sup>, see Supplementary section A.6.2). Night-time lights reflect the existence of potential alternative sources of income, since off-farm activities often require access to power sources. The distance to the nearest city determines travel costs and, more generally, degree of accessibility of markets.

## 3.3 Statistical identification

I estimated a pixel-year fixed effect regression of the log of deforestation level.

$$x_{it} = \alpha_i + \gamma_t + \beta \cdot Z_{it} + \mu_{it} \quad (1)$$

where  $x_{it}^*$  is the log of deforestation in ares<sup>7</sup> in pixel  $i$  and year  $t$ ;  $\alpha_i$  is the specific effect of pixel  $i$ ;  $\gamma$  is the generalised annual shock of year  $t$ ;  $Z_{it}$  is the vector of explanatory variables (i.e. the dummy variable that indicates the presence of a bad rainy season in  $t$ ) and  $\mu_{it}$  is the error term. Choosing to work with the log transformation of the independent variable is justified by the highly positively skewed and lognormal shape of the deforestation distribution.  $\beta$  is the parameter estimated for every combination of proportion of crop area and remoteness.

Current rainfall shocks are largely exogenous to tree cover losses (see Supplementary section C.1), and linked to agricultural income, which represents a large proportion of rural household income. Therefore, I can assert that what we are looking at is a causal relationship between income shocks and tree cover losses. Although there may be multiple transmission channels, the selected estimation charts the short-term impact of seasonal rainfall shocks and deforestation at fine scales of time and space.

Using pixel fixed effects in our panel regressions enables to control for all time invariant pixel specifics. Using year fixed effects in our panel regressions enables to control for

<sup>7</sup> $\log(X+1)$ ,  $X$  being the deforestation in ares observed in pixel  $i$  and year  $t$ . Deforestation is the average forest cover in 2000 of 1 arc-second (approximately 30m at the equator) pixels lying in 0.05 degree pixels considered as cleared in year  $t$  by Hansen et al. (2013)'s algorithm. See Appendix A.2 of Supplementary material for further details about deforestation estimates.

all global annual variations observable in all pixels. Every regression is clustered at the second highest sub-national administrative level (see Appendix A.4 of Supplementary material), corresponding to regions lying under the province level.

The fixed effect estimator requires that there be no dependence of the error term across pixels and across time periods. It is possible for spatial correlation to occur in this framework, as the fine spatial unit renders spatial dependence highly probable, as a result of leakage, community level trade-offs (Alix-Garcia, 2007), neighbour effects or simply because deforestation creates routes of access, increasing the accessibility of remote locations in dense forests. In addition, although annual weather indices are independent from past realisations, serial correlation may occur in the case of deforestation, depending on potential accumulation of past weather events or simply on past deforestation shocks. In order to address these issues, I check that the main results are robust to a nonparametric standard errors estimation, allowing for both cross-sectional spatial correlation and location-specific serial correlation (Conley, 1999; Hsiang et al., 2011). In the Supplementary Tables of Section D.5, I allow for spatial correlation of 50 kilometres and serial correlation over 7 consecutive years, using recent Stata routine *acreg* developed by Colella et al. (2019), and based on Hsiang (2010) and Conley (1999). Robust standard errors are such that, in most of the cases, our coefficients of interest are still statistically significant for the main specifications.

## 4 Main results: heterogeneous but significant impacts of a bad rainy season on deforestation

### 4.1 Weather shocks and deforestation in the 4 eco-climatic zones

When one examines the impact of a bad rainy season on the whole sample, one finds that season length and cumulative rainfall over the rainy season have opposite impacts. Indeed, a short rainfall season increases deforestation while a drought tends to reduce deforestation (Supplementary Table 4). Moreover, by interacting annual rainfall shocks and long-term average cumulative rainfall, I find that both impacts are largely driven by pixels with a humid climate. Because this impact largely depends on average cumulative rainfall, I present the results for each eco-climatic zones hereafter.

The impacts of bad rainy seasons on deforestation are displayed in Figure 3. Impacts heterogeneity across eco-climatic zones can be due heterogeneities in: water resource availability, land use, anthropogenic pressure and/or economic development. Except for the dryer (Sahelian) eco-climatic zone, the two rainfall indices considered also have distinct and significant impacts. Indeed, a short rainy season tends to increase deforestation while a lack of seasonal rainfall reduces deforestation.

The impacts of a bad rainy season can account for a significant proportion of observed deforestation in the Guinean and Guineo-congolian eco-climatic zones: during a year with an exceptionally short rainy season, deforestation is approximately 15% higher. Inversely, during drought years, deforestation is 10% lower. In the most humid eco-climatic zone (Guineo-congolian), this negative impact is driven by pixels with a




Figure 3: Impact of a bad rainy season on deforestation.

Regressions are displayed in Supplementary Tables 5, 6, 7 and 8.

significant crop area. Deforestation is reduced by 40% during drought years in pixels where cropping exceeds 25% of the area in 2000 (left panels of Figure 3).

## 4.2 Robustness checks

Considering other forest definitions, corresponding to different canopy cover thresholds, demonstrates the robustness of previous results to forest definition. Regressions were run without any threshold<sup>8</sup> and for higher (25 and 30%) canopy cover (see Section D.2), in accordance with global studies using the same data and considering different tree cover types (Hansen et al., 2010; Heino et al., 2015). No significant impact is found in the Sahelian eco-climatic zone with a 20% canopy cover threshold (previous Section, 4.1). This may be explained by the fact that wooded landscapes are scarce and mainly consist in savannahs, and is consistent with the significant impact found without any limitations of canopy cover.

<sup>8</sup>In the absence of any threshold, any vegetation taller than 5m in height is considered, which is compatible with the Sahelian zone, where woody cover mainly consist in woody savannah or shrubland.

1  
2  
3  
4  
5  
6 When considering both more extreme and more common events, by setting the SPI  
7 threshold to -0.5, -0.7 and -1.5 (as compared to -1 in the previous section), results seem  
8 to hold in all eco-climatic zones (see Section D.3 of Supplementary material).

9 When considering alternative definitions of eco-climatic zones, the results also hold  
10 in all zones (see section D.1 of Supplementary material).

11 The main results hold when allowing for spatial correlation within 50 kilometres and  
12 serial correlation over 7 consecutive years (see section D.5 of Supplementary material).

### 13 14 15 **4.3 Deforestation and remoteness**

16 The impact of an income shock on deforestation may depend on the elasticity of the  
17 demand for agricultural products, which may itself be influenced by remoteness. The  
18 more connected an area is, the lower transport costs are for importing food or exporting  
19 non-agricultural products, and the greater off-farm income is to smooth consumption  
20 and cope with shocks. As a result, the impact of bad rainy seasons on land use decisions  
21 could be lower in connected areas.

22  
23 Moreover, the proportion of cultivated land is a good proxy for agricultural pro-  
24 duction, which can potentially both relax the subsistence constraint and increase the  
25 pressure on land use. If one considers that households are trapped in a subsistence and  
26 isolated economy when they live in a remote location, it makes sense that the subsistence  
27 constraint is alleviated by a large crop production, thereby leading to a more elastic de-  
28 mand for crop production. Additionally, the higher the pressure on land use, the less  
29 room there is for extension of the cultivated area. The elasticity of the supply would thus  
30 be lower in places where there is already a large proportion of land dedicated to culti-  
31 vation. Both inelastic supply and elastic demand leading to lower cropland expansion,  
32 the high propensity to deforest in remote areas is expected to be inhibited in pixels with  
33 high crop area.

34  
35 By focusing on the eco-climatic zones in which the impact of a bad rainy season  
36 on deforestation is significant (all zones except the Sahelian eco-climatic zone), I test  
37 the hypothesis that the impact of a bad rainy season depends upon the presence of  
38 alternative income sources or on local agricultural production. On average, every pixel  
39 experience about 3 droughts and 3 short rainy seasons during the study period<sup>9</sup>. Both  
40 short rainy seasons and droughts represent negative income shocks. However, they seem  
41 to have an opposite short-term effect on deforestation. By taking advantage of the  
42 opposite short-term impact of rainfall shocks on deforestation shocks (depending on  
43 rainy season specifics), it is possible to test whether the occurrence of positive and  
44 negative deforestation shocks depends on the pixel time invariant characteristics.

45  
46 Figure 4 shows that additional deforestation due to negative income shocks is clearly  
47 driven by unconnected pixels, i.e. pixels with low population density (a), pixels far away  
48 from markets (b) or pixels far away from energy sources (c). To put it in another  
49 way, after a short rainy season, positive deviations to the average long term (19 years)

50  
51  
52 <sup>9</sup>The probability of return of the events studied is 15%, meaning that the indices exceed one standard  
53 deviation less than once every 6 years.




Figure 4: Impact of a bad rainy season on deforestation in Sudanian, Guinean and Guineo-congolian eco-climatic zones, as a function of Relative remoteness in terms of population density (a) and Relative remoteness in terms of time-distance to cities (b) Relative remoteness in terms of distance to night-time lights (c).

Pixels with high (low) population density correspond to values higher (lower) than the median value of the log population density (2.6); pixels far (near) from the nearest market are at more (less) than 90 minutes from the nearest city and pixels far (near) from powered settlements are at more (less) than 13.5 km from a detectable night-time light. Regressions are displayed in Supplementary tables 9, 10 and 11.



1  
2  
3  
4  
5  
6 deforestation in each pixel are of higher amplitude in remote or poorly connected areas.  
7 This additional deforestation in years with short rainy seasons reaches 20% in remote  
8 areas with a low share of crop areas. This significant impact is robust to the choice of  
9 the proxy used to estimate the remoteness (Figure 4) and to the sample of eco-climatic  
10 zones considered (see Supplementary Figure 17 in section D.4, showing the results on a  
11 sample restricted to the Guinean and Guineo-congolian eco climatic zones).

12  
13 Inversely, the negative impact of a drought does not seem to be significantly influ-  
14 enced by given pixel's remoteness or share of crop area.

15 To summarise the results, a negative income shock would thus have an ambiguous  
16 short-term impact on land use, depending on intra-seasonal rainfall timing, but positive  
17 impact on deforestation mainly depends on relative remoteness and pressure on land use  
18 in the vicinity. These results are consistent with existing smallholder subsistence land  
19 use theories and with neo-classical economic theories of land rent (Meyfroidt et al., 2018),  
20 and could potentially reconcile the puzzling ambivalent empirical evidence relating to  
21 the short-term impact of income shocks on deforestation.  
22  
23

## 24 5 Discussion

25  
26 The complexity of ecosystem management, especially in the presence of relationships  
27 between humans and the ecosystems, generates wicked problems (DeFries and Nagendra,  
28 2017). The existence of climate feedbacks on land use via human-related activities  
29 support the hypothesis that there may be vicious circles in inter- and intra-annual climate  
30 variability and land use changes dynamics. Thinking about poverty alleviation in a  
31 broader context, in which the ecological and social consequences of shocks are integrated,  
32 should help to increase smallholders resilience in the long term (Lade et al., 2017).  
33

34 This article shows that in Western and Central Africa, deforestation depends on  
35 the quality of the rainy season. Short rainy seasons lead to greater deforestation and  
36 a low amount of seasonal rainfall leads to reduced deforestation. This difference may  
37 be explained by the relative predictability of these extreme events. Since the length of  
38 the rainy season largely depends on its onset (the offset being much less variable, see  
39 Supplementary Figure 6), it is easy for farmers to anticipate a short rainy season and  
40 to plan farm work outside the crop calendar. However, a lack of rainfall may occur  
41 throughout the rainy season, making this event unpredictable. Finally, it may also  
42 be that, under humid climates (Guinean and Guineo-congolian), a short rainy season  
43 coincides with a longer period, during which forests accessibility is increased.  
44

45 These impacts are large in amplitude. Indeed, in the most humid eco-climatic zones,  
46 deforestation is 5 to 20% higher during years with a short rainy season. This suggests  
47 that tree cover loss may be linked to economic, and more precisely, agricultural outcomes.  
48 Therefore, if we consider the way that rainfall shocks influence deforestation, we could  
49 improve the design of policies created to fight deforestation in poor regions in which  
50 income data availability is scarce.  
51

52 The positive impacts of bad rainy seasons on deforestation are more salient in remote  
53 areas with low pressure on land use. It is possible to elaborate development and conser-  
54  
55

1  
2  
3  
4  
5  
6 vation programmes in such way that they simultaneously support conservation efforts  
7 and adaptation of smallholder agriculture. As the impacts depend on both the quality  
8 of the rainy season and the remoteness of considered locations, smart adaptation policies  
9 could be designed to target remote areas to meet environmental and development ob-  
10 jectives. Given that the onset of the rainy season is much more variable than the offset  
11 in all eco-climatic zones, policies could focus on this indicator, already recognised in the  
12 Sudano-Sahelian zone as a good indicator of seasonal rainfall quality for agricultural pro-  
13 duction (Dodd and Jolliffe, 2001; Marteau et al., 2011). For example, drought insurance  
14 compensating farmers after the late onset of a rainfall season could significantly reduce  
15 the pressure on the remaining forest landscapes of remote areas, in addition to providing  
16 a safety net for smallholders. Alternatively, conservation programme enforcement could  
17 be designed to target years with a short rainfall season in remote areas with low pressure  
18 on land use to increase conservation effectiveness.

19  
20 Further research could help to identify the transmission channels of these feedbacks.  
21 Forest clearing is not only driven by the need to extend cultivation areas. Results  
22 provide evidence of an immediate impact of a bad rainy season on deforestation, though  
23 there also exist many potential indirect and long-term impacts, such as migration and  
24 urbanisation (Ruf et al., 2015; Henderson et al., 2017), that may interact with this  
25 immediate impact. Furthermore, adaptation to weather variability by using crop mix,  
26 improved seed varieties or physical infrastructure such as dams and irrigation systems  
27 may inhibit these feedbacks. Finally, future work could also be dedicated to extending  
28 the analysis to other regions, on a greater scale or globally.

## 31 32 6 Data and code availability

33  
34 The data and code that support the findings of this study are openly available at  
35 [doi.org/10.5281/zenodo.4266325](https://doi.org/10.5281/zenodo.4266325).

## 36 37 38 References

- 39  
40 (2017): “ESA Land Cover CCI Product User Guide Version 2. Tech.  
41 Rep.” =[http://maps.elie.ucl.ac.be/CCI/viewer/download/ESACCI-LC-Ph2-](http://maps.elie.ucl.ac.be/CCI/viewer/download/ESACCI-LC-Ph2-PUGv2.2.0.pdf)  
42 [PUGv2.2.0.pdf](http://maps.elie.ucl.ac.be/CCI/viewer/download/ESACCI-LC-Ph2-PUGv2.2.0.pdf), european Space Agency and Climate Change Initiative.  
43  
44 AGARWAL, B. (1991): “Social security and the family: coping with seasonality and  
45 calamity in rural India,” in *E. Ahmad, J. Dreze, J. Hills, A. Sen (ed.), Social Security*  
46 *in Developing Countries*, Oxford: Clarendon Press, pp. 17-24.  
47  
48 ALEIXO, I., D. NORRIS, L. HEMERIK, A. BARBOSA, E. PRATA, F. COSTA, AND  
49 L. POORTER (2019): “Amazonian rainforest tree mortality driven by climate and  
50 functional traits,” *Nature Climate Change*, 9, 384.  
51  
52 ALIX-GARCIA, J. (2007): “A spatial analysis of common property deforestation,” *Jour-*  
53 *nal of Environmental Economics and Management*, 53, 141–157.  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6 ALIX-GARCIA, J., C. MCINTOSH, K. R. SIMS, AND J. R. WELCH (2013): “The ecological footprint of poverty alleviation: evidence from Mexico’s Oportunidades program,”  
7 *Review of Economics and Statistics*, 95, 417–435.  
8  
9  
10 ANDEREGG, W. R., J. M. KANE, AND L. D. ANDEREGG (2013): “Consequences  
11 of widespread tree mortality triggered by drought and temperature stress,” *Nature*  
12 *climate change*, 3, 30.  
13  
14 ANGELSEN, A. AND S. WUNDER (2002): “Exploring the forest-poverty link,” *CIFOR*  
15 *Occasional Paper*, 40.  
16  
17 ARAUJO, C., J.-L. COMBES, AND J. G. FERES (2019): “Determinants of Amazon  
18 deforestation: the role of off-farm income,” *Environment and Development Economics*,  
19 24, 138–156.  
20  
21 ARCHIBALD, S. (2016): “Managing the human component of fire regimes: lessons from  
22 Africa,” *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371,  
23 20150346.  
24  
25 ASSUNÇÃO, J., C. GANDOUR, R. ROCHA, AND R. ROCHA (2019): “The Effect of Rural  
26 Credit on Deforestation: Evidence from the Brazilian Amazon Effect of Rural Credit  
27 on Deforestation,” *The Economic Journal*.  
28  
29 BALAND, J. AND P. FRANCOIS (2005): “Commons as insurance and the welfare impact  
30 of privatization,” *Journal of Public Economics*, 89, 211–231.  
31  
32 BALAND, J.-M., P. BARDHAN, S. DAS, D. MOOKHERJEE, AND R. SARKAR (2010):  
33 “The environmental impact of poverty: evidence from firewood collection in rural  
34 Nepal,” *Economic Development and Cultural Change*, 59, 23–61.  
35  
36 BARON, C., B. SULTAN, M. BALME, B. SARR, S. TRAORE, T. LEBEL, S. JANICOT,  
37 AND M. DINGKUHN (2005): “From GCM grid cell to agricultural plot: scale issues  
38 affecting modelling of climate impact,” *Philosophical Transactions of the Royal Society*  
39 *B: Biological Sciences*, 360, 2095–2108.  
40  
41 BASTIN, J.-F., N. BERRAHMOUNI, A. GRAINGER, D. MANIATIS, D. MOLLICONE,  
42 R. MOORE, C. PATRIARCA, N. PICARD, B. SPARROW, E. M. ABRAHAM, ET AL.  
43 (2017): “The extent of forest in dryland biomes,” *Science*, 356, 635–638.  
44  
45 BIASUTTI, M. (2019): “Rainfall trends in the African Sahel: Characteristics, processes,  
46 and causes,” *Wiley Interdisciplinary Reviews: Climate Change*, e591.  
47  
48 BICHET, A. AND A. DIEDHIOU (2018): “West African Sahel has become wetter during  
49 the last 30 years, but dry spells are shorter and more frequent,” *Climate Research*, 75,  
50 155–162.  
51  
52 BLUFFSTONE, R. A. (1995): “The effect of labor market performance on deforestation  
53 in developing countries under open access: an example from rural Nepal,” *Journal of*  
54 *Environmental Economics and Management*, 29, 42–63.  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6 BOYARD-MICHEAU, J., P. CAMBERLIN, N. PHILIPPON, AND V. MORON (2013):  
7 “Regional-scale rainy season onset detection: a new approach based on multivariate  
8 analysis,” *Journal of Climate*, 26, 8916–8928.
- 9  
10 BRANDT, M., C. TUCKER, A. KARIRYAA, K. RASMUSSEN, C. ABEL, L. V. RAS-  
11 MUSSEN, AND P. HIERNAUX (2020): “Civil conflicts are associated with the global  
12 climate,” *Nature*, 1476–4687.
- 13  
14 BRANDT, M., J.-P. WIGNERON, J. CHAVE, T. TAGESSON, J. PENUELAS, P. CIAIS,  
15 K. RASMUSSEN, F. TIAN, C. MBOW, A. AL-YAARI, ET AL. (2018): “Satellite passive  
16 microwaves reveal recent climate-induced carbon losses in African drylands,” *Nature*  
17 *ecology & evolution*, 2, 827.
- 18  
19 BRODRIBB, T. J., J. POWERS, H. COCHARD, AND B. CHOAT (2020): “Hanging by a  
20 thread? Forests and drought,” *Science*, 368, 261–266.
- 21  
22 COLELLA, F., R. LALIVE, S. O. SAKALLI, AND M. THOENIG (2019): “Inference with  
23 arbitrary clustering,” .
- 24  
25 CONLEY, T. G. (1999): “GMM estimation with cross sectional dependence,” *Journal*  
26 *of econometrics*, 92, 1–45.
- 27  
28 COTILLON, S. E. AND G. G. TAPPAN (2016): “Landscapes of West Africa: A win-  
29 dows on a changing world,” [https://eros.usgs.gov/westafrika/land-cover/land-use-and-](https://eros.usgs.gov/westafrika/land-cover/land-use-and-land-cover-trends-west-africa)  
30 [land-cover-trends-west-africa](https://eros.usgs.gov/westafrika/land-cover/land-use-and-land-cover-trends-west-africa).
- 31  
32 CURTIS, P. G., C. M. SLAY, N. L. HARRIS, A. TYUKAVINA, AND M. C. HANSEN  
33 (2018): “Classifying drivers of global forest loss,” *Science*, 361, 1108–1111.
- 34  
35 DEFRIES, R. AND H. NAGENDRA (2017): “Ecosystem management as a wicked prob-  
36 lem,” *Science*, 356, 265–270.
- 37  
38 DESBUREAUX, S. AND R. DAMANIA (2018): “Rain, forests and farmers: Evidence of  
39 drought induced deforestation in Madagascar and its consequences for biodiversity  
40 conservation,” *Biological Conservation*, 221, 357–364.
- 41  
42 DIACONESCU, E. P., P. GACHON, J. SCINOCCA, AND R. LAPRISE (2015): “Evalu-  
43 ation of daily precipitation statistics and monsoon onset/retreat over western Sahel in  
44 multiple data sets,” *Climate Dynamics*, 45, 1325–1354.
- 45  
46 DODD, D. E. AND I. T. JOLLIFFE (2001): “Early detection of the start of the wet season  
47 in semiarid tropical climates of western Africa,” *International Journal of Climatology:*  
48 *A Journal of the Royal Meteorological Society*, 21, 1251–1262.
- 49  
50 DUNNING, C. M., E. BLACK, AND R. P. ALLAN (2018): “Later wet seasons with  
51 more intense rainfall over Africa under future climate change,” *Journal of Climate*,  
52 31, 9719–9738.
- 53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6 DUNNING, C. M., E. C. BLACK, AND R. P. ALLAN (2016): “The onset and cessation  
7 of seasonal rainfall over Africa,” *Journal of Geophysical Research: Atmospheres*, 121,  
8 11–405.
- 9  
10 ETONGO, D., I. N. S. DJENONTIN, M. KANNINEN, K. FOBISSIE, K. KORHONEN-  
11 KURKI, AND H. DJOUDI (2015): “Land tenure, asset heterogeneity and deforestation  
12 in Southern Burkina Faso,” *Forest Policy and Economics*, 61, 51–58.
- 13  
14 FAO (2007): “Sahel weather and crop situation report,” [Http://www.fao.org/3/ah870e.pdf](http://www.fao.org/3/ah870e.pdf).
- 15  
16  
17 ——— (2020): “Global Forest Resources Assessment 2020,”  
18 [Http://www.fao.org/3/CA8753EN/CA8753EN.pdf](http://www.fao.org/3/CA8753EN/CA8753EN.pdf).
- 19  
20 FERRARO, P. J. AND R. SIMORANGKIR (2020): “Conditional cash transfers to alleviate  
21 poverty also reduced deforestation in Indonesia,” *Science Advances*, 6, eaaz1298.
- 22  
23 FUNK, C., P. PETERSON, M. LANDSFELD, D. PEDREROS, J. VERDIN, S. SHUKLA,  
24 G. HUSAK, J. ROWLAND, L. HARRISON, A. HOELL, ET AL. (2015): “The climate  
25 hazards infrared precipitation with stations—a new environmental record for moni-  
26 toring extremes,” *Scientific data*, 2, 150066.
- 27  
28 HANSEN, M. C., P. V. POTAPOV, R. MOORE, M. HANCHER, S. TURUBANOVA,  
29 A. TYUKAVINA, D. THAU, S. STEHMAN, S. GOETZ, T. LOVELAND, ET AL. (2013):  
30 “High-resolution global maps of 21st-century forest cover change,” *Science*, 342, 850–  
31 853.
- 32  
33 HANSEN, M. C., S. V. STEHMAN, AND P. V. POTAPOV (2010): “Quantification of  
34 global gross forest cover loss,” *Proceedings of the National Academy of Sciences*, 107,  
35 8650–8655.
- 36  
37 HEINO, M., M. KUMMU, M. MAKKONEN, M. MULLIGAN, P. H. VERBURG,  
38 M. JALAVA, AND T. A. RÄSÄNEN (2015): “Forest loss in protected areas and in-  
39 tact forest landscapes: a global analysis,” *PloS one*, 10, e0138918.
- 40  
41 HENDERSON, J. V., A. STOREYGARD, AND U. DEICHMANN (2017): “Has climate  
42 change driven urbanization in Africa?” *Journal of development economics*, 124, 60–  
43 82.
- 44  
45 HESS, S., D. JAIMOVICH, AND M. SCHÜNDELN (2019): “Environmental Effects of  
46 Development Programs: Experimental Evidence from West African Dryland Forests,”  
47 Tech. rep., Frankfurt University.
- 48  
49 HSIANG, S. M. (2010): “Temperatures and cyclones strongly associated with eco-  
50 nomic production in the Caribbean and Central America,” *Proceedings of the National  
51 Academy of Sciences*, 107, 15367–15372.
- 52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6 HSIANG, S. M., K. C. MENG, AND M. A. CANE (2011): “Civil conflicts are associated  
7 with the global climate,” *Nature*, 476, 438–441.
- 8  
9 JAYACHANDRAN, S., J. DE LAAT, E. F. LAMBIN, C. Y. STANTON, R. AUDY, AND  
10 N. E. THOMAS (2017): “Cash for carbon: A randomized trial of payments for ecosys-  
11 tem services to reduce deforestation,” *Science*, 357, 267–273.
- 12  
13 JIANG, Y., L. ZHOU, C. J. TUCKER, A. RAGHAVENDRA, W. HUA, Y. Y. LIU, AND  
14 J. JOINER (2019): “Widespread increase of boreal summer dry season length over the  
15 Congo rainforest,” *Nature Climate Change*, 9, 617–622.
- 16  
17 KOSMOWSKI, F., A. LEBLOIS, AND B. SULTAN (2016): “Perceptions of recent rainfall  
18 changes in Niger: a comparison between climate-sensitive and non-climate sensitive  
19 households,” *Climatic change*, 135, 227–241.
- 20  
21 LADE, S. J., L. J. HAIDER, G. ENGSTRÖM, AND M. SCHLÜTER (2017): “Resilience  
22 offers escape from trapped thinking on poverty alleviation,” *Science Advances*, 3,  
23 e1603043.
- 24  
25 LAWRENCE, D. AND K. VANDECAR (2015): “Effects of tropical deforestation on climate  
26 and agriculture,” *Nature Climate Change*, 5, 27.
- 27  
28 LEITE-FILHO, A. T., V. Y. DE SOUSA PONTES, AND M. H. COSTA (2019): “Effects  
29 of deforestation on the onset of the rainy season and the duration of dry spells in  
30 southern Amazonia,” *Journal of Geophysical Research: Atmospheres*.
- 31  
32 LIEBMANN, B., I. BLADÉ, G. N. KILADIS, L. M. CARVALHO, G. B. SENAY, D. AL-  
33 LURED, S. LEROUX, AND C. FUNK (2012): “Seasonality of African precipitation from  
34 1996 to 2009,” *Journal of Climate*, 25, 4304–4322.
- 35  
36 LIEBMANN, B. AND J. MARENGO (2001): “Interannual variability of the rainy season  
37 and rainfall in the Brazilian Amazon Basin,” *Journal of Climate*, 14, 4308–4318.
- 38  
39 MARTEAU, R., B. SULTAN, V. MORON, A. ALHASSANE, C. BARON, AND S. B.  
40 TRAORÉ (2011): “The onset of the rainy season and farmers’ sowing strategy for  
41 pearl millet cultivation in Southwest Niger,” *Agricultural and forest meteorology*, 151,  
42 1356–1369.
- 43  
44 MAYAUX, P., F. ACHARD, AND J.-P. MALINGREAU (1998): “Global tropical forest  
45 area measurements derived from coarse resolution satellite imagery: a comparison  
46 with other approaches,” *Environmental Conservation*, 25, 37–52.
- 47  
48 MEYFROIDT, P., R. R. CHOWDHURY, A. DE BREMOND, E. C. ELLIS, K.-H. ERB,  
49 T. FILATOVA, R. GARRETT, J. M. GROVE, A. HEINIMANN, T. KUEMMERLE, ET AL.  
50 (2018): “Middle-range theories of land system change,” *Global Environmental Change*,  
51 53, 52–67.
- 52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6 MOLINARIO, G., M. HANSEN, P. POTAPOV, A. TYUKAVINA, AND S. STEHMAN (2020):  
7 “Contextualizing Landscape-Scale Forest Cover Loss in the Democratic Republic of  
8 Congo (DRC) between 2000 and 2015,” *Land*, 9, 23.
- 9  
10 NOACK, F., M.-C. RIEKHOF, AND S. DI FALCO (2019): “Droughts, Biodiversity, and  
11 Rural Incomes in the Tropics,” *Journal of the Association of Environmental and Re-*  
12 *source Economists*, 6, 823–852.
- 13  
14 ODEKUNLE, T. (2004): “Rainfall and the length of the growing season in Nigeria,”  
15 *International Journal of Climatology: A Journal of the Royal Meteorological Society*,  
16 24, 467–479.
- 17  
18 PANTHOU, G., T. LEBEL, T. VISCHÉL, G. QUANTIN, Y. SANE, A. BA, O. NDIAYE,  
19 A. DIONGUE-NIANG, AND M. DIOPKANE (2018): “Rainfall intensification in tropical  
20 semi-arid regions: The Sahelian case,” *Environmental Research Letters*, 13, 064–013.
- 21  
22 PATRIARCA, C., M. BAKO, A. BRANTHOMME, T. S. FRESCINO, F. F. HADDAD,  
23 A. H. HAMID, A. MARTUCCI, H. O. CHOUR, P. L. PATTERSON, N. PICARD,  
24 ET AL. (2019): “Trees, forests and land use in drylands: The first global assessment,”  
25 *FAO Forestry Paper No. 184. Rome, Italy: Food and Agriculture Organization of the*  
26 *United Nations. 184 p.*
- 27  
28 PATTANAYAK, S. AND E. SILLS (2001): “Do tropical forests provide natural insurance?  
29 The microeconomics of non-timber forest product collection in the Brazilian Amazon,”  
30 *Land Economics*, 77, 595–612.
- 31  
32 PHILLIPS, O. L., L. E. ARAGÃO, S. L. LEWIS, J. B. FISHER, J. LLOYD, G. LÓPEZ-  
33 GONZÁLEZ, Y. MALHI, A. MONTEAGUDO, J. PEACOCK, C. A. QUESADA, ET AL.  
34 (2009): “Drought sensitivity of the Amazon rainforest,” *Science*, 323, 1344–1347.
- 35  
36 ROUDIER, P., B. SULTAN, P. QUIRION, AND A. BERG (2011): “The impact of future  
37 climate change on West African crop yields: what does the recent literature say?”  
38 *Global Environmental Change*, 21, 1073–1083.
- 39  
40 RUF, F., G. SCHROTH, AND K. DOFFANGUI (2015): “Climate change, cocoa migrations  
41 and deforestation in West Africa: What does the past tell us about the future?”  
42 *Sustainability Science*, 10, 101–111.
- 43  
44 SEXTON, J. O., P. NOOJIPADY, X.-P. SONG, M. FENG, D.-X. SONG, D.-H. KIM,  
45 A. ANAND, C. HUANG, S. CHANNAN, S. L. PIMM, ET AL. (2016): “Conservation  
46 policy and the measurement of forests,” *Nature Climate Change*, 6, 192–196.
- 47  
48 SHIVELY, G. E. AND S. PAGIOLA (2004): “Agricultural intensification, local labor mar-  
49 kets, and deforestation in the Philippines,” *Environment and Development Economics*,  
50 241–266.
- 51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- SOMORIN, O. A., H. C. P. BROWN, I. J. VISSEREN-HAMAKERS, D. J. SONWA, B. ARTS, AND J. NKEM (2012): “The Congo Basin forests in a changing climate: policy discourses on adaptation and mitigation (REDD+),” *Global Environmental Change*, 22, 288–298.
- STAAL, A., B. M. FLORES, A. P. D. AGUIAR, J. H. BOSMANS, I. FETZER, AND O. A. TUINENBURG (2020): “Feedback between drought and deforestation in the Amazon,” *Environmental Research Letters*, 15, 044024.
- SVOBODA, M., M. HAYES, AND D. WOOD (2012): “Standardized precipitation index user guide,” *World Meteorological Organization Geneva, Switzerland*.
- TATEM, A. J. (2017): “WorldPop, open data for spatial demography,” *Scientific data*, 4.
- TAYLOR, C. M., D. BELUŠIĆ, F. GUICHARD, D. J. PARKER, T. VISCHEL, O. BOCK, P. P. HARRIS, S. JANICOT, C. KLEIN, AND G. PANTHOU (2017): “Frequency of extreme Sahelian storms tripled since 1982 in satellite observations,” *Nature*, 544, 475.
- THE WORLD BANK (2020): “Agriculture, forestry, and fishing, value added (% of GDP),” .
- TSCHARNTKE, T., Y. CLOUGH, T. C. WANGER, L. JACKSON, I. MOTZKE, I. PERFECTO, J. VANDERMEER, AND A. WHITBREAD (2012): “Global food security, biodiversity conservation and the future of agricultural intensification,” *Biological conservation*, 151, 53–59.
- TYUKAVINA, A., M. C. HANSEN, P. POTAPOV, D. PARKER, C. OKPA, S. V. STEHMAN, I. KOMMAREDDY, AND S. TURUBANOVA (2018): “Congo Basin forest loss dominated by increasing smallholder clearing,” *Science advances*, 4, eaat2993.
- UNEP, A. (2001): “An Assessment of the Status of the World’s Remaining Closed Forests.” Tech. rep., UNEP/DEWA/TR01-2, Division of Earth Warming and Assessment, UNEP.
- VERBESSELT, J., N. UMLAUF, M. HIROTA, M. HOLMGREN, E. H. VAN NES, M. HEROLD, A. ZEILEIS, AND M. SCHEFFER (2016): “Remotely sensed resilience of tropical forests,” *Nature Climate Change*, 6, 1028.
- WEISS, D. J., A. NELSON, H. GIBSON, W. TEMPERLEY, S. PEEDELL, A. LIEBER, M. HANCHER, E. POYART, S. BELCHIOR, N. FULLMAN, ET AL. (2018): “A global map of travel time to cities to assess inequalities in accessibility in 2015,” *Nature*, 553, 333.
- WOLFF, N. H., Y. J. MASUDA, E. MEIJAARD, J. A. WELLS, AND E. T. GAME (2018): “Impacts of tropical deforestation on local temperature and human well-being perceptions,” *Global environmental change*, 52, 181–189.



1  
2  
3  
4  
5  
6 WUNDER, S. (2001): “Poverty alleviation and tropical forests: what scope for syner-  
7 gies?” *World Development*, 29, 1817–1833.

8  
9 ZAVERI, E., J. RUSS, AND R. DAMANIA (2020): “Rainfall anomalies are a significant  
10 driver of cropland expansion,” *Proceedings of the National Academy of Sciences*, 117,  
11 10225–10233.

12  
13 ZEMP, D. C., C.-F. SCHLEUSSNER, H. M. BARBOSA, M. HIROTA, V. MONTADE,  
14 G. SAMPAIO, A. STAAL, L. WANG-ERLANDSSON, AND A. RAMMIG (2017): “Self-  
15 amplified Amazon forest loss due to vegetation-atmosphere feedbacks,” *Nature com-  
16 munications*, 8, 14681.

17  
18 ZHANG, W., M. BRANDT, F. GUICHARD, Q. TIAN, AND R. FENSHOLT (2017): “Using  
19 long-term daily satellite based rainfall data (1983-2015) to analyze spatio-temporal  
20 changes in the sahelian rainfall regime,” *Journal of hydrology*, 550, 427–440.

## 21 22 23 **7 Appendix**

### 24 25 **A Data**

#### 26 27 **A.1 Tree cover**

28  
29 Hansen et al. (2013) tree cover data for year 2000 is available at 1 arc-second resolution:  
30 about 30m at equator. The data considers pixels with more than 1% of forest cover  
31 (vegetation taller than 5m height) in 2000. The forest and more generally tree cover of  
32 the region studied is very diverse, and largely follow the rainfall gradient (see Figure 1  
33 and Figure 5).

#### 34 35 **A.2 Deforestation**

36  
37 Hansen et al. (2013) estimates tree cover loss annually over the 2001-2019 period (version  
38 1.7, available at 1 arc-second resolution), defined as a stand-replacement disturbance, or  
39 a change from a forest to non-forest state. It provides a year of tree cover loss for every  
40 pixel that is estimated to endure a loss of more than 50% of the 2000 forest cover, using  
41 an underlying algorithm trained in Central Africa (more precisely RDC in the Congo  
42 Basin).

43  
44 I have considered that the whole pixel (30x30m) tree cover was reduced to zero  
45 when losses occurs, and forest degradation, for example selective removals from within  
46 forested stands that do not lead to a non-forest state, is not included in the change  
47 characterisation. Moreover, the data does not allow to distinguish quality of the canopy  
48 and select every vegetation higher than 5m, potentially leading to consider secondary  
49 forest loss as deforestation.

50  
51 Bastin et al. (2017) showed that drylands, covering about two fifth of the Earth’s  
52 land surface, had more than 10% tree-cover and also comprise forests. Although the  
53  
54  
55




Figure 5: 500 mm isohyets (average cumulative annual precipitations, in mm.) in the study region, according to CHIRPS 2000-2019 daily data.

terms of forest in this recent study is disputed, Brandt et al. (2020) also recently shows there is much more trees in the West African Sahara and Sahel than expected. Tree cover of the region studied, is diversified and included shrubland, woodland and wooded savannas. I used a 20% tree canopy cover threshold in the main text of the paper to define whether the 30m resolution grid cells are classified as forest or non-forest in 2000. I finally compute the average losses tree cover and 3 definition of forest cover (20, 25 and 30% of canopy cover), at a 0.05 decimal degree resolution (corresponding to about 5.4 km at the equator) for robustness checks (see section 4.2).

### A.3 Daily rainfall estimates

Western Africa is characterised by a rainfall gradient that largely follows latitude (see Figure 5), with the presence of different biomes depending on these isohyets.

Rainfall indices are computed using the daily CHIRPS climate data (Funk et al., 2015) available at  $0.05^\circ$ , downloaded in June 2020 (version 2.0), incorporating satellite imagery with in-situ station data to create gridded rainfall time series for trend analysis and seasonal drought monitoring.

#### A.3.1 Annual rainfall season quality indices

I retain two main annual indices that reflect the quality of the rainy season:

- the length of the rainy season in days (offset minus onset).
- the bounded seasonal cumulative rainfall. I only consider significant daily rainfall (superior to 1 mm of daily rainfall, assuring that it would not be entirely evaporated, following Odekunle (2004)), with a maximum daily bound fixed to 30mm, considering the potential runoff, following Baron et al. (2005). I restrict rainfall to seasonal rainfall, considering that the sowing and harvesting of crops largely follows the onset and offset of the rainy season in the region. Such index tries to consider

only rainfall that are able to be used by crops, and reflect the quality of the rainfall season for agricultural activities.

Both indices are standardised at the pixel level, i.e. rescaled to have a mean of 0 and a standard deviation of 1, using 20 years of data. SPIs are fitted on a normal distribution, by standardizing 20 year series of these two indices, computed from daily rainfall data (CHIRPS estimates, described in this Supplementary material, section A.3) for every pixel.

### A.3.2 Definition of the rainfall season

I only consider rainfall during the rainy season, i.e. after the onset and before the offset, that are useful for agricultural activities. Outside the cropping calendar and agricultural activities, rainfall accumulation is not necessarily critical in order to estimate the impact of an agricultural income shock, on which this article focuses.

I use the definition of the rainy season (onset and offset) of Liebmann and Marengo (2001), that allows to only consider the rainfall useful to crops. This definition has been calibrated on and applied to the Brazilian forest but also proved to be useful in the Sahel (Liebmann et al., 2012; Diaconescu et al., 2015) and all over Africa (Boyard-Micheau et al., 2013; Dunning et al., 2018). It has the advantage of being compatible with the presence of two rainy seasons during a single year, which occurs in the southern part of the study region (humid eco-climatic zones, as defined by Figure 2(b)).

It is based on the daily rainfall accumulation quantity (A):

$$A(\text{day}) = \sum_1^{\text{day}} R(n) - \bar{R} \times \text{day} \quad (2)$$

where  $R(n)$  is the daily climatological (or that of a particular year) rainfall as a function of day of year, and  $\bar{R}$  is the annual mean daily rainfall. The onset is the day which has the minimum value of this rainfall accumulation quantity over the year, for each pixel. Symmetrically, the offset corresponds to the day of the year with the maximum value of A.

Figure 6 shows the heterogeneity of onset and offset of the rainy seasons in different eco-climatic zones.

## A.4 Administrative boundaries

Every regression is clustered at the second highest international administrative level (see Figure 7), with comparable population levels. These correspond to Local Government Areas (LGAs) in Nigeria, Districts in Equatorial Guinea, Gambia, Ghana, Liberia and Sierra Leone, prefectures in Guinea, autonomous sectors in Guinea-Bissau and *Départements* in countries previously under French colonial influence<sup>10</sup> except in

<sup>10</sup>i.e. Benin, Burkina Faso, Cameroon, Central African Republic (RCA) Chad, Congo, Democratic Republic of the Congo (DRC), Gabon, Ivory Coast, Niger, Mali, Mauritania and Senegal.




Figure 6: Distribution of onset, offset date (day of the year) and length (in days) of the rainy season, by eco-climatic zone.

Mali where they correspond to *Cercles* and in the Central African Republic where they correspond to Sub-Prefectures. Political borders data, from the *GADM world political borders* was downloaded on November 2015

### A.5 Land cover classification

I use the European Space Agency (ESA) Climate Change Initiative (CCI) Land use (ESA CCI LC ESA (2017)) data, available annually from 1992 to 2019, to characterize the year 2000 land cover and compute the crop area in each pixel. Only the crop area for the year 2000 is considered, in order to limit endogeneity issues: while later land use may interfere with our interest variable (tree and forest cover) it might thus scramble the causal link.

4 land types of the original land cover classification were aggregated into 3 groups (Table 1) to assess the overall cropland cover. This classification allows to compute a final value for crop area, which is the proportion of cropland, plus 0.5 times the proportion of cropland (majoritary) and tree cover plus 0.25 times the proportion of tree (majoritary) and cropland cover.




Figure 7: 1st administrative level (blue lines) and 2nd administrative level (grey lines), the latter is used for clustering all statistical regressions. Source: GADM world political borders (<https://gadm.org/>).

Table 1: ESA land cover classification

Original (ESA) class	<b>Cropland cover</b>	Definition
10	Cropland	Cropland, rainfed
20	Cropland	Cropland, irrigated or post-flooding
40	Majoritary	Mosaic natural vegetation (tree, shrub, herbaceous cover) (>50%) / cropland (<50%)
30	Minoritary	Mosaic cropland (>50%) / natural vegetation (tree, shrub, herbaceous cover) (<50%)




Figure 8: Night-time lights data (presence of lights during the period, at less than approximately 8.1 km, pixels in yellow, and 13.5 km, in red).

## A.6 Relative remoteness

### A.6.1 Night-time lights: distance to powered settlements

Night-time lights original data are composed of 30 arc-second grids cloud-free nighttime lights composites<sup>11</sup> for the year 2000. Figure 8 shows the areas situated at less than 8.1 (one pixel, in yellow) and less than 13.5 km (two pixels, in red) of a detectable night-time light, for at least one year during the period considered.

On average, 30% of the pixels considered are at less than 8.5 km of a persistent Night-time light, while more than 45% are at less than 13.5 km of a persistent Night-time light (see zone specific summary statistics in B.2 for eco-climatic zones specific average).

### A.6.2 Accessibility: travel time to cities

Figure 9 shows the travel time (in minutes), in log scale, to the nearest city in the study region in 2015 (Weiss et al., 2018)<sup>12</sup>. Cities are defined as pixels with more than 1500 inhabitants (original resolution: 1 km).

<sup>11</sup> Version 4 DMSP-OLS Nighttime Lights Time Series, available at: <https://ngdc.noaa.gov/eog/dmsp/downloadV4composites.html>

<sup>12</sup>The data can be accessed and visualised at the following link access map in 2015.




Figure 9: Log time-distance (minutes) to cities in 2015 (of pixels of 1500 inhabitants per  $\text{km}^2$ , initial resolution: 1 km).

## B Descriptive statistics

### B.1 Whole sample

The following Tables display the descriptive statistics of the sample used in the study, i.e. pixels with at least 1 hectare of forest cover in 2000 and that have a minimum of 100 mm of rainfall per year on average during the period: which correspond to 140 418 pixels followed during 19 years (2001-2019).

Table 2 shows the descriptive statistics of the time invariant characteristics for the sample of pixels followed for 19 years. Probability of being deforested over the period is 62%, with an average of about 60 ha deforested.

Table 2: Time invariant summary statistics (N = 140 418 pixels)

Variable	Mean	Std. Dev.	Min.	Max.	N
Total tree cover loss over the period ( $\text{km}^2$ )	0.649	1.481	0	21.377	140 418
Total deforestation over the period ( $\text{km}^2$ )	0.598	1.483	0	21.377	140 418
Probability of deforestation over the period	0.626	0.484	0	1	140 418
Crop area (%)	0.313	0.345	0	1	140 409
Population density (inhab. $\text{km}^2$ )	45.076	205.34	0.003	18 149	140 418
Time-distance to the nearest city (min.)	213.495	282.402	0	3 156.4	140 418
Near (< 13.5 km) a night-time light in 2000	0.289	0.454	0	1	140 418

Table 3: Summary statistics: whole sample

Variable	Mean	Std. Dev.	Min.	Max.	N
<b>Whole sample</b>					
Cumulative annual rainfall (mm)	1 276.826	564.52	101.54	4165.3	2 644 412
Length of the rainy season (days)	154.577	43.43	25	326	2 644 412
Seasonal cumulative rainfall (mm)	1060.414	458.043	47.749	4226.1	2 644 412
Short rainfall season (year t)	0.155	0.362	0	1	2 644 412
Drought (year t)	0.151	0.358	0	1	2 644 412
Tree cover loss in ha	3.421	13.269	0	2096.4	2 644 412
Deforestation in ha	3.149	13.152	0	2096.3	2 644 412
Annual probability of deforestation	0.576	0.494	0	1	2 644 412
<b>Among which:</b>					
<i>Sahelian zone</i>					
Cumulative annual rainfall (mm)	645.454	160.875	101.54	899.99	689 339
Length of the rainy season (days)	118.152	19.894	36	279	689 339
Seasonal cumulative rainfall (mm)	598.229	169.225	47.749	1252.4	689 339
Short rainfall season (year t)	0.158	0.365	0	1	689 339
Drought (year t)	0.154	0.361	0	1	689 339
Tree cover loss in ha	0.03	0.491	0	128.13	689 339
Deforestation in ha	0.004	0.305	0	113.14	689 339
Annual probability of deforestation	0.114	0.318	0	1	689 339
<i>Sudanian zone</i>					
Cumulative annual rainfall (mm)	1005.525	55.341	900.01	1099.9	412 632
Length of the rainy season (days)	154.465	31.369	45	286	412 632
Seasonal cumulative rainfall (mm)	904.358	128.948	252.9	1473.5	412 632
Short rainfall season (year t)	0.154	0.361	0	1	412 632
Drought (year t)	0.159	0.366	0	1	412 632
Tree cover loss in ha	0.968	4.339	0	320.71	412 632
Deforestation in ha	0.474	3.552	0	320.37	412 632
Annual probability of deforestation	0.614	0.487	0	1	412 632
<i>Guinean zone</i>					
Cumulative annual rainfall (mm)	1402.256	184.672	1100	1699.9	1 110 791
Length of the rainy season (days)	171.778	44.39	25	326	1 110 791
Seasonal cumulative rainfall (mm)	1143.356	241.401	249.78	2223	1 110 791
Short rainfall season (year t)	0.156	0.363	0	1	1 110 791
Drought (year t)	0.147	0.355	0	1	1 110 791
Tree cover loss in ha	3.874	13.355	0	2096.4	1 110 791
Deforestation in ha	3.474	13.21	0	2096.3	1 110 791
Annual probability of deforestation	0.743	0.437	0	1	1 110 791
<i>Guineo-congolian zone</i>					
Cumulative annual rainfall (mm)	2221.686	424.39	1700.1	4165.3	431 650
Length of the rainy season (days)	168.591	42.837	35	326	431 650
Seasonal cumulative rainfall (mm)	1733.957	487.917	365.62	4226.1	431 650
Short rainfall season (year t)	0.146	0.353	0	1	431 650
Drought (year t)	0.145	0.352	0	1	431 650
Tree cover loss in ha	10.016	23.089	0	935.58	431 650
Deforestation in ha	9.894	23.022	0	935.25	431 650
Annual probability of deforestation	0.846	0.361	0	1	431 650



## B.2 Eco-climatic zones

Summary statistics of the data, a sample limited to pixels with at least one hectare of tree cover in 2000 and 100 mm of average cumulative rainfall and disaggregate descriptive statistics by eco-climatic zones, are shown in Table 3.

## C Statistical analysis

### C.1 Transmission channels: impacts mainly driven by human activities

The objective of the paper is to assess the impact of agricultural income shocks on deforestation, via human-induced activities. One may argue that direct impacts of rainfall shocks on forest stands could hamper the statistical analysis by introducing a bias in the estimated relation between rainfall and deforestation.

Many recent studies show that climate factors may influence forest health (Jiang et al., 2019; Aleixo et al., 2019; Zemp et al., 2017; Anderegg et al., 2013). However, the algorithm behind Hansen et al. (2013)'s data is designed to assess forest stand loss and not canopy disturbance. This algorithm detects only stand-replacement disturbances, or a change from a forest to non-forest state, and only considers 30m side pixels with forest cover in 2000 entirely cleared during the 2001-2019 period. Thanks to this feature of the algorithm, the estimation of bad rainy season impacts provided in this article are not attributable to tree mortality. In addition, because mortality occurs at least two years after the weather event (Aleixo et al., 2019), it is ruled out from our statistical estimations, which only consider bad rainy seasons in years  $t$ .

Another potential concern is the role of forest fires, which may be triggered by a dry or a short season and directly impact tree cover. A large proportion of existing forest fires are triggered and controlled by humans, and this proportion may increase in Africa in the future as result of increased population densities and urbanisation (Archibald, 2016). By only considering rainfall during the rainy season, most of the impacts of naturally triggered forest fires (that usually occur during the dry season) are excluded from consideration. However, slash-and-burn forest clearing methods are used in the study area and these human triggered fires should be considered. Yet, due to a lack of data relating to the causes of forest fires, it is impossible to completely rule out the hypothesis that a bad rain season may directly increase tree cover loss as a result of naturally caused fires.

### C.2 Whole study region

The average impact of a short rainy season on deforestation in the whole study region is positive (9%) and the average impact of a drought, i.e. a lack of seasonal rainfall, on deforestation is negative (8.5%)

### C.3 Results by eco-climatic zones

Table 4: Drivers of tree cover loss, zone

	(1)	(2)
	Log of deforestation (ares)	Log of deforestation (ares)
Short rainfall season (year t)	0.0913*** (0.0208)	0.0165 (0.0291)
Drought (year t)	-0.0854*** (0.0237)	0.0114 (0.0458)
av. annual cum. rainfall $\times$ Short rainfall season (year t)		0.0000631** (0.0000301)
av. annual cum. rainfall $\times$ Drought (year t)		-0.0000803* (0.0000453)
Observations	2687335	2687335
$R^2$	0.786	0.786
Adjusted $R^2$	0.774	0.774

Standard errors in parentheses, robust to clustering at the 2nd administrative level.

\*  $p < .1$ , \*\*  $p < .05$ , \*\*\*  $p < .01$

Table 5: Drivers of deforestation, Sahelian zone

	(1)	(2)	(3)	(4)
Log of deforestation (ares)	All pixels	Crop area > 1%	Crop area > 5%	Crop area > 25%
Short rainfall season (year t)	0.000604 (0.00100)	0.000287 (0.000943)	0.000526 (0.000976)	0.000118 (0.000801)
Drought (year t)	0.00127 (0.00160)	0.000466 (0.00154)	0.000476 (0.00138)	-0.000660 (0.000812)
Observations	695381	651852	618260	514976
$R^2$	0.448	0.429	0.444	0.486
Adjusted $R^2$	0.418	0.398	0.413	0.458

Standard errors in parentheses, robust to clustering at the 2nd administrative level.

\*  $p < .1$ , \*\*  $p < .05$ , \*\*\*  $p < .01$

Table 6: Drivers of deforestation, Sudanian zone

	(1)	(2)	(3)	(4)
Log of deforestation (ares)	All pixels	Crop area > 1%	Crop area > 5%	Crop area > 25%
Short rainfall season (year t)	0.0629** (0.0246)	0.0686*** (0.0239)	0.0598*** (0.0206)	0.0414*** (0.0159)
Drought (year t)	-0.0551** (0.0232)	-0.0330* (0.0193)	-0.0200 (0.0156)	-0.00647 (0.0115)
Observations	420669	322382	289284	226527
$R^2$	0.729	0.767	0.788	0.823
Adjusted $R^2$	0.714	0.754	0.776	0.813

Standard errors in parentheses, robust to clustering at the 2nd administrative level.

\*  $p < .1$ , \*\*  $p < .05$ , \*\*\*  $p < .01$

Table 7: Drivers of deforestation, **Guinean zone**

Log of deforestation (ares)	(1) All pixels	(2) Crop area > 1%	(3) Crop area > 5%	(4) Crop area > 25%
Short rainfall season (year t)	0.146*** (0.0343)	0.0893*** (0.0279)	0.0815*** (0.0295)	0.0939*** (0.0320)
Drought (year t)	-0.0925** (0.0432)	-0.0437 (0.0300)	-0.0252 (0.0277)	-0.0291 (0.0311)
Observations	1124633	587799	476170	315193
$R^2$	0.718	0.840	0.855	0.870
Adjusted $R^2$	0.702	0.831	0.847	0.862

Standard errors in parentheses, robust to clustering at the 2nd administrative level.

\*  $p < .1$ , \*\*  $p < .05$ , \*\*\*  $p < .01$

Table 8: Drivers of deforestation, **Guineo-congolian zone**

Log of deforestation (ares)	(1) All pixels	(2) Crop area > 1%	(3) Crop area > 5%	(4) Crop area > 25%
Short rainfall season (year t)	0.170*** (0.0522)	0.124*** (0.0427)	0.0852** (0.0384)	0.0225 (0.0471)
Drought (year t)	-0.0728 (0.0621)	-0.181*** (0.0660)	-0.265*** (0.0719)	-0.457*** (0.0914)
Observations	446652	269846	213168	123962
$R^2$	0.649	0.700	0.714	0.718
Adjusted $R^2$	0.629	0.683	0.698	0.702

Standard errors in parentheses, robust to clustering at the 2nd administrative level.

\*  $p < .1$ , \*\*  $p < .05$ , \*\*\*  $p < .01$

## C.4 Deforestation and remoteness

Table 9: Drivers of deforestation in Sudanian, Guinean and Guineo-congolian zones, depending on **population density** and crop area

	(1)	(2)	(3)	(4)
	Log of def. (ares)	Log of def. (ares)	Log of def. (ares)	Log of def. (ares)
Remoteness (popul. density)	Low	Low	High	High
Crop area	<25%	≥25%	<25%	≥25%
Short rainfall season (year t)	0.221*** (0.0451)	0.138*** (0.0482)	0.111*** (0.0287)	0.0454** (0.0219)
Drought (year t)	-0.0878 (0.0544)	-0.121* (0.0654)	-0.0920*** (0.0337)	-0.151*** (0.0373)
Observations	867032	62843	427403	597623
$R^2$	0.627	0.897	0.782	0.879
Adjusted $R^2$	0.606	0.891	0.770	0.872

Standard errors in parentheses, robust to clustering at the 2nd administrative level.

\*  $p < .1$ , \*\*  $p < .05$ , \*\*\*  $p < .01$

Low (high) population density is under the median level of log population density (2.6) corresponding to less (more) than 13.5 people by km<sup>2</sup>.

Table 10: Drivers of deforestation in Sudanian, Guinean and Guineo-congolian zones, depending on **time-distance to the nearest market** and crop area

	(1)	(2)	(3)	(4)
	Log of def. (ares)	Log of def. (ares)	Log of def. (ares)	Log of def. (ares)
Remoteness (nearest market)	Near	Near	Far	Far
Crop area	<25%	≥25%	<25%	≥25%
Short rainfall season (year t)	0.210*** (0.0403)	0.0293 (0.0302)	0.0897** (0.0388)	0.0682*** (0.0217)
Drought (year t)	-0.0796 (0.0513)	-0.126** (0.0507)	-0.107** (0.0456)	-0.149*** (0.0376)
Observations	988632	211980	305803	448486
$R^2$	0.658	0.906	0.764	0.866
Adjusted $R^2$	0.638	0.900	0.751	0.858

Standard errors in parentheses, robust to clustering at the 2nd administrative level.

\*  $p < .1$ , \*\*  $p < .05$ , \*\*\*  $p < .01$

Far (near) from the nearest markets correspond to pixels that are at more (less) than 90 minutes of time-distance from the nearest market.

Table 11: Drivers of deforestation in Sudanian, Guinean and Guineo-congolian zones, depending on **proximity to powered settlements** and crop area

	(1)	(2)	(3)	(4)
	Log of def. (ares)	Log of def. (ares)	Log of def. (ares)	Log of def. (ares)
Remoteness (powered settlements)	Near	Near	Far	Far
Crop area	<25%	≥25%	<25%	≥25%
Short rainfall season (year t)	0.205*** (0.0402)	-0.0268 (0.0237)	0.123*** (0.0375)	0.106*** (0.0273)
Drought (year t)	-0.104** (0.0493)	-0.181*** (0.0524)	-0.0260 (0.0400)	-0.111*** (0.0338)
Observations	1056210	305689	238225	354777
$R^2$	0.673	0.905	0.721	0.857
Adjusted $R^2$	0.654	0.900	0.705	0.849

Standard errors in parentheses, robust to clustering at the 2nd administrative level.

\*  $p < .1$ , \*\*  $p < .05$ , \*\*\*  $p < .01$

Far (near) from powered settlements correspond to pixels that are at more (less) than 13.5 km from the nearest powered settlement.

## D Robustness checks

### D.1 Considering USGS bioclimatic regions instead of FAO eco-climatic zones

Using the USGS definition of bioclimatic regions<sup>13</sup> rather than the FAO eco-climatic zones, the results hold, as shown in Figure 10. Under this definition, the Sahelian region, corresponds to pixels with more than 100 mm and less than 600 mm of annual rainfall, the Sudano-Sahelian region lies between 600 and 1200 mm, the Guinean region between 1200 and 1700 mm, the Guineo-Congolian region between 1700 mm and 2200 and the Congolian region over 2200 mm annually. Only 204 257 observations lies in the Congolian bioclimatic region and this may explain that results for that zones are found less significant.

### D.2 Forest definition: thresholds of canopy cover

I consider here different canopy cover, corresponding to different forest definition. Figure 11 shows the impact of a bad rainy season on tree cover loss, Figure 12 shows the impact of a bad rainy season on Forests defined by a 25% canopy cover and Figure 13 the impact of a bad rainy season on Forests defined by a 30% canopy cover.

The opposite effect of a short rainy season (negative impact of rainy season length) and a lack of seasonal rainfall (positive impact of drought) in the dryer eco-climatic (Figure 11) zone must be put into perspective in the light of the scarce presence of trees in this zone (see Table 3).

<sup>13</sup><https://eros.usgs.gov/westafrika/node/147>




Figure 10: Impact of a bad rainy season on deforestation, using the USGS definition of bioclimatic regions.




Figure 11: Impact of a bad rainy season on tree cover loss (no threshold on canopy cover).




Figure 12: Impact of a bad rainy season on deforestation (**threshold: 25%** of canopy cover).




Figure 13: Impact of a bad rainy season on deforestation (**threshold: 30%** of canopy cover).

### D.3 Rainfall shocks severity: SPI thresholds

The results are robust to different severity of rainfall shocks, corresponding to different SPI threshold. I consider lower thresholds for mild and moderate rainfall shocks (.5, .7) and a higher threshold for a severe rainfall shock (1.5), respectively in Figures 14, 15 and 16. The probability of return of a mild rainfall shock is 23.4% for droughts and 23.9% for short rainy seasons (every 4 to 5 years), the probability of return of a mild rainfall shock is 30.1% for droughts and 31.6% for short rainy seasons (every 3 to 4 years) and the probability of return of a severe rainfall shock is 5.1% for droughts and 6.5% for short rainy seasons (every 15 to 20 years).




Figure 14: Impact of a bad rainy season on deforestation (**mild** droughts and short rainy seasons, both defined by SPI inferior to **-.5**).




Figure 15: Impact of a bad rainy season on deforestation (**moderate** droughts and short rainy seasons, both defined by SPI inferior to **-.7**).

Extremely severe droughts (SPI inferior to -2) correspond to events only occurring every fifty year, which seem to small considering the time span of the study period. Similarly, heavy droughts are rare, it may explain why they are found to have a less






Figure 16: Impact of a very bad rainy season on deforestation (**heavy** droughts and very short rainy seasons, both defined by SPI inferior to **-1.5**).

significant impacts on deforestation in spite of a similar or higher amplitude (see Figure 16).

#### D.4 Deforestation and remoteness: Guinean and Guineo-congolian eco-climatic zones

Here, I show the robustness of heterogeneous results depending on the remoteness and the proportion of crop area, when restricting the sample to the two most humid eco-climatic zones.

#### D.5 Standard error adjustment for spatial correlation and serial correlation in panel data

I consider here an alternative estimation method of standard errors that is robust to controlling for spatial correlation and serial correlation in panel data. I employ the recent Stata routine `acreg` developed by Colella et al. (2019) based on Hsiang (2010) and Conley (1999). Details about the statistical regression method are given at the following links:

<http://www.fight-entropy.com/2010/06/standard-error-adjustment-ols-for.html>.

Table 12 shows the standard errors allowing for spatial correlation of 50 kilometres, and for a serial correlation over 7 consecutive years.




Figure 17: Impact of a bad rainy season on deforestation in Guinean and Guineo-congolian eco-climatic zones, depending on the population density (a) and relative remoteness in terms of: time-distance to cities (b) and distance to night-time lights (c).

Table 12: Drivers of deforestation: robust to spatial and serial correlation.

<b>Sahelian zone</b>				
	(1)	(2)	(3)	(4)
Log of deforestation (ares)	All pixels	Crop area > 1%	Crop area > 5%	Crop area > 25%
Short rainfall season (year t)	0.000604 (0.00127)	0.000287 (0.00126)	0.000526 (0.00123)	0.000118 (0.000985)
Drought (year t)	0.00127 (0.00168)	0.000466 (0.00153)	0.000476 (0.00134)	-0.000660 (0.000982)
Observations	695381	651852	618260	514976
<b>Sudanian zone</b>				
Short rainfall season (year t)	0.0629*** (0.0187)	0.0686*** (0.0193)	0.0598*** (0.0175)	0.0414** (0.0172)
Drought (year t)	-0.0551*** (0.0192)	-0.0330* (0.0192)	-0.0200 (0.0177)	-0.00647 (0.0183)
Observations	420669	322382	289284	226527
<b>Guinean zone</b>				
Short rainfall season (year t)	0.146*** (0.0353)	0.0893** (0.0369)	0.0815** (0.0402)	0.0939* (0.0488)
Drought (year t)	-0.0925** (0.0436)	-0.0437 (0.0452)	-0.0252 (0.0489)	-0.0291 (0.0603)
Observations	1124634	587799	476170	315193
<b>Guineo-congolian zone</b>				
Short rainfall season (year t)	0.169*** (0.0626)	0.124* (0.0705)	0.0852 (0.0781)	0.0225 (0.0944)
Drought (year t)	-0.0725 (0.0673)	-0.181** (0.0774)	-0.265*** (0.0869)	-0.457*** (0.109)
Observations	446481	269808	213130	123924

Conley (1999) standard errors in parentheses, allowing for spatial correlation within a 50 km radius and for 7 years serial correlation.

\*  $p < .1$ , \*\*  $p < .05$ , \*\*\*  $p < .01$