



**HAL**  
open science

# Critical analysis of methods to estimate the fraction of absorbed or intercepted photosynthetically active radiation from ground measurements: Application to rice crops

Wenjuan Li, Hongliang Fang, Shanshan Wei, Marie Weiss, Frédéric Baret

## ► To cite this version:

Wenjuan Li, Hongliang Fang, Shanshan Wei, Marie Weiss, Frédéric Baret. Critical analysis of methods to estimate the fraction of absorbed or intercepted photosynthetically active radiation from ground measurements: Application to rice crops. *Agricultural and Forest Meteorology*, 2020, 297, 10.1016/j.agrformet.2020.108273 . hal-03122813

**HAL Id: hal-03122813**

**<https://hal.inrae.fr/hal-03122813>**

Submitted on 15 Dec 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Critical analysis of methods to estimate the fraction of absorbed or intercepted**  
2 **photosynthetically active radiation from ground measurements: application to rice**  
3 **crops**

4  
5 Wenjuan Li<sup>a,b,c\*</sup>, Hongliang Fang<sup>a,b</sup>, Shanshan Wei<sup>a,b</sup>, Marie Weiss<sup>c</sup> and Frédéric Baret<sup>c</sup>

6 <sup>a</sup>LREIS, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy  
7 of Sciences, Beijing 100101, China,

8 <sup>b</sup>College of Resources and Environment, University of Chinese Academy of Sciences, Beijing  
9 100049, China

10 <sup>c</sup>INRAE, Avignon Université, UMR 1114 EMMAH, UMT CAPTE, F-84000, Avignon,  
11 France

12

13

14 \*Corresponding author

15

16

17

18 **Highlights**

19

- 20 • fIPAR is a good proxy of fAPAR when the background is dark as in most rice crops.
- 21 • Impact of illumination conditions and non-green components was analyzed.
- 22 • Green fAPAR can be estimated from canopy fAPAR and the GAI/PAI ratio.
- 23 • Downward DHP is recommended when estimating green fIPAR.

25 **Abstract**

26

27 Continuous and accurate ground measurements of the fraction of absorbed (fAPAR) or  
28 intercepted (fIPAR) photosynthetically active radiation by green canopy components is  
29 important to monitor canopy functioning. fAPAR and fIPAR are sensitive to illumination  
30 conditions and non-green components during the senescence stage. While several methods  
31 have been developed to estimate fAPAR or fIPAR in the field from different methods  
32 including AccuPAR, LAI-2200 and Digital Hemispheric Photograph Photography (DHP), the  
33 differences among these methods still need more investigations. The principles on which they  
34 are based are first reviewed with due attention to the assumptions used and approximations  
35 made. Two field campaigns conducted in 2012 and 2013 in northeastern China over paddy  
36 rice fields were then used to compare fAPAR and fIPAR measured using AccuPAR, DHP and  
37 LAI-2200. Results demonstrated that considering only canopy light transmittance (fIPAR),  
38 measured with AccuPAR, DHP or LAI-2200, is a good proxy of fAPAR which is computed  
39 from AccuPAR measurements of the four fluxes of the radiation balance. However, when  
40 canopy is senescing, downward looking DHP method is recommended since it is the only  
41 method that directly measures the light intercepted by green elements. Methods based on  
42 upward looking (DHP upward, AccuPAR, LAI-2200) cannot distinguish between the green  
43 and senescent vegetation elements. Corrections based on independent measurements of the  
44 ratio of the green area index (GAI) to the plant area index (PAI) (GAI/PAI) need to be used in  
45 this case, while assuming that green and senescent elements are well mixed in the canopy  
46 volume. Downward looking DHP appears to be the preferred method for relatively short and

47 dense canopies such as rice since it does not disturb the canopy, it is sensitive to the green  
48 elements only and allows to simulate fIPAR for any illumination conditions.

49

50 **Keywords**

51 fAPAR; fIPAR; Green fAPAR; Green fIPAR; Paddy rice; Diffuse fraction

52 **Nomenclature**

$I_t^\downarrow$	Incoming downward flux measured at the top of the canopy
$I_t^\uparrow$	Upward flux reflected by the canopy
$I_b^\downarrow$	Downward fluxes measured at the bottom of the canopy
$I_b^\uparrow$	Upward fluxes measured at the bottom of the canopy
$R_c$	Canopy reflectance
$R_c^{bs}$	Black-sky canopy reflectance under direct illumination conditions
$R_c^{ws}$	White-sky canopy reflectance under diffuse illumination conditions
$T$	Canopy transmittance
$T^{bs}$	Black-sky transmittance under direct illumination conditions
$T^{ws}$	White-sky transmittance under diffuse illumination conditions
$R_s$	Soil background reflectance
$R_s^{bs}$	Black-sky soil reflectance under direct illumination conditions
$R_s^{ws}$	White-sky soil reflectance under diffuse illumination conditions
$R_\infty$	Canopy reflectance for very dense foliage
$R_{sen}$	Reflectance of senescent layer
$P$	Canopy gap fraction
$\theta$	Zenith angle
$G(\theta)$	Leaf projection function
$\Omega(\theta)$	Canopy clumping index
DOY	Day of Year
ESU	Elementary Sampling Units
DHP	Digital Hemispherical Photography
PAR	Photosynthetically Active Radiation

$f$	Fraction of diffuse PAR in total PAR
fAPAR	Fraction of Absorbed PAR
fAPAR <sub>T</sub>	fAPAR measured from the two-stream method using transmittance only
fAPAR <sup>bs</sup>	Black-sky fAPAR under direct illumination conditions
fAPAR <sup>ws</sup>	White-sky fAPAR under diffuse illumination conditions
fAPAR <sub>T</sub> <sup>bs</sup>	Black-sky fAPAR measured from the two-stream method
fAPAR <sub>T</sub> <sup>ws</sup>	White-sky fAPAR measured from the two-stream method
fIPAR	Fraction of Intercepted PAR
fIPAR(LAI-2200)	fIPAR measured from LAI-2200
fIPAR <sup>ws</sup> (LAI-2200)	White-sky fIPAR measured from LAI-2200
fIPAR(DHP <sub>up</sub> )	fIPAR measured from the upward DHP
fIPAR <sup>bs</sup> (DHP <sub>up</sub> )	Black-sky fIPAR from the upward DHP
fIPAR <sup>ws</sup> (DHP <sub>up</sub> )	White-sky fIPAR from the upward DHP
GfIPAR(DHP <sub>down</sub> )	fIPAR of green canopy components measured from the downward DHP
GfIPAR <sup>bs</sup>	Black-sky GfIPAR measured from the downward DHP
GfIPAR <sup>ws</sup>	White-sky GfIPAR measured from the downward DHP
GfAPAR	fAPAR of canopy green components
GfAPAR <sub>top</sub>	GfAPAR corrected from canopy fAPAR using Eq. (14)
GfAPAR <sub>mix</sub>	GfAPAR corrected from canopy fAPAR using Eq. (15)
GF	Green Fraction
GAI	Green Area Index
GLAI	Green Leaf Area Index
PAI	Plant Area Index

## 53 **1 Introduction**

54

55 The fraction of photosynthetically active radiation (PAR, 400-700nm) absorbed by green  
56 vegetation elements (fAPAR) is closely linked to canopy functioning processes such as  
57 photosynthesis and transpiration. It also quantifies the incoming radiation available at the soil  
58 level that is mandatory for modeling soil temperature and evaporation. It is thus a key variable  
59 required in many ecosystems and crop functioning models to simulate photosynthesis and  
60 primary production (Goward and Huemmrich, 1992; McCallum et al., 2010; Monteith, 2015).  
61 fAPAR is listed as an essential climate variable (ECV) by the Global Climate Observing  
62 System (GCOS, 2016). It is often approximated by the fraction of intercepted PAR (fIPAR)  
63 because the vegetation pigments present a strong absorption in this spectral domain and the  
64 reflectivities from background are usually small for well-developed canopies (Gower et al.,  
65 1999).

66

67 Several methods have been developed to estimate fAPAR and fIPAR from ground  
68 measurements. Handheld optical devices, such as AccuPAR (Meter Group, Inc., USA),  
69 provide an efficient way to measure fAPAR under different illumination conditions  
70 (Steinberg et al., 2006). AccuPAR measures the downward and upward PAR fluxes at the top  
71 and bottom of the canopy by placing the probes above and below the canopy. Other methods  
72 such as Digital Hemispherical Photography (DHP) measure the gap fraction (upward looking)  
73 or green fraction (downward looking) to derive fIPAR in all directions. Pixel classification of  
74 the RGB images is mainly based on color contrast between leaves and the sky for the upward  
75 looking DHP to get the gap, and between green leaves and non-green elements including the  
76 background to get the green pixels for the downward looking DHP (Baret et al., 1993;  
77 Demarez et al., 2008; Leblanc et al., 2005). However, image segmentation may be affected by

78 the illumination conditions, especially when shadows or specular reflection are observed  
79 (Fang et al., 2014a, 2018; Ye et al., 2015). LAI-2200 (LI-COR Inc., Lincoln, Nebraska, USA)  
80 measures the transmittance in the blue wavelength domain in five zenithal directions from  
81 which fIPAR can be estimated. However LAI-2200 measurements are also sensitive to the  
82 illumination conditions (Asner et al., 1998; Kobayashi et al., 2013; Leblanc and Chen, 2001).  
83 A thorough intercomparison of these instruments is still lacking.

84

85 fAPAR depends on solar zenith angle and illumination conditions, e.g., overcast or clear sky  
86 condition. The instantaneous fAPAR is highly sensitive to variations of the solar zenith angle  
87 and presents diurnal variations under clear sky conditions (Fensholt et al., 2004; Rahman et  
88 al., 2015; Zhao et al., 2018), while it shows a much smaller diurnal variations under cloudy  
89 conditions (Nouvellon, 2000; Thomas et al., 2006). The daily integrated fAPAR, which is a  
90 variable used by many canopy functioning models, has been demonstrated to be smaller under  
91 clear sky as compared to overcast conditions (Gower et al., 1999; Thomas et al., 2006).  
92 Therefore, it is required to compare the fAPAR quantities measured by different instruments  
93 under a range of illumination conditions and solar zenith angles. However, direct comparison  
94 between instruments is not always feasible due to the intrinsic properties of each device. As  
95 an example, the fAPAR measured by AccuPAR accounts for the diffuse fraction, while  
96 devices based on gap fraction measurements (DHP) may account both for the direct sunlight  
97 and the diffuse illumination. To facilitate the comparison between those different instruments,  
98 we used the decomposition proposed by Martonchik et al. (2000): fAPAR is considered as the  
99 sum of a black-sky and a white-sky components, weighted by the PAR diffuse fraction. The  
100 black-sky fAPAR,  $fAPAR^{bs}$ , corresponds to the direct component (collimated beam irradiance  
101 in the sun direction only) while the white-sky fAPAR,  $fAPAR^{ws}$ , corresponds to diffuse  
102 illumination conditions generally assumed perfectly isotropic (GCOS, 2016). Although the



103 impact of diffuse fraction on fAPAR has been investigated (Gu et al., 2002; Jongschaap et al.,  
104 2006; Lizaso et al., 2005), few studies focused on the estimation of the black-sky and white-  
105 sky components of fAPAR or fIPAR in crops (Cohen et al., 1997; Hanan and Bégué, 1995)  
106 and none of them have intercompared the ability of the current instruments to well measure  
107 these quantities.

108

109 Since only the green photosynthetically active elements contribute directly to key processes  
110 such as photosynthesis and transpiration, green vegetation elements should be isolated to  
111 estimate fAPAR (Huemmrich et al., 2005; Pinter, 1993; Weiss et al., 2007; Xiao, 2004; Zhang  
112 et al., 2005). The presence of senescent leaves during late crop growth stages have a  
113 significant impact on fAPAR, and the relationship between fAPAR and vegetation indices (Di  
114 Bella et al., 2004; Rahman et al., 2019; Viña and Gitelson, 2005). The ground measured  
115 canopy fAPAR can be partitioned into fAPAR of green components and non-green  
116 components. Among optical instruments listed above, only downward looking DHPs allow to  
117 separate the green from the non-green elements to estimate the corresponded fraction of  
118 intercepted light. Upward looking DHPs should not be used for such a purpose since  
119 senescence often starts from the bottom layer of the crop, while the light penetrates from the  
120 top of the canopy (Baret et al., 2010). The other upward looking techniques, such as  
121 AccuPAR and LAI-2200, do not allow distinguishing between green and non-green elements.  
122 Some corrections have been proposed to consider only the green elements depending on the  
123 canopy type, either assuming that the green elements are located at the top of the canopy  
124 (Chen, 1996), or assuming that green and non-green elements are well mixed in the canopy  
125 volume (Viña and Gitelson, 2005).

126

127 The objective of this study is to compare the several methods proposed and evaluate the  
128 impact of the presence of non-green vegetation elements during the senescence phase, under  
129 different illumination conditions. For this purpose, a dedicated experiment was conducted in  
130 2012 and 2013 where AccuPAR, DHP and LAI-2200 devices were concurrently used over  
131 paddy rice fields in northeastern China.

## 132 **2 Methods**

### 133 **2.1 Theoretical background**

#### 134 **2.1.1 Derivation of canopy fAPAR and fIPAR**

135

136 fAPAR is calculated from the radiation balance in the PAR domain:

137

$$138 \text{ fAPAR} = \frac{I_t^\downarrow - I_t^\uparrow - (I_b^\downarrow - I_b^\uparrow)}{I_t^\downarrow} \quad (1)$$

139

140 where  $I_t^\downarrow$  and  $I_t^\uparrow$  are the downward and upward fluxes measured at the top of the canopy.  $(I_b^\downarrow -$   
141  $I_b^\uparrow)$  is the radiation absorbed by the soil background calculated as the difference between the  
142 downward ( $I_b^\downarrow$ ) and upward ( $I_b^\uparrow$ ) fluxes measured at the bottom of the canopy. Note that the net  
143 horizontal PAR fluxes are considered negligible as we focus on rice crops which are short  
144 canopies that do not present major heterogeneity at the scale investigated corresponding to  
145 few square meters located in an homogeneous field (Widlowski, 2010). Eq. (1) can be  
146 expressed more simply as:

147

$$148 \text{ fAPAR} = 1 - R_c - T(1 - R_s) \quad (2)$$

149

150 where  $R_c = \frac{I_t^\uparrow}{I_t^\downarrow}$  is the canopy reflectance,  $T = \frac{I_b^\downarrow}{I_t^\downarrow}$  is the canopy transmittance, and  $R_s = \frac{I_b^\uparrow}{I_b^\downarrow}$  is the  
151 soil background reflectance in the PAR spectral domain. For short canopies such as paddy  
152 rice, it is usually difficult to measure the upward flux at the bottom of the canopy because of  
153 the short distance between the sensors and the soil surface and the large spatial heterogeneity  
154 of this flux. However, the soil background reflectance can be estimated from other  
155 independent measurements in the laboratory or over bare soils at nearby locations.

156

157 In the PAR domain, the canopy reflectance can be approximated as a linear decomposition of  
158 soil and foliage reflectance:

159

$$160 \quad R_c \approx TR_s + (1 - T)R_\infty \quad (3)$$

161

162 where  $R_\infty$  is the reflectance for very dense foliage. Combining Eqs. (2) and (3), fAPAR can  
163 be approximated using two terms:

164

$$165 \quad fAPAR \approx (1 - T)(1 - R_\infty) \quad (4)$$

166

167 For dense vegetation,  $R_\infty$  is very small ( $R_\infty \approx 0.04$ ) because of the strong absorption by  
168 chlorophyll pigments in the PAR domain (Weiss et al., 2018). Therefore, Eq. (4) can be  
169 further simplified as:

170

$$171 \quad fAPAR \approx fAPAR_T = 1 - T \quad (5)$$

172

173 The accuracy of this simplification depends on the fluxes reflected by the canopy and the soil  
174 background, which vary with canopy structure, illumination conditions, and background

175 properties (Widlowski, 2010). If leaves are considered opaque, the fraction of intercepted  
176 PAR (fIPAR) can be calculated from the gap fraction  $P$  (Eq. (6)). In these conditions,  $P$  is  
177 closely approximated by canopy transmittance ( $T$ ) and  $fIPAR \approx fAPAR_T$ .

178

$$179 \quad fIPAR = 1 - P \approx 1 - T \approx fAPAR_T \quad (6)$$

180

### 181 **2.1.2 Estimation of the canopy fAPAR and fIPAR under different illumination** 182 **conditions**

183

184 At a given time of the day, the total canopy fAPAR is the sum of the black-sky and white-sky  
185 fAPAR, weighted by the fraction of the incoming diffuse PAR radiation ( $f$ ):

186

$$187 \quad fAPAR = (1 - f) \cdot fAPAR^{bs} + f \cdot fAPAR^{ws} \quad (7)$$

188

189 The same black-sky and white-sky components are also defined for the fIPAR quantities.

190 During a day, if clear-sky  $fAPAR(\theta)$  and white-sky observations,  $fAPAR^{ws}$ , are measured,  
191 instantaneous black-sky fAPAR ( $fAPAR^{bs}$ ) can be estimated based on Eq. (8):

192

$$193 \quad fAPAR^{bs}(\theta) = \frac{fAPAR(\theta) - f(\theta) \cdot fAPAR^{ws}}{1 - f(\theta)} \quad (8)$$

194

195 Similarly, transmittance measured in the five directions by the LAI-2200 allows to compute  
196 the black-sky fIPAR,  $fIPAR^{bs}(\theta)$  for  $\theta < 68^\circ$  by linear interpolation between the five crowns.

197

198 The fraction of intercepted black-sky PAR ( $fIPAR^{bs}(\theta)$ ) was calculated from the green  
 199 fraction (GF) for downward looking DHP or gap fraction (P) for upward looking DHP after  
 200 classifying the green (downward) or sky (upward) pixels:

201

$$202 \begin{cases} FIPAR^{bs}(\theta) = GF(\theta) & \text{for downward DHP} \\ FIPAR^{bs}(\theta) = 1 - P(\theta) & \text{for upward DHP} \end{cases} \quad (9)$$

203

204 For each zenith direction,  $\theta$ , with  $\theta < 60^\circ$ , the green or gap fraction is averaged across all  
 205 azimuthal directions from all images in an ESU to compute  $GF(\theta)$  or  $P(\theta)$  (Weiss and Baret,  
 206 2010). Data for  $\theta > 60^\circ$  were not considered because of the large uncertainties in the green  
 207 fraction estimation due to the degraded resolution for these directions.

208

209 White-sky fIPAR ( $fIPAR^{ws}$ ) for LAI-2200 and DHP devices can be derived by integrating  
 210  $FIPAR^{bs}$  over the hemisphere (Weiss and Baret, 2010):

211

$$212 FIPAR^{ws} = 2 \int_0^{\pi/2} (FIPAR^{bs}(\theta)) \cos \theta \sin \theta d\theta \quad (10)$$

213

214 For  $\theta > 60^\circ$  (DHP) or  $\theta > 68^\circ$  (LAI-2200), the term  $(FIPAR^{bs}(\theta)) \cos \theta \sin \theta$  was  
 215 approximated by linear interpolation between  $\theta = 60^\circ$  or  $\theta = 68^\circ$  and  $\theta = 90^\circ$  with  
 216  $(FIPAR^{bs}(90^\circ)) \cos 90^\circ \sin 90^\circ = 0$ .

217

### 218 **2.1.3 Derivation of the green fAPAR and fIPAR (GfAPAR and GfIPAR)**

219

220 Assuming that all canopy elements are randomly distributed in the canopy volume, the  
 221 canopy transmittance can be derived using the Poisson model (Nilson, 1971):

222

$$223 \quad T = e^{-G(\theta) \cdot PAI \cdot \Omega(\theta) / \cos \theta} \quad (11)$$

224

225 where  $G(\theta)$  is the projection function that depends on the leaf inclination distribution and  
226 direction ( $\theta$ ), and  $\Omega(\theta)$  is the canopy clumping index. It is here assumed that  $G(\theta)$  and  $\Omega(\theta)$   
227 values are the same for the green and non-green elements. The four-stream fAPAR (Eq. (2))  
228 can then be approximated as:

229

$$230 \quad fAPAR = 1 - R_c - (1 - R_s) \cdot e^{-G(\theta) \cdot PAI \cdot \Omega(\theta) / \cos \theta} \quad (12)$$

231

232 When there are no senescent elements, GfAPAR=fAPAR. Conversely, for canopies having  
233 senescent elements, GfAPAR can be estimated from fAPAR measurements using an  
234 independent estimate of GAI/PAI and assumptions about the distribution of the senescent  
235 elements in the canopy. When the green leaves are located at the top of the canopy above the  
236 senescent elements, Chen (1996) proposed to estimate GfAPAR using the following  
237 formulation:

238

$$239 \quad GfAPAR^{top} = 1 - R_c - (1 - R_{sen}) \cdot e^{-G(\theta) \cdot GAI \cdot \Omega(\theta) / \cos \theta} \quad (13)$$

240

241 where  $R_{sen}$  is the reflectance of the senescent layer above the soil background. It plays the  
242 same role as  $R_s$  in Eq. (12) when there is no senescent element. Finally, GAI in Eq. (13) can  
243 be replaced by PAI in Eq. (12) using the GAI/PAI ratio:

244

$$245 \quad GfAPAR^{top} = 1 - R_c - (1 - R_{sen}) \cdot e^{\frac{GAI}{PAI} \ln\left(\frac{1-R_c-fAPAR}{1-R_s}\right)} \quad (14)$$

246

247 Conversely, Viña and Gitelson (2005) assumed that the green and non-green elements are  
 248 well mixed within the canopy volume, proposed the following formulation of the green  
 249 fAPAR as a function of the total canopy fAPAR and the GAI/PAI ratio:

250

$$251 \quad GfAPAR^{mix} = fAPAR \cdot GAI/PAI \quad (15)$$

252

253 Based on the same considerations, Eqs. (14) and (15) can be applied to fIPAR values derived  
 254 from upward DHP and LAI-2200 devices to get the corresponding green fIPAR, GfIPAR:

255

$$256 \quad GfIPAR^{top} = 1 - e^{\frac{GAI}{PAI} \cdot \ln(1-fIPAR)} \quad (16)$$

257

$$258 \quad GfIPAR^{mix} = fIPAR \cdot GAI/PAI \quad (17)$$

259

260 Table 1 lists the fAPAR and fIPAR quantities derived from the several instruments and the  
 261 associated notations and equations used. All these quantities can be computed for both black-  
 262 sky and white-sky conditions.

263

264 **Table 1.** Quantities estimated from AccuPAR, DHP, and LAI-2200.  $R_c$ ,  $R_s$ ,  $R_{sen}$  and  $T$   
 265 represent the canopy reflectance, the background soil and senescent layer reflectance, and the  
 266 canopy transmittance, respectively.  $P$  is the canopy gap fraction and  $GF$  is the green fraction.

267

Instruments	Notation	Equation	Eq. #
AccuPAR	$fAPAR(AccuPAR)$	$1 - R_c - T(1 - R_s)$	(2)
	$fAPAR_T(AccuPAR)$	$1 - T$	(5)

	$GfAPAR^{top}(AccuPAR)$	$1 - R_c - (1 - R_{sen})e^{\frac{GAI}{PAI} \ln(\frac{1-R_c-fAPAR}{1-R_s})}$	(14)
	$GfAPAR^{mix}(AccuPAR)$	$fAPAR \cdot GAI/PAI$	(15)
Downward DHP	$GfIPAR(DHP_{down})$	$GF$	(9)
	$fIPAR(DHP_{up})$	$1 - P$	(9)
Upward DHP	$GfIPAR^{top}(DHP_{up})$	$1 - e^{\frac{GAI}{PAI} \ln(1-fIPAR)}$	(16)
	$GfIPAR^{mix}(DHP_{up})$	$fIPAR \cdot GAI/PAI$	(17)
	$fIPAR(LAI - 2200)$	$1 - P$	(9)
LAI-2200	$GfIPAR^{top}(LAI - 2200)$	$1 - e^{\frac{GAI}{PAI} \ln(1-fIPAR)}$	(16)
	$GfIPAR^{mix}(LAI - 2200)$	$fIPAR \cdot GAI/PAI$	(17)

268

## 269 2.2 Study area

270

271 The study area is located at the Honghe Farm (47.65° N, 133.52° E) in the Heilongjiang  
 272 Province, China. The area is subjected to a humid continental monsoon climate with long and  
 273 cold winter and warm, short, and humid summer. The water and soil are frozen from late  
 274 October to April and thaw in late April. A single rice cultivar (*Longjing 29*) is grown in flat  
 275 fields sharing the same soil properties and where the same cropping practices are applied.  
 276 Rice crops are grown once a year from May to September (Fig. A1). The fields are flooded  
 277 during most of the growing season.

278

279 A total of 55 Elementary Sampling Units (ESUs) of about 20 × 20 m<sup>2</sup> each were selected in  
 280 five fields closely located and chosen to be homogeneous and similar in terms of soils and  
 281 management practices. This allows to consider each ESU as representative of all the other  
 282 ESUs. All ESUs were located at least at 1.5 m from the field border to limit potential edge



283 effects. More details about the site and sampling strategy can be found in Fang et al. (2014a,  
284 2014b).


285

### 286 **2.3 Ground measurements**

287

288 Ground measurements were carried out frequently from June 11 to September 17 in 2012, and  
289 from June 22 to August 29 in 2013 (Fig. 1). We used the “moving ESU strategy” as described  
290 by Fang et al. (2014a), considering that the measurements achieved in one ESU at a given  
291 date are representative of all the other ESUs. This allows to prevent disturbances caused by  
292 the handheld measurements along the growing season and makes destructive measurements  
293 possible. In 2012, all the measurements were taken close to sunset or under overcast  
294 conditions to estimate the white-sky fAPAR. In 2013, white-sky fAPAR was also measured  
295 near sunset or under overcast conditions and completed the same day by black-sky fAPAR  
296 measurements when the sky was clear in the morning (9:30 to 10:30 am). The field  
297 measurement dates and the corresponding solar zenith angles and diffuse fraction are shown  
298 in Fig. 1 for the several instruments considered in this study.

299



300

301 **Fig. 1.** Measurement dates for AccuPAR (blue diamond),  $DHP_{down}$  (green square),  $DHP_{up}$   
 302 (pink asterisk) and LAI-2200 (cyan triangle) in (a) 2012 and (b) 2013 under cloudy (filled  
 303 marker) and clear (open marker, 2013) conditions. Black filled and open downward-pointing  
 304 triangles represent solar zenith angles for cloudy and clear conditions (first right y-axis). Gray  
 305 dashed line with open circles in (b) indicate the diffuse fraction measured for clear sky  
 306 conditions in 2013 (second right y-axis).

307

308 Decagon's AccuPAR LP-80 PAR/LAI Ceptometer measures PAR using 80 individual sensors  
 309 with a  $180^\circ$  field of view on a 1-m probe (Huemmrich et al., 2005; Senna, 2005; Steinberg et  
 310 al., 2006; Thomas et al., 2006). The downward and reflected PAR fluxes at the top of canopy  
 311 were measured by placing the probe approximately 1.5 m above the canopy, facing upward  
 312 and downward, respectively. The canopy transmitted PAR was measured by placing the probe  
 313 below the canopy looking upward. The below-canopy measurements were repeated four times  
 314 in different directions to account for the row effect (Campos et al., 2017; Timlin et al., 2014;  
 315 Zhong et al., 2015). The soil reflected PAR was measured twice in two different rows by

316 placing the probe approximately 5 cm above the ground looking downward. Prior to each  
317 measurement, the AccuPAR was calibrated when the above canopy PAR was  $> 600 \mu\text{mol}/\text{m}^2\text{s}$   
318 as recommended in the user manual (Decagon Devices, 2010). Under clear skies in 2013, the  
319 diffuse PAR was measured by blocking the direct solar illumination with a black board placed  
320 0.5 m from the sensor. The diffuse fraction was then computed as the ratio of the diffuse to  
321 the total downward PAR. The measurement was repeated three times within one minute  
322 before, during, and after fAPAR measurements. Because the three replicates were generally  
323 consistent, their average value was considered as the diffuse fraction at the time of the fAPAR  
324 measurements.

325

326 The DHP images were taken using a Nikon D5100 camera equipped with a 4.5 mm F2.8 EX  
327 DC fisheye convertor. The DHP camera was calibrated before measurements following the  
328 CAN-EYE manual (Weiss and Baret, 2010) to obtain the optical center and the projection  
329 function of the camera and fish-eye system. The total height of the camera, including the lens,  
330 was about 16.5 cm. Two bubble levels were attached to the camera to keep it horizontal for  
331 both downward and upward measurements. In each ESU, 15 to 20 DHPs were acquired for  
332 both downward and upward directions (Fang et al., 2014a). The downward images were taken  
333 by holding the camera 0.8–1.5 m above the canopy. When the rice was higher than 70 cm,  
334 upward images were taken by placing the camera right above the background soil or water in  
335 the row. All DHP images were processed using the CAN-EYE version 6.3.3 software  
336 (<https://www6.paca.inrae.fr/can-eye>). Green pixels were manually separated from senescent  
337 and background pixels for the downward images during the classification step. This step was  
338 performed by the same operator throughout the season.

339

340 LAI-2200 measures the blue radiation in 5 concentric rings centered at 7°, 23°, 38°, 53° and  
341 68°. LAI-2200 measurements were conducted always under diffuse conditions. Each  
342 measurement was repeated twice, with one above and four below canopy readings along  
343 diagonal transects between the rows. For the below canopy readings, the instrument was held  
344 about 5 cm above the background. Throughout the season, a 270° view cap was used to shield  
345 the operator. The four measurements over an ESU were averaged to obtain the mean  
346 transmittance (Fang et al., 2014a, 2014b). All AccuPAR, DHP, and LAI-2200 measurements  
347 were made within a maximum time difference of 10 minutes.

348  
349 In addition to the optical measurements, canopy green area index (GAI) and plant area index  
350 (PAI) were measured in 2012 using a destructive method (Fang et al., 2014a, 2014b). Five  
351 plants were randomly harvested in the ESU and the area of green and non-green leaves, stems  
352 and ears were measured using a LI-3100C Area Meter (LI-COR, Lincoln, NE, USA). Leaf,  
353 stem, and ear area are the sum of the corresponding green and non-green measured areas. The  
354 corresponding area indices were then computed using the plant density to get the area of  
355 elements per unit ground area. GLAI corresponds to the green leaves only, while LAI  
356 includes green and non-green parts. GAI corresponds to the area of all green elements, while  
357 PAI includes the senescent parts as well.

358


## 359 **3 Results**

### 360 **3.1 Dynamics of LAI, GLAI, GAI and PAI**

361

362 During the rice green-up stage from sowing to the end of July (Day of year (DOY) 210), no  
363 senescence is observed: GAI and PAI are equal (Fig. 2). When the senescence starts to

364 progress, some leaves disappear, and both PAI and GAI decrease gradually after DOY 210.  
 365 Once the stems and ears are fully developed around DOY 220, their total area keep about  
 366 constant. However, senescence is also progressing gradually up to almost full senescence at  
 367 maturity, i.e. DOY 265 (Fig. 2). Conversely, senescence of leaves stops on DOY 240: LAI  
 368 and GLAI and PAI keep about constant up to maturity, while GAI still decreases because of  
 369 the senescing stems and ears. The high consistency observed between measurements across  
 370 time demonstrates that the spatial variability among the several ESUs sampled was very small.




371  
 372  
 373 **Fig. 2.** Seasonal variation of plant, leaf, stem and ear area index measured by destructive  
 374 method in 2012. The corresponding area of the green parts are indicated by the dashed lines.  
 375 The black circles represent the actual measurements days.

377 **3.2 fAPAR from AccuPAR**

378  
 379 Results show that for both 2012 and 2013, canopy reflectance ( $R_c$ ) is slightly higher in the  
 380 beginning when the soil background is not fully covered by the vegetation, and at the end of

381 season after the ears and senescent components began to appear. When the canopy is fully  
382 covering the soil,  $R_c$  keeps about to a low and stable value with  $R_c \approx 0.04$  (Fig. 3). Soil  
383 background reflectance ( $R_s$ ) shows little variation during the growing season and is low  
384 because the soil was always wet or covered by water. Canopy transmittance ( $T$ ) decreases  
385 continuously from the beginning of the season until DOY 210 and then increases slightly  
386 during the senescent stage (Fig. 3) since part of the leaves are dead while another part of them  
387 show a decrease in chlorophyll, leading to an increase in leaf reflectance and transmittance in  
388 the PAR domain. Accordingly, canopy fAPAR increases from the beginning of the season up  
389 to DOY 210 and decreases during the senescent stage (Fig. 3a). The influence of the  
390 illumination conditions on the different components can be analyzed in 2013 (Fig. 3b).  
391 Canopy and soil reflectance are little impacted and remain stable. Conversely, the canopy  
392 transmittance depends on the illumination conditions mostly before DOY 210 when the  
393 canopy is not fully covering the soil. The black-sky transmittance is higher than its white-sky  
394 counterpart, and consequently the black-sky fAPAR is smaller than the white-sky fAPAR.  
395 After DOY 210, the difference between black-sky and white-sky values for both  
396 transmittance and fAPAR becomes very small due to the saturation of the canopy  
397 transmittance.

398




399

400 **Fig. 3.** Seasonal variation of canopy reflectance ( $R_c$ ), soil reflectance ( $R_s$ ), canopy  
 401 transmittance ( $T$ ), and fAPAR measured in 2012 (a) and 2013 (b) with AccuPAR. The solid  
 402 and dashed lines represent the black-sky (with superscript <sup>'bs'</sup>) and white-sky (with superscript  
 403 <sup>'ws'</sup>) conditions. The filled circles on lines represent the actual measurement days.

404

405 Our experimental results (Fig. 4) show that  $fAPAR_T(\text{AccuPAR})$  estimated from the two-  
 406 stream assumption (Eq. (5)) agrees very well with the reference four-stream fAPAR,  
 407  $fAPAR(\text{AccuPAR})$  (Eq. (2)) under both black-sky and white-sky conditions ( $R^2 = 0.94\sim 1$ ,  
 408  $RMSE = 0.03\sim 0.08$ ). These two fAPAR quantities differ from less than 0.03 (4%) under  
 409 black-sky conditions, the differences being larger when  $fAPAR(\text{AccuPAR})$  is higher than 0.7  
 410 and under white-sky conditions.

411



412

413 **Fig. 4.** Comparison of four-stream fAPAR(AccuPAR) (Eq. (2)) and the two-stream fAPAR<sub>T</sub>  
 414 (AccuPAR)(Eq. (5)) values derived from AccuPAR measurements in 2012 and 2013 under  
 415 both black (magenta) and white-sky (blue) conditions.

416

### 417 3.3 fAPAR and fIPAR of different instruments


418

419 White-sky fAPAR and fIPAR values rapidly increase until DOY 210 in 2012 and 2013 (Fig.  
 420 5). As expected, fAPAR<sub>T</sub>(AccuPAR), fIPAR(DHP<sub>up</sub>) and fIPAR(LAI-2200) and fAPAR  
 421 (AccuPAR) are very close together during the entire season. Conversely, GfIPAR(DHP<sub>down</sub>) is  
 422 slightly higher than fAPAR(AccuPAR) during the early development stages and is much  
 423 lower than the other quantities during the later stages: White-sky GfIPAR(DHP<sub>down</sub>) decreases  
 424 sharply after DOY 210. In contrast, the other quantities remain stable from DOY 210 to DOY  
 425 250 and slightly decrease after DOY 250. In 2013 where both black-sky and white-sky values  
 426 were measured (Fig. 5b), the black-sky values are substantially smaller than the white-sky



427 counterparts. However black-sky  $GfIPAR(DHP_{down})$  is higher than the white-sky values at the  
 428 end of the season (Fig. 5b).

429



430

431 **Fig. 5.** Seasonal variation of fAPAR and fIPAR in 2012 (a) and 2013 (b). The solid and  
 432 dashed lines represent the black-sky (with superscript 'bs') and white-sky (with superscript 'ws')  
 433 conditions, respectively. In 2012, only white-sky conditions are presented.

434

435 We will focus here on the first growth period (before DOY 210) where senescence is  
 436 marginal (Fig. 2) and  $GAI=PAI$ . As a consequence,  $GfAPAR=fAPAR$  and  $GfIPAR=fIPAR$ .


437 We will therefore use here only the terms fAPAR and fIPAR except for  $GfIPAR(DHP_{down})$   
 438 for which only the green elements are accessible (Table 1). The comparison between fAPAR

439 and fIPAR will be made using fAPAR<sub>T</sub>(AccuPAR) as a reference since we demonstrated  
 440 previously that fAPAR(AccuPAR)≈0.96×fAPAR<sub>T</sub>(AccuPAR) (Fig. 4).

441 GfIPAR(DHP<sub>down</sub>) shows a high agreement with fAPAR<sub>T</sub>(AccuPAR) under white-sky  
 442 conditions (Fig. 6a) ( $R^2 = 0.82$ ) with almost no bias. A strong correlation is also observed  
 443 under black-sky conditions (Fig. 6a) with however a systematic overestimation (Bias=0.13).

444 The correlation between fAPAR<sub>T</sub>(AccuPAR) and the fIPAR(DHP<sub>up</sub>) is weak both for the  
 445 white-sky and black-sky values (Fig. 6b). fIPAR(LAI-2200) shows a high agreement with  
 446 fAPAR<sub>T</sub>(AccuPAR) (Fig. 6c), particularly under white-sky conditions.

447



448

449

450 **Fig. 6.** Comparison between fAPAR<sub>T</sub>(AccuPAR) used as a reference and GfIPAR(DHP<sub>down</sub>),  
 451 fIPAR(DHP<sub>up</sub>) and fIPAR(LAI-2200). Data from the first period (before DOY 210) when no  
 452 senescent elements are present. Black-sky (pink circles) and white-sky illumination  
 453 conditions (blue crosses) are presented.

454


### 455 3.4 GfAPAR and GfIPAR during the senescence stage

456

457 We focus on the period starting after DOY 210 when senescence increases up to the maturity  
 458 stage (Fig. 2). As a consequence, the GAI/PAI ratio decreases regularly with time (Fig. 7).

459 The canopy fAPAR measured by AccuPAR shows small variations due to saturation when  
 460 PAI is generally higher than 4.0. Conversely, the green fIPAR derived from downward  
 461 looking DHP, which can be taken as the best proxy of GfIPAR, decreases swiftly from 0.9 to  
 462 0.3 (Fig. 7). Assuming that green and non-green elements are mixed within the canopy (Viña  
 463 and Gitelson, 2005), GfAPAR measured by AccuPAR (Eq. (15)) and GfIPAR measured by  
 464 LAI-2200 (Eq. (17)) show a temporal profile close to the reference GfIPAR from downward  
 465 looking DHP. Conversely, all green quantities derived with Chen (1996), (Eq. (14)) (e.g.  
 466 assuming that the green elements are distributed at the top of canopy) are systematically  
 467 higher than the reference GfIPAR from downward looking DHP.  $GfIPAR^{mix}$  estimated from  
 468 LAI-2200 and upward looking DHP are similar and higher than  $GfAPAR^{mix}$  derived from  
 469 AccuPAR.

470




471

472 **Fig. 7.** Seasonal variation of fAPAR and fIPAR quantities considered in Table 1 during the  
 473 senescent stage (after DOY 210). GAI/PAI is the ratio of GAI to PAI (right y-axis). All  
 474 measurements were performed under white-sky illuminations in 2012.

475

476 Fig. 8 shows that GfAPAR(AccuPAR), GfIPAR(DHP<sub>up</sub>) and GfIPAR(LAI-2200) are well  
 477 correlated with GfIPAR(DHP<sub>down</sub>) considered as the reference. However, significant biases  
 478 are observed. Under the assumption that the green and non-green elements are mixed in the  
 479 canopy (Viña and Gitelson, 2005, Eq. (15)), GfAPAR<sup>mix</sup>(AccuPAR) is closer to the reference  
 480 GfIPAR(DHP<sub>down</sub>) (Bias = 0.02, Fig. 8a), while the GfIPAR<sup>mix</sup>(DHP<sub>up</sub>) and GfIPAR<sup>mix</sup>(LAI-  
 481 2200) are larger by around 0.1 (Fig. 8b and 8c). Conversely, assuming that the green elements  
 482 are distributed at the top of canopy as proposed by Chen (1996), GfAPAR<sup>top</sup>(AccuPAR),  
 483 GfIPAR<sup>top</sup>(DHP<sub>up</sub>) and GfIPAR<sup>top</sup>(LAI-2200) are systematically higher by 0.13 ~ 0.25 than  
 484 reference GfIPAR(DHP<sub>down</sub>).

485



486

487

488 **Fig. 8.** After DOY 210 (senescence), from left to right: comparison of GfAPAR derived from  
 489 AccuPAR, GfIPAR by upward DHP and LAI-2200 with the GfIPAR from downward DHP  
 490 used as a reference. “top” and “mix” refers to the assumptions used to derive the green  
 491 fAPAR, e.g. the senescence is occurring from the top of the canopy (Chen, 1996) or is  
 492 randomly distributed within the canopy (Viña and Gitelson, 2005).

493

## 494 **4 Discussion**

### 495 **4.1 Four-stream fAPAR versus two-stream fAPAR<sub>T</sub> estimation from AccuPAR**

496

497 AccuPAR is appropriate to measure the fAPAR based on the four-stream approach (Eq. (2)).  
498 However, application of the four-stream assumption to compute fAPAR requires measuring  
499 simultaneously canopy reflectance and transmittance, together with the background  
500 reflectance. Measurement of the background reflectance is difficult since it requires setting  
501 the sensors close to the background which may disturb the canopy and influence the  
502 measurement. Furthermore, the spatial representativeness may also be an issue considering  
503 the high local spatial variability of the radiation field at the bottom of the canopy, due to the  
504 row spacing and canopy cover (Timlin et al., 2014). Conversely, the two-stream assumption  
505 (Eq. (5)) based on the sole measurement of canopy transmittance is appealing to estimate  
506 fAPAR.

507

508 The high consistency between fAPAR(AccuPAR) and fAPAR<sub>T</sub>(AccuPAR) (Fig. 4) is mainly  
509 due to the small values of canopy and soil reflectance (Fig. 3). Furthermore, both terms are  
510 partly counterbalancing each other: in Eq. (2), canopy reflectance ( $R_c$ ) varies between  $R_s$  for  
511 PAI=0 to  $R_\infty$  for very large PAI values. Conversely, the term  $T_{R_s}$  varies between  $R_s$  for  
512 PAI=0 to 0 for large PAI values. These experimental results are consistent with that of other  
513 studies (Gallo and Daughtry, 1986; Gobron et al., 2006; Gower et al., 1999; Kukul and Irmak,  
514 2020). However, as shown by Eq. (4), the measured transmittance includes the contribution  
515 from multiple scattering between the bottom of the canopy and the ground, leading to an  
516 overestimation of the actual transmittance and thus on fAPAR<sub>T</sub> (Eklundh et al., 2011). Closer  
517 inspection of the values shows that fAPAR<sub>T</sub>(AccuPAR) is systematically higher than  
518 fAPAR(AccuPAR), particularly for the well-developed canopies fAPAR<sub>T</sub>(AccuPAR)≈1 when

519  $fAPAR(AccuPAR) \approx 1 - R_{\infty} \approx 0.96$  as expected from Eq. (4) since  $R_{\infty} \approx 0.04$  (Fig. 3). We also  
520 computed the actual transmittance which is smaller than the measured one by -0.78% to -  
521 0.14% under cloudy conditions and -0.41% to -0.01% under clear sky conditions. Similarly,  
522  $fAPAR_T$  computed when considering multiple scattering is slightly larger than the  $fAPAR_T$   
523 we estimated by 0.22% to 3.3% under cloudy conditions and 0.2% to 3.09% under clear  
524 conditions. This small uncertainty is mainly due to low background reflectance of paddy rice.  
525 Nevertheless, higher uncertainties may occur for canopies with brighter backgrounds (Asner  
526 et al., 1998; Gower et al., 1999; Widlowski, 2010).

527

## 528 **4.2 Comparison of fAPAR and fIPAR measured from different instruments during the** 529 **green-up stage**

530

531 The overestimations observed between  $GfIPAR^{bs}(DHP_{down})$  and  $fAPAR^{bs}(AccuPAR)$  under  
532 black-sky conditions are mostly due to the limited spatial sampling when considering only the  
533 sun direction. In case of the black-sky conditions, AccuPAR measurements provide a better  
534 spatial sampling with the 80 sensors set along the 1 m long device. Conversely, for white-sky  
535 conditions,  $GfIPAR^{ws}(DHP_{down})$  results from the integration of the black-sky values over all  
536 the directions (Eq. (10)) which provides to a much larger area sampled.

537

538  $fIPAR(DHP_{up})$  has a weak correlation with  $fAPAR_T(AccuPAR)$ . This is mostly explained by  
539 the limited range of variation of  $fAPAR_T$  points available. DHP measurements looking  
540 upward requires to set the camera at the bottom of the canopy. When the back of the camera is  
541 laying on the ground, the focal point of the lens is at about 16.5 cm above the ground. It is  
542 therefore not possible to use this technique for the early growth stages when the canopy is too  
543 short. This explains why no points are available for the low values of fAPAR or fIPAR (Fig.

544 6b). Further, only part of the vegetation elements are seen by the camera looking upward,  
545 resulting in possible underestimation of  $fIPAR(DHP_{up})$ . In addition, setting the camera on the  
546 ground disturbs canopy architecture and may also bias the spatial sampling since it is not  
547 possible to set the camera at the position of the row. Finally, the area sampled by the camera  
548 looking upward from the bottom of the canopy is lower than in the case of  $fIPAR(DHP_{down})$ :  
549 the distance between the camera and the top of the canopy (upward looking DHP) is shorter  
550 than the distance from the camera to the ground (downward looking DHP). This explains why  
551 significant scattering of data is observed between  $fIPAR(DHP_{up})$  and  $fAPAR_T(AccuPAR)$ . It  
552 is therefore recommended to use a very small camera and to improve the spatial sampling by  
553 taking more images. Nevertheless,  $fIPAR(DHP_{up})$  should be used mostly for relatively high  
554 and sparse canopies such as maize crops to limit both the disturbances when taking the  
555 pictures and the parts not sampled at the bottom of the canopy because of the height of the  
556 lens above the ground.

557

558 The small discrepancies observed between both quantities demonstrate that the spatial  
559 sampling was sufficient for LAI-2200 (8 points per ESU), although more limited than that of  
560 the AccuPAR (4 readings of the 80 PAR sensors set along the 1m probe). Under black-sky  
561 conditions, only three matching pairs were available because the LAI-2200 was only  
562 performed under cloudy conditions and the large sun zenith angles prevent the black-sky  
563  $fIPAR^{bs}(LAI-2200)$  calculations.

564

565 Among the three methods investigated ( $DHP_{down}$ ,  $DHP_{up}$  and LAI2200),  $DHP_{down}$  shows  
566 obvious advantages: it provides a good agreement with  $fAPAR_T$ , while not disturbing canopy  
567 architecture since the camera is placed above the canopy. However, in the case of deriving  
568 black-sky  $fIPAR$  values, more samples should be taken to compensate the small footprint of

569 the camera in the sun direction. Further, great care should be taken when segmenting the  
570 image which is more difficult and uncertain for dense canopies and sunny illumination  
571 conditions (Garrigues et al., 2008). Indeed, more advanced classification method is necessary  
572 to improve the DHP data processing (Duveiller and Defourny, 2010; Jonckheere et al., 2017).

573

#### 574 **4.3 Impacts of illumination conditions on fAPAR and fIPAR estimations**

575

576 fAPAR and fIPAR present diurnal variations due to variations of the solar zenith angle and  
577 the proportion of diffuse PAR in the total downwelling radiation. These variations have a  
578 significant impact on the photosynthetic efficiency and on the canopy light regime (Aikman,  
579 1989; Grant, 1999; Wang et al., 2006). We therefore compared the ability of instruments to  
580 retrieve the black-sky and white-sky fAPAR components. Our results show that instantaneous  
581 fAPAR and fIPAR under white-sky conditions are slightly higher than under black-sky  
582 conditions, which is consistent with previous results based on both model simulation and  
583 ground measurements (Li and Fang, 2015; Nouvellon, 2000; Thomas et al., 2006). The  
584 resulting daily integrated fAPAR can be more or less affected depending on the variation of  
585 the diffuse PAR fraction throughout the day. Therefore, except for AccuPAR, accurate daily  
586 fAPAR estimation requires auxiliary measurements of the PAR diffuse fraction or specific  
587 development such as proposed by Hanan and Bégué (1995) for LAI-2200.

588

#### 589 **4.4 Estimations of green fAPAR and fIPAR during the senescence period**

590

591 During the senescence period, both green and senescent elements contribute to fAPAR at the  
592 canopy level (Asner et al., 1998; Di Bella et al., 2004; Huemmrich et al., 2005; Rahman et al.,  
593 2019). Since only the green components are used for photosynthesis and transpiration, the



594 green fAPAR should be the quantity to be considered. Downward DHP is the only method  
595 that provides a direct estimate of green fIPAR because it minimizes problems due to  
596 senescent elements generally located at the bottom of the canopy (Baret et al., 2010). Green  
597 fIPAR from downward DHP is therefore used as the reference method. Conversely, the green  
598 fAPAR cannot be directly measured by the other methods since the instruments are looking  
599 from the bottom of the canopy and green and non-green components cannot be easily  
600 distinguished. We evaluated two methods to derive green fAPAR or green fIPAR from  
601 canopy fAPAR and fIPAR measured quantities using the GAI/PAI ratio, based on different  
602 assumptions on the spatial distribution of green and non-green elements. In paddy rice crops,  
603 the senescence happens right after the ear appearance, and is observed at leaf tips and at the  
604 bottom of the canopy. The ears, distributed mainly at the top layer and mixed with green  
605 leaves, become yellow and brown, and the senescent leaves at the bottom layer grow upward  
606 and mix with other green stems and leaves. This behavior is thus closer to the random mixing  
607 hypothesis of Viña and Gitelson (2005) than to the Chen (1996) assumption that green  
608 elements are concentrated in the top layer. However, these two correction methods developed  
609 to get the green fAPAR or fIPAR from the canopy fAPAR or fIPAR requires the  
610 measurement of the GAI/PAI ratio during the senescence period.

611

## 612 **5 Conclusion**

613

614 The main objective of this study was to compare several methods and instruments for fAPAR  
615 or fIPAR estimates over paddy rice and investigate the impact of canopy senescence under  
616 different illumination conditions. Results showed that using only canopy transmittance  
617 ( $fAPAR_T(\text{AccuPAR})$ ) measured by AccuPAR provides a good proxy of the four-stream

618 reference fAPAR(AccuPAR). This allows to simplify the AccuPAR measurements over  
619 paddy rice fields while keeping a high degree of accuracy.

620

621 Canopy transmittance can also be measured using DHP looking upward or downward and  
622 LAI-2200, resulting respectively into fIPAR(DHP<sub>up</sub>), GfIPAR(DHP<sub>down</sub>) and  
623 fIPAR(LAI2200). Our results demonstrated that fIPAR(DHP<sub>up</sub>) was leading to uncertainties  
624 mostly because of the dimensions of the camera used, disturbing canopy architecture when  
625 placed at the bottom of the crop and missing also a significant fraction of the vegetation  
626 elements located below the lens of the camera. For these reasons, downward looking DHP  
627 (GfIPAR(DHP<sub>down</sub>)), AccuPAR (fAPAR<sub>T</sub>(AccuPAR)) and LAI-2200 (fIPAR(LAI-2200)) are  
628 better suited for rice crops that are dense and relatively short. However, the spatial sampling  
629 should be adapted to the actual footprint of each instrument. Three AccuPAR, four LAI-2200  
630 or 15 to 20 DHPs seems sufficient to get precise estimates of white-sky fAPAR or fIPAR over  
631 an area of  $\approx 100\text{m}^2$  of homogeneous rice crops. This minimum sampling appears also  
632 sufficient under black-sky conditions, except for DHPs for which the footprint is very small in  
633 the sun direction. To avoid taking more images in order to improve the area sampled, it is  
634 advised to integrate canopy transmittance over all the compass directions as done for LAI-  
635 2200. Nevertheless, the daily integrated green fAPAR and fIPAR are required in many  
636 vegetation functional models (Baret and Guyot, 1991; Gower et al., 1999; Weiss et al., 2007).  
637 The daily integrated fAPAR and fIPAR values can be derived from the DHP images, which  
638 will also result in a much larger area sampled. Note that DHPs appear the best suited method  
639 to estimate daily variation and daily integrated values of fIPAR since a single image taken  
640 during the day allows to derive canopy transmittance for all possible incoming light directions,  
641 assuming that canopy architecture keeps stable during the day. This assumption seems

642 reasonable for rice crops, but not realistic for heliotropic species and species presenting leaf  
643 rolling reaction to water stresses (Baret et al., 2018).

644

645 Downward looking DHPs is the only method that measures directly GfIPAR, the fraction of  
646 incoming light intercepted by the green photosynthetically active parts of the vegetation. This  
647 offers a great advantage over the other instruments when a significant part of the organs are  
648 senescing as observed over rice crops after flowering. AccuPAR and LAI-2200 are measuring  
649 canopy transmittance from the bottom of the canopy and are not able to distinguish between  
650 the green and non-green parts. Corrections are proposed for these instruments, based on  
651 independent measurement of the GAI/PAI ratio. Measuring the GAI/PAI ratio is generally  
652 done by destructive methods, which is laborious, time consuming, and not well suited for crop  
653 monitoring. Further, the corrections need assumptions on the vertical distribution of the  
654 senescing parts. For rice crops, we demonstrated that the method proposed by Viña and  
655 Gitelson (2005), assuming that green and non-green elements are well mixed, provides the  
656 best agreement with GfIPAR(DHP<sub>down</sub>) considered as the reference method.

657

658 Downward looking DHPs appears thus to be the best method to estimate GfIPAR under  
659 relatively short canopies. It is currently used intensively over a number of crops (Camacho et  
660 al., 2013; Li et al., 2015; Weiss et al., 2007). For taller canopies that prevents easy  
661 characterization from the top, fAPAR<sub>T</sub>(AccuPAR), fIPAR(DHP<sub>up</sub>) and fIPAR(LAI-2200)  
662 should be preferred. Exploitation of DHPs requires images with good resolution and acquired  
663 under favorable illumination conditions. As a matter of facts, sunny conditions are not ideal  
664 since the distinction between green and non-green parts (background and senescent elements)  
665 is difficult in the shadows because of the small dynamics of the pixel values as well as in the  
666 specularly reflected areas where colors are lost. Using HDR (High Dynamic Range) features

667 and applying a gamma factor should partly solve the problem. Nevertheless, image  
668 segmentation to identify the green pixels is still not fully automatic which is the main  
669 limitation of the DHP downward looking method as compared to AccuPAR and LAI-2200.  
670 Additional work is therefore required to develop algorithms capable of identifying  
671 automatically the green pixels in the images with a high degree of accuracy.

672

## 673 **6 Acknowledgements**

674

675 This study was supported by the National Natural Science Foundation of China (41171333,  
676 H.F.). We would like to thank the local farmers for allowing us to take experiment on their  
677 fields and the students who helped in the field campaigns in 2012 and 2013. We also would  
678 like to thank Dr. Chongya Jiang and Dr. Tao Sun in the preparation of field measurements.  
679 Anonymous reviewers are thanked for valuable comments.

680

681

682

683

684

685

686

687

688

689

690

691

692

693 **7 Appendix A. Rice field pictures during growing season**

694



695

696 **Fig. A1.** Rice field pictures of Plot B from end of July to September, 2012.

697

698

699

700 **8 References**

- 701 Aikman, D.P., 1989. Potential Increase in Photosynthetic Efficiency from the Redistribution of Solar  
702 Radiation in a Crop. *J Exp Bot* 40, 855–864. <https://doi.org/10.1093/jxb/40.8.855>
- 703 Asner, G.P., Wessman, C.A., Archer, S., 1998. Scale Dependence of Absorption of Photosynthetically  
704 Active Radiation in Terrestrial Ecosystems. *Ecological Applications* 8, 1003–1021.
- 705 Baret, F., Andrieu, B., Steven, M., 1993. Gap frequency and canopy architecture of sugar beet and  
706 wheat crops. *Agricultural and Forest Meteorology* 65, 261–279. [https://doi.org/10.1016/0168-](https://doi.org/10.1016/0168-1923(93)90008-6)  
707 [1923\(93\)90008-6](https://doi.org/10.1016/0168-1923(93)90008-6)
- 708 Baret, F., de Solan, B., Lopez-Lozano, R., Ma, K., Weiss, M., 2010. GAI estimates of row crops from  
709 downward looking digital photos taken perpendicular to rows at 57.5° zenith angle:  
710 Theoretical considerations based on 3D architecture models and application to wheat crops.  
711 *Agricultural and Forest Meteorology* 150, 1393–1401.  
712 <https://doi.org/10.1016/j.agrformet.2010.04.011>
- 713 Baret, F., Guyot, G., 1991. Potentials and limits of vegetation indices for LAI and APAR assessment.  
714 *Remote Sensing of Environment* 35, 161–173. [https://doi.org/10.1016/0034-4257\(91\)90009-U](https://doi.org/10.1016/0034-4257(91)90009-U)
- 715 Baret, F., Madec, S., Irfan, K., Lopez, J., Comar, A., Hemmerlé, M., Dutartre, D., Praud, S., Tixier,  
716 M.H., 2018. Leaf-rolling in maize crops: from leaf scoring to canopy-level measurements for  
717 phenotyping. *Journal of Experimental Botany* 69, 2705–2716.  
718 <https://doi.org/10.1093/jxb/ery071>
- 719 Camacho, F., Cernicharo, J., Lacaze, R., Baret, F., Weiss, M., 2013. GEOV1: LAI, FAPAR essential  
720 climate variables and FCOVER global time series capitalizing over existing products. Part 2:  
721 Validation and intercomparison with reference products. *Remote Sensing of Environment* 137,  
722 310–329. <https://doi.org/10.1016/j.rse.2013.02.030>
- 723 Campos, I., Neale, C.M.U., Calera, A., 2017. Is row orientation a determinant factor for radiation  
724 interception in row vineyards?: Row direction influences light capture in vineyard. *Australian*  
725 *Journal of Grape and Wine Research* 23, 77–86. <https://doi.org/10.1111/ajgw.12246>

726 Chen, J., 1996. Canopy Architecture and Remote Sensing of the Fraction of Photosynthetically Active  
727 Radiation Absorbed by Boreal Conifer Forests. *IEEE Transactions on Geoscience and Remote*  
728 *Sensing* 34, 1353–1368.

729 Cohen, S., Rao, R.S., Cohen, Y., 1997. Canopy transmittance inversion using a line quantum probe for  
730 a row crop. *Agricultural and Forest Meteorology* 86, 225–234.

731 Decagon Devices, 2010. AccuPAR PAR/LAI ceptometer model LP-80 Operator's Manual Version 10  
732 [WWW Document]. URL  
733 [http://www.cen.ulaval.ca/nordicanad/donnees/n\\_45561/v582105/file/supp/Decagon\\_Accupar\\_](http://www.cen.ulaval.ca/nordicanad/donnees/n_45561/v582105/file/supp/Decagon_Accupar_LP80_Web.pdf)  
734 [LP80\\_Web.pdf](http://www.cen.ulaval.ca/nordicanad/donnees/n_45561/v582105/file/supp/Decagon_Accupar_LP80_Web.pdf)

735 Demarez, V., Duthoit, S., Baret, F., Weiss, M., Dedieu, G., 2008. Estimation of leaf area and clumping  
736 indexes of crops with hemispherical photographs. *Agricultural and Forest Meteorology* 148,  
737 644–655. <https://doi.org/10.1016/j.agrformet.2007.11.015>

738 Di Bella, C.M., Paruelo, J.M., Becerra, J.E., Bacour, C., Baret, F., 2004. Effect of senescent leaves on  
739 NDVI-based estimates of *f* APAR: Experimental and modelling evidences. *International*  
740 *Journal of Remote Sensing* 25, 5415–5427. <https://doi.org/10.1080/01431160412331269724>

741 Duveiller, G., Defourny, P., 2010. Batch processing of hemispherical photography using object-based  
742 image analysis to derive canopy biophysical variables. Presented at the GEOBIA 2010-  
743 Geographic Object-Based Image Analysis, Ghent University, Ghent, Belgium, p. 5.

744 Eklundh, L., Jin, H., Schubert, P., Guzinski, R., Heliasz, M., 2011. An Optical Sensor Network for  
745 Vegetation Phenology Monitoring and Satellite Data Calibration. *Sensors* 11, 7678–7709.  
746 <https://doi.org/10.3390/s110807678>

747 Fang, H., Li, W., Wei, S., Jiang, C., 2014a. Seasonal variation of leaf area index (LAI) over paddy rice  
748 fields in NE China: Intercomparison of destructive sampling, LAI-2200, digital hemispherical  
749 photography (DHP), and AccuPAR methods. *Agricultural and Forest Meteorology* 198–199,  
750 126–141. <https://doi.org/10.1016/j.agrformet.2014.08.005>

751 Fang, H., Li, W., Wei, S., Sun, T., Jiang, C., 2014b. Paddy Rice Experiment in the Sanjiang Plain  
752 (PRESP) Field Measurement Report. Institute of Geographic Sciences and Natural Resources  
753 Research, Chinese Academy of Sciences, Beijing, China.

754 Fang, H., Ye, Y., Liu, W., Wei, S., Ma, L., 2018. Continuous estimation of canopy leaf area index  
755 (LAI) and clumping index over broadleaf crop fields: An investigation of the PASTIS-57  
756 instrument and smartphone applications. *Agricultural and Forest Meteorology* 253–254, 48–  
757 61. <https://doi.org/10.1016/j.agrformet.2018.02.003>

758 Fensholt, R., Sandholt, I., Rasmussen, M.S., 2004. Evaluation of MODIS LAI, fAPAR and the relation  
759 between fAPAR and NDVI in a semi-arid environment using in situ measurements. *Remote*  
760 *Sensing of Environment* 91, 490–507. <https://doi.org/10.1016/j.rse.2004.04.009>

761 Gallo, K.P., Daughtry, C.S.T., 1986. Techniques for Measuring Intercepted and Absorbed  
762 Photosynthetically Active Radiation in Corn Canopies <sup>1</sup>. *Agron. J.* 78, 752–756.  
763 <https://doi.org/10.2134/agronj1986.00021962007800040039x>

764 Garrigues, S., Shabanov, N.V., Swanson, K., Morisette, J.T., Baret, F., Myneni, R.B., 2008.  
765 Intercomparison and sensitivity analysis of Leaf Area Index retrievals from LAI-2000,  
766 AccuPAR, and digital hemispherical photography over croplands. *Agricultural and Forest*  
767 *Meteorology* 148, 1193–1209. <https://doi.org/10.1016/j.agrformet.2008.02.014>

768 GCOS, 2016. The Global Observing System for Climate: Implementation Needs (GCOS- 200). World  
769 Meteorological Organization.

770 Gobron, N., Pinty, B., Ausedat, O., Chen, J.M., Cohen, W.B., Fensholt, R., Gond, V., Huemmrich,  
771 K.F., Lavergne, T., Mélin, F., Privette, J.L., Sandholt, I., Taberner, M., Turner, D.P.,  
772 Verstraete, M.M., Widlowski, J.-L., 2006. Evaluation of fraction of absorbed  
773 photosynthetically active radiation products for different canopy radiation transfer regimes:  
774 Methodology and results using Joint Research Center products derived from SeaWiFS against  
775 ground-based estimations. *J. Geophys. Res.* 111, D13110.  
776 <https://doi.org/10.1029/2005JD006511>

777 Goward, S.N., Huemmrich, K.E., 1992. Vegetation Canopy PAR Absorptance and the Normalized  
778 Difference Vegetation Index: An Assessment Using the SAIL Model. *Remote Sensing of*  
779 *Environment* 39, 119–140.



780 Gower, S.T., Kucharik, C.J., Norman, J.M., 1999. Direct and Indirect Estimation of Leaf Area Index,  
781 fAPAR, and Net Primary Production of Terrestrial Ecosystems. *Remote Sensing of*  
782 *Environment* 70, 29–51. [https://doi.org/10.1016/S0034-4257\(99\)00056-5](https://doi.org/10.1016/S0034-4257(99)00056-5)

783 Grant, R.H., 1999. Ultraviolet-B and photosynthetically active radiation environment of inclined leaf  
784 surfaces in a maize canopy and implications for modeling. *Agricultural and Forest*  
785 *Meteorology* 95, 187–201. [https://doi.org/10.1016/S0168-1923\(99\)00023-4](https://doi.org/10.1016/S0168-1923(99)00023-4)

786 Gu, L., Baldocchi, D., Verma, S.B., Black, T.A., Vesala, T., Falge, E.M., Dowty, P.R., 2002.  
787 Advantages of diffuse radiation for terrestrial ecosystem productivity: advantages of diffuse  
788 radiation. *J. Geophys. Res.* 107, ACL 2-1-ACL 2-23. <https://doi.org/10.1029/2001JD001242>

789 Hanan, N.P., Bégué, A., 1995. A method to estimate instantaneous and daily intercepted  
790 photosynthetically active radiation using a hemispherical sensor. *Agricultural and Forest*  
791 *Meteorology* 74, 155–168. [https://doi.org/10.1016/0168-1923\(94\)02196-Q](https://doi.org/10.1016/0168-1923(94)02196-Q)

792 Huemmrich, K.F., Privette, J.L., Mukelabai, M., Myneni, R.B., Knyazikhin, Y., 2005. Time-series  
793 validation of MODIS land biophysical products in a Kalahari woodland, Africa. *International*  
794 *Journal of Remote Sensing* 26, 4381–4398. <https://doi.org/10.1080/01431160500113393>

795 Jonckheere, I.G.C., Macfarlane, C., Walter, J.-M.N., 2017. Image Analysis of Hemispherical  
796 Photographs, Algorithms and Calculations, in: *Hemispherical Photography in Forest Science:*  
797 *Theory, Methods, Applications.* Springer Netherlands, Dordrecht, pp. 115–151.

798 Jongschaap, R.E.E., Dueck, T.A., Marissen, N., Hemming, S., Marcelis, L.F.M., 2006. Simulating  
799 seasonal patterns of increased greenhouse crop production by conversion of direct radiation  
800 into diffuse radiation. *Acta Hort.* 315–322. <https://doi.org/10.17660/ActaHortic.2006.718.36>

801 Kobayashi, H., Ryu, Y., Baldocchi, D.D., Welles, J.M., Norman, J.M., 2013. On the correct estimation  
802 of gap fraction: How to remove scattered radiation in gap fraction measurements? *Agricultural*  
803 *and Forest Meteorology* 174–175, 170–183. <https://doi.org/10.1016/j.agrformet.2013.02.013>

804 Kukal, M.S., Irmak, S., 2020. Light interactions, use and efficiency in row crop canopies under  
805 optimal growth conditions. *Agricultural and Forest Meteorology* 284, 107887.  
806 <https://doi.org/10.1016/j.agrformet.2019.107887>

807 Leblanc, S.G., Chen, J.M., 2001. A practical scheme for correcting multiple scattering effects on  
808 optical LAI measurements. *Agricultural and Forest Meteorology* 15.

809 Leblanc, S.G., Chen, J.M., Fernandes, R., Deering, D.W., Conley, A., 2005. Methodology comparison  
810 for canopy structure parameters extraction from digital hemispherical photography in boreal  
811 forests. *Agricultural and Forest Meteorology* 129, 187–207.  
812 <https://doi.org/10.1016/j.agrformet.2004.09.006>

813 Li, W., Fang, H., 2015. Estimation of direct, diffuse, and total FPARs from Landsat surface  
814 reflectance data and ground-based estimates over six FLUXNET sites: Landsat direct, diffuse  
815 and total FPARs. *J. Geophys. Res. Biogeosci.* 120, 96–112.  
816 <https://doi.org/10.1002/2014JG002754>

817 Li, W., Weiss, M., Waldner, F., Defourny, P., Demarez, V., Morin, D., Hagolle, O., Baret, F., 2015. A  
818 generic algorithm to estimate LAI, FAPAR and FCOVER variables from SPOT4\_HRVIR and  
819 Landsat sensors: Evaluation of the consistency and comparison with ground measurements.  
820 *Remote Sensing* 7, 15494–15516.

821 Lizaso, J.I., Batchelor, W.D., Boote, K.J., Westgate, M.E., Rochette, P., Moreno-Sotomayor, A., 2005.  
822 Evaluating a Leaf-Level Canopy Assimilation Model Linked to CERES-Maize. *Agron. J.* 97,  
823 734–740. <https://doi.org/10.2134/agronj2004.0172>

824 Martonchik, J.V., Bruegge, C.J., Strahler, A.H., 2000. A review of reflectance nomenclature used in  
825 remote sensing. *Remote Sensing Reviews* 19, 9–20.  
826 <https://doi.org/10.1080/02757250009532407>

827 McCallum, I., Wagner, W., Schmullius, C., Shvidenko, A., Obersteiner, M., Fritz, S., Nilsson, S.,  
828 2010. Comparison of four global FAPAR datasets over Northern Eurasia for the year 2000.  
829 *Remote Sensing of Environment* 114, 941–949. <https://doi.org/10.1016/j.rse.2009.12.009>

830 Monteith, J.L., 2015. Light Interception and Radiative Exchange in Crop Stands, in: Eastin, J.D.,  
831 Haskins, F.A., Sullivan, C.Y., van Bavel, C.H.M. (Eds.), *Physiological Aspects of Crop Yield*.  
832 American Society of Agronomy, Crop Science Society of America, Madison, WI, USA, pp.  
833 89–111. <https://doi.org/10.2135/1969.physiologicalaspects.c9>

834 Nilson, T., 1971. A theoretical analysis of the frequency of gaps in plant stands. *Agricultural*  
835 *Meteorology* 8, 25–38. [https://doi.org/10.1016/0002-1571\(71\)90092-6](https://doi.org/10.1016/0002-1571(71)90092-6)

836 Nouvellon, Y., 2000. PAR extinction in shortgrass ecosystems: effects of clumping, sky conditions  
837 and soil albedo. *Agricultural and Forest Meteorology* 105, 21–41.  
838 [https://doi.org/10.1016/S0168-1923\(00\)00194-5](https://doi.org/10.1016/S0168-1923(00)00194-5)

839 Pinter, P.J., 1993. Solar angle independence in the relationship between absorbed PAR and remotely  
840 sensed data for alfalfa. *Remote Sensing of Environment* 46, 19–25.  
841 [https://doi.org/10.1016/0034-4257\(93\)90029-W](https://doi.org/10.1016/0034-4257(93)90029-W)

842 Rahman, M.M., Lamb, D.W., Samborski, S.M., 2019. Reducing the influence of solar illumination  
843 angle when using active optical sensor derived NDVIAOS to infer fAPAR for spring wheat  
844 (*Triticum aestivum* L.). *Computers and Electronics in Agriculture* 156, 1–9.  
845 <https://doi.org/10.1016/j.compag.2018.11.007>

846 Rahman, M.M., Lamb, D.W., Stanley, J.N., 2015. The impact of solar illumination angle when using  
847 active optical sensing of NDVI to infer fAPAR in a pasture canopy. *Agricultural and Forest*  
848 *Meteorology* 202, 39–43. <https://doi.org/10.1016/j.agrformet.2014.12.001>

849 Senna, M.C.A., 2005. Fraction of photosynthetically active radiation absorbed by Amazon tropical  
850 forest: A comparison of field measurements, modeling, and remote sensing. *J. Geophys. Res.*  
851 110, G01008. <https://doi.org/10.1029/2004JG000005>

852 Steinberg, D.C., Goetz, S.J., Hyer, E.J., 2006. Validation of MODIS F/sub PAR/ products in boreal  
853 forests of Alaska. *IEEE Trans. Geosci. Remote Sensing* 44, 1818–1828.  
854 <https://doi.org/10.1109/TGRS.2005.862266>

855 Thomas, V., Finch, D.A., McCaughey, J.H., Noland, T., Rich, L., Treitz, P., 2006. Spatial modelling  
856 of the fraction of photosynthetically active radiation absorbed by a boreal mixedwood forest  
857 using a lidar–hyperspectral approach. *Agricultural and Forest Meteorology* 140, 287–307.  
858 <https://doi.org/10.1016/j.agrformet.2006.04.008>

859 Timlin, D.J., Fleisher, D.H., Kemanian, A.R., Reddy, V.R., 2014. Plant Density and Leaf Area Index  
860 Effects on the Distribution of Light Transmittance to the Soil Surface in Maize. *Agronomy*  
861 *Journal* 106, 1828–1837. <https://doi.org/10.2134/agronj14.0160>

862 Viña, A., Gitelson, A.A., 2005. New developments in the remote estimation of the fraction of  
863 absorbed photosynthetically active radiation in crops: REMOTE ESTIMATION OF FAPAR  
864 IN CROPS. *Geophys. Res. Lett.* 32. <https://doi.org/10.1029/2005GL023647>

865 Wang, Xiping, Guo, Y., Li, B., Wang, Xiyong, Ma, Y., 2006. Evaluating a three dimensional model of  
866 diffuse photosynthetically active radiation in maize canopies. *Int J Biometeorol* 50, 349–357.  
867 <https://doi.org/10.1007/s00484-006-0032-0>

868 Weiss, M., Baret, F., 2010. CAN-EYE V6.1 user manual [WWW Document]. URL  
869 <https://www6.paca.inrae.fr/can-eye>

870 Weiss, M., Baret, F., Garrigues, S., Lacaze, R., 2007. LAI and fAPAR CYCLOPES global products  
871 derived from VEGETATION. Part 2: validation and comparison with MODIS collection 4  
872 products. *Remote Sensing of Environment* 110, 317–331.  
873 <https://doi.org/10.1016/j.rse.2007.03.001>

874 Weiss, M., Baret, F., Li, W., Fang, H., 2018. Differences in FAPAR definitions for the validation of  
875 satellite products against ground measurements. Presented at the Workshop on Land Product  
876 Validation and Evolution.

877 Widlowski, J.-L., 2010. On the bias of instantaneous FAPAR estimates in open-canopy forests.  
878 *Agricultural and Forest Meteorology* 150, 1501–1522.  
879 <https://doi.org/10.1016/j.agrformet.2010.07.011>

880 Xiao, X., 2004. Modeling gross primary production of temperate deciduous broadleaf forest using  
881 satellite images and climate data. *Remote Sensing of Environment* 91, 256–270.  
882 <https://doi.org/10.1016/j.rse.2004.03.010>

883 Ye, M., Cao, Z., Yu, Z., Bai, X., 2015. Crop feature extraction from images with probabilistic  
884 superpixel Markov random field. *Computers and Electronics in Agriculture* 114, 247–260.  
885 <https://doi.org/10.1016/j.compag.2015.04.010>

886 Zhang, Q., Xiao, X., Braswell, B., Linder, E., Baret, F., Mooreiii, B., 2005. Estimating light absorption  
887 by chlorophyll, leaf and canopy in a deciduous broadleaf forest using MODIS data and a  
888 radiative transfer model. *Remote Sensing of Environment* 99, 357–371.  
889 <https://doi.org/10.1016/j.rse.2005.09.009>

890 Zhao, L., Liu, Z., Xu, S., He, X., Ni, Z., Zhao, H., Ren, S., 2018. Retrieving the Diurnal FPAR of a  
891 Maize Canopy from the Jointing Stage to the Tasseling Stage with Vegetation Indices under  
892 Different Water Stresses and Light Conditions. *Sensors* 18, 3965.  
893 <https://doi.org/10.3390/s18113965>

894 Zhong, W.W., Liu, J.Q., Zhou, X.B., Chen, Y.H., Bi, J.J., 2015. Row spacing and irrigation effect on  
895 radiation use efficiency of winter wheat. *J. Anim. Plant Sci.* 25, 448–455.

896