

The repertoire of vertebrate STAT transcription factors: Origin and variations in fish

Pierre Boudinot, Steve Bird, Louis Du Pasquier, Bertrand Collet

▶ **To cite this version:**

Pierre Boudinot, Steve Bird, Louis Du Pasquier, Bertrand Collet. The repertoire of vertebrate STAT transcription factors: Origin and variations in fish. Developmental & Comparative Immunology, 2021, 116, pp.103929. 10.1016/J.DCI.2020.103929 . hal-03129753

HAL Id: hal-03129753
https://hal.inrae.fr/hal-03129753

Submitted on 15 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Version of Record: https://www.sciencedirect.com/science/article/pii/S0145305X20304845
Manuscript_a732fde7198a0371c89fc0166eeff8c0

The repertoire of vertebrate STAT transcription factors: origin and variations in fish

Pierre Boudinot[1], Steve Bird[2], Louis Du Pasquier[3] and Bertrand Collet[1]

[1] Université Paris-Saclay, INRAE, UVSQ, VIM, 78350, Jouy-en-Josas, France.

[2] Biomedical Unit, School of Science, University of Waikato, Hamilton 3240, New Zealand

[3] Zoology and Evolutionary Biology, University of Basel, 4051 Basel, Switzerland

Correspondence:

Pierre Boudinot phone: 00331 34652585; email: Pierre.Boudinot@inrae.fr
Bertrand Collet phone: 00331 34652637; email: Bertrand.Collet@inrae.fr
Université Paris-Saclay, INRAE, UVSQ, VIM, 78350, Jouy-en-Josas, France

Abbreviations:
CCD: coiled coil domain; CBP: CREB-binding protein**;** GAS: Gamma interferon activation site; ISRE: interferon-sensitive responsive element; SH2: Src homology 2 domain; STAT: signal transducer and activator of transcription; TAD: C terminal transactivation domain; TAZ: Transcription Adaptor putative Zinc finger; WGD: whole genome duplication;

Keywords: STAT, Interferon signalling, gene duplication, comparative immunology, evolution, vertebrates

© 2020 published by Elsevier. This manuscript is made available under the CC BY NC user license
https://creativecommons.org/licenses/by-nc/4.0/

Abstract

The *stat* gene family diversified during early vertebrate evolution thanks to two rounds of whole genome duplication (WGD) to produce a typical repertoire composed of 6 STAT factors (named 1-6). In contrast, only one or two *stat* genes have been reported in *C. elegans* and in *D. melanogaster*. The main types of STAT found from bony fish to mammals are present in Agnathan genomes, but a typical STAT1-6 repertoire is only observed in jawed vertebrates. Comparative syntenies showed that STAT6 was the closest to the ancestor of the family. An extensive survey of *stat* genes across fish including polyploid species showed that whole genome duplications did not lead to a uniform expansion of *stat* genes. While 2 to 5 *stat*1 are present in salmonids, whose genome duplicated about 35My ago, only one copy of *stat*2 and *stat*6 is retained. In contrast, common carp, with a recent whole genome duplication (5-10My), possesses a doubled *stat* repertoire indicating that the elimination of *stat*2 and *stat*6 additional copies is not immediate. Altogether our data shed light on the multiplicity of evolutionary pathways followed by key components of the canonical cytokine receptor signalling pathway, and point to differential selective constraints exerted on these factors.

1. Introduction

Animals have evolved a number of efficient strategies to combat a large diversity of pathogens. In mammals, complex immune mechanisms are orchestrated and regulated by a network of cytokines acting through cognate ligand/receptor on multiple specialised immunocytes [1]. In mammals, a large number of these cytokines signal through the JAK/STAT signalling factors [2] composed of a particular combination of four Janus kinases (JAK1-3, Tyrosine Kinase TYK2) and one of 7 Signal Transducer and Activator of Transcription

(STAT1-4, 5A, 5B, 6) [3]. The pathway involves a cascade of phosphorylation reactions [4], multimeric complex formation and nuclear translocation [5] resulting in the induction of a particular set of genes responsible for a specific cellular response [6]. Gamma interferon activation site (GAS) is the core genomic motif targeted by STAT1 homodimers [7-9]. STATs heterodimers, associated with additional transcription factors, can bind variants of GAS motifs such as interferon-sensitive responsive element (ISRE) resulting in transcriptional regulation of large gene sets. Such gene sets leading to particular immune responses were associated to different STAT-dependant signalling. In addition, variations in the epigenetic status of genomic elements and in the type of immune cells involved explain, at least in part, the "specificity paradox" of the JAK/STAT signalling pathway, namely, how a 7-member protein family can ensure the specificity of response of dozens of cytokines [10]. The human STAT repertoire is composed of 7 transcription factors encoded by genes located on 3 chromosomes: *STAT1* and *STAT4* closely linked on chromosome 2, *STAT2* and *STAT6* on chromosome 12, and *STAT3*, *STAT5A* and *STAT5B* closely linked on chromosome 17.

All these proteins share four domains: a N-terminal Protein interaction domain ("STAT-i"), a coiled coil domain ("STAT a", CCD), a DNA-binding domain ("STAT b", DBD) and a Src homology 2 domain ("STAT-SH2"). Additionally, STAT1 and STAT2 comprise a C terminal transactivation (TAD) domain: the STAT1 transactivation domain (IPR022752) binds selectively to the Transcription Adaptor putative Zinc finger (TAZ)2 domain of C CREB-binding protein (CBP)/p300, while the STAT2 transactivation domain (IPR022756) binds to the TAZ1 domain of this protein [16-18] (Figure 1A). This domain confers to STAT1 and STAT2 an additional capacity to regulate gene expression since CBP and P300 are histone acetyltransferases that control acetylation of histones in nucleosomes, thus regulating chromatin remodelling and gene transcription.

The vertebrate *stat* repertoire emerged from an ancestral sequence present in the common ancestor of protostomes and deuterostomes with all STAT typical domains [11], through WGD, tandem duplication and dispersion [12]. Liongue et al. [13], proposed that vertebrate *stat* genes originated from a set of two paralogs produced by local duplication, subsequently duplicated "en bloc" by the two rounds of WGD that occurred during early vertebrate evolution, leading to four copies of this cluster. Three of these copies (*STAT3-STAT5, STAT2-STAT6,* and *STAT1-STAT4*) have been retained in human and most vertebrates. In zebrafish, additional copies of *stat1* and *stat5* were found, likely due to the additional, teleost-specific WGD [13].

In this work, we revisited the origins and the evolutive dynamic of the vertebrate *stat* gene repertoire. To find out whether duplicated *stat* copies were retained or lost, we focused on groups and species in which additional WGD occurred. We thus focused on ray finned fish because their genomes were subjected to several WGD events including a teleost-specific WGD event ("3R") that occurred at the root of this lineage about 350 million years ago (Myr) and more recent events for example in salmonids 50-60 Myr ago [14-15] and carps 5-10 Myr ago. In addition, salmonid fish such as Atlantic salmon and Rainbow trout are the most relevant species for the fish farming industry in Europe and worldwide, and their genomes are among the best characterized in teleost fish. We also characterized *stat* genes from Chondrichthyans, Agnathans and non-vertebrate deuterostomians to clarify how these transcription factors evolved during the emergence of vertebrates.

2. Results and discussion

111 *The repertoire of stat genes is well conserved across tetrapods*

112 A fundamental repertoire of six *stat* genes is well-conserved across all tetrapod
113 classes and in the coelacanth, as illustrated in Figure 1B (see also Table S1). One-to-one
114 orthology relationships between tetrapod and coelacanth genes are also supported by
115 conserved synteny groups comprising several markers flanking all *stat* gene clusters (as
116 shown for *stat2* in Figure 1C).

117

118 *Loss and retention of stat genes after WGD during fish evolution reveal contrasted*
119 *constraints on different stat subtypes.*

120 In ray-finned fishes, the *stat* repertoire comprises the same types as in the
121 coelacanth and tetrapods, with *stat1, stat2, stat3, stat4, stat5* and *stat6* present in all
122 species across teleosts. After a WGD occurred some 350 Myr during the early evolution of
123 this group, two copies of each *stat* gene should have been generated [19-21]. This *stat*
124 repertoire has been reshaped by further duplication and gene loss.

125 In fish groups that did not undergo additional WGD, such as herring (*Clupea*
126 *harengus*), pike (*Esox lucius*), zebrafish (*Danio rerio*), stickleback (*Gasterosteus aculeatus*)
127 and the marine species fugu (*Fugu rubripes*) and sea bream (*Sparus aurata)* (Figure 2 and
128 Table S1), *stat*2-6 could be found as single copy in contrast with two or more, *stat1*
129 paralogs. One *stat1* paralog (named "b") is always linked to *stat4* as observed across
130 tetrapods, while the other copy (named "a") is located on another chromosome. This was
131 also the case of Atlantic cod (*Gadus morhua)*, a gadiform species with a particular immune
132 system lacking CD4 and a functional MHC class II pathway. In some cases, an additional
133 s*tat1* can be found like in herring on a third chromosome (Figure 2). In zebrafish, a *stat1*
134 pseudogene has been described close to *stat1b*, [13] but is not present in the last genome

assembly. Only one copy of *stat3, 4* and *5* was generally present in these species with a few exceptions as a double *stat5* in zebrafish, produced by a local duplication. In contrast, the retention of two functional *stat1* genes across multiple families of ray-finned fish suggests that different types of selection pressures may affect this gene, compared to other stat family members.

Multiple stat1 paralogs are also retained in tetraploid Salmonids

To further test this hypothesis, we then focused on tetraploid species in which larger *stat* repertoires have been produced by an additional WGD, providing the opportunity to test their evolutionary fate.

We first performed a comprehensive survey of *stat* genes in salmonids, a fish family tetraploidized by an additional WGD that occurred about 50-60 Myr ago. In these species, we typically found two blocks *stat3+5*, two blocks *stat4+1,* four or five copies of *stat1,* but only one *stat2* and one *stat6* gene (as for rainbow trout in Figure 2). A comprehensive characterization of *stat* genes across salmonids is presented in table 1. Among the two genera *Oncorhynchus* and *Salmo,* we analysed six species for which high quality genomes were available: Sockeye salmon *O. nerka*, rainbow trout *O. mykiss*, chinook salmon *O. tshawytscha*, Coho salmon *O. kisutch*, brown trout *S. trutta* and Atlantic salmon *S. salar*. A total of 16 *stat* loci were found in these six salmonid genomes (Table 1, Figure 2). They were located on 9 chromosomes corresponding to 6 chromosomes in zebrafish (Figure 3), a diploid cyprinid. Linkage analyses showed that *stat1a1-3, stat1b-4* and *stat3-5b* duplicated blocks generated by the salmonid-specific WGD were retained (Figure 3), while there was no evidence of multiple copies of *stat2* and *stat6* (not even pseudogenes).

158 Phylogenetic and synteny block analyses across species provided consistent insights
159 into the origin of these *stat* genes (Figure 3 and Figure S1) and allowed unambiguous
160 identification and annotation. For example, all *stat1a* were linked to *ccr4not* and *ftcd* – as
161 zebrafish *stat1a* – while *stat1b* genes were associated to *stat4* and *slc40* genes. A number of
162 sequences encoding ORFs with size lower than 50% of the average size of STATs proteins
163 were additionally found in the rainbow trout, brown trout and Atlantic salmon (Table 2).
164 These, which likely are assembly artefacts or pseudogenes, were not included in the
165 phylogenetic analysis.

166 In salmonids, an additional (fifth) *stat1* gene that we named *stat1a3* was found
167 immediately downstream of *stat1a2,* suggesting it was generated by local duplication.
168 Interestingly, in the rainbow trout, brown trout and Coho salmon, the STAT1A3 protein is
169 twice the size of the normal size of the STAT1. These long STAT1 proteins contain twice the
170 typical set of domains in tandem [STATi- STATa- STATb- STATSH2-CTD- STATi- STATa- STATb-
171 STATSH2-CTD] and seem to be due to a local duplication-fusion of two *stat1* ORFs. The
172 double *stat1a3* was confirmed in the rainbow trout by the EST CA361350 covering the
173 junction area between the end of the putative first *stat1* and the beginning of the second,
174 which excludes that *stat1a3* has been produced by an assembly error. Further functional
175 studies are required to determine the function of the encoded protein, its potential
176 intramolecular dimerization and GAS elements binding abilities.

177 While the ancient WGD that in early teleost fish has left two *stat1* but only one of
178 the other *stat* paralogs in diploid species, the more recent salmonid-specific WGD resulted
179 in five *stat1* being retained. In contrast, only one copy of *stat2* and *stat6* were kept, either
180 due to an early complete loss post-WGD or because of consistent selection pressures in
181 favour of a single copy.

182

183 *Up-regulation of salmonid stat genes during antiviral responses*

184 Figure 4 shows the expression profile of all chinook salmon *stat* genes from an
185 RNAseq experiment carried out on the EC cell line [22]. We checked whether the salmonid
186 *stat1* and *stat2* genes were induced by type I IFN in a manner consistent with zebrafish
187 where, in zebrafish larva, recombinant IFNϕ1 induces a robust up-regulation of *stat1b* and
188 *stat2,* but not of *stat1a* [23]. In the chinook salmon cell line EC [22] *stat1b1* and *stat2* were
189 induced with a FC>1.5 following stimulation by salmonid recombinant type I IFN. *Stat1a1*
190 was also induced to some extent (Figure 4). However, *stat1b2* was not up-regulated. At
191 steady state, s*tat1a* paralogs were more expressed than *stat1b,* as in zebrafish, a pattern
192 consistent with a functional constitutive expression of *stat1a* genes [22].

193 Thus, there is no strict conservation of the *stat1/2* genes inducibility between
194 salmonids and zebrafish, although the most upregulated genes are *stat1b* and *stat2* in both
195 species. Overall, the paralogs of a given genes may be expressed at low levels in healthy
196 cells, but can reach much higher levels after stimulation, offering opportunities for complex
197 regulations. Whether this profile is different in other cells or under different stimulation
198 conditions remains to be clarified. Similar variations of steady state expression levels were
199 also observed for *stat5: stat5.1* and *stat5.2* were detected at low levels, while *stat5.3*
200 transcripts were at least 10 times more abundant (Figure 4).

201 Duplicated genes in polyploid species are expected to be eliminated by
202 deletion/accumulation of mutations, if they do not acquire new functions (neo/sub-
203 functionalization) or are not kept by selection for gene dosage [24]. Our data about
204 zebrafish and salmonid multiple *stat1* paralogs strongly suggest that they were indeed

subjected to neofunctionalization. More functional work will be necessary to establish if this is also true for salmonid *stat3, 4* and *5* paralogs.

Classification and nomenclature of stat genes in tetraploid species based on the example of salmonids.

The survey of the *stat* gene cluster in salmonid fish highlighted a nomenclature issue for *stat* genes in polyploid species. The current annotation of such complex duplicated genomes is often misleading because of assembly errors. Some annotations inherited the nomenclature used at the time of the first and often single gene discovery by homology cloning and lack consistency with annotation in other fish species. Regarding salmonid *stat* genes, the rainbow trout *stat1a1* and *stat1a2* were annotated *stat1-1* and *stat1*-2 with no reference to the *stat1a* group defined previously in non-salmonid teleost such as zebrafish. The *stat1a3* was left annotated as "uncharacterized protein" whereas phylogeny and blast against the mammalian protein database allocated it to the *stat1* group. Based on our results from phylogeny and synteny conservation, we therefore established a coherent nomenclature (Figure 3, table S1). A similar approach may be followed in other groups of tetraploid vertebrates for example in Amphibians.

Other tetraploid genomes tell more about stat evolutionary dynamics.

We also studied the *stat* genes from the common carp (*Cyprinus carpio,* Ensembl 100: German_Mirror_carp_1.0), an allotetraploid teleost due to a recent WGD that occurred relatively recently 5-10 Myr ago. In this species, all duplicated loci were retained, with exactly twice as many genes as in the diploid cyprinid zebrafish with 4 *stat1*, 2 *stat2*, 2 *stat3*, 2 *stat4*, 4 *stat5* and 2 *stat6* (Table S1). The phylogenetic tree and the distribution of these

genes in contigs indicate that they correspond to a duplication of the blocks typically found in zebrafish and other diploid teleosts (Table S1, Figure 4A).

Polyploid species also originate by allopolyploidization, *i.e.* by genome association due to hybridization among different species. The availability of the genome for the frog *Xenopus laevis* (2n=36) offers an opportunity to estimate the effect of evolution of the two subgenomes of an allotetraploid species that were combined about 17-18 Myr ago, on the diversification of the *stat* gene family [25]. In parallel, we analysed the *stat* repertoire from the genome of *Xenopus tropicalis* (2n=20), which is not made of obvious pairs of homoeologous chromosomes [26]. Thirteen (13) and seven (7) *stat* genes were identified in the genome of *X. laevis* and *X. tropicalis*, respectively. *X. laevis* shows an almost perfect duplication, with the exception of the loss of *stat4.S* (Table S1; Figure 4B), while 8.3% and 31.5% of *X. laevis* genes with clear 1:1 or 2:1 orthologs in *X. tropicalis* were lost, respectively, from L and S subgenomes [25].

Interestingly, these observations in the common carp *Cyprinus carpio* and the African clawed frog *Xenopus laevis* show that additional *stat* genes in polyploid species are not rapidly eliminated, maybe because different copies can get specialized functions easily and quickly. The presence of two functional *stat2* and *sta6* genes is tolerated in both cases, and the loss of one copy is not necessarily immediate after duplication. Furthermore, the pattern of evolution of duplicated genomes in salmonids suggest some selection pressures possibly associated to viral subversions strategies [27].

Agnathan-specific stat genes shed light on the origin of vertebrate STAT transcription factors

The repertoire of *stat* genes is generally more diverse in Vertebrates than in other Metazoans [13], likely due to the two cycles of WGD that occurred in the early evolution of

this lineage. To get insight into the early steps of *stat* evolution in vertebrates, we analysed genomes from cartilaginous fish (*i.e.*, Chondrichthyans) and Agnathans.

Orthologs of all vertebrate *stat* were found in cartilaginous fish (Table 3, Figure 5A). S*tat1, 2, 3, 5* and *6* have been annotated in most species of sharks and rays for which a genome is available (Table S1). A typical *stat4* genomic sequence was not detected in shark genomes except in the whale shark *Rhincodon typus* (Genbank ID XP_020376005). An EST was also found in the dogfish shark *Squalus acanthias* (Genbank ID EE627912). Phylogenetic analysis confirmed that *stat* genes from Chondrichthyans have human orthologs (Figure 5A).

In contrast, the list of *stat* genes was different in Agnathans: in two species, the sea lamprey *Petromyzon marinus* and the hagfish *Eptatretus burgeri,* phylogenetic analysis identified orthologs of human *STAT3* and *STAT5* (Figure 5A). Two other *stat* sequences clustered with group1-4 (later referred as "*stat1-4"*) and group5-6 (later referred as "*stat5-6"*) respectively but could not be assigned to a particular set, suggesting that the *stat* repertoire of "modern" vertebrates was consolidated and standardized in Gnathostomes. Additionally, the genomic neighborhood of agnathan *stat* did not fit the well-conserved synteny blocks observed in jawed vertebrates (Figure 5B). These regions contain markers located close to *stat* genes in vertebrates, such as in *ab1, gls, myo1b, cavin1, tmeff2, slc39A10, dnah7.* However, these markers do not seem to be associated consistently with *stat* sequences in agnathans and jawed vertebrates, suggesting that these regions were produced by several duplications of an ancestral segment followed by extensive gene loss, making the reconstitution of the history of this region difficult. Markers have been best conserved in the regions encoding tetrapod *stat1, stat2, stat3 and stat4* and lamprey *stat1-4* (Figure 5B). While two *stat* genes closely linked on lamprey scaffold 5 are most similar to *stat3* and *stat5*/6 respectively, the markers found at close proximity do not match with

genes located close to human *stat3* and *5*: in human, *nab1*, *gls* and *myo1* homologs are located on chromosome 2 close to *stat2* and *stat4*. Moreover, the lamprey *stat5* is linked to *cavin*, a marker associated to human *stat*3/5, but also to *timeless* which is found close to human *stat2* on chromosome 12.

Thus, all vertebrates seem to possess genes from both *stat1-4* and *stat5-6* groups, encoded in genomic blocks inherited from an ancestral region containing *nab1, gls, myo1b, cavin1, tmeff2, slc39A10,* and *dnah7* genes. However, the standardized *stat* repertoire found in human was apparently established later in early gnathostomes. Further assemblies of agnathan genomes will help to better understand the evolution of this region.

Conserved linkages indicate that stat6 is a genomic environment closest to the ancestral stat gene.

We then analysed genomes from other deuterostomians. In these species, the repertoire of *stat* genes was significantly smaller compared to vertebrates (Table 4, Figure S2): three *stat* sequences were found in the cephalochordate lancelet *Branchiostoma floridae* and in the tunicate *Ciona intestinalis,* and one in the appendicularia *Oikopleura dioica*, in the hemichordate *Saccoglossus kowalevskii* and in the sea urchin *Strongylocentrotus purpuratus.* Most sequences clustered in phylogenetic trees with STAT5 and STAT6 (data not shown), as reported previously for non-vertebrate STAT sequences [13]. Only one sequence from the lancelet was more similar to the STAT1-4 group (Figure S2). STAT5 and STAT6 have pleiotropic roles and are involved as transcription factors in the biology of different cell types including epithelial and haematopoietic as well as immune cells. Such critical functions in a wide range of contexts are consistent with a primordial

status of these genes within the family. Overall, these results confirmed that a complete STAT1-6 repertoire could not be found in these species.

As published previously [11], protostomians genomes also contain typical *stat* genes, sometimes with multiple copies such as in the annelids *Helobdella* and *Capitella* (Table S1). These sequences were most similar to vertebrate *stat5* and *stat6* as previously reported [6]. However, we were not able to find any *stat* synteny blocks shared between these species and vertebrates. In contrast, linkage groups with *stat* genes from the placozoan *Trichoplax adhaerens* stand out as an intriguing exception (Figure 6), which reminds of our previous report about MHC [28]. In this species, *stat* genes were mainly located close to each other on scaffold 2. Seven genes flanking this cluster were homologous to 7 markers located on human chromosome 12, most of them in the close neighborhood of *STAT6* and *STAT2*, to 3 markers on human chromosome 2 close to *STAT1*, and to one marker on human chromosome 17 close to *STAT5* and *STAT3*. Interestingly, the best conserved set of linkages involved the region of *stat6*, which appears to be most closely related to the ancestral *stat* in phylogenetic analyses. These observations are consistent with the idea that both vertebrate and *Trichoplax* genomes evolved relatively slowly while those of Protostomes were subjected to extensive rearrangements. It also establishes a link between vertebrate *stat* genes and basal bilaterians.

Conclusions

The canonical signalling pathway “cytokine receptor - JAK/STAT“ contributes to many functions in invertebrates, as illustrated by in depth studies in Drosophila. In this species, this axis in involved in embryonic segmentation, in stem cell proliferation, in growth as well as in immunity [29-36].. This repertoire of types of STAT transcription factors was remarkably stable during tetrapod evolution. We found the same types of STAT in

Chondrichthyans but not in Agnathans, showing that this repertoire was likely standardized with the emergence of Gnathostomes. In ray finned fish, successive WGD offered multiple opportunities of further functional diversification and specialization. Our work shows that only *stat1* paralogs were retained after the R3 WGD, with one being constitutive and the other strongly induced by IFN. Focusing on Salmonids, we found several *stat-1*, *-3*, *-4* and *-5* due to the most recent WGD, while one copy of the *stat2*/*6* block has been retained. With 5 paralogs and a remarkable long version with additional domains, *stat1* stands out as the only member of the family prone to expansion and diversification. We have already reported that chinook salmon cells in which *stat1a1* and *stat1a2* (with constitutive expression) have been disrupted, completely lost type I IFN responsiveness [22]. Further work is needed to dissect the specialized functions of these multiple *stat1* in various cell types and infectious contexts. This evolutionary trend seems to be supported by the high number of *stat1* genes in cyprinids which have been subjected to an independent WGD. Overall, our work shows that the kinetics of *stat* loss is consistently variable across the members of the family (Figure 7). Hence, the *stat* gene family is particularly suited to study the fate of recently duplicated genes and in particular, loss-of function (or pseudogeneization), dosage effect and neofunctionalization aspects [40]. Contrasted inducibility of *stat* paralogs, which is a key mechanism of *stat* mediated immune responses, provide a fast and efficient pathway towards neo/sub-functionalization for these critical factors.

Material and methods

Identification of stat sequences

Genomes analyses were carried out using the Ensembl (Release 100) and NCBI web interfaces. tBlastn and delta blast searches on the Refseq genomes, and genome annotations searches were combined to pull out all the members of the *stat* gene family. The NCBI genomes released version are as follows: Omyk_1.0 for *Oncorhynchus mykiss*, Okis_V2 for *O. kisucht*, Otsh_v1.0 for *O. tshawytscha*, Oner_1.0 for *O. nerka*, fSalTru1.1 for *Salmo trutta*, ICSASG_v2 for *S. salar*, fSpaAur1.1 for *Sparus aurata*, gadMor3.0 for *Gadus morhua*, GRCz11 for *Danio rerio*, UCB_Xtro_10.0 for *Xenopus tropicalis*, Xenopus_laevis_v2 for *X. laevis*, GRCg6a for *Gallus gallus* and GRCh38.p13 for *Homo sapiens*. The domain structure of the proteins encoded by *stat* genes was checked using SMART and pfam to look for assembly problems and fragmentary sequences. MegaX software was used to carry out phylogenetic analyses and confirm the homology relationships between sequences. The evolutionary history was inferred by using the Maximum Likelihood method and JTT matrix-based model. The bootstrap consensus tree inferred from 1000 replicates was taken to represent the evolutionary history of the taxa analysed. Initial tree(s) for the heuristic search were obtained automatically by applying Neighbor-Joining and BioNJ algorithms to a matrix of pairwise distances estimated using a JTT model, and then selecting the topology with superior log likelihood value.

Microsynteny analysis

Synteny were retrieved from Genomicus (version 100) and the orthology/paralogy relationships available in Ensembl, and complemented by visual examination of the graphical interface in both Ensembl and NCBI. A linkage was considered a conserved microsynteny only when three or more such genes were linked in such a way in two species.

Expression analysis

RNAseq transcriptome analysis on the chinook salmon STAT2-KO GS2 and CHSE-EC cell lines was described by [22]. Briefly, STAT2 KO or control CHSE-EC cells were stimulated (or not) in EMEM medium supplemented with 250 ng/ml of recombinant *O. mykiss* IFNA2. Three biological replicates (Flask 1-3) were used for library construction for each group, and RNA-Seq libraries were prepared using TruSeq Stranded mRNA Sample Preparation Kit (Illumina) according to the manufacturer's instructions. Libraries were validated for quality on Agilent DNA1000 Kit, pooled in equimolar amounts and sequenced in pair-ends 2x75 bp on Illumina NextSeq 500/550. For each library, a depth 20 M reads were generated. Reads were then spliced-aligned to 47,898 genes (47,022 Gnomon, 876 RefSeq, GCF_002163495.1_Omyk_1.0_genomic.gff from the NCBI).

Acknowledgements

We thank Luc Jouneau for help with transcriptome analysis and annotation. This work was funded in part by the VetBioNet project (EU Grant Agreement INFRA-2016-1 731014 and by institutional grant from INRAE. BC, PB, SB and LDP carried out *in silico* analyses. BC, PB and LDP wrote the manuscript. All authors read and approved the final manuscript

Data availability

All data generated or analysed during this study are included in this published article and its supplementary information files

Conflict of interests disclosure

The authors declare no commercial or financial conflict of interest.

References

1. Aaronson, D.S., Horvath, C.M., 2002. A road map for those who don't know JAK-STAT. Science (80-.). https://doi.org/10.1126/science.1071545

2. Kiu, H., Nicholson, S.E., 2012. Biology and significance of the JAK/STAT signalling pathways. Growth Factors 30, 88–106. https://doi.org/10.3109/08977194.2012.660936

3. Villarino, A. V, Kanno, Y., O'Shea, J.J., 2017. Mechanisms and consequences of Jak–STAT signaling in the immune system. Nat. Immunol. 18, 374–384. https://doi.org/10.1038/ni.3691

4. Decker, T., Kovarik, P., 2000. Serine phosphorylation of STATs. Oncogene 19, 2628–2637. https://doi.org/10.1038/sj.onc.1203481

5. Reich, N.C., 2013. STATs get their move on. JAK-STAT 2, e27080. https://doi.org/10.4161/jkst.27080

6. Liongue, C., Sertori, R., Ward, A.C., 2016. Evolution of Cytokine Receptor Signaling. J. Immunol. 197, 11–18. https://doi.org/10.4049/jimmunol.1600372

7. Nast, R., Staab, J., Meyer, T., 2019. Gene Activation by the Cytokine-Driven Transcription Factor STAT1, in: Gene Regulation. IntechOpen. https://doi.org/10.5772/intechopen.82699

8. Stark, G.R., Darnell, J.E., 2012. The JAK-STAT Pathway at Twenty. Immunity. https://doi.org/10.1016/j.immuni.2012.03.013

9. Decker, T., Kovarik, P., Meinke, A., 1997. GAS elements: A few nucleotides with a major impact on cytokine-induced gene expression. J. Interf. Cytokine Res. 17, 121–134. https://doi.org/10.1089/jir.1997.17.121

10. Lin, J.-X., Leonard, W.J., 2019. Fine-Tuning Cytokine Signals. Annu. Rev. Immunol. 37, 295–324. https://doi.org/10.1146/annurev-immunol-042718-041447

11. Wang, Y., Levy, D.E., 2012. Comparative evolutionary genomics of the STAT family of transcription factors. JAK-STAT 1, 23–36. https://doi.org/10.4161/jkst.19418

12. Copeland, N.G., Gilbert, D.J., Schindler, C., Zhong, Z., Wen, Z., Darnell, J.E., Mui, A.L.-F., Miyajima, A., Quelle, F.W., Ihle, J.N., Jenkins, N.A., 1995. Distribution of the mammalian stat gene family in mouse chromosomes. Genomics 29, 225–228. https://doi.org/10.1006/geno.1995.1235

13. Liongue, C., O'Sullivan, L.A., Trengove, M.C., Ward, A.C., 2012. Evolution of JAK-STAT pathway components: Mechanisms and role in immune system development. PLoS One 7, e32777. https://doi.org/10.1371/journal.pone.0032777

14. Lien, S., Koop, B.F., Sandve, S.R., Miller, J.R., Kent, M.P., Nome, T., Hvidsten, T.R., Leong, J.S., Minkley, D.R., Zimin, A., Grammes, F., Grove, H., Gjuvsland, A., Walenz, B., Hermansen, R.A., Von Schalburg, K., Rondeau, E.B., Di Genova, A., Samy, J.K.A., Olav Vik, J., Vigeland, M.D., Caler, L., Grimholt, U., Jentoft, S., Inge Våge, D., De Jong, P., Moen, T., Baranski, M., Palti, Y., Smith, D.R., Yorke, J.A., Nederbragt, A.J., Tooming-Klunderud, A., Jakobsen, K.S., Jiang, X., Fan, D., Hu, Y., Liberles, D.A., Vidal, R., Iturra, P., Jones, S.J.M., Jonassen, I., Maass, A., Omholt, S.W., Davidson, W.S., 2016. The Atlantic salmon genome provides insights into rediploidization. Nature 533, 200–205. https://doi.org/10.1038/nature17164

15. Pasquier, J., Cabau, C., Nguyen, T., Jouanno, E., Severac, D., Braasch, I., Journot, L., Pontarotti, P., Klopp, C., Postlethwait, J.H., Guiguen, Y., Bobe, J., 2016. Gene evolution and gene expression after whole genome duplication in fish: the PhyloFish database. BMC Genomics 17, 368. https://doi.org/10.1186/s12864-016-2709-z

16. Wojciak, J.M., Martinez-Yamout, M.A., Dyson, H.J., Wright, P.E., 2009. Structural basis for recruitment of CBP/p300 coactivators by STAT1 and STAT2 transactivation domains. EMBO J. 28, 948–958. https://doi.org/10.1038/emboj.2009.30

17. Bhattacharya, S., Eckner, R., Grossman, S., Oldread, E., Arany, Z., D'Andrea, A., Livingston, D.M., 1996. Cooperation of Stat2 and p300/CBP in signalling induced by interferon- α. Nature 383, 344–347. https://doi.org/10.1038/383344a0

18. Zhang, J.J., Vinkemeier, U., Gu, W., Chakravarti, D., Horvath, C.M., Darnell, J.E., 1996. Two contact regions between Stat1 and CBP/p300 in interferon γ signaling. Proc. Natl. Acad. Sci. U. S. A. 93, 15092–15096. https://doi.org/10.1073/pnas.93.26.15092

19. Van de Peer, Y., 2004. Tetraodon genome confirms Takifugu findings: Most fish are ancient polyploids. Genome Biol. https://doi.org/10.1186/gb-2004-5-12-250

20. Jaillon, O., Aury, J.M., Brunet, F., Petit, J.L., Stange-Thomann, N., Maucell, E., Bouneau, L., Fischer, C., Ozouf-Costaz, C., Bernot, A., Nicaud, S., Jaffe, D., Fisher, S., Lutfalla, G., Dossat, C., Segurens, B., Dasilva, C., Salanoubat, M., Levy, M., Houdet, N., Castellano, S., Anthouard, V., Jubin, C., Castelli, V., Katinka, M., Vacherie, B., Blémont, C., Skalli, Z., Cattolico, L., Poulain, J., De Berardinis, V., Cruaud, C., Dupart, S., Brottler, P., Coutanceau, J.P., Gouzy, J., Parra, G., Lardier, G., Chapple, C., McKernan, K.J., McEwan, P., Bosak, S., Kellis, M., Volff, J.N., Gulgó, R., Zody, M.C., Mesirov, J., Lindblad-Toh, K., Birren, B., Nusbaum, C., Kahn, D., Robinson-Rechavi, M., Laudet, V., Schachter, V., Quétler, F., Saurin, W., Scarpeill, C., Wincker, P., Lander, E.S., Weissenbach, J., Roest Crollius, H., 2004. Genome duplication in the teleost fish Tetraodon nigroviridis reveals the early vertebrate proto-karyotype. Nature 431, 946–957. https://doi.org/10.1038/nature03025

21. Christoffels, A., Brenner, S., Venkatesh, B., 2006. Tetraodon genome analysis provides further evidence for whole-genome duplication in the ray-finned fish lineage. Comp. Biochem. Physiol. - Part D Genomics Proteomics 1, 13–19. https://doi.org/10.1016/j.cbd.2005.06.001

22. Dehler, C.E., Lester, K., Della Pelle, G., Jouneau, L., Houel, A., Collins, C., Dovgan, T., Machat, R., Zou, J., Boudinot, P., Martin, S.A.M., Collet, B., 2019. Viral Resistance and IFN Signaling in STAT2 Knockout Fish Cells. J. Immunol. 203, 465–475. https://doi.org/10.4049/jimmunol.1801376

23. Levraud, J.-P., Jouneau, L., Briolat, V., Laghi, V., Boudinot, P., 2019. IFN-Stimulated Genes in Zebrafish and Humans Define an Ancient Arsenal of Antiviral Immunity. J. Immunol. 203, 3361–3373. https://doi.org/10.4049/jimmunol.1900804

24. Levasseur, A., Pontarotti, P., 2011. The role of duplications in the evolution of genomes highlights the need for evolutionary-based approaches in comparative genomics. Biol. Direct. https://doi.org/10.1186/1745-6150-6-11

25. Session, A.M., Uno, Y., Kwon, T., Chapman, J.A., Toyoda, A., Takahashi, S., Fukui, A., Hikosaka, A., Suzuki, A., Kondo, M., Van Heeringen, S.J., Quigley, I., Heinz, S., Ogino, H., Ochi, H., Hellsten, U., Lyons, J.B., Simakov, O., Putnam, N., Stites, J., Kuroki, Y., Tanaka, T., Michiue, T., Watanabe, M., Bogdanovic, O., Lister, R., Georgiou, G., Paranjpe, S.S., Van Kruijsbergen, I., Shu, S., Carlson, J., Kinoshita, T., Ohta, Y., Mawaribuchi, S., Jenkins, J., Grimwood, J., Schmutz, J., Mitros, T., Mozaffari, S. V., Suzuki, Y., Haramoto, Y., Yamamoto, T.S., Takagi, C., Heald, R., Miller, K., Haudenschild, C., Kitzman, J., Nakayama, T., Izutsu, Y., Robert, J., Fortriede, J., Burns, K., Lotay, V., Karimi, K., Yasuoka, Y., Dichmann, D.S., Flajnik, M.F., Houston, D.W., Shendure, J., Dupasquier, L., Vize, P.D., Zorn, A.M., Ito, M., Marcotte, E.M.,

Wallingford, J.B., Ito, Y., Asashima, M., Ueno, N., Matsuda, Y., Veenstra, G.J.C., Fujiyama, A., Harland, R.M., Taira, M., Rokhsar, D.S., 2016. Genome evolution in the allotetraploid frog Xenopus laevis. Nature 538, 336–343. https://doi.org/10.1038/nature19840

26. Uno, Y., Nishida, C., Takagi, C., Ueno, N., Matsuda, Y., 2013. Homoeologous chromosomes of Xenopus laevis are highly conserved after whole-genome duplication. Heredity (Edinb). 111, 430–436. https://doi.org/10.1038/hdy.2013.65

27. Nan, Y., Wu, C., Zhang, Y.J., 2017. Interplay between Janus kinase/signal transducer and activator of transcription signaling activated by type I interferons and viral antagonism. Front. Immunol. https://doi.org/10.3389/fimmu.2017.01758

28. Suurväli, J., Jouneau, L., Thépot, D., Grusea, S., Pontarotti, P., Du Pasquier, L., Rüütel Boudinot, S., Boudinot, P., 2014. The Proto-MHC of Placozoans, a Region Specialized in Cellular Stress and Ubiquitination/Proteasome Pathways. J. Immunol. 193, 2891–2901. https://doi.org/10.4049/jimmunol.1401177

29. Perrimon, N., Mahowald, A.P., 1986. l(1)hopscotch, a larval-pupal zygotic lethal with a specific maternal effect on segmentation in Drosophila. Dev. Biol. 118, 28–41. https://doi.org/10.1016/0012-1606(86)90070-9

30. Zeidler, M.P., Perrimon, N., Strutt, D.I., 1999. Polarity determination in the Drosophila eye: A novel role for unpaired and JAK/STAT signaling. Genes Dev. 13, 1342–1353. https://doi.org/10.1101/gad.13.10.1342

31. Gregory, L., Came, P.J., Brown, S., 2008. Stem cell regulation by JAK/STAT signaling in Drosophila. Semin. Cell Dev. Biol. https://doi.org/10.1016/j.semcdb.2008.06.003

512 32. Morin-Poulard, I., Vincent, A., Crozatier, M., 2013. The Drosophila JAK-STAT pathway in
513 blood cell formation and immunity. JAK-STAT 2, e25700.
514 https://doi.org/10.4161/jkst.25700
515 33. Rajan, A., Perrimon, N., 2012. Drosophila cytokine unpaired 2 regulates physiological
516 homeostasis by remotely controlling insulin secretion. Cell 151, 123–137.
517 https://doi.org/10.1016/j.cell.2012.08.019
518 34. Vanha-aho, L.M., Valanne, S., Rämet, M., 2016. Cytokines in Drosophila immunity.
519 Immunol. Lett. https://doi.org/10.1016/j.imlet.2015.12.005
520 35. Stepkowski, S.M., Chen, W., Ross, J.A., Nagy, Z.S., Kirken, R.A., 2008. STAT3: An
521 important regulator of multiple cytokine functions. Transplantation.
522 https://doi.org/10.1097/TP.0b013e3181739d25
523 36. Stepkowski, S.M., Chen, W., Ross, J.A., Nagy, Z.S., Kirken, R.A., 2008. STAT3: An
524 important regulator of multiple cytokine functions. Transplantation.
525 https://doi.org/10.1097/TP.0b013e3181739d25
526 37. Whelan, S., Goldman, N., 2001. A general empirical model of protein evolution derived
527 from multiple protein families using a maximum-likelihood approach. Mol. Biol. Evol.
528 18, 691–699. https://doi.org/10.1093/oxfordjournals.molbev.a003851
529 38. Felsenstein, J., 1985. Confidence limits on phylogenies: An approach using the bootstrap.
530 Evolution (N. Y). 39, 783–791. https://doi.org/10.1111/j.1558-5646.1985.tb00420.x
531 39. Kumar, S., Stecher, G., Li, M., Knyaz, C., Tamura, K., 2018. MEGA X: Molecular
532 evolutionary genetics analysis across computing platforms. Mol. Biol. Evol. 35, 1547–
533 1549. https://doi.org/10.1093/molbev/msy096

40. Lan, X., Pritchard, J.K., 2016. Coregulation of tandem duplicate genes slows evolution of subfunctionalization in mammals. Science (80-.). 352, 1009–1013. https://doi.org/10.1126/science.aad8411

Abbreviations

STAT Signal Transducers and Activators of Transcription **GMCSF** Granulocyte-Macrophage Colony–Stimulating Factor **R1-4 WGD** Round 1-4 of Whole Genome Duplication **IFN** Interferon **GAS** IFN-γ Activation Sequence **Myr** Million Years **CCD** Coiled coil domain **DBD** DNA-binding Domain **SH2** Src Homology 2 **CTD** C-Terminal Domain **CREB** C-AMP Response Element-Binding **CCRNOT** Carbon Catabolite Repression—Negative On TATA-less **FTCD** Formimidoyl Transferase CycloDeaminase **EST** Expressed Sequence Tag

Correspondence:

B. Collet and P. Boudinot

Figure legends

Figure 1. **Evolutionary history of STAT transcription factors across tetrapods. A**. Domain structure of vertebrate STAT proteins. **B**. Maximum likelihood phylogenetic tree of STAT amino-acid sequences from human *Homo sapiens* (hosa), chicken *Gallus gallus* (gaga), *Anolis carolinensis* anole (anco), clawed frog *Xenopus tropicalis* (xetr) and *Latimeria chalumnae* coelacanth (lach). Bootstrap values (in %) of key nodes are indicated. Bootstrap values lower than 60% are not indicated. All sequences and sequence ID are provided in Table S1. **C**. Conservation of genomic neighborhood of *stat2* genes in the same tetrapod species (based on genome assemblies from Ensembl release 100).

Figure 2. **Repertoires of fish STAT amino-acid sequences and *stat* genes chromosomic distribution**. Data from genome assembly Omyk_1.0 (Oncorhynchus mykiss, rainbow trout, RefSeq GCF_002163495.1) and for other species from Ensembl release 100.

Figure 3. *stat* genes from genomes of Salmonidae (4 *Oncorhynchus* species and 2 *Salmo species*): synteny block conservation analysis. Data based on NCBI genome assemblies: Okis_V2, *Oncorhynchus kisutch* (coho salmon): GCF_002021735.2 ; Oner_1.0, *Oncorhynchus nerka* (sockeye salmon): GCF_006149115.1; Otsh_v1.0, *Oncorhynchus tshawytscha* (Chinook salmon): GCF_002872995.1; Oket_V1, *Oncorhynchus keta* (chum salmon): GCF_012931545.1 ; Omyk_1.0, *Oncorhynchus mykiss* (rainbow trout): GCF_002163495.1; fSalTru1.1, *Salmo trutta* (river trout): GCF_901001165.1 ; ICSASG_v2, *Salmo salar* (Atlantic salmon): GCF_000233375.1.

Figure 4. **Expression levels of *stat* genes (basal and induced by recombinant type I interferon) determined by RNAseq in CHSE-EC cell *O. tshawytscha* [22].** Data are on a log scale and represent the average + Standard deviation (N = 3). When the induction is statistically significant (*** p<0.001), the Fold Change is indicated

Figure 5. STAT repertoires in other polyploid species. **A.** Phylogenetic tree of STAT proteins from common carp *Cyprinus carpio* (Cyca) and zebrafish *Danio rerio* (Dare). The evolutionary history was inferred using the Maximum likelihood method (number of bootstrap tests :1000 replicates). Bootstrap values (in %) of key nodes are indicated. Bootstrap values lower than 60% are not shown. All ambiguous positions were removed for each sequence pair (pairwise deletion option). The chromosome (for zebrafish) or the scaffold (for carp) are indicated. **B.** Phylogenetic tree of STAT proteins from *Xenopus tropicalis* (Xetr), *Xenopus laevis* (Xela) and *Gallus gallus* (gaga). The gene ID are indicated and refer to Table S1. The evolutionary history was inferred using the Maximum likelihood method as for A.

Figure 6. **STAT amino-acid sequences from Chondrichthyans and Agnathans**. A. Maximum likelihood phylogenetic tree of STAT amino-acid sequences from human, elephant shark (a chimera), spiny dogfish and zebra bullhead sharks, the little skate (a ray) and sea lamprey and inshore hagfish (Agnathans). Bootstrap values (in %) of key nodes are indicated. Bootstrap values lower than 60% are not indicated. All sequences and sequence ID are provided in Table S1. B. Genomic context of *stat* genes in human and sea lamprey based on data from Ensembl release 100 (Human GRCh38.p13 and Sea Lamprey Pmarinus_7.0).

Figure 7. **Conserved genomic neighborhood between human STAT genes located on chromosomes 2, 12 and 17, and *stat* genes found in *Trichoplax adhaerens***. The location of markers is indicated besides gene names between brackets when relevant. Data from genome assemblies in Ensembl release 100 (Human GRCh38.p13 and *Trichoplax adhaerens* ASM15027v1).

Figure 8. Evolutionary pathways of stat genes in Deuterostomians. WGD are indicated by red "X", and the date indicated for the most recent events. Salmonid *stat* genes for which paralogs have not been retained are boxed in red.


Figure S1. Phylogenetic tree of salmonids STAT amino-acid sequences.


The human and zebrafish sequences were included in the analysis as reference and basis for nomenclature. The evolutionary history was inferred by using the Maximum Likelihood method and JTT matrix-based model. The bootstrap consensus tree inferred from 500 replicates is taken to represent the evolutionary history of the taxa analysed. This analysis involved 95 amino acid sequences (hs: *Homo sapiens*; dr: *Danio rerio*, zebrafish; on: *Oncorhynchus nerka*, Sockeye salmon; om: *Oncorhynchus mykiss*, rainbow trout, ot: *Oncorhynchus tshawytscha*, chinook salmon; ok: *Oncorhynchus kisutch*, Coho salmon; st: *Salmo trutta*, brown trout and ss: *Salmo salar*, Atlantic salmon). There were a total of 1742 positions in the final dataset. All sequences and sequence ID are provided in Table S1 and are based on the NCBI accession number except for ss stat1a3, only annotated as a translated transcript in ENSEMBL Evolutionary analyses were conducted in MEGA X [39].

Figure S2. Phylogenetic tree of human and non-vertebrate deuterostomian STAT amino-acid sequences.

The evolutionary history was inferred by using the Maximum Likelihood method and JTT matrix-based model. (Hosa: human; Brfl: lancelet, *Branchiostoma lanceolatum*; Ciin: *Ciona intestinalis*; Sako: *Sacchoglossus kowalevsky*; Oidi: *Oikopleura dioica*; Stpu: sea urchin, *Strongylocentrotus purpuratus*). All sequences and sequence ID are provided in Table S1. Evolutionary analyses were conducted in MEGA X [39].

Figure 1


A
intD
CCD
DBD
SH2
CTD
STAT1
STAT2
STAT3
STAT4
STAT5
STAT6
STAT i, protein interaction domain
STAT a, stat CCD (Coiled-coil domain)
STAT b, stat DBD (DNA-binding domain)
STAT SH2 (Src homology 2 domain)
CTD (C terminal domain)


B
anco_STAT1
xetr_STAT1
lach_STAT1
gaga_STAT1
hosa_STAT1
lach_STAT4
xetr_STAT4
xetr_STAT4b
gaga_STAT4
anco_STAT4
hosa_STAT4
lach_STAT3
gaga_STAT3
anco_STAT3
hosa_STAT3
xetr_STAT3
xetr_STAT3b
lach_STAT2
xetr_STAT2
hosa_STAT2
gaga_STAT2
anco_STAT2
lach_STAT6
xetr_STAT6
gaga_STAT6
anco_STAT6
hosa_STAT6
lach_STAT5a
lach_STAT5b
xetr_STAT5b
gaga_STAT5
anco_STAT5
hosa_STAT5a
hosa_STAT5b
STAT1
STAT4
STAT3
STAT2
STAT6
STAT5
0.2



C
human Chr12
Chicken Chr33
Anole Chr2
Frog Chr2
Coelacanth Scaff_JH126563

Figure 2.




Human
2
STAT1
STAT4
12
STAT2
17
STAT3
STAT5A/B
STAT6
Herring
21
stat1a
15
stat1
5
stat2
1
stat3
stat5
2
stat1b
stat4
4
stat6
Pike
22
stat1a
17
stat2
11
stat3
stat5a
16
stat1b
stat4
12
stat6
Rainbow trout
18
stat1a1
22
stat1b1
3
stat1b2
stat4a
12
stat3a
stat5a2
7
stat1a2/a3
17
stat2
22
stat1b1
stat4b
13
stat3b
16
stat5a1
stat6
stat5b
Zebrafish
22
stat1a
6
stat2
3
stat3
stat5a
9
stat1b
stat4
12
stat5b
23
stat6
Fugu
8
stat1a
19
stat2
5
stat3
stat5a
1
stat1b
stat4
3
stat6
Stickleback
I
stat1a
scaff27
stat2
XI
stat3
stat5a
XVI
stat1b
stat4
XII
stat6

Figure 3



Stat1a2 absent in ss, is a pseudogene in st
Stat1a3 has two copies in om, ok, sf
Stat5a1 and *stat5a2* are pseudogenes in ot



Level of expression (Average corrected counts + Standard deviation, N=3)
1000000
100000
10000
1000
100
10
1
***
FC 1.4
***
FC 20.2
***
FC 1.53
stat1a1
stat1a2
stat1a3
stat1b1
stat1b2
stat2
stat3a
stat3b
stat4a
stat4b
stat5a1
stat5a2
stat5b
stat6
gapdh
megfp
ncas9n
Cont
IFN

Figure 5




A.
Cyca_STAT1a2_sca292
Cyca_STAT1b2_sca68
Cyca_STAT1b1_sca28
dare_STAT1b_chr9
Cyca_STAT1a1_sca173
dare_STAT1a_chr22
Cyca_STAT4_sca28
Cyca_STAT4_sca68
dare_STAT4
Cyca_STAT3_sca270
Cyca_STAT3_sca251
dare_STAT3_chr3
Cyca_STAT2_sca26
Cyca_STAT2_sca11
dare_STAT2_chr6
dare_STAT6_ch
Cyca_STAT5a_sca251
dare-STAT5a
Cyca_STAT5b_sca97
dare_STAT5b_chr12
0.2
B.
Xetr_780397
Xela_398324
Xela_108703082
Gaga_STAT1_424044
STAT1
Gaga_STAT4_768406
Xetr_100496013
Xela_108701493
STAT4
Gaga_STAT3_420027
Xela_779100
Xela_108702168
Xetr_100498278
Xetr_100498547
Xela_379264
Xela_108702458
STAT3
Gaga_STAT2_100858732
Xela_108709841
Xetr_100170174
Xela_108708573
STAT2
Xetr_100495240
Xela_399176
Xela_414637
Gaga_STAT5_395556
STAT5
Gaga_STAT6_100859196
Xela_108709852
Xetr_100485184
Xela_495683
STAT6
0,20

Figure 6


A.
Human_STAT1_ENSG00000115415
Elephant_Shark_STAT1_JW863248
Spiny_Dogfish_STAT1_HAGW01107643.1
Zebra_bullhead_shark_STAT1_GGGL01517949.1
Little_Skate_STAT1_Contig91042/13782
Sea Lamprey_STAT1 GL482783(ENSPMAT00000003171)
Inshore Hagfish_STAT1 FYBX02009374(ENSEBUG00000007506)
Human_STAT3_ENSG00000168610
Elephant_Shark_STAT3_JW863591.1
Zebra_bullhead_shark_STAT3_GGGL01454852.1
Little_Skate_STAT3_Contig90098
Spiny_Dogfish_STAT3_HAGW01031230.1
Sea Lamprey STAT3 GL477977 (ENSPMAG00000000244)
Inshore_Hagfish_STAT_ENSEBUG00000002998
Human_STAT4_ENSG00000138378
Whale shark_STAT4_XP_020376005
Human_STAT2_ENSG00000170581
Little_Skate_STAT2_Contig10504
Spiny_Dogfish_STAT2_HAGV01033278
Zebra_bullhead_shark_STAT2_GGGL01522981
Human_STAT6_ENSG00000166888
Little_Skate_STAT6_Contig11267
Spiny_Dogfish_STAT6_HAGT01087885.1
Zebra_bullhead_shark_STAT6_GGGL01255670.1
Elephant_Shark_STAT6_JW864710.1
Human_STAT5A_ENSG00000126561
Human_STAT5B_ENSG00000173757
Little_Skate_STAT5B_Contig89644
Elephant_Shark_STAT5B_JW866184.1
Spiny_Dogfish_STAT5B_HAGV01114964.1/HAGV01114961.1
Zebra_bullhead_shark_STAT5B_GGGL01392334
Sea Lamprey STAT5 GL477207(ENSPMAG00000009008)
Inshore Hagfish STAT5 FYBX02010131(ENSEBUG00000013827)
Sea Lamprey STAT5B GL477086(ENSPMAG00000006622)
Inshore_Hagfish_STAT_ENSEBUG00000006280
STAT1
STAT3
STAT4
STAT2
STAT6
STAT5
Group1-4
Group5-6
100%
99%
80%
93%
90%
95%
76%
0.3



B.
hChr2
HIBCH MFSD6 NAB1 STAT1 MYO1B CAVIN2 SLC39A10 STK17B
INPP1 NEMP2 GLS STAT4 NABP1 TMEFF2 DNAH7
hChr12
slc39a5 coq10a cnpy2 il23a apof mip gls2 rdh16 nemp1 nab2
ankrd52 cs pan2 stat2 timeless spryd4 rbms2p myo1a stat6
hChr17
kat2a rab5c hcrt stat5b stat3 atp6v0a1 hsd17b1 psmc3ip
hspb9 kcnh4 ghfc stat5a cavin1 naglu coasy
Lamprey Scaf_4
pms1 cdk4 nemp1 stat1-4 stom cavin-like atp6v0a1 dnah7
zswim2 naglu nab1 myo1b nabp1 tmeff2 slc39a10
Lamprey Scaf_5
heximl calcrl dsc-3 stat56 stat3 blvra npy zbed8
vps50 psdm3 nab1-like gls myo1b ngfr mpp6
Lamprey Scaf_28
hoxb13 agps gdf11 sarnp timeless pmel umodl1 gcg
atp5mc3 dnajc14 abitram stat5 cavin lrp1b nell2

Figure 7




H. sapiens
hChr 17
STAT5
STAT3
CAVIN1
ATP6VOA1
hChr 2
NEMP2
NAB1
STAT1
MYO1B
CAVIN2
hChr 12
STAT2
TIM1
RBMS2
RDH16
MYO1A
NEMP1
STAT6
NAB2
SHMT2
ATP6VOA2
(57.2)
(123)
(56.3)
Scaff 2
TRIADG53559 (5.8)
TRIADG63669 (6.0)
TRIADG21854 (6.8)
TRIADG20417 (7.2)
TRIADG20799(7.9)
TRIADG53910-
12-14-18-20-47
TRIADG53906 (8.8)
TRIADG20682 (9.2)
T. adhaerens
Scaff 1
TRIADG52438 (9.8)

Figure 8



	Stat1-4	*Stat5-6*	*Stat1*	*Stat2*	*Stat3*	*Stat4*	*Stat5*	*Stat6*
Sea urchin		1						
Accorn worm		2						
Lancelet	1	1						
Sea squirt		3						
Oikopleura		1						
Agnathans								
Lamprey	1	1			1		1	
Chondrichthyans								
Leucoraja erinacea			1	1	1		1	1
Squalus acanthias			1	1	1		1	1
Rhincodon typus						1		
Bony fishes								
Zebrafish			2	1	1	1	2	1
Common carp			4	2	2	2	4	2
Atlantic Salmon			5	1	2	2	3	1
Rainbow trout			5	1	2	2	3	1
Chinook Salmon			5	1	2	2	3	1
Amphibians								
Xenopus tropicalis			1	1	2	1	1	1
Xenopus laevis			2	2	4	1	2	2
Other tetrapods			1	1	1	1	1 or 2	1

Table 1: List of potentially functional stat genes in five species of salmonids fish. The nomenclature is extended from *Danio rerio*.

	Oncorhynchus				*Salmo*	
	nerka	*mykiss*	*tshawytscha*	*Kisutch***	*truttae*	*salar*
stat 1a1	115129673 (5/+)	100136755 (18/-)	112266551 (14/-)[a]	109906681 (16/-)	115195984 (6/+)	100136558 (16/+)
stat 1a2	115133971 (2/-)	100137016 (7/-)	112253897 (7/-)[b]	109890649 (5/-)	115179269 (39/-)[##]	
stat1a3	115133986 (2/+)	110527523 (7/+)[*,d,§]	112253898 (7/+)[c]	109890433 (5/+)[d]	115179399 (39/+)[d]	106575214[#] (17/-)
stat 1b1	115138513 (1/-)	110501544 (22/+)	112244575 (Un/+)	109871062 (26/+)	115156118 (20/+)	100196256 (21/-)[e]
stat 1b2	115108041 (3/-)	110520020 (3/-)	112235369 (3/-)	109865728 (2/+)	115160666 (24/-)	106586142 (25/-)
stat 2	115143174 (15/+)	110494323 (17/+)	112217577 (2/+)	109895769 (1/+)	115207827 (14/-)	100270812 (12/+)
stat3a	115110281 (26/-)	110538194 (12/-)	112225989 (27/-)[***]	116374454 (6/-)	115171116 (32/-)	106601025 (3/-)
stat3b	115104128 (21/-)	100136756 (13/-)	112258660 (9/-)	109898422 (10/-)	115197994 (1/-)	106607297 (6/+)
stat4a	115108086 (3/-)	110520023 (3/-)	112235400 (3/-)	109865765 (2/-)	115160669 (24/-)	106586145(25/-)
stat4b	115138591 (1/-)	110501546 (22/+)	112225151 (26/-)[***]	109870786 (26/+)	115156112 (20/+)	100380385 (21/-)
stat5a1	115103495 (21/-)	100135887 (13/-)[**]	112258659 (9/-)[##]	109897291 (10/-)	115198021 (1/-)	106607295 (6/+)[**]
stat5a2	115110283 (26/-)	110538192 (12/-)	112225727 (27/-)[##]	109893324 (6/-)	115171115 (32/-)	100380532 (3/-)
stat5b	115106643 (23/+)	110491683 (16/+)	112223777 (24/+)	109865530 (20/+)	115161213 (2/+)	106579144 (19/+)
stat6	115102666 (20/+)	110491929 (16/+)	112221662 (22/+)	109869625 (24/+)	115167799 (30/+)	106567004 (13/+)

[*] annotated as "uncharacterised protein", [**] RefSeq status indicated as provisional, [***] annotated as "low quality protein", [#] annotated as pseudogene in NCBI but as coding for ENSSSAT00000091945 in Ensembl, [##] annotated as pseudogene, [a] duplicated due to genome assembly errors, identical to 112253955, [b] duplicated due to genome assembly errors, identical to 112253778, [c] duplicated due to genome assembly errors, identical to 112253779, [d] double size, [e] small size (not included in phylogenetic analysis).

** The chinook genome still contains some assembly errors resulting in artificially duplicated regions in particular between the chromosomes 7 and 14 and within the chromosome 7. In this species, only one stat5 was annotated as functional in contrast to 3 stat5 genes found in the five other species analysed, named stat5.1-3.

§ stat1a3 is doubled in NCBI but separate in Ensembl as ENSOMYG00000034815 and ENSOMYG00000035706, both annotated as stat1a.

Table 2. Additional genes encoding short ORF with significant hit to signal transducer and activator of transcription. None could be identified in *O. tshawytscha* nor in *O. kisutch.*

Species	GeneID (chromosome/orientation)	Size of the largest isoform (aa)
O. mykiss	110503197 (24/+)	243
S. truttae	115188323 (Un/+)	184
	115189982 (Un/+)	249
	115190002 (Un/-)	136
	115156120 (20/+)	249
	115156122 (20/+)	186
	115156123 (20/+)	179
	115156124 (20/+)	147
	115156138 (20/+)	186
	115156139 (20/+)	184
S. salar	106583229 (22/-)	314

Table 3. Presence and number of *stat* genes in genomes of Chondrichthyans and Agnathans

Phylum	Species	Gene number	Gene name	Conserved domains
Chondrichthyans	Elephant shark *Callorhincus milli*	>=4	ENSCMIG00000003696 (3) ENSCMIG00000003757 (5b) ENSCMIG00000010732 (1) ENSCMIG00000015418 (1)	STATi-STATa-STATb-SH2 STATi-STATa-STATb-SH2 STATi-STATa-STATb-SH2 STATi-STATa-STATb-SH2**
	Whale shark *Rhyncodon typus*		XP_020376005	STATi-STATa-STATb-SH2
Agnathans	Lamprey *Petromyzon marinus*	4	ENSPMAG00000000244 ENSPMAG00000002770 ENSPMAG00000006622 ENSPMAG00000009008	STATi-STATa-STATb-SH2 STATi-STATa-STATb-SH2 … STATa-STATb… STATi-STATa-STATb-SH2
	Hagfish *Eptatretus burgeri*	4/5	ENSEBUG00000002998& ENSEBUG00000003439 &ENSEBUG00000006280 ENSEBUG00000007506 ENSEBUG00000013827	… STATb-SH2 …-SH2 (separated by gls) STATi-STATa-STATb-SH2 STATi-STATa-STATb

** there are 2 STAT genes in this entry !!

Table 4. Presence and number of *stat* genes in genomes of other deuterostomians.

Phylum	Species	Gene number	Gene name	Conserved domains
Non vertebrate Deuterostomians	Sea urchin *Strongylocentrotus purpuratus*	1	SP-STAT	STATi-STATa-STATb-SH2
	Saccoglossus kowalevskii (Hemichordates)	1	XP_006814941	STATi-STATa-STATb-SH2
	Branchiostoma floridae (Cephalochordates)	2	XP_019630041/BL18533 XP_002594129BL09530	STATi-STATa-STATb-SH2 STATi-STATa-STATb-SH2
	Oikopleura dioica (Appendicularia)	1	AAS21327	STATi-STATa-STATb-SH2
	Ciona intestinalis (Tunicates)	3	ENSCING00000004044 ENSCING00000010308 ENSCING00000024295	STATi-STATa-STATb-SH2 …STATa-STATb-SH2 STATi-STATa