



HAL
open science

Spatial distribution and activity patterns as welfare indicators in response to water quality changes in European sea bass, *Dicentrarchus labrax*

Sébastien Alfonso, Bastien Sadoul, Xavier Cousin, Marie-Laure Bégout

► To cite this version:

Sébastien Alfonso, Bastien Sadoul, Xavier Cousin, Marie-Laure Bégout. Spatial distribution and activity patterns as welfare indicators in response to water quality changes in European sea bass, *Dicentrarchus labrax*. *Applied Animal Behaviour Science*, 2020, 226, pp.1-11. 10.1016/j.applanim.2020.104974 . hal-03135465

HAL Id: hal-03135465

<https://hal.inrae.fr/hal-03135465v1>

Submitted on 22 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Spatial distribution and activity patterns as welfare indicators in response to water quality**
2 **changes in European sea bass, *Dicentrarchus labrax***

3

4 Sébastien Alfonso^{1,2}, Bastien Sadoul^{2,3}, Xavier Cousin^{2,4} and Marie-Laure Bégout^{1,2}

5

6 ¹ Ifremer, Laboratoire Ressources Halieutiques, Station de La Rochelle, Place Gaby Coll, 17137

7 L'Houmeau, France.

8 ² MARBEC, Univ. Montpellier, CNRS, IRD, Ifremer, 34250 Palavas-les-flots, France.

9 ³ INRAE, UR1037 Fish physiology and Genomics, 35000 Rennes, France

10 ⁴ Université Paris-Saclay, AgroParisTech, INRAE, GABI, 78350 Jouy-en-Josas, France.

11

12 * Corresponding author: Sébastien Alfonso

13 Phone : +39 (0) 809 246 190; e-mail: Sebastien.alfonso1@gmail.com

14 Present address: COISPA Tecnologia & Ricerca, Via dei trulli 18, 70126 Torre a Mare (Bari), Italy.

15

16 **Abstract**

17 In aquaculture, fish are exposed to unavoidable stressors that can be detrimental for their health and
18 welfare. However, welfare in farmed fish can be difficult to assess, and, so far, no standardized test
19 has been universally accepted as a welfare indicator. This work contributes to the establishment of
20 behavioural welfare indicators in a marine teleost in response to different water quality acute stressors.
21 Groups of ten fish were exposed to high Total Ammonia Nitrogen concentration (High TAN, 18 mg.L⁻¹),
22 Hyperoxia (200 % O₂ saturation), Hypoxia (20 % O₂ saturation), or control water quality (100% O₂
23 saturation and TAN < 2.5 mg.L⁻¹) over 1 hour. Fish were then transferred in a novel environment for a
24 group behaviour test under the same water quality conditions over 2 hours. Videos were recorded to
25 assess thigmotaxis, activity and group cohesion. After this challenge, plasma cortisol concentration
26 was measured in a subsample, while individual behavioural response was measured in the other fish
27 using novel tank diving test. Prior to this study, the novel tank diving test was validated as a
28 behavioural challenge indicative of anxiety state, by using nicotine as anxiolytic drug. Overall, all
29 stress conditions induced a decrease in activity, thigmotaxis and group cohesion while only fish
30 exposed to Hypoxia and High TAN conditions displayed elevated plasma cortisol concentrations. In
31 *post*-stress condition, activity was still affected but normal behaviour was recovered within the 25
32 minutes of the test duration. Our work suggests that the activity, thigmotaxis and group cohesion are
33 good behavioural indicators of exposure to degraded water quality, and could be used as standardized
34 measures to assess fish welfare.

35 **Keywords:** Fish; Welfare; Water quality; Behaviour; Stress.

36 1. Introduction

37 Fish production has expanded importantly during the last decades, both because of the world's
38 diminishing natural wild resources and the increase in demand for fish products (FAO, 2018).
39 Aquaculture represented 53 % of the total fish production (including non-food uses) in 2016 (FAO,
40 2018) and is now recognized as a major food production industry. Thus, as well as for terrestrial
41 farming industry, concerns about sustainability, environmental issues and animal welfare in
42 aquaculture are increasing (Conte, 2004; Ashley, 2007; Martins et al., 2010; Martins et al., 2012;
43 Hixson, 2014; FAO, 2018; Lembo et al., 2019). It is common that under aquaculture conditions, and in
44 every fish husbandry system, variations of water quality variables such as temperature, pH, oxygen
45 (O₂), carbon dioxide (CO₂) or Total Ammonia Nitrogen concentrations (TAN) occur. Such variations
46 when they reach a certain threshold, depending on the species *preferendum*, could be considered as
47 stress factors (stressor) and therefore deleterious for fish health and welfare. The exposure to stressors,
48 such as degraded water quality may mobilize fish energy for coping with the stressor hereby
49 decreasing the available energy allocated to growth and reproduction, or directly causes death if the
50 magnitude of stress is too high (Barton, 2002; Sneddon et al., 2016). Therefore, it can finally affect
51 fish production and has economic consequences for farmers (Conte, 2004; Lembo et al., 2019).

52 Exposure to stress factors triggers a cascade of biological events within an organism to cope with these
53 factors. In fish, the hypothalamo-pituitary-interrenal axis (HPI) is involved in the production and
54 release of cortisol into circulation acting as an activator of the physiological and behavioural responses
55 (Sumpter, 1997; Sadoul and Vijayan, 2016; Schreck and Tort, 2016). Among the previously cited
56 variables, oxygen and ammonia concentrations are known to activate the HPI axis when they vary,
57 leading to the stimulation of cortisol release and the triggering of behavioural adaptive responses
58 (Knoph and Olsen, 1994; van Raaij et al., 1996; Espmark and Baeverfjord, 2009).

59 Thus, behavioural measurements have proven to be sensitive indicators of the complex existing
60 biochemical and physiological changes that occur in response to stress (Schreck, 1990; Scherer, 1992;
61 Schreck et al., 1997; Martins et al., 2012). Behaviours, such as changes in food-anticipatory activity,

62 feed intake, ventilation rate, individual and group swimming activity are commonly used as welfare
63 indicators (Huntingford et al., 2006; Martins et al., 2012; Huntingford and Kadri, 2014; Carbonara et
64 al., 2015; Carbonara et al., 2019). Group swimming behaviour is defined as the spatial distribution and
65 swimming activity of the group of fish held within an aquaculture production unit and it covers shoal
66 structure, the horizontal and vertical distribution of the group, their swimming speed and direction
67 (Martins et al., 2012). For instance, exposure to negative *stimuli*, such as poor water quality, is known
68 to lead to rapid escape movements (Stien et al., 2007; Bratland et al., 2010) or to alter group cohesion
69 (Domenici et al., 2002; Espmark and Baeverfjord, 2009; Sadoul et al., 2014; Sadoul et al., 2017).
70 Thus, group swimming behaviour appears to be a sensitive welfare indicator even if it is still lacking
71 calibration efforts to be precisely translated into an operational welfare indicator; nevertheless some
72 examples exist (Papandroulakis et al., 2014; Pettersen et al., 2014). Moreover, the appraisal of
73 negative or positive stimuli and, hence, the psychological dimension of stress as defined for fish by
74 Galhardo and Oliveira (2009) is seldom tackled in welfare research. There exists however a
75 complementary measure which is the individual behavioural responses to novel environment and in
76 particular the novel tank diving test which is worldwide used along with the measure of stereotypies,
77 such as thigmotaxis to assess anxiety in zebrafish (*Danio rerio*) in ecotoxicology and pharmacology
78 research (Levin et al., 2007; Egan et al., 2009; Vignet et al., 2014; Macaulay et al., 2015; Alfonso et
79 al., 2019a). In further details, the novel tank diving test was validated as a tool for evaluating anxiety
80 by using drugs, such as nicotine. Short exposure to nicotine is known to reduce anxiety in fish, through
81 its action on nicotinic acetylcholine receptors as demonstrated by the use of specific inhibitors (Levin
82 et al., 2007; Bencan & Levin, 2008). In the context of novel tank diving test, nicotine-exposure
83 (bathing) has been shown to be anxiolytic by triggering change in fish space utilization, such as higher
84 time spent in the top area of the novel tank which translate a relief from bottom dwelling behaviour
85 that fish would express under predator threat for example. The novel tank diving test could thus be a
86 helpful non-invasive tool to monitor farmed fish anxiety state *post* stress exposure hereby assessing
87 psychological stress and contributing to the assessment of positive or negative emotions and, hence
88 better welfare state determination.

89 Overall, the objectives of the present study were to further contribute to the establishment of
90 behavioural welfare indicators including the psychological dimension of stress in a model marine
91 teleost in response to different water quality stressors. Firstly, the novel tank diving test outcome was
92 validated as a behavioural indicator of anxiety in European sea bass *Dicentrarchus labrax*, using
93 nicotine as an anxiolytic reference drug. Secondly, behavioural responses of fish group in response to
94 a novel environment under acute and severe water quality deterioration, including Total Ammonia
95 Nitrogen (High TAN) increase (18 mg.L⁻¹), Hyperoxia (200 % O₂ saturation) and Hypoxia (20 % O₂
96 saturation) were evaluated along with cortisol measurement. Finally, individual behaviour expressed
97 following the same water quality exposures were assessed using the novel tank diving test translated
98 from ecotoxicology studies.

99 **2. Material and methods**

100 Experiments were authorized by ethics committee agreement APAFIS#7098 and all procedures
101 involving animals were in accordance with the ethical standards of the institution and followed the
102 recommendations of Directive 2010/63/EU.

103 **2.1. Fish rearing**

104 Juvenile European sea bass were hatched and reared at Ifremer Palavas-les-flots research station
105 (France, 34250) until 280 days *post* fertilization (dpf) according to sea bass rearing standard (Chatain,
106 1994). They were then transferred to Ifremer L’Houmeau (France, 17137). No mortality occurred
107 during the transfer between the two facilities. They were then randomly separated in groups of 100
108 fish into 4 tanks of 400 L (90x90x50 cm). Tanks shared a recirculating system with a flow rate of 4 m³
109 per hour and water was renewed at a rate of 20 % per day. Water temperature was maintained at 21.5
110 ± 1°C, oxygen around 100 % saturation, and salinity and pH were respectively set to 20.5 ± 1 and 8.3.
111 The light regime was 13:11 L/D. Total Ammonia Nitrogen (TAN) concentration was < 2.5 mg.L⁻¹
112 (equivalent to 0.1 mg.L⁻¹ [NH₃]). The fish were hand-fed using commercial diet from Le Gouessant
113 (France) once a day each morning at 9:00 at 1 % of biomass. Fish were reared in L’Houmeau for 3
114 months before the first experiment.

2.2. Translating and validating the novel tank diving test in European sea bass

The novel tank diving test assessing position choice along the vertical dimension has been previously validated as a metric of adaptation to a novel environment and a proxy of the anxiety level of individual fish (Levin et al., 2007; Egan et al., 2009). The first objective of this work was to adapt the test protocol (observation duration, tank size relative to fish size) to European sea bass, and then to validate the measure of anxiety using nicotine as a reference anxiolytic drug.

Two groups of fish (385 dpf, n=24 per group) were randomly selected and transferred from the home rearing tank to two 50 L tanks into the experimental room. After a 1-hour acclimation period, control fish (n=24) were transferred one by one in a 3L tank with the same water quality for 8 min (24.5 x 15 x 13.5 cm, AquaBox 3; Aqua Schwarz GmbH). The same protocol was followed for the nicotine-bathed fish (n=24), except they were bathed one at a time during five min in a 3L tank containing 5 mg.L⁻¹ nicotine solution (Pestanal®, Sigma Aldrich) then placed in normal plain water for the next three minutes. After a 1-hour acclimation period, control fish (n=24) were transferred one by one in a 3L tank (24.5 x 15 x 13.5 cm, AquaBox 3; Aqua Schwarz GmbH) filled with normal plain water for five min and then transferred to a second 3L tank (same water quality) for an extra three min. The same protocol was followed for the nicotine-bathed fish (n=24), except they were first bathed one at a time during 5 min in a 3L tank containing 5 mg.L⁻¹ nicotine solution (Pestanal®, Sigma Aldrich) added to normal water and then transferred in plain water for the next three minutes. This two steps bathing protocol was applied to ensure correct elimination of nicotine (at least in the gills) and to ensure similar handling for both conditions. For both treatment, directly after the bathing period, the individual fish was gently transferred in a novel tank containing normal plain water (29 x 21 x 17 cm, 10 L trapezoid tank from Aquatic Habitat. Inc.), and a video was recorded in side view during 25 minutes. For space occupancy analysis, the tank was virtually separated into two areas according to Egan et al. (2009): top area including one half of the volume and bottom area including the other half. Time spent in top area (s) and latency to enter top area (s), variables which are both indicative of anxiety level, were measured. In addition, distance travelled (cm) and number of transitions between

141 areas, indicative of fish activity level were also measured. Variables were recorded in each frame and
142 they were summed over periods of 5 min or over the whole test duration (*i.e.* 25 min).

143 **2.3. Exposure to stress condition and water quality characterization**

144 In the morning prior to the experiment (*i.e.* exposure to stress condition), a group of 10 fish was gently
145 randomly caught from the rearing tank and transferred to the behavioural room where they were
146 maintained in a tank (70 L, height 48 cm, diameter 49.5 cm) filled with 60 L of the same water as in
147 their home tank (**Figure 1**). Fish were kept under standard condition during 1 h for tank acclimation
148 and then, during the next hour, one of the four following conditions was applied *i.e.* Control, High
149 Total Ammonia Nitrogen (TAN) concentration, Hyperoxia and Hypoxia. Experiments were performed
150 in triplicates (n=10 fish x 3 run per condition).

151 For the control condition, fish were maintained under the same standard condition than in the rearing
152 room (*i.e.*, at $21.5 \pm 1^\circ\text{C}$, 100 % O_2 saturation and $\text{TAN} < 2.5 \text{ mg.L}^{-1}$). For high TAN condition, 8.5 g
153 of ammonium chloride NH_4Cl (Fluka 09711, Sigma Aldrich) dissolved in 0.5 L of water was added
154 twice at 30 min interval to reach a targeted TAN concentration of 18 mg.L^{-1} which corresponds to 1.6
155 mg.L^{-1} of NH_3 . Three water samples (50 mL) were taken to quantify *a posteriori* the TAN
156 concentration: (i) before adding ammonium chloride, (ii) at the beginning and (iii) at the end of the
157 novel environment in group. Samples were stored at -22°C before further analysis. Then, TAN
158 concentration was quantified using a spectrophotometer with continuous flow (Alliance Integral
159 Futura, Frépillon, France) and analytical method is described below.

160 Samples were filtered using GF/F $0.7\mu\text{m}$ filter (Whatman, Maidstone, United Kingdom). TAN
161 concentration was quantified using a spectrophotometer with continuous flow (Alliance Integral
162 Futura, Frépillon, France) using colorimetric method. The solutions used for the calibration analyses
163 that were performed the same day came from stock solutions from 0.1g.L^{-1} to 1g.L^{-1} of ammoniacal
164 nitrogen (NH_4^+) stocked at a temperature of 8°C . The calibration curve of ammoniacal nitrogen
165 showed a high R^2 validating the procedure ($\text{NH}_4^+=1.439 \times \text{OD} - 0.002$; $R^2 = 0.99$, where OD is the
166 optical density measured using the spectrophotometer).

167 The NH₃ concentration was determined using the following equation as described by Johansson and
168 Wedborg (1980):

169
$$[NH_3] = \frac{[NH_4^+]}{K_1 \times [H^+]}$$

170 Where $[H^+] = 10^{-pH}$ and $\log K_1 = -0.467 + 0.00113 \times \text{salinity} \times 2887.9/\text{temperature (K)}$ according to
171 Johansson and Wedborg (1980).

172 For Hypoxia condition, oxygen concentration was slowly lowered to reach 20 % of O₂ saturation using
173 nitrogen bubbling. For Hyperoxia condition, oxygen concentration was slowly increased to reach 200
174 % of O₂ saturation using oxygen bubbling. Oxygen concentration was monitored every 5 min during
175 the entire experiment using an Oxygen probe (Oxi 3310, WTW, Xylem Analytics Germany Sales
176 GmbH & Co. KG). Oxygen concentration, NH₄⁺ and NH₃ concentrations recorded over the
177 experimental duration are presented below.

178 For control condition, water oxygen concentration was maintained at 100% saturation (at 21.5 ± 1°C)
179 and TAN concentration was < 2.5 mg.L⁻¹ during both bathing and novel environment in group. For
180 High TAN condition, oxygen concentration was above 100 % and TAN concentrations were 15.3 ±
181 4.2 and 18.8 ± 7.6 mg.L⁻¹ at the beginning and at the end of the test respectively, corresponding to
182 UIA-N concentrations of 1.34 ± 0.37 and 1.64 ± 0.67 mg.L⁻¹. For both Hyperoxia and Hypoxia
183 conditions, TAN concentration was maintained at less than 2.5 mg.L⁻¹. For the hyperoxia condition,
184 oxygen concentration increased slowly during the bathing period to reach 208 ± 5.6 % of saturation,
185 while for the hypoxia it slowly decreased to reach 21.3 ± 3.8 % of saturation before the start of the
186 novel environment challenge in group. All conditions were then maintained during two hours until the
187 end of the group test (**Figure 2**). In the high TAN condition, the concentration of NH₃ represents half
188 of the LC50 concentration reported for European sea bass after 96 h of exposure under similar
189 hydrological conditions (i.e. temperature=17.5°C, salinity=34, and pH=8.15) than in our experiment
190 (Person-Le Ruyet et al., 1995).

191 After the initial bathing period, the fish group was challenged in a novel environment with the same
192 water quality (see section 2.4.1). After 2 hours, fish from two replicates were then monitored
193 individually into novel tank diving test in plain system water (see section 2.4.2, n=20 for control, n=20

194 for High TAN and n=18 for Hyperoxia and Hypoxia). The third replicate was used to quantify plasma
195 cortisol concentration following the challenge in group (n=10 per condition, see section 2.5). Fish
196 were randomly selected for blood sampling or for performing novel tank diving test. After blood
197 sampling (n=9 or 10 fish per condition) or novel tank diving test, all fish were measured for weight (g,
198 to the nearest mg) and standard length (cm, to the nearest mm) under the same anaesthesia conditions
199 (see section 2.5).

200 **2.4. Behavioural procedures**

201 All behavioural experiments were performed in a dedicated room where environmental parameters
202 were identical to rearing conditions. All videos were recorded at 25 frame.s⁻¹ with an analogue camera
203 ICD-48E (Ikegami) and a 2.1-13.5 lens (Fujinon) linked to a computer with an acquisition card and
204 EthoVision XT 10.0 software (Noldus, The Netherlands). Data extraction and analyses were
205 performed using EthoVision XT 13.1 software. Swaps between individuals during novel environment
206 in group were manually corrected using the track editor module (Noldus, The Netherlands).

207 *2.4.1. Novel environment in group*

208 After the bathing period, the entire group of fish (n=10) was gently transferred into a novel arena (110
209 cm x 110 cm x 6 cm, 70 L). After 1 min, video recording started for 2 hours in top view. For each
210 condition, water parameters were maintained similar to those obtained at the end of the bathing period
211 for the respective condition following the same procedure explained in section 2.2.2.

212 For space occupancy, the visualization of heat maps was produced for each 5-min period using
213 Ethovision XT 13 (Noldus The Netherlands). For further analysis, the arena was separated into two
214 areas: Centre area including one half of the volume and periphery area including the other half; time
215 spent in periphery (s), indicative of thigmotaxis behaviour (Ferrari et al., 2014), was recorded.
216 Distance travelled by each fish (cm), indicative of individual fish activity, and the interindividual
217 distances (cm), indicative of group cohesion (Buske and Gerlai, 2011), were also recorded. Variables
218 were recorded for each frame and they were summed (for Distance travelled) or averaged (for time

219 spent in periphery and interindividual distances) over period of 5 min every 30 minutes (*i.e.*, four
220 sampling periods: 0-5 min, 30-35 min, 60-65 min and 90-95 min).

221 2.4.2. *Novel tank diving test post-stress condition*

222 After novel environment challenge in group, fish were gently transferred individually into a novel tank
223 (same as the one used for method validation with nicotine, see above) containing plain system water.
224 Fish swimming activity and vertical position was monitored during 25 min to assess anxiety state and
225 behavioural recovery following stressful conditions exposure. For space occupancy and fish
226 swimming activity analyses, the same procedure as presented before in section 2.2 was followed.

227 2.5. Cortisol measurement

228 Immediately after the end of the observation period in the novel environment challenge, fish were
229 gently caught and transferred into a 10 L tank which contained 500 $\mu\text{L.L}^{-1}$ of a benzocaine stock
230 solution (50 g.L^{-1} in 100 % ethanol; Benzocaine Sigma-Aldrich, Saint-Quentin Fallavier, France).
231 Blood samples were obtained within 3 minutes from the venous sinus with heparinised syringes.
232 Thereafter, blood was centrifuged (5 min at 4000 g) to obtain plasma samples which were stored at -
233 22°C until further analyses. Plasma cortisol concentration was determined by ELISA (RE52061, IBL
234 International, Hamburg, Germany) using a Synergy-HT (BioTek Instruments, Winooski, VT, USA)
235 following manufacturer instructions.

236 At the end of the experiment, all fish were euthanized and sexed using an overdose of benzocaine (1
237 mL.L^{-1} from the stock solution described above), following the recommendations of Directive
238 2010/63/EU.

239 2.6. Statistical analyses

240 Statistical analyses were performed using Statistica 9.0 software (StatSoft, USA). All statistical
241 analyses were carried out at a 5 % level of significance and values are represented as mean \pm SEM
242 except where otherwise mentioned. Normality and homoscedasticity were tested *a priori* using
243 Shapiro-Wilks test and when sample sizes were too small non parametric statistics were used.

244 Fish weight and length were compared between conditions using one-way ANOVA to ensure no bias
245 existed for behavioural and physiological responses interpretation.

246 For validation of the Novel tank diving test, time spent in top area and distance travelled were
247 compared between fixed factors (Nicotine bathed *vs.* Control) with a repeated-measures ANOVA
248 (with 5 periods, *i.e.* 5-min, 10-min, 15-min, 20-min and 25-min) and a Tukey HSD *post-hoc* test. A
249 Mann-Whitney U-test was also performed to compare between conditions, total time spent in top area,
250 latency to enter top area, total distance travelled by fish and number of transitions between top and
251 bottom areas.

252 For Novel environment challenge in group, distance travelled, interindividual distances and time spent
253 in periphery area were compared between conditions with a Factorial ANOVA (with 4 periods, *i.e.* 0-
254 5-min, 30-35-min, 60-65min, 90-95min) followed by Tukey HSD *post-hoc* test.

255 For novel tank diving test *post-stress* condition, time spent in top area and distance travelled were
256 compared between conditions with a repeated-measures ANOVA (with 5 periods, *i.e.* 5-min, 10-min,
257 15-min, 20-min and 25-min) and a Tukey HSD *post-hoc* test. A One-way ANOVA followed by Tukey
258 HSD *post-hoc* test was performed to compare between conditions the total time spent in top area,
259 latency to enter top area, total distance travelled by fish and number of transitions

260 Finally, since sex had no significant effect on plasma cortisol concentration, a Mann-Whitney U-test
261 was performed to compare cortisol values between control and each condition.

262 **3. Results**

263 Overall, 168 fish were used for the different experiments: 48 for the validation of the novel tank diving
264 test (body weight: 50.8 ± 0.6 g, standard length: 15.4 ± 0.9 cm, $n=24$ per group); 120 for the stress
265 condition exposures (51 ± 1 g, 15.3 ± 0.2 cm, $n=30$ per group), batches were homogeneous between
266 conditions ($F=1.5$, $df=3$, $p=0.23$ and $F=2$, $df=3$, $p=0.13$ for body weight and standard length
267 respectively).

268 **3.1. Validation of novel tank diving test in European sea bass**

269 Irrespective of the condition (Nicotine bathed vs. Control) fish progressively explored the top area of
270 the novel tank ($F=6.0$, $df=4$, $p<0.001$). Nicotine bathed-fish spent significantly more time in the top
271 area than control ones during each 5-min period of the test ($F=31.8$, $df=1$, $p<0.001$; **Figure 3.A**) as
272 well as over the total duration of the test ($Z=4.8$, $p<0.001$; **Figure 3.B**). Fish also progressively
273 travelled more distance during the experiment whatever the condition ($F=13.3$, $df=4$, $p<0.001$) but
274 nicotine bathed-fish travelled significantly more distance than control ones during each 5-min period
275 of the test ($F=15.9$, $df=1$, $p<0.001$; **Figure 3.C**) as well as over the total duration of the test ($Z=3.5$,
276 $p<0.001$; **Figure 3.D**). Nicotine bathed-fish swam more between the two areas and entered quicker in
277 the top area than control ones ($Z=4.3$, $p<0.001$ and $Z=-5.0$, $p<0.001$ respectively; **Figure 3.E,F**).

278 **3.2. Novel environment in group**

279 Heatmaps, representing the mean location frequency of fish from different conditions and during
280 different periods of the novel environment challenge in group, are plotted in **Figure 4**. For control
281 condition, fish responded to the group test by staying mainly in the periphery of the arena during the
282 first five minutes of the test. Progressively, they explored the centre of the arena. On the contrary,
283 other groups of fish (*i.e.* High TAN, Hyperoxia and Hypoxia), were located in the centre area from the
284 beginning of the test. In addition, fish dispersion increased over the test duration for exposed animals,
285 while fish from the control condition stayed aggregated during the entire experiment.

286 Under stress conditions, distance travelled by fish differed between conditions ($F=212.7$, $df=3$,
287 $p<0.001$), and periods ($F=33.2$, $df=3$, $p<0.001$) with a significant interaction observed over time
288 ($F=10.3$, $df=9$, $p<0.001$). Control fish travelled progressively more distance over the test duration;
289 3178 ± 268 cm during period 0-5 min vs. 5148 ± 254 cm during period 90-95 min ($p<0.001$). Fish
290 from High TAN, Hyperoxia and Hypoxia conditions showed a lower distance travelled compared to
291 control fish during the entire test ($p<0.05$ for all conditions) and it was stable over time (**Figure 5.A**).

292 Interindividual distances, indicative of group cohesion, differed between conditions ($F=449.3$, $df=3$,
293 $p<0.001$), periods ($F=17.4$, $df=3$, $p<0.001$) with a significant interaction between conditions and
294 periods ($F=29.1$, $df=9$, $p<0.001$). Fish from control condition progressively decreased their

295 interindividual distances during the challenge; 48.3 ± 1.4 cm during period 0-5 min vs. 41.9 ± 1.1 cm
296 during period 90-95 min ($p < 0.001$). During the 0-5 min period, fish from control and Hypoxia
297 conditions showed similar interindividual distances ($p = 0.06$), whereas fish from High TAN and
298 Hyperoxia conditions swam closer to each other ($p < 0.001$ for both conditions). However, starting from
299 the 60-65 min to the 90-95 min period, fish from both High TAN and Hypoxia conditions showed
300 higher dispersion than control fish ($p < 0.001$) while fish from Hyperoxia condition continued to display
301 lower dispersion than control fish from the beginning to the end of the testing period ($p < 0.001$; **Figure**
302 **5.B**).

303 Time spent in periphery area also differed between conditions ($F = 15.3$ $df = 3$, $p < 0.001$), not between
304 periods ($F = 0.5$, $df = 3$, $p = 0.66$) but the interaction between conditions and periods was significant
305 ($F = 10$, $df = 9$, $p < 0.001$). Fish from control condition spent more time in the centre area from the second
306 time period (30-35 min) until the end (90-95 min) compared to the first 0-5 min period ($p = 0.003$,
307 $p = 0.003$ and $p < 0.001$ for 30-35 min, 60-65 min and 90-95 min respectively). During the first 0-5 min
308 period of the test, fish from High TAN, Hyperoxia and Hypoxia conditions spent less time in the
309 peripheric area than control fish ($p < 0.001$ for both conditions) whereas they spent the same time in the
310 periphery during the rest of the test, with the exception of High TAN which still spent more time in
311 Centre than Control during 30-35 period (**Figure 5.C**).

312 **3.3. Novel tank diving test *post*-stress condition**

313 Irrespective of the condition, fish progressively explored the top area of the novel tank ($F = 5.9$, $df = 4$,
314 $p < 0.001$). There was an effect of condition ($F = 11.8$, $df = 3$, $p < 0.001$) and an interaction between
315 conditions and period were also found ($F = 4.4$, $df = 12$, $p < 0.001$; **Figure 6.A**). The total time spent in
316 top area differed according to conditions ($F = 11.8$, $df = 3$, $p < 0.001$; **Figure 6.B**) with High TAN fish
317 spending significantly more time in the top area than the three other conditions ($p < 0.001$).

318 Distance travelled by fish varied in relation to the duration of the test ($F = 13.0$, $df = 4$, $p < 0.001$),
319 conditions ($F = 15$, $df = 3$, $p < 0.001$) and interactions between period and conditions were significant
320 ($F = 3$, $df = 12$, $p < 0.001$; **Figure 6.C**). Total distance travelled by fish also differed between conditions

321 (F=15, df=3, p<0.001; **Figure 6.D**). High TAN, Hyperoxia and Hypoxia fish travelled less distance
322 than control fish during the novel tank test (p<0.001 for all conditions). No difference was observed
323 for the number of transitions between the two areas between conditions (F=2.3, df=3, p=0.8; **Figure**
324 **6.E**) as well as for the latency to enter into the top area (F=0.5, df=3, p=0.7; **Figure 6.F**).

325 **3.4. HPI axis response**

326 Plasma cortisol concentrations for fish from Hyperoxia condition were not different from control fish
327 values (Z=0.1, p=0.93, **Figure 7**). On the contrary, plasma cortisol concentrations were higher for both
328 High TAN and Hypoxia conditions compared to control values (Z=-3.5, p<0.001 and Z=-2.1, p=0.04
329 respectively).

330 **4. Discussion**

331 Overall, this study demonstrates that commonly used behavioural tests in neurobiology or
332 ecotoxicology on model species are relevant for evaluating anxiety in a model marine farmed fish and
333 this opens new opportunities to evaluate psychological stress following environmental perturbation.
334 Moreover, we have shown that activity, group cohesion and thigmotaxis are relevant behavioural
335 indicators revealing acute exposures to High Total Ammonia Nitrogen (TAN), Hyperoxia and
336 Hypoxia in a group situation. Individual bottom dwelling behaviour was also assessed in response to
337 the same stressors using the novel tank diving test and proved to be sensitive.

338 Fish display typical swimming features in a new environment, e.g. bottom dwelling, which have been
339 used to define a suitable procedure to measure anxiety or anxiety-like responses. Common protocols
340 exist for using these behavioural features for anxiety phenotyping in zebrafish (Levin et al., 2007) and
341 they were successfully transferred to other small model species such as three-spine stickleback and
342 fathead minnow (Margiotta-Casaluci et al., 2014; Thompson et al., 2016). To our knowledge, our
343 study represents the first attempt to adapt this test for a larger fish of commercial and ecological
344 importance such as European Sea bass (Vandeputte et al., 2019). Since this species is known to be a
345 highly stress responding fish (Levin et al., 2007; Fanouraki et al., 2011; Alfonso et al., 2019b), we
346 extended the test duration to 25 minutes. Despite this longer test period (usually between 5 and 10

347 minutes for other species) the time spent in the upper part of the tank remained limited to
348 approximately 10%.

349 Nicotine has been found to reduce anxiety in zebrafish through activation of acetylcholine nicotinic
350 receptors (Levin et al., 2007; Bencan & Levin, 2008). In the context of novel tank diving test,
351 nicotine-exposure has been shown to be anxiolytic by triggering change in fish space utilization. In the
352 present study, acute treatment with nicotine induced the same relief of anxiety as observed in zebrafish
353 (Levin 2007). It is to note however that the concentration we used, 5 mg.L⁻¹, was much lower than the
354 one efficient in zebrafish (100 mg.L⁻¹) and a concentration of 50 mg.L⁻¹ was not able to induce change
355 in bottom dwelling behaviour in zebrafish (Levin et al., 2007). Indeed, in zebrafish, exposure to
356 nicotine results in a release from bottom dwelling behaviour (i.e. swimming in the top area of the
357 novel tank) over test duration and increases the locomotor activity (Levin et al., 2007; Bencan and
358 Levin, 2008). In the present study, after exposure to nicotine, sea bass similarly spent more time in the
359 top area and were more active (i.e., travelled more distance and showed a higher transitions number
360 between tank areas) compared to control. Altogether, these results advocate that anxiolytic effects of
361 nicotine are behaviourally observable in European sea bass, as in zebrafish. These results firstly
362 suggest conservation of the sensitivity to nicotine and the associated behavioural response in
363 phylogenetically relatively distant fish species. This is supported by the high conservation of nicotinic
364 acetylcholine receptors in fishes as recently shown (Pedersen et al., 2019) and the fact that we have
365 found sequence coding for protein sharing high similarity with several nicotinic receptors in the
366 genome of European sea bass (not shown). Secondly, this study supports the fact that the novel tank
367 diving test is a promising tool for monitoring anxiety in marine teleost such as European sea bass.

368 The four experimental conditions had very distinct water qualities. In the control condition, oxygen
369 saturation was at 100 % and TAN concentration was under 2.5 mg.L⁻¹ while TAN concentration was
370 approximately 8 fold higher in High TAN condition and oxygen saturation was 2 fold higher and 5
371 fold lower in Hyperoxia and Hypoxia conditions respectively. TAN concentrations in the High TAN
372 condition corresponded to a concentration of NH₃ of 1.6 mg.L⁻¹ (half of the LC50, Person-Le Ruyet et
373 al., 1995). Oxygen saturations chosen during the Hyperoxia and Hypoxia conditions were previously

374 described to affect fish physiology and behaviour during chronic exposure without being lethal
375 (Chapman and Mckenzie, 2009; Espmark and Baeverfjord, 2009; Rimoldi et al., 2016). Both
376 concentrations of oxygen and TAN used in the present study were quite extreme but can occasionally
377 occur (or even co-occur) in aquaculture conditions in case of technical failures.

378 In the novel environment in group, control fish displayed thigmotaxis behaviour and spent most of the
379 time in the periphery area during the first 5 minutes of the test. Indeed, handling fish from bathing tank
380 to a novel environment is known to induce a typical thigmotaxis behavioural response. This
381 thigmotaxis behaviour has been shown to indicate an anxiety state (Prut and Belzung, 2003; Schnorr et
382 al., 2012). Interestingly, fish from High TAN, Hyperoxia and Hypoxia conditions spent less time in
383 the periphery area than control fish, which suggest that they were not displaying anxiety-like
384 behaviour contrary to control fish in response to handling. High TAN, Hyperoxia and Hypoxia are
385 known to directly affect fish survival (Magaud et al., 1997; Person-Le Ruyet and Boeuf, 1998; Shimps
386 et al., 2005; Rimoldi et al., 2016). Indeed, fish survival can be impacted in many ways following their
387 exposure to one of these stressors. For instance, in hypoxic conditions, the oxygen quantity is limited,
388 therefore fish have to adapt their physiology and behaviour to maximize oxygen uptake and avoid
389 death by asphyxia (Chapman & Mckenzie, 2009). Then, Hyperoxia may alter the equilibrium of ions
390 in the gill, causes a decrease in the ventilatory frequency inducing oxidative damages or triggers
391 diseases and thus can cause fish death (Dejours et al. 1977; Liepelt et al.1995; Brauner et al. 2000).
392 Finally, Ammonia exposure can affect osmoregulation, represses the immune system, or causes
393 asphyxiation leading to hyperventilation, convulsions and death (Randall and Tsui, 2002; Eddy, 2005;
394 Camargo and Alonso, 2006). By contrast to these stressors, handling is mostly documented for its
395 effects on the HPI axis, through cortisol release (Barton 2002). To our knowledge, handling does not
396 have direct effects on survival except if fish are injured which was not the case in our experiment.
397 Therefore, we hypothesise that when fish are coping with stressors affecting survival, the thigmotaxis
398 expression in response to a psychological stressor is overruled, possibly through endorphin release.
399 Further mechanistic studies are needed to better understand what changes within the fish are related to
400 these behavioural adaptations under multi-stressors exposure. Furthermore, during the test in group

401 situation, control fish travelled progressively more distance over the test duration whereas High TAN,
402 Hyperoxia and Hypoxia conditions showed lower and stable distance travelled. Such lower swimming
403 activity under high TAN was described before in rainbow trout (Shingles et al., 2001) and under
404 hypoxia conditions in common sole (*Solea solea*; Dalla Via et al. (1998)) and dogfish (*Scyliorhinus*
405 *canicula*; Metcalfe and Butler (1984)) whereas hyperoxia was shown to lead to higher variability of
406 the activity pattern in Atlantic salmon (Espmark and Baeverfjord, 2009). It is, however, important to
407 note that behavioural responses under hypoxia were found to be clearly dependant on the intensity and
408 duration of the hypoxia challenge and the species (Chapman and Mckenzie, 2009). NH₃ is known to
409 cause asphyxiation by reducing the blood oxygen-carrying capacity and hence alter the swimming
410 performances reported above (Shingles et al., 2001; Camargo and Alonso, 2006). Indeed, decrease in
411 oxygen or increase in TAN concentration both lead to reduced active metabolic rate (Muusze et al.,
412 1998; Shingles et al., 2001) and may explain why exposed fish are being less active than controls. It
413 seems that fish adopted a “wait and see” strategy consisting in minimizing energy expenditure and
414 waiting for an improvement in water quality (van Raaij et al., 1996; Clingerman et al., 2007). Finally,
415 concerning interindividual distances, during the 0-5 min period of novel environment in group, fish
416 from Control and Hypoxia conditions expressed the same group cohesion whereas fish from high TAN
417 and Hyperoxia conditions were closer to each other. Fish from Hyperoxia condition stayed closer
418 during all test duration while both fish from High TAN and Hypoxia conditions displayed lower group
419 cohesion than control fish from 60-65 to 90-95 min periods. Interestingly, Domenici et al. (2002)
420 reported the same larger dispersion in Atlantic herring under O₂ saturation of 20 % and same results
421 were observed in Atlantic salmon under O₂ saturation of 150 % (Espmark and Baeverfjord, 2009).
422 Lower group cohesion was also observed in rainbow trout under other stress factor such as
423 hypercapnia by Sadoul et al. (2017). When given the possibility, it is well known that fish are able to
424 avoid stressful environment such as hypoxic waters, high ammonia or high CO₂ concentrations
425 (Richardson et al., 2001; Clingerman et al., 2007; Skjæraasen et al., 2008; Herbert et al., 2010).
426 Therefore we suggest that fish interindividual distance increases to maximize oxygen uptake
427 (Domenici et al., 2017) and thus could be used as a relevant welfare indicator (Juell, 1995; Espmark
428 and Baeverfjord, 2009). This behavioural strategy can also be perceived as a decrease in attention

429 and/or used for saving energy for a better coping with stress favouring a return to a homeostatic state.
430 Additional experiments are however required to fully understand the observed disruptions of the shoal
431 cohesion but it overall fits well with the on-growing research effort about the effects of abiotic factors
432 on fish group behaviour (Weetman et al., 1999; Domenici et al., 2002; Espmark and Baeverfjord,
433 2009; Colchen et al., 2017; Domenici et al., 2017; Sadoul et al., 2017; Colson et al., 2019).

434 Concerning the HPI axis responsiveness, plasma cortisol concentrations are always high for European
435 sea bass upon stress exposure since this species is known to be high cortisol responder (Rotllant et al.,
436 2003; Samaras et al., 2016; Alfonso et al., 2019b). In undisturbed conditions, plasma cortisol values
437 are expected to be around 100 ng.L⁻¹ but it can reach a value 10 fold higher upon stress (Samaras et al.,
438 2018; Alfonso et al., 2019b). In addition to the confinement stress in the bathing tank, fish also
439 experienced two handling stresses, one transfer between the rearing and the bathing tanks and then
440 between the bathing and the experimental tanks explaining the high cortisol values measured in all
441 conditions. Measured plasma cortisol concentrations were similar between fish exposed to Hyperoxia
442 and control fish but were higher in fish exposed to Hypoxia and High TAN conditions than control. A
443 similar increase in cortisol release was previously reported in Atlantic salmon following chronic
444 exposure to ammonia (0.15 mg.L⁻¹, (Knoph and Olsen, 1994) and hyperoxia (150 % O₂ saturation,
445 (Espmark and Baeverfjord, 2009) or acute exposure to hypoxia (25 % of O₂ saturation) in rainbow
446 trout (van Raaij et al., 1996). Behavioural disruptions observed in Hyperoxia treated fish did not
447 translate in terms of cortisol values compared to control fish, suggesting the possible implication of
448 other physiological mechanisms, such as plasma catecholamines as previously shown in rainbow trout
449 (van Raaij et al., 1996; Gesto et al., 2013; 2015). Interestingly, within a species, individuals are known
450 to differ from each other in terms of amplitude of primary stress response (e.g. cortisol and
451 catecholamines) and, along with divergent behavioural responses, that defines individual coping styles
452 (Castanheira et al., 2017). Therefore, it would be interesting in future studies to include
453 complementary coping style characterizations to investigate the link between coping style, behavioural
454 responses within a group and primary responses upon stress exposure. This would help in assessing
455 whether different strategies co-exist to cope with these kinds of stressors within a fish group.

456 After 2 h-exposure to any of the tested conditions (High TAN, Hyperoxia and Hypoxia), fish travelled
457 less distance in novel tank diving test than control fish during the first 5 minutes in system water.
458 Thereafter all fish reached the same activity level. Such behavioural recovery from hypoxia in
459 European sea bass is rapid and similar to that reported in Nile Tilapia (*Oreochromis niloticus*, Xu et al.
460 (2006)). To our knowledge, it is the first time that behavioural recovery from Hyperoxia and High
461 ammonia concentration has been monitored. In addition to distance travelled, fish previously exposed
462 to High TAN explored the top area more than control fish and decreased progressively time spent in
463 top to reach control's level after 20 min in system water. On the contrary, the two other conditions
464 (*i.e.*, Hyperoxia and Hypoxia) explored progressively the top area in the same way than control fish.
465 The fast release from bottom dwelling (or the search for access to water surface) observed in High
466 TAN exposed fish resembled the surface swimming observed for nicotine-exposed fish. As explained
467 before for the novel environment test in group situation, the lower anxiety-like behaviour observed in
468 High TAN fish could also be the result of fish setting priority to cope with TAN instead of expressing
469 psychological stress effects *i.e.* a relief of anxiety.

470 **Conclusion**

471 In conclusions, our results showed that thigmotaxis, swimming activity, group cohesion and bottom
472 dwelling behaviour are reliable behavioural indicators of health and welfare status in European sea
473 bass. Moreover, this study reports the fact that it is important to distinguish between stressors that
474 could affect survival and psychological stressors such as handling or confinement alone. Besides
475 cortisol, other molecular factors should be sought for to fully understand individual stress responses
476 depending on the type and/or duration of stressors. Finally, the novel tank diving test seems to be
477 sensitive for screening recovery state and thus better evaluate welfare status in farmed fish.

478 **Acknowledgments**

479 This work was supported by the French National Research Agency, project ERANET ANIHWa Win-
480 Fish (ANR-14-ANWA-0008). SA received a PhD grant from Ifremer to conduct this research. Lucette
481 Joassard, Christine Jarc, Didier Leguay, Mathieu Auvinet, Maxime Petitjean and Thibault Geoffroy

482 from Ifremer technical staff are acknowledged for their help in conducting the experiments. We also
483 thank the two anonymous reviewers for helpful comments on the previous version of the manuscript.

484 **References**

485 Alfonso, S., Blanc, M., Joassard, L., Keiter, S.H., Munsch, C., Loizeau, V., Begout, M.L., Cousin,
486 X., 2019a. Examining multi- and transgenerational behavioral and molecular alterations resulting from
487 parental exposure to an environmental PCB and PBDE mixture. *Aquat Toxicol* 208, 29-38.

488 Alfonso, S., Sadoul, B., Gesto, M., Joassard, L., Chatain, B., Geffroy, B., Begout, M.L., 2019b.
489 Coping styles in European sea bass: The link between boldness, stress response and neurogenesis.
490 *Physiol Behav* 207, 76-85.

491 Ashley, P.J., 2007. Fish welfare: Current issues in aquaculture. *Applied Animal Behaviour Science*
492 104, 199–235.

493 Barton, B., 2002. Stress in fishes: a diversity of responses with particular reference to changes in
494 circulating corticosteroids. *Integ. and Comp. Biol* 42, 517-525.

495 Bencan, Z., Levin, E.D., 2008. The role of alpha 7 and alpha 4 beta 2 nicotinic receptors in the
496 nicotine-induced anxiolytic effect in zebrafish. *Physiology & Behavior* 95, 408-412.

497 Bratland, S., Stien, L.H., Braithwaite, V.A., Juel, J.E., Folkedal, O., Nilsson, J., Oppedal, F.,
498 Fosseidenge, J.E., Kristiansen, T.S., 2010. From fright to anticipation: using aversive light stimuli to
499 investigate reward conditioning in large group of Atlantic salmon (*Salmo salar*). *Aquaculture*
500 *International* 18, 991-1001.

501 Buske, C., Gerlai, R., 2011. Early embryonic ethanol exposure impairs shoaling and the dopaminergic
502 and serotonergic systems in adult zebrafish. *Neurotoxicol Teratol* 33, 698-707.

503 Brauner CJ, Seidelin M, Madsen SS, Jensen F.B. 2000. Effects of freshwater hyperoxia and
504 hypercapnia and their influences on subsequent seawater transfer in Atlantic salmon *Salmo*
505 *salar* smolts. *Can J Fish Aquat Sci* 57:2054–2064.

506 Camargo, J.A., Alonso, A., 2006. Ecological and toxicological effects of inorganic nitrogen pollution
507 in aquatic ecosystems: A global assessment. *Environ Int* 32, 831-849.

508 Carbonara, P., Scolamacchia, M., Spedicato, M.T., Zupa, W., McKinley, R.S., Lembo, G., 2015.
509 Muscle activity as a key indicator of welfare in farmed European sea bass (*Dicentrarchus labrax* L.
510 1758). *Aquaculture research* 46(9), 2133-2146.

511 Carbonara, P., Alfonso, S., Zupa, W., Manfrin, A., Fiocchi, E., Pretto, T., Spedicato, M.T. & Lembo,
512 G. (2019). Behavioral and physiological responses to stocking density in sea bream (*Sparus aurata*):
513 Do coping styles matter?. *Physiol Behav* 212, 112698.

514 Castanheira, M. F., Conceição, L. E. C., Millot, S., Rey, S., Bégout, M. L., Damsård, B., Kristiansen,
515 T., E, H., Øverli, Ø. & Martins, C. I. M. (2017). Coping styles in farmed fish: consequences for
516 aquaculture. *Reviews in Aquaculture* 9, 23-41.

517 Chapman, L.J., Mckenzie, D.J., 2009. Chapter 2 Behavioral Responses and Ecological Consequences.
518 *Fish physiology Hypoxia* (Ed Richard J.G, Farrell A.P and Brauner C.J), 27, 25-77.

519 Chatain, B., 1994. Estimation et amélioration des performances zootechniques de l'élevage larvaire de
520 *Dicentrarchus labrax* et de *Sparus auratus*.

521 Clingerman, J., Bebak, J., Mazik, P.M., Summerfelt, S.T., 2007. Use of avoidance response by
522 rainbow trout to carbon dioxide for fish self-transfer between tanks. *Aquacultural Engineering* 37,
523 234-251.

524 Colchen, T., Teletchea, F., Fontaine, P., Pasquet, A., 2017. Temperature modifies activity, inter-
525 individual relationships and group structure in a fish. *Current Zoology* 63, 175-183.

526 Conte, F.S., 2004. Stress and the welfare of cultured fish. *Applied Animal Behaviour Science* 86, 205–
527 223.

528 Colson, V., Mure A., Valotaire, C., Le Calvez, J.M., Goardon, Labbe, L., Leguen I., Prunet P., 2019.
529 A novel emotional and cognitive approach to welfare phenotyping in rainbow trout exposed to poor
530 water quality. *Applied animal behaviour science* 210, 103-112.

531 Dalla Via, J., Van Den Thillart, G., Cattani, O., Cortesi, P., 1998. Behavioural responses and
532 biochemical correlates in *Solea solea* to gradual hypoxic exposure. *Canadian Journal of Zoology* 76,
533 2108-2113.

534 Dejours, P., Toulmond, A., Truchot, J.P., 1977. The effect of hyperoxia on the breathing of
535 marine fishes. *Comp. Biochem. Physiol A* 58, 409-411.

536 Domenici, P., Ferrari, R.S., Steffensen, J.F., Batty, R.S., 2002. The effect of progressive hypoxia on
537 school structure and dynamics in Atlantic herring *Clupea harengus*. *Proc Biol Sci* 269, 2103-2111.

538 Domenici, P., Steffensen, J.F., Marras, S., 2017. The effect of hypoxia on fish schooling. *Philos Trans*
539 *R Soc Lond B Biol Sci* 372.

540 Eddy, F.B., 2005. Ammonia in estuaries and effects on fish. *J. Fish Biol* 67(6), 1495-1513.

541 Egan, R.J., Bergner, C.L., Hart, P.C., Cachat, J.M., Canavello, P.R., Elegante, M.F., Elkhayat, S.I.,
542 Bartels, B.K., Tien, A.K., Tien, D.H., Mohnot, S., Beeson, E., Glasgow, E., Amri, H., Zukowska, Z.,
543 Kalueff, A.V., 2009. Understanding behavioral and physiological phenotypes of stress and anxiety in
544 zebrafish. *Behavioural Brain Research* 205, 38-44.

545 Espmark, A.M., Baeverfjord, G., 2009. Effects of hyperoxia on behavioural and physiological
546 variables in farmed Atlantic salmon (*Salmo salar*) parr. *Aquaculture International* 17, 341-353.

547 Fanouraki, E., Mylonas, C.C., Papandroulakis, N., Pavlidis, M., 2011. Species specificity in the
548 magnitude and duration of the acute stress response in Mediterranean marine fish in culture. *General*
549 *and Comparative Endocrinology* 173, 313-322.

550 FAO, 2018. The state of world fisheries and aquaculture., Food and Organisation of the United
551 Nations, p. 227 pp.

552 Ferrari, S., Benhaim, D., Colchen, T., Chatain, B., Begout, M.L., 2014. First links between self-
553 feeding behaviour and personality traits in European seabass, *Dicentrarchus labrax*. *Applied Animal*
554 *Behaviour Science* 161, 131-141.

555 Galhardo, L., Oliveira, R.F., 2009. Psychological stress and welfare in fish. *Annual review of*
556 *Biomedical Sciences*, 1-20.

557 Gesto, M., Lopez-Patino, M.A., Hernandez, J., Soengas, J.L., Miguez, J.M., 2013. The response of
558 brain serotonergic and dopaminergic systems to an acute stressor in rainbow trout: a time course study.
559 *Journal of Experimental Biology* 216, 4435-4442.

560 Gesto, M., Lopez-Patino, M.A., Hernandez, J., Soengas, J.L., Miguez, J.M., 2015. Gradation of the
561 stress response in rainbow trout exposed to stressors of different severity: the role of brain
562 serotonergic and dopaminergic systems. *J Neuroendocrinol* 27, 131-141.

563 Herbert, N.A., Skjæraasen, J.E., Nilsen, T., Salvanes, A.G.V., Steffensen, J.F., 2010. The hypoxia
564 avoidance behaviour of juvenile Atlantic cod (*Gadus morhua L.*) depends on the provision and
565 pressure level of an O₂ refuge. *Marine Biology* 158.

566 Hixson, S.M., 2014. Fish Nutrition and Current Issues in Aquaculture: The Balance in Providing Safe
567 and Nutritious Seafood, in an Environmentally Sustainable Manner. *Journal of Aquaculture Research*
568 *Development* 5, 234.

569 Huntingford, F.A., Adams, C., Braithwaite, V.A., Kadri, S., Pottinger, T.G., Sandøe, P., Turnbull, J.F.,
570 2006. Current issues in fish welfare. *Journal of Fish Biology* 68, 332-372.

571 Huntingford, F.A., Kadri, S., 2014. Defining, assessing and promoting the welfare of farmed fish.
572 *Revue Scientifique Et Technique-Office International Des Epizooties* 33, 233-244.

573 Johansson, O., Wedborg, M., 1980. Ammonia-Ammonium Equilibrium in Seawater at Temperatures
574 between 5 and 25-Degrees-C. *Journal of Solution Chemistry* 9, 37-44.

575 Juell, J.E., 1995. The behaviour of Atlantic salmon in relation to efficient cage-rearing. *Reviews in*
576 *Fish Biology and Fisheries* 5, 320-335.

577 Knoph, M.B., Olsen, Y.A., 1994. Subacute Toxicity of Ammonia to Atlantic Salmon (*Salmo-Salar L*)
578 in Seawater - Effects on Water and Salt Balance, Plasma-Cortisol and Plasma Ammonia Levels.
579 *Aquatic Toxicology* 30, 295-310.

580 Lembo, G., Carbonara, P., Fabris, A., Manfrin, A., Zupa, W., 2019. Welfare issues and veterinary
581 treatments.

582 Levin, E.D., Bencan, Z., Cerutti, D.T., 2007. Anxiolytic effects of nicotine in zebrafish. *Physiol Behav*
583 90, 54-58.

584 Liepelt, A., Karbe, L., Westendorf, J. 1995. Induction of DNA strand breaks in rainbow trout
585 *Oncorhynchus mykiss* under hypoxic and hyperoxic conditions. *Aquat. Toxicol* 33, 177-181.

586 Macaulay, L.J., Bailey, J.M., Levin, E.D., Stapleton, H.M., 2015. Persisting effects of a PBDE
587 metabolite, 6-OH-BDE-47, on larval and juvenile zebrafish swimming behavior. *Neurotoxicol Teratol*
588 52, 119-126.

589 Magaud, H., Migeon, B., Morfin, P., Garric, J., Vindimian, E., 1997. Modelling fish mortality due to
590 urban storm run off: interacting effects of hypoxia and un-ionized ammonia. *Water research* 31, 211-
591 218.

592 Margiotta-Casaluci, L., Owen, S.F., Cumming, R.I., de Polo, A., Winter, M.J., Panter, G.H., Rand-
593 Weaver, M., Sumpter, J.P., 2014. Quantitative Cross-Species Extrapolation between Humans and
594 Fish: The Case of the Anti-Depressant Fluoxetine. *Plos One* 9.

595 Martins, C.I., Eding, E.H., Verdegem, M.C.J., Heinsbroek, L.T.N., Schneider, O., Blancheton, J.P.,
596 Roque d'Orbcastel, E., Verreth, J.A.J., 2010. New developments in recirculating aquaculture systems
597 in Europe: A perspective on environmental sustainability. *Aquacultural Engineering* 43, 83-93.

598 Martins, C.I., Galhardo, L., Noble, C., Damsgard, B., Spedicato, M.T., Zupa, W., Beauchaud, M.,
599 Kulczykowska, E., Massabuau, J.C., Carter, T., Planellas, S.R., Kristiansen, T., 2012. Behavioural
600 indicators of welfare in farmed fish. *Fish Physiol Biochem* 38, 17-41.

601 Metcalfe, J.D., Butler, P.J., 1984. Changes in activity and ventilation in response to hypoxia in
602 unrestrained, unoperated dogfish (*Scyliorhinus canicula* L.). *J Exp Biol* 108, 411-418.

603 Muusze, B., Marcon, J., van den Thillart, G., Almeida-Val, 1998. Hypoxia tolerance of Amazon fish
604 Respirometry and energy metabolism of the cichlid *Astronotus Ocellatus*. *Comparative Biochemistry*
605 *and Physiology Part A* 120, 151-156.

606 Papandroulakis, N., Lika, K., Kristiansen, T.S., Oppedal, F., Divanach, P., Pavlidis, M., 2014.
607 Behaviour of European sea bass, *Dicentrarchus labrax* L., in cages - impact of early life rearing
608 conditions and management. *Aquaculture Research* 45, 1545-1558.

609 Pedersen, J.E., Bergqvist, C.A., Larhammar, D., 2019. Evolution of vertebrate nicotinic acetylcholine
610 receptors. *BMC Evol Biol* 19(1):38.

611 Person-Le Ruyet, J., Boeuf, G., 1998. L'Azote ammoniacal, un toxique potentiel en élevage de
612 poissons : le cas du turbot. *Bulletin Francais de la Peche et de la Pisciculture* 350-351, 393-412.

613 Person-Le Ruyet, J., Chartois, H., Quemener, L., 1995. Comparative acute ammonia toxicity in marine
614 fish and plasma ammonia response. *Aquaculture* 136, 181-194.

615 Pettersen, J.M., Bracke, M.B.M., Midtlyng, P.J., Folkedal, O., Stien, L.H., Steffenak, H., Kristiansen,
616 T.S., 2014. Salmon welfare index model 2.0: an extended model for overall welfare assessment of

617 caged Atlantic salmon, based on a review of selected welfare indicators and intended for fish health
618 professionals. *Reviews in Aquaculture* 6, 162-179.

619 Prut, L., Belzung, C., 2003. The open field as a paradigm to measure the effects of drugs on anxiety-
620 like behaviors: a review. *Eur J Pharmacol* 463, 3-33.

621 Randall, D.J., Tsui, T.K.N., 2002. Ammonia toxicity in fish. *Marine Pollution Bulletin* 45(1-
622 12), 17-23.

623 Richardson, J., Williams, E.E., Hickey, W., 2001. Avoidance behaviour of freshwater fish and shrimp
624 exposed to ammonia and low dissolved oxygen separately and in combination. *New Zealand Journal*
625 *of Marine and Freshwater Research* 35, 625-633.

626 Rimoldi, S., Terova, G., Zaccone, G., Parker, T., Kuciel, M., Dabrowski, K., 2016. The Effect of
627 Hypoxia and Hyperoxia on Growth and Expression of Hypoxia-Related Genes and Proteins in Spotted
628 Gar *Lepisosteus oculatus* Larvae and Juveniles. *J Exp Zool B Mol Dev Evol* 326, 250-267.

629 Rotllant, J., Ruane, N.M., Caballero, M.J., Montero, D., Tort, L., 2003. Response to confinement in
630 sea bass (*Dicentrarchus labrax*) is characterised by an increased biosynthetic capacity of interrenal
631 tissue with no effect on ACTH sensitivity. *Comp Biochem Physiol A Mol Integr Physiol* 136, 613-
632 620.

633 Sadoul, B., Evouana Mengues, P., Friggens, N.C., Prunet, P., Colson, V., 2014. A new method for
634 measuring group behaviours of fish shoals from recorded videos taken in near aquaculture conditions.
635 *Aquaculture* 430, 179-187.

636 Sadoul, B., Friggens, N.C., Valotaire, C., Labbe, L., Colson, V., Prunet, P., Leguen, I., 2017.
637 Physiological and behavioral flexibility to an acute CO₂ challenge, within and between genotypes in
638 rainbow trout. *Comp Biochem Physiol A Mol Integr Physiol* 209, 25-33.

639 Sadoul, B., Vijayan, M.M., 2016. Stress and growth. *Fish physiology Biology of stress in fish*,
640 academic Press, 167-205.

641 Samaras, A., Dimitroglou, A., Sarropoulou, E., Papaharisis, L., Kottaras, L., Pavlidis, M., 2016.
642 Repeatability of cortisol stress response in the European sea bass (*Dicentrarchus labrax*) and
643 transcription differences between individuals with divergent responses. *Sci Rep* 6, 34858.

644 Samaras, A., Esperito Santo, C., Papandroulakis, N., Mitrizakis, N., Pavlidis, M., Höglund E.,
645 Pelgrim, T.N.M., Zetholf, J., Spanings F.A.T., Vindas, M.A., Ebbesson, L.O.E., Flik, G., Gorissen,
646 M., 2018. Allostatic Load and Stress Physiology in European Seabass (*Dicentrarchus labrax* L.) and
647 Gilthead Seabream (*Sparus aurata* L.). *Front Endocrinol* 9, 451.

648 Scherer, E., 1992. Behavioural responses as indicators of environmental alterations: approaches,
649 results, developments. *J. Appl. Ichthyol.* 8, 122-131.

650 Schnorr, S.J., Steenbergen, P.J., Richardson, M.K., Champagne, D.L., 2012. Measuring thigmotaxis in
651 larval zebrafish. *Behaviour Brain Research* 228, 367-374.

652 Schreck, C.B., 1990. Physiological, behavioral, and performance indicators of stress. *Am. Fish. Soc.*
653 *Symp.* 8, 29-37.

654 Schreck, C.B., Olla B., L., Davis, M.W., 1997. Behavioral responses to stress, in: Iwama, G.K.,
655 Pickering, A.D., Sumpter, J.P., Schreck, C.B. (Eds.), *Fish stress and Health in Aquaculture*,
656 Cambridge University Press, Cambridge, U.K., pp. 145-170.

657 Schreck, C.B., Tort, L., 2016. The concept of stress in fish. *Fish physiology Biology of stress in fish*,
658 academic Press, 1-34.

659 Shimps, E.L., Rice, J.A., Osborne, J.A., 2005. Hypoxia tolerance in two juvenile estuary-dependent
660 fishes. *Journal Experimental of Marine Biology and Ecology* 325, 146-162.

661 Shingles, A., McKenzie, D.J., Taylor, E.W., Moretti, A., Butler, P.J., Ceradini, S., 2001. Effects of
662 sublethal ammonia exposure on swimming performance in rainbow trout (*Oncorhynchus mykiss*). *J*
663 *Exp Biol* 204, 2691-2698.

664 Skjæraasen, J.E., Nilsen, T., Meager, J.J., Herbert, N.A., Moberg, G.P., Tronci, V., Johansen, T.,
665 Salvanes, A.G.V., 2008. Hypoxic avoidance behaviour in cod (*Gadus morhua* L.): The effect of
666 temperature and haemoglobin genotype. *Journal Experimental of Marine Biology and Ecology* 358,
667 70-77.

668 Sneddon, L.U., Wolfenden, D.C.C., J.S., T., Tort, L., 2016. Stress management and welfare. *Fish*
669 *physiology Biology of stress in fish*, academic Press, 463-539.

670 Stien, L.H., Bratland, S., Austevoll, I., Oppedal, F., Kristiansen, T.S., 2007. A video analysis
671 procedure for assessing vertical fish distribution in aquaculture tanks. *Aquacultural Engineering* 37,
672 115-124.

673 Sumpter, J.P., 1997. The endocrinology of stress. *Fish Stress and Health in Aquaculture*, Cambridge
674 University Press, Cambridge Society for Experimental Biology, Seminar series:62, 95-118.

675 Thompson, R.R.J., Paul, E.S., Radford, A.N., Purser, J., Mendl, M., 2016. Routine handling methods
676 affect behaviour of three-spined sticklebacks in a novel test of anxiety. *Behavioural Brain Research*
677 306, 26-35.

678 van Raaij, M.T., Pit, D.S., Balm, P.H., Steffens, A.B., van den Thillart, G.E., 1996. Behavioral
679 strategy and the physiological stress response in rainbow trout exposed to severe hypoxia. *Horm*
680 *Behav* 30, 85-92.

681 Vandeputte, M., Gagnaire, P.A., Allal, F., 2019. The European sea bass: a key marine fish model in
682 the wild and in aquaculture. *Anim Genet* 50, 195-206.

683 Vignet, C., Devier, M.H., Le Menach, K., Lyphout, L., Potier, J., Cachot, J., Budzinski, H., Begout,
684 M.L., Cousin, X., 2014. Long-term disruption of growth, reproduction, and behavior after embryonic
685 exposure of zebrafish to PAH-spiked sediment. *Environ Sci Pollut Res Int* 21, 13877-13887.

686 Weetman, D., Atkinson, D., Chubb, J.S., 1999. Water temperature influences the shoaling decisions of
687 guppies, *Poecilia reticulata*, under predation threat. *Animal behavior* 58, 735-741.

688 Xu, J., Liu, Y., Cui, S., Miao, X., 2006. Behavioral responses of tilapia (*Oreochromis niloticus*) to
689 acute fluctuations in dissolved oxygen levels as monitored by computer vision. *Aquacultural*
690 *Engineering* 35, 207-217.

691

692 **Captions**

693 **Figure 1.** Experimental protocol followed for fish exposure to different water qualities (stress
694 condition) and behavioural and physiological measurements.

695 **Figure 2.** Mean \pm SEM of TAN and UIA-N ($\text{mg}\cdot\text{L}^{-1}$) and oxygen saturation ($\% \text{O}_2$) for High TAN,
696 Hyperoxia and Hypoxia conditions during bathing period (at 0, 30 and 60 min) and at the start and end
697 of novel environment challenge in group.

698 **Figure 3.** Swimming characteristics in the novel tank diving test. Mean \pm SEM of **(A)** time spent in
699 top area in relation to the observation period (s); **(B)** total time spent in top area (s); **(C)** distance
700 travelled in relation to the observation period (cm); **(D)** total distance travelled (cm); **(E)** number of
701 transitions between top and bottom areas and **(F)** latency to enter in top area (s) in nicotine-bathed and
702 control-conditions (n=24 per condition). **(A, C)** Tukey HSD *post-hoc*: *: $p < 0.05$; **: $p < 0.01$; ***,
703 $p < 0.001$. **(B,D,E,F)** Mann-Whitney U-test: ***: $p < 0.001$.

704 **Figure 4.** Spatial location of fish during the novel environment in group. Averaged heatmaps of fish
705 spatial location (n=30 per condition) over the different time periods. From low (blue colour) to high
706 location frequency (red colour).

707 **Figure 5.** Swimming characteristics in the novel environment in group challenge. Mean \pm SEM of **(A)**
708 distance travelled (cm); **(B)** interindividual distance (cm); **(C)** proportion of time spent in different
709 areas (%) for each condition (n=30 control, High TAN, Hyperoxia and Hypoxia). Tukey HSD *post-*
710 *hoc*: different letters indicate significant differences between conditions within observation periods.

711 **Figure 6.** Swimming characteristics in the novel tank diving test. Mean \pm SEM of **(A)** time spent in
712 the top area (s); **(B)** total time spent in the top area (s); **(C)** distance travelled (cm); **(D)** total distance
713 travelled (cm); **(E)** number of transitions between the two areas and **(F)** latency to enter in the top area
714 (s) *post-stress* for all conditions (n=20 control, n=20 High TAN, n=18 Hyperoxia; n=18 Hypoxia).
715 Tukey HSD *post-hoc*: different letters indicate significant differences between conditions.

716 **Figure 7.** Plasma cortisol concentration *post*-stress in the novel environment group challenge (Mean ±
717 SD) for each condition (n=9 control, n=9 High TAN, n=9 Hyperoxia; n=10 Hypoxia). Mann &
718 Whitney U-test: *: p<0.05; ***: p<0.001

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7

