



**HAL**  
open science

## Management of grasslands by mowing versus grazing – impacts on soil organic matter quality and microbial functioning

Aliia Gilmullina, Cornelia Rumpel, Evgenia Blagodatskaya, Abad Chabbi

### ► To cite this version:

Aliia Gilmullina, Cornelia Rumpel, Evgenia Blagodatskaya, Abad Chabbi. Management of grasslands by mowing versus grazing – impacts on soil organic matter quality and microbial functioning. *Applied Soil Ecology*, 2020, 156, pp.1-9. 10.1016/j.apsoil.2020.103701 . hal-03140069

**HAL Id: hal-03140069**

**<https://hal.inrae.fr/hal-03140069>**

Submitted on 22 Aug 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Management of grasslands by mowing versus grazing – impacts on soil organic matter**  
2 **quality and microbial functioning**

3 Gilmullina Aliia<sup>1,2</sup>, Rumpel Cornelia<sup>3</sup>, Blagodatskaya Evgenia<sup>4,5</sup>, Chabbi Abad<sup>1,2\*</sup>

4 <sup>1</sup> UMR P3F, INRA, Lusignan, France

5 <sup>2</sup> UMR ECOSYS, INRA, Thiverval-Grignon, France,

6 <sup>3</sup> Institute of Ecology and Environmental Sciences- Paris (iEES-Paris) UMR CNRS, INRA,

7 UPMC, Thiverval-Grignon, France

8 <sup>4</sup> Agro-Technology Institute, RUDN University, Moscow, Russia

9 <sup>5</sup> Dept. of Soil Ecology, Helmholtz Centre for Environmental Research – UFZ, Halle (Saale),

10 Germany

11 \* - corresponding author, abad.chabbi@inra.fr

12

13

14 **Keywords**

15 Grassland, grazing, mowing, non-cellulosic polysaccharides, lignin, microbial activity

16

17

18 **Highlights**

19 • Grazing and mowing have contrasting effects on soil biogeochemical properties

20 • Grazing promoted more efficient microbial functioning

21 • Mowing leads to more degraded lignin than grazing

22 • Only microbial properties were sensitive to treatment effects in subsurface soil

23

24        **Abstract**

25        Although 30% of the European surface area is covered with grasslands, little is known  
26 about the effect of their management on soil quality and biogeochemical cycling. Here, we  
27 analysed soil from an experimental site in Western France, which had been under either  
28 grazing or mowing regime for 13 years. We aimed to assess the effect of the two  
29 management practices on the biogeochemical functioning of soil system. To this end we  
30 compared soil organic matter (SOM) composition and microbial properties at two soil depths.  
31 We analysed for elemental, lignin and non-cellulosic polysaccharide content and composition  
32 and for microbial biomass, soil microbial respiration and enzyme activities. Our results  
33 showed higher soil organic carbon (SOC) and nitrogen contents in the surface soil under  
34 grazing as compared to mowing. Soil biogeochemical properties differed between grazing  
35 and mowing treatments. In particular, soil under grazing showed lower lignin and higher  
36 microbial biomass. Despite the similar non-cellulosic polysaccharide content under both  
37 treatments, microbial community under mowing was characterised by higher enzyme  
38 production per microbial biomass, leading to more degraded SOM in the mowing system as  
39 compared to grazing. We conclude that grazing and mowing regimes impact differently the  
40 biogeochemical soil functioning. Higher and more diverse carbon input under grazing  
41 compared to mowing may lead to enhanced substrate availability and thus more efficient  
42 microbial functioning, which could favour SOC sequestration through formation of microbial  
43 products.

44

45

46

## 47 **1. Introduction**

48 Dangerous climate change can only be avoided if we succeed to remove CO<sub>2</sub> from the  
49 atmosphere with negative emission technologies (IPCC, 2018). Soil organic carbon (SOC)  
50 sequestration is a nature-based negative emission technology, which may be achievable at  
51 scale through the introduction of sustainable management practices with permanent soil cover  
52 (Rumpel et al., 2018). Permanent grasslands, which in Europe, occupy about 30% of the  
53 agricultural area (Ec.europa.eu, 2018), are responsible for many ecosystem services including  
54 forage for animal production and SOC storage (Havstad et al., 2007; Rumpel et al., 2015).  
55 Biogeochemical cycling in grassland soils can be influenced by a variety of management  
56 practices (Rumpel et al., 2015). The impact of these management practices on processes  
57 impacting soil biogeochemical cycling via soil-plant interactions are poorly understood  
58 (Dignac et al., 2017). These interactions result in contrasting effects of grassland  
59 management on SOC storage potential (Post and Kwon, 2000; Rumpel et al., 2015; Smith et  
60 al., 2008).

61 Grazing and mowing are the most frequently used grassland management practices.  
62 Both practices lead to defoliation (removal of plant aboveground tissue). Defoliation alters  
63 root exudation and C allocation in plants but the direction of these changes was found to be  
64 contrasting (Bazot et al., 2005; Gavrichkova et al., 2010; Medina-Roldán and Bardgett,  
65 2011), related to different climatic and pedological conditions (Pineiro *et al.*, 2010).

66 Defoliation under grazing management is caused by herbivores during several days  
67 (Senapati et al., 2014). This process plays an important role in terms of carbon and nutrient  
68 return (Soussana et al., 2006), because about 50-70% of the ingested biomass is returned to  
69 soil in the form of excreta. In mowing systems, plant biomass is removed in a day with up to  
70 20% of all cut biomass remaining as green litter in form of harvesting losses (Sanaullah et al.,

71 2010). In order to compensate for nutrient exportation during defoliation events, mineral  
72 fertilisers are applied in mowing systems.

73 Due to the different types of biomass returned in the two systems, the quality of  
74 biomass input also varies. Mowing systems receive only plant residues while input in grazing  
75 systems comprises additionally animal depositions. Dung and urine inputs are characterised  
76 by lower C:N ratio, higher amount of easily available compounds (Dungait et al., 2009) and  
77 relatively stable compounds, such as crude proteins and fats (Dungait et al., 2005; Ngo et al.,  
78 2011). Moreover, in grazing systems, there is a return of senescent brown litter, which  
79 contains less N and less soluble compounds compared to the green litter returned as  
80 harvesting losses in mowing systems (Sanaullah et al., 2010).

81 These differences may affect belowground processes (Wilson et al., 2018), SOC  
82 formation and storage (Cotrufo et al., 2015; Rumpel et al., 2015) through their effect on the  
83 soil microbial biomass and its activity (Liang et al., 2017). We therefore hypothesised that the  
84 two management systems may lead to contrasting soil microbial functioning and affect  
85 differently biogeochemical cycling. The effect of management has been analysed up to now  
86 mainly in the first few centimetres of soil, although it has been shown that management can  
87 affect SOC stored down to 2 m depth (Tautges et al., 2019). We thus hypothesised that  
88 grassland management affects SOC below the first centimetres.

89 We focused on an experimental site with grazing and mowing as two contrasting  
90 management practices under similar soil and climatic conditions. We aimed to evaluate the  
91 differences in biogeochemical cycling in soil under the two different management practices at  
92 two depths. To this end we analysed C and N contents, molecular signatures of  
93 polysaccharide and lignin monomers. These variables were compared to the functioning of  
94 the soil microbial communities, assessed by the analyses of soil microbial respiration, growth  
95 kinetic parameters and activity of 9 enzymes as well as microbial biomass C and N.

96

## 97 2. Materials and methods

### 98 2.1. Site description and soil sampling

99 The field experiment is located in Lusignan (southwest of France, 46°25'12,91"N;  
100 0°07'29,35"E) at the national long-term experimental observatory SOERE ACBB  
101 (Agroecosystems, Biogeochemical Cycles and Biodiversity). The mean annual temperature  
102 and precipitation for the period 2006–2010 were 11.2°C and 773 mm, (Senapati et al., 2014).  
103 The landscape is flat. The soil is classified as a Dystric Cambisol with loamy texture (Chabbi  
104 et al., 2009).

105 The current study is focused on two permanent sown grasslands (each of about 3 ha in  
106 size), which were established in 2005 by sowing a mixture of three plant species (*Lolium*  
107 *perenne*, *Festuca arundinacea*, *Dactylis glomerata* L.) in both treatments. In the grazing  
108 system, legume *Trifolium repens* was included in the species mixture but covered only 5% of  
109 grazed paddock in 2017. The mown grassland was cut four times per year with biomass  
110 exported. To replace the exported nutrients, nitrogen (N) fertilizer was applied at rates  
111 between 170 and 380 kg N ha<sup>-1</sup> year<sup>-1</sup> (Puche et al., 2019). Grazing in the grazed paddock  
112 took place from March to December with 50 days per year using 15 to 20 livestock units per  
113 hectare. Grazed grasslands received less nitrogen fertilization (60-150 kg N ha<sup>-1</sup> year<sup>-1</sup>, Puche  
114 et al., 2019) because nitrogen losses were additionally returned by dung and urine and  
115 through the presence of the leguminous species. In order to compare the treatments at similar  
116 N status, fertilizer application rates were adjusted to maintain the Nitrogen Nutrition Index  
117 between 0.9 and 1.0 for both treatments, close to non-limiting nitrogen nutrition to near  
118 maximum plant production (Senapati et al., 2016). Moreover, both sites were limed regularly  
119 in order to neutralize acid pH.

120 Due to the large land requirements (3 ha for plots with cows), it was not possible to  
121 establish and maintain a completely replicated field experiment including grazing treatment  
122 for several decades. Limitations to generalization of the treatment effects due to the absence  
123 of replication of the experiments were limited by choosing homogenous flat areas in close  
124 proximity with similar land use history, climate, and soil type. Moreover, we carried out  
125 baseline measurements, in form of geostatistical evaluation of the soils SOC and N contents  
126 and included initial SOC stocks as a co-variate. These data showed that both plots were  
127 significantly different in initial SOC and N contents (n=28). The SOC contents on mowing  
128 plots varied between 9.9 and 13.7 mg g<sup>-1</sup> (average 12.0 ± 1.0 mg g<sup>-1</sup>), while under grazing it  
129 was between 11.9 and 19.1 mg g<sup>-1</sup> (average 14.8 ± 1.5 mg g<sup>-1</sup>). N contents varied between 1.0  
130 and 1.4 mg g<sup>-1</sup> (average 1.2 ± 0.1 mg g<sup>-1</sup>) under mowing, while under grazing the values  
131 ranged between 1.2 and 1.9 mg g<sup>-1</sup> (average 1.5 ± 0.1 mg g<sup>-1</sup>). These previous analyses  
132 indicated on average non-significant differences in SOC stock changes between grazing and  
133 mowing after nine years of treatment (Crème et al., 2020). The study also showed  
134 partitioning of the field into different zones with SOC gain and loss (Crème et al., 2020; Fig  
135 S1, Supplementary materials).

136

137 Five replicated soil samples were taken from each of the two zones, giving a total of 10  
138 replicated field samples per plot. Sampling took place in November 2017, 2 weeks and 5  
139 months after the last grazing and mowing events, accordingly. The shortest distance between  
140 samples was 25 m. Soil samples were collected with a mechanical auger (5cm Ø, 30cm) at  
141 two depths: 0-10 cm (surface soil) and 20-30 cm (subsurface soil) giving in total 40 samples.  
142 All samples were sieved through a 2-mm mesh. Thereafter, half of the samples was air-dried  
143 and ground for measurements of physicochemical analysis and the other half was stored at  
144 4°C before microbial analyses. Because of dry field conditions prior to measurements of



145 microbiological analysis, soil samples were moistened by distilled water to adjust 50% of  
146 WHC and pre-incubated at 22 °C for 7 days.

## 147 2.2. *Soil general properties*

148 Soil pH (H<sub>2</sub>O) was measured in a soil:water suspension (1:2.5 weight/volume). Soil  
149 organic carbon (SOC), nitrogen (N) and stable isotopes (<sup>13</sup>C and <sup>15</sup>N) contents were measured  
150 using a CHN auto-analyser (Flash EA, Thermo Electron Corporation, Bremen, Germany)  
151 coupled with an isotope ratio mass spectrometer. The isotopic ratios were calculated relative  
152 to the Pee Dee Belemnite Standard (PDB) for C and relative to atmospheric N<sub>2</sub> for nitrogen.

153

## 154 2.3. *Soil chemical properties*

155 Lignin was analysed by the alkaline cupric oxide (CuO) oxidation method (Hedges and  
156 Ertel, 1982; Kögel and Bochter, 1985). Briefly, oxidation was carried out under alkaline  
157 conditions (2M NaOH) at 172 °C for 4 hours using 500 mg of air-dried soil, 250 mg of CuO,  
158 50 mg of ammonium ferrous hexahydrate and 50 mg of glucose. After cooling, samples were  
159 acidified with 5 M HCl and left overnight for humic acid precipitation. Removal of humic  
160 acids was conducted through centrifugation (10 min at 10000 rpm) and followed by  
161 extraction of phenolic oxidation products with C18 reversed phase columns. The phenols  
162 were derivatized with BSTFA and quantified as trimethylsilyl derivatives by gas  
163 chromatography with a HP gas chromatograph (HP GC 6890) equipped with a flame  
164 ionization detector and a SGE BPX-5 column (50 m length, 0.25 mm inner diameter, 0.32 µm  
165 coating). Samples were injected in split mode (1:10). The GC oven temperature was  
166 programmed at 100 °C for 2 min, then increased from 100 to 172 °C at a heating rate of 8 °C  
167 min<sup>-1</sup>, from 172 to 184 °C at 4 °C min<sup>-1</sup>, and from 184 to 300 °C at a rate of 10°C min<sup>-1</sup>.

168 The internal standard ethylvanillin was added before the purification step to quantify  
169 lignin recovery and the quantification standard phenylacetic acid was added before GC  
170 analyses.

171 The total lignin content ( $\text{mg g}^{-1}$  dry soil) of the sample was determined as the sum of  
172 phenolic oxidation products: vanillyl (V), syringyl (S) and p-coumaryl (C) in their acid (Ac),  
173 aldehyde (Al) and ketone forms. Lignin content was also expressed as lignin content per SOC  
174 ( $\text{mg g}^{-1}$  SOC). Lignin decomposition was assessed by the ratios of S, C to V and (Ac/Al)  
175 ratios of V and S, which generally indicate decomposition state (Thevenot et al., 2010).

176 Non-cellulosic polysaccharides of plant and microbial origin (Kögel-Knabner, 2002)  
177 were determined by gas chromatography after trifluoroacetic acid (TFA) hydrolysis and  
178 reduction-acetylation using a method introduced by Rumpel and Dignac (2006) and modified  
179 by Eder et al. (2010). The analysis was performed using 700 mg of soil samples. Briefly,  
180 hydrolysis of non-cellulose polysaccharides was carried out at  $105^{\circ}\text{C}$  for 4 h with 10 ml of 4  
181 M TFA. Thereafter, Myo-inositol was added as quantification standard to account for the  
182 losses during the purification procedure. Removal of soil was performed by filtration through  
183 glass fibre filters (Whatman GF/C  $0.45\ \mu\text{m}$ ). Afterwards, TFA was evaporated using a  
184 centrifugal Evaporator EZ-2 ENVI at  $35^{\circ}\text{C}$  for 4 hours and dry samples were left overnight in  
185 the freezer. Thereafter, dry samples were dissolved in 0.5 ml of  $\text{H}_2\text{O}$  followed by the addition  
186 of 0.9 EDTA in order to avoid co-precipitation of organic material with metal oxides and  
187 hydroxides (Eder et al., 2010). One mL sodium borohydride ( $\text{NaBH}_4$ ) in dimethylsulfoxide  
188 ( $20\ \text{g L}^{-1}$ ) was added for reduction of polysaccharide monomers into alditol forms and kept  
189 at  $40^{\circ}\text{C}$  for 1.5 hours. Then, acetylation was conducted by addition of 0.2 mL acetic acid, 2  
190 mL of acetic anhydride and 0.2 mL Methylimidazole. Acetylated alditols were extracted by 1  
191 ml of dichloromethane and quantified with a HP GC 6890 gas chromatograph equipped with  
192 a flame ionization detector. Separation was achieved with a 60 m fused silica capillary

193 column (SGE BPX 70, 0.32 mm internal diameter, 0.25 mm film thickness) under the  
194 following temperature program: 170 to 250 °C at 8 °C.min<sup>-1</sup>, followed by 12 min at 250 °C  
195 (isothermal). Helium was used as the carrier gas at a flow rate of 1.0 mL min<sup>-1</sup>. The injector  
196 was kept at 250 °C and the detector at 260 °C. The non-cellulosic polysaccharides content of  
197 soil samples was determined as the sum of monosaccharides: C5 (pentoses: xylose, ribose  
198 and arabinose), C6 (hexoses: glucose, galactose and mannose), and desoxyC6  
199 (desoxyhexoses: fucose and rhamnose) (Kögel-Knabner, 2002). A higher C6/C5 ratio  
200 generally indicates higher contribution of microbial sugars.

#### 201 2.4. Soil microbial properties

202 Microbial biomass C (MBC) and nitrogen (MBN) were determined by the chloroform  
203 fumigation-extraction method (Vance et al., 1987). Dissolved organic C and N in fumigated  
204 and non-fumigated soil samples were extracted in 0.05 M K<sub>2</sub>SO<sub>4</sub> and were measured using a  
205 multi C/N analyzer (multi C/N analyser 2100S, Analytic Jena). MBC and MBN were  
206 calculated with a conversion factor of 0.45 (Jenkinson et al., 2004). For measuring soil  
207 microbial respiration (SMR) a half gram of soil sample was placed in 2 ml Eppendorf tubes.  
208 The CO<sub>2</sub> efflux was trapped in 3 ml of 0.1 M NaOH and determined by conductometry. The  
209 metabolic quotient (qCO<sub>2</sub>), reflecting decomposition activity (Anderson, 2003; Anderson and  
210 Domsch, 1993), was calculated as soil microbial respiration expressed per gram of microbial  
211 biomass carbon:  $qCO_2 = SMR/MBC$  (μg CO<sub>2</sub>-C g<sup>-1</sup> MBC h<sup>-1</sup>).

212 We used microbial growth kinetics technique as an approach to estimate microbial  
213 biomass activity state (Blagodatskaya and Kuzyakov, 2013). This approach is based on soil  
214 respiratory response to unlimited nutrient amendments (Panikov and Sizova, 1996). For this  
215 purpose, soil samples were treated with a solution (0.1 ml per g of dw soil) containing per g  
216 soil: 10 mg glucose, 1.9 mg (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 3.8 mg MgSO<sub>4</sub>\*7H<sub>2</sub>O, 0.11 mg K<sub>2</sub>HPO<sub>4</sub> and 1.68  
217 mg KH<sub>2</sub>PO<sub>4</sub> for surface soil samples and 10 mg glucose, 1.9 mg (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 3.8 mg

218 MgSO<sub>4</sub>\*7H<sub>2</sub>O, 0.53 mg K<sub>2</sub>HPO<sub>4</sub> and 1.35 mg KH<sub>2</sub>PO<sub>4</sub> for subsurface soil samples. The  
219 amount of mineral salts was preliminary selected in order to avoid soil pH change of more  
220 than 0.1 units after addition. For active microbial biomass (AMB) and specific growth rate  
221 calculation, the results of substrate induced respiration rate were fitted with a model proposed  
222 by Panikov and Sizova (Panikov and Sizova, 1996; Wutzler et al., 2012):

$$223 \quad CO_2(t) = A + B * \exp(\mu * t) \quad (1)$$

224 In order to estimate catabolic (decomposition) activity in regards to specific substrates  
225 in soil, we measured extracellular enzyme activity using the fluorometric technique (Koch et  
226 al., 2007; Marx et al., 2005; Razavi et al., 2015). Nine types of fluorogenic substrates based  
227 on 4-methylumbelliferone (MUF) and 7-amino-4-methylcoumarin (AMC) were used: (1)  
228 MUF- $\alpha$ -D-glucopyranoside for  $\alpha$ -glucosidase, (2) MUF- $\beta$ -D-glucopyranoside for  $\beta$ -  
229 glucosidase, (3) MUF- $\beta$ -D-galactopyranoside for  $\beta$ -galactosidase, (4) MUF- $\beta$ -D-  
230 xylopyranoside for  $\beta$ -xylosidase, (5) MUF- $\beta$ -D-cellobioside for  $\beta$ -cellobiohydrolase, (6)  
231 MUF-N-acetyl- $\beta$ -D-glucosamide for chitinase, (7) Leucine-AMC for leucine aminopeptidase,  
232 (8) MUF-heptanoate for lipase and (9) MUF-phosphate for phosphatase. Saturation  
233 concentrations of fluorogenic substrates were determined in preliminary experiments and  
234 comprised 20  $\mu$ mol g<sup>-1</sup> soil for all enzymes except lipase with 60  $\mu$ mol g<sup>-1</sup> soil. Briefly, a  
235 water extract of soil (1:10) was homogenised by low-energy sonication (40 J s<sup>-1</sup> output  
236 energy) for 60 s. Thereafter 50 ml of the soil suspension were added to 150 ml of each  
237 substrate solution in a 96-well microplate. Fluorescence was measured at an excitation  
238 wavelength of 355 nm and an emission wavelength of 460 nm (Victor3 1420-050 Multilabel  
239 Counter, PerkinElmer, USA).

240

## 241 2.5. *Statistical analysis*

242 All results are presented as arithmetic means with standard error. The statistical  
243 analyses were conducted by using R (Studio Version 1.1.447). We used analyses of  
244 covariance (ANCOVA) to test treatment effect, depth effect and their interactions using  
245 chemical and microbial variables with initial SOC stock as a covariate. The initial SOC  
246 stocks data was obtained from exactly the same sampling points based on the geostatistical  
247 evaluation before the beginning of the experiment. This procedure allowed us to account for  
248 the lack of field replication by taking into account the original difference between the grazed  
249 and mowed plots. In order to obtain better understanding of treatment and depth effects, non-  
250 transformed data (except C and N contents) were subjected to Principal Component Analysis  
251 (PCA) and the results were also tested by ANCOVA with initial SOC stock as a covariate.  
252 The equations were fitted by non-linear regression, using Model Maker-3 software (SB  
253 technology Ltd.).

254

## 255 **3. Results**

### 256 3.1. *Soil properties*

257 Soil physicochemical properties are presented in Table 1. The pH was not controlled by  
258 initial SOC stock ( $P=0.70$ ). Lower pH was found for both treatments in surface soil compared  
259 to subsurface soil, although, the lowest pH value was recorded in surface soils under mowing  
260 treatment. SOC and N contents were nearly twice as high in the surface soil compared to the  
261 subsurface soil under both treatments. Even if SOC and N contents were dependant on initial  
262 SOC stock ( $P=0.03$  and  $0.02$ , respectively), there were still significant effects of depth  
263 ( $P<0.001$ ) and treatment ( $P<0.001$ ) after correction by using it as covariate. C:N ratio differed  
264 only between soil depths ( $P<0.001$ ) showing slightly higher C:N ratios in surface soils as

265 compared to subsurface soils.  $\delta^{13}\text{C}$  followed the same pattern as SOC content and the highest  
266 enrichment was recorded for the surface soil of the grazing treatment (depth effect  $P < 0.001$   
267 and treatment effect  $P = 0.002$ ). The  $\delta^{15}\text{N}$  did not differ between the treatments and was  
268 enriched in surface soils compared to subsurface soils.

269

### 270 3.2. Specific SOM compounds

271 Non-cellulosic polysaccharide (NCP) content was not affected by initial SOC stock  
272 ( $P = 0.52$ ) and there was treatment  $\times$  depth interaction (Table 2,  $P < 0.001$ ). Grazing resulted in  
273 higher NCP content in both depths compared to mowing. The NCP content per SOC ( $\text{mg g}^{-1}$   
274 soil C) was affected only by depth ( $P = 0.002$ ). Concerning the NCP monomers ratio, C6/C5  
275 and Man/Xyl ratios were controlled by initial SOC stock ( $P = 0.03$  and  $0.04$ , respectively),  
276 consequently, after ANCOVA application the treatment effect was vanished while depth  
277 effect remained significant (Table 2,  $P < 0.001$ ). All NCP monomers ratios were higher in  
278 subsurface soil compared to surface soil under both treatments.

279 Lignin content was not affected by initial SOC stock correction ( $P = 0.82$ ), so the effects  
280 of depth ( $P < 0.001$ ), treatment ( $P < 0.001$ ) and their interactions ( $P = 0.04$ ) remained significant  
281 (Table 3). Lignin content was higher in surface soils than in subsurface soils and was higher  
282 under grazing compared to mowing as well. Correcting for initial SOC stock caused the  
283 elimination of all effects on lignin content per SOC content. The C/V ratio was affected only  
284 by depth ( $P = 0.006$ ) showing higher values in surface soils than in subsurface soils. The S/V  
285 ratio was greater under grazing treatment than under mowing treatment at both depths even  
286 after correction by initial SOC stock (Table 3,  $P = 0.01$ ). Based on the presence of treatment  $\times$   
287 depth interaction  $(\text{Ac/Al})_V$  and  $(\text{Ac/Al})_S$  ratios were lower in the surface soil of grazing  
288 treatment as compared to mowing treatment ( $P < 0.001$ ). In contrast to surface soils, treatments  
289 did not show any effects on these lignin ratios in subsurface soils.

290

### 291 3.3. Soil microbial properties

292 The soil microbial respiration (SMR) ranged between 0.2 and 0.7  $\mu\text{g CO}_2 \text{ -C g}^{-1} \text{ h}^{-1}$   
293 with highest values in the surface soil under grazing treatment (Fig. 1A). After correcting for  
294 initial SOC stock, treatment  $\times$  depth interaction effect on SMR was significant (Table S1,  
295 Supplementary materials,  $P < 0.001$ ). Soil microbial respiration per SOC was around 33%  
296 higher in the surface soil under grazing as compared to mowing (Fig. 1B). In contrast, it was  
297 greater in the subsurface soil under mowing than under grazing treatment. Including initial  
298 SOC stock as covariate resulted only in significant effect of treatment  $\times$  depth interaction on  
299 soil microbial respiration per SOC (Table S1, Supplementary materials,  $P = 0.004$ ).

300 MBC per SOC was highest in the surface soil under grazing (20  $\mu\text{g C mg}^{-1} \text{ SOC}$ , Fig.  
301 1C). Mowing treatment resulted in two times lower MBC per SOC in the surface soil  
302 compared to grazing treatment. After correction for initial SOC stock, treatment ( $P < 0.001$ )  
303 and their interaction ( $P < 0.001$ ) showed significant effects on  $q\text{CO}_2$ . Mowing treatment  
304 resulted in higher  $q\text{CO}_2$  at both depths as compared to grazing treatment (Fig. 1D,  $P = 0.02$ ).

305 Microbial C:N ratio ranged between 4.9 and 6.4. It was affected by treatments in all  
306 depths showing higher values under mowing (Fig. 2A). After taking into account initial SOC  
307 stock, the treatment effect was still significant (Table S1, Supplementary materials,  $P < 0.001$ ).  
308 Active microbial biomass was also higher under mowing at both depths compared to grazing  
309 treatment (Fig. 2B,  $P = 0.02$ ). The highest specific microbial growth rate (Fig. 2C) was  
310 recorded in subsurface soils without difference between treatments. But in surface soils, the  
311 specific microbial growth rate was higher under grazing than under mowing (Fig. 2C).  
312 However, ANCOVA with initial SOC stock as covariate decreased the significance treatment  
313 effects ( $P = 0.05$ ) on specific microbial growth rate but increased the depth effect Table S1,  
314 Supplementary materials,  $P < 0.001$ ).

315 Treatment effect on absolute enzyme activities is presented only for leucine  
316 aminopeptidase in surface soil and chitinase and phosphatase in subsurface soil (Fig S2,  
317 Supplementary materials). When the initial SOC stock was used as covariate, treatment  
318 differences between enzyme activities per MBC were observed for all enzymes (except  
319 leucine aminopeptidase) in surface soil. Soil under mowing treatment showed 2-2.5 times  
320 higher enzyme activity per MBC under mowing compared to soil under grazing (Fig. 3). The  
321 differences between treatments were more pronounced in surface soil for activities of  
322 chitinase,  $\beta$ -galactosidase,  $\beta$ -glucosidase and phosphatase (Table S1, Supplementary  
323 materials).

324

#### 325 3.4. *Principal component analysis*

326 Principal component analysis based on SOC normalised data of all soil properties  
327 showed that the first two factors explained 54.4% of the variation (Fig. 4). The first  
328 component (Dim1) was related to microbial functioning, as it was strongly associated with  
329 the soil microbial properties MBC and MBN per SOC in negative direction. The positive  
330 direction was related to the lipase activity per MBC. The second component (Dim2) was  
331 explained by variables related to polysaccharides. It was positively correlated with enzymes  
332 participating in polysaccharide degradation and negatively with polysaccharide ratios. The  
333 clustering of samples allowed to separate surface soil and subsurface soil samples along both  
334 axes, while surface soil samples were additionally separated by treatments along the first axes  
335 (Fig. 4). Subsurface soil samples were differentiated from surface soil by high neutral  
336 polysaccharide monomer ratio, low enzymes activities per MBC, MBC and MBN per SOC.  
337 Treatments in surface soil were separated by C- and N-cycle enzyme activity and MBC and  
338 MBN per SOC. We also applied ANCOVA with initial SOC stock as a covariate on new  
339 PCA coordinates which resulted in significant effects of treatment, depth and their



340 interaction. Treatment effect was more pronounced on Dim1, while Dim2 was more affected  
341 by depth.

342

## 343 **4. Discussion**

### 344 *4.1. Effect of grazing and mowing on chemical properties of surface soil*

345 Since the primary factor of SOM formation is organic matter input (Fujisaki et al.,  
346 2018; Kögel-Knabner, 2002), higher SOC and N contents in the surface soil under grazing  
347 system might be explained by greater C input compared to mowing systems. This was shown  
348 through ecosystem flux measurements at these plots (Senapati et al., 2014). Moreover, dung  
349 return comprising about 50-80% of plant biomass could also favour higher SOC and N  
350 content under grazing (Soussana et al., 2006). Even if mowing leads to some biomass input in  
351 the form of plant material lost during grass removal (Sanaullah et al., 2010), the amount is  
352 not enough to reach a similar input level than under grazing. Additionally, the lower pH  
353 under mowing could contribute to indirect losses of SOC via changing C cycle and microbial  
354 functioning (Kemmitt et al., 2006). Consequently, our results suggest that temperate loamy  
355 soil under grazing is more prone to higher SOC contents when compared to mowing.

356 With regards to the biogeochemical composition of SOC, we did not find any  
357 differences in non-cellulosic polysaccharide concentrations. These results are in agreement  
358 with other studies showing that the soils' polysaccharide content is more or less stable and  
359 even plant removal does not have a strong effect on the total polysaccharide concentrations  
360 (Marchus et al., 2018). Soil lignin content, in contrast, was lower under grazing than mowing.  
361 As lignin is a biomarker for plant-derived organic matter and more difficult to decompose,  
362 because it requires a specific enzyme system (Buswell et al., 1987; Thevenot et al., 2010),  
363 lower exportation of plant biomass and lignin input via dung deposition in soil under grazing  
364 would suggest the opposite trend. However, dung contains only small amounts of lignin

365 (Dungait et al., 2005), which is relatively instable being degraded during one year (Dungait et  
366 al., 2008). All lignin parameters (except the C/V ratio) suggested that lignin was less  
367 degraded in the grazing than the mowing system. More acid pH in fertilised mowing systems  
368 could have favoured the activity of lignin-degrading fungi (Couto et al., 2006). In mowing  
369 systems microbial activity is fuelled exclusively by plant litter, whereas in grazing systems  
370 organic matter input is supplied also by animal depositions. We hypothesise that this could  
371 lead to contrasting quantitative lignin inputs, but could also impact its decomposition. Our  
372 data show that lignin degradation in the mowing system is slower and less complete than in  
373 the grazing system, leading to accumulation of partially degraded lignin molecules (Filley et  
374 al., 2006). Therefore, lignin in the mowing system was characterised by a higher state of  
375 degradation and at the same time its contribution to SOC was higher as compared to the  
376 grazing system.

#### 377 4.2. *Effect of grazing treatment on biological properties of surface soil*

378 Higher maturity and sustainability of the grazing system was shown by higher MBC per  
379 SOC together with a lower  $qCO_2$  (Anderson and Domsch, 2010). Higher  $qCO_2$  in the mowing  
380 system indicates that the microbial communities were less efficient and respired more C to  
381 maintain metabolic activity as compared to those under grazing (Anderson, 2003).  
382 Microorganisms are the main SOM decomposers leading to release of greenhouse gases and  
383 nutrients in natural as well as in managed soils (Bardgett et al., 2008; Gougoulias et al.,  
384 2014). This is particularly relevant for grazed pastures. Higher soil microbial respiration and  
385 microbial  $CO_2$  –C per unit SOC (soil microbial respiration per SOC) in the grazing system  
386 was probably related to dung input with a huge amount of easily available compounds (Chu  
387 et al., 2007; Marinari et al., 2000).

388 Contrary to our expectations, absolute enzyme activity did not differ among the  
389 treatments, even after normalisation by SOC. A treatment effect was only observed after

390 normalisation by MBC, which expresses microbial activity in terms of enzyme production.  
391 The enzymatic activities per MBC were higher in the mowing system as compared to the  
392 grazing one, indicating that microorganisms in mowed soil produced enzymes more actively  
393 than those under grazing. Microbial communities in the mowing system stayed active and  
394 were investing in enzyme production probably to adapt to less decomposable organic  
395 materials with higher lignin contents (see above). This maintenance of active state requires a  
396 lot of energy, consequently, it could change C-cycling rates and decomposition of SOM  
397 (Schimel and Schaeffer, 2012; Wang et al., 2014).

398         Microbial communities in the mowed soil are probably characterised by a higher  
399 contribution of fungi than those of the grazed soil because we recorded a higher C:N ratio of  
400 the microbial biomass (Joergensen and Emmerling, 2006) and more acid pH. Lower specific  
401 growth rates in the mowing system may indicate relative domination of K-strategists in the  
402 microbial community, which are more adapted to nutrient poor conditions (Strickland and  
403 Rousk, 2010; Xu et al., 2017) and the decomposition of specific substances, such as plant  
404 material containing high amounts of biopolymers (Fontaine et al., 2003). As illustrated by  
405 lower enzyme activity per MBC, microorganisms in the grazing system invested less energy  
406 for the degradation of complex compounds than those of the mowing system, most probably  
407 because of higher availability of easily decomposable substrates. These conditions favour r-  
408 strategists (Fierer et al., 2007; Xu et al., 2017) and thus stimulate microbial activity, as shown  
409 by higher MBC per SOC and higher soil microbial respiration under the grazing as compared  
410 to mowing system. As a consequence, the biogeochemical soil functioning under the two  
411 management practices is quite different. This may affect significantly SOM formation, which  
412 is favoured in systems with intensive microbial processing of C input (Kallenbach et al.,  
413 2016; Liang et al., 2017) thus corroborating the high SOC contents observed under grazing.

414 4.3. *Less pronounced treatment effects in subsurface soil*

415 Treatment effects on soil properties were less pronounced in subsurface soil compared  
416 to surface soil. Enhanced leaching and activity of soil fauna (Bohlen et al., 2004; Rumpel and  
417 Kögel-Knabner, 2011) promote nutrient transport to subsurface soil under grazing which  
418 resulted in higher SOC and N contents in subsurface soil under grazing than the one under  
419 mowing. Treatment effects in the subsurface soil were neither observed for non-cellulosic  
420 polysaccharide content and origin nor for lignin content or its degradation status. Since  
421 lignins are typical indicators of plant input (Kögel-Knabner, 2002), this could indicate that  
422 grazing and mowing have only small effects on plant rooting behaviour at lower depths.

423 On the other hand, the treatment effects on MBC and MBN was also observable in  
424 subsurface soil. Soil microbial respiration did not differ between the treatments but microbial  
425 CO<sub>2</sub> –C per SOC and qCO<sub>2</sub> were higher in the subsurface soil under mowing indicating that  
426 the microbial communities used C inefficiently, similarly to surface soil. Higher  
427 galactosidase activity in the subsurface soil of the mowing treatment is related to higher  
428 contribution of galactose monomers in grass roots compared to grass leaves (Schädel et al.,  
429 2010). As lipase is hydrolysing triglycerides, higher lipase activity in the subsurface soil  
430 indicates accumulation of lipid compounds at depth, which probably serve as C source for  
431 microorganisms under C-limiting conditions (Heitkötter et al., 2017).

432 The absence of treatment separation for the subsurface soils on the PCA plot might  
433 indicate that in deeper soil probably more time is required to make treatment effects  
434 observable. It was interesting to note that chemical properties related to SOM composition  
435 were not sensitive to treatment effects in the subsurface soil, whereas microbial properties  
436 were. This is in agreement with other studies, which showed that microbial properties are  
437 most sensitive to changes introduced by management activities (Allison and Martiny, 2008;  
438 Bending et al., 2004).

439

## 440 **5. Conclusions**

441 In this study we investigated the effect of grazing and mowing treatments on soil  
442 biogeochemical and microbial properties. Our data indicated significant differences in the soil  
443 organic matter composition as well as microbial functioning of the two treatments. Both plots  
444 were also characterized by contrasting SOC contents and pH values. The grazing system was  
445 characterized by (1) more efficient microbial community and (2) less decomposed organic  
446 matter as compared to the mowing system. We conclude that the harvesting regime by  
447 grazing or mowing affects the biogeochemical functioning of grassland soils. Even though  
448 both systems are favorable to SOC storage, grazing might be preferable to mowing because it  
449 leads to better substrate quality and more efficient microbial functioning. Although SOM  
450 changes were only evident in surface soil, microbial properties suggest that these processes  
451 are also occurring in subsurface soil.

452

## 453 **6. Conflicts of interests**

454 We state that there is no conflict of interests.

455

## 456 **7. Acknowledgements**

457 The research leading to these results has received funding principally from the New  
458 Zealand Government to support the objectives of the Livestock Research Group of the Global  
459 Research Alliance on Agricultural Greenhouse Gases - Grants SOW12-GPLER-LCR-PM  
460 (Proposal ID 16949-15 LCR), the ANR (ANR-11-INBS-0001), the CNRS-INSU and the  
461 INRA/Region Nouvelle-Aquitaine. We are grateful to Xavier Charrier, Jean-François  
462 Bouhiron, Jerome Chargelegue, Marie-Laure Decau for help with site access, organization,  
463 and field sampling and to Karin Schmidt and Valerie Pouteau for laboratory assistance. Two

464 anonymous reviewers are acknowledged for their valuable comments on an earlier version of  
465 the manuscript.

466

## 467 **8. References**

- 468 Abdalla, M., Hastings, A., Chadwick, D.R., Jones, D.L., Evans, C.D., Jones, M.B., Rees,  
469 R.M., Smith, P., 2018. Critical review of the impacts of grazing intensity on soil organic  
470 carbon storage and other soil quality indicators in extensively managed grasslands.  
471 *Agric. Ecosyst. Environ.* 253, 62–81. <https://doi.org/10.1016/j.agee.2017.10.023>
- 472 Allison, S.D., Martiny, J.B.H., 2008. In the Light of Evolution, In the Light of Evolution.  
473 National Academies Press, Washington, D.C. <https://doi.org/10.17226/12501>
- 474 Anderson, T.-H., 2003. Microbial eco-physiological indicators to asses soil quality. *Agric.*  
475 *Ecosyst. Environ.* 98, 285–293. [https://doi.org/10.1016/S0167-8809\(03\)00088-4](https://doi.org/10.1016/S0167-8809(03)00088-4)
- 476 Anderson, T.-H., Domsch, K.H., 2010. Soil microbial biomass: The eco-physiological  
477 approach. *Soil Biol. Biochem.* 42, 2039–2043.  
478 <https://doi.org/10.1016/j.soilbio.2010.06.026>
- 479 Anderson, T.-H., Domsch, K.H., 1993. The metabolic quotient for CO<sub>2</sub> (qCO<sub>2</sub>) as a specific  
480 activity parameter to assess the effects of environmental conditions, such as ph, on the  
481 microbial biomass of forest soils, *Soil Biology and Biochemistry*. Pergamon.  
482 [https://doi.org/10.1016/0038-0717\(93\)90140-7](https://doi.org/10.1016/0038-0717(93)90140-7)
- 483 Bardgett, R.D., Freeman, C., Ostle, N.J., 2008. Microbial contributions to climate change  
484 through carbon cycle feedbacks. *ISME J.* 2, 805–814.  
485 <https://doi.org/10.1038/ismej.2008.58>
- 486 Bazot, S., Mikola, J., Nguyen, C., Robin, C., 2005. Defoliation-induced changes in carbon  
487 allocation and root soluble carbon concentration in field-grown *Lolium perenne* plants:  
488 Do they affect carbon availability, microbes and animal trophic groups in soil? *Funct.*  
489 *Ecol.* 19, 886–896. <https://doi.org/10.1111/j.1365-2435.2005.01037.x>
- 490 Bending, G.D., Turner, M.K., Rayns, F., Marx, M.-C., Wood, M., 2004. Microbial and  
491 biochemical soil quality indicators and their potential for differentiating areas under  
492 contrasting agricultural management regimes. *Soil Biol. Biochem.* 36, 1785–1792.  
493 <https://doi.org/10.1016/J.SOILBIO.2004.04.035>
- 494 Blagodatskaya, E., Kuzyakov, Y., 2013. Active microorganisms in soil: Critical review of  
495 estimation criteria and approaches. *Soil Biol. Biochem.*

496 <https://doi.org/10.1016/j.soilbio.2013.08.024>

497 Bohlen, P.J., Pelletier, D.M., Groffman, P.M., Fahey, T.J., Fisk, M.C., 2004. Influence of  
498 Earthworm Invasion on Redistribution and Retention of Soil Carbon and Nitrogen in  
499 Northern Temperate Forests. *Ecosystems* 7, 13–27. [https://doi.org/10.1007/s10021-003-](https://doi.org/10.1007/s10021-003-0127-y)  
500 [0127-y](https://doi.org/10.1007/s10021-003-0127-y)

501 Buswell, J.A., Odier, E., Kirk, T.K., 1987. Lignin Biodegradation. *Crit. Rev. Biotechnol.* 6,  
502 1–60. <https://doi.org/10.3109/07388558709086984>

503 Chabbi, A., Kögel-Knabner, I., Rumpel, C., 2009. Stabilised carbon in subsoil horizons is  
504 located in spatially distinct parts of the soil profile. *Soil Biol. Biochem.* 41, 256–261.  
505 <https://doi.org/10.1016/j.soilbio.2008.10.033>

506 Chu, H., Lin, X., Fujii, T., Morimoto, S., Yagi, K., Hu, J., Zhang, J., 2007. Soil microbial  
507 biomass, dehydrogenase activity, bacterial community structure in response to long-term  
508 fertilizer management. *Soil Biol. Biochem.* 39, 2971–2976.  
509 <https://doi.org/10.1016/J.SOILBIO.2007.05.031>

510 Cotrufo, M.F., Soong, J.L., Horton, A.J., Campbell, E.E., Haddix, M.L., Wall, D.H., Parton,  
511 W.J., 2015. Formation of soil organic matter via biochemical and physical pathways of  
512 litter mass loss. *Nat. Geosci.* 8, 776–779. <https://doi.org/10.1038/ngeo2520>

513 Couto, S.R., Moldes, D., Sanromán, M.A., 2006. Optimum stability conditions of pH and  
514 temperature for ligninase and manganese-dependent peroxidase from *Phanerochaete*  
515 *chrysosporium*. Application to in vitro decolorization of Poly R-478 by MnP. *World J.*  
516 *Microbiol. Biotechnol.* 22, 607–612. <https://doi.org/10.1007/s11274-005-9078-0>

517 Dignac, M.-F., Derrien, D., Barré, P., Barot, S., Cécillon, L., Chenu, C., Chevallier, T.,  
518 Freschet, G.T., Garnier, P., Guenet, B., Hedde, M., Klumpp, K., Lashermes, G., Maron,  
519 P.-A., Nunan, N., Roumet, C., Basile-Doelsch, I., 2017. Increasing soil carbon storage:  
520 mechanisms, effects of agricultural practices and proxies. A review. *Agron. Sustain.*  
521 *Dev.* 37, 14. <https://doi.org/10.1007/s13593-017-0421-2>

522 Dungait, J.A.J., Bol, R., Bull, I.D., Evershed, R.P., 2009. Tracking the fate of dung-derived  
523 carbohydrates in a temperate grassland soil using compound-specific stable isotope  
524 analysis. *Org. Geochem.* 40, 1210–1218.  
525 <https://doi.org/10.1016/j.orggeochem.2009.08.001>

526 Dungait, J.A.J., Bol, R., Evershed, R.P., 2005. Quantification of dung carbon incorporation in  
527 a temperate grassland soil following spring application using bulk stable carbon isotope  
528 determinations. *Isotopes Environ. Health Stud.* 41, 3–11.  
529 <https://doi.org/10.1080/10256010500053516>

530 Dungait, J.A.J., Stear, N.A., van Dongen, B.E., Bol, R., Evershed, R.P., 2008. Off-line  
531 pyrolysis and compound-specific stable carbon isotope analysis of lignin moieties: a  
532 new method for determining the fate of lignin residues in soil. *Rapid Commun. Mass*  
533 *Spectrom.* 22, 1631–1639. <https://doi.org/10.1002/rcm.3454>

534 Ec.europa.eu, 2018. Share of main land types in utilised agricultural area (UAA) by NUTS 2  
535 regions [WWW Document]. URL <https://ec.europa.eu/eurostat/data/database> (accessed  
536 1.23.19).

537 Eder, E., Spielvogel, S., Kölbl, A., Albert, G., Kögel-Knabner, I., 2010. Analysis of  
538 hydrolysable neutral sugars in mineral soils: Improvement of alditol acetylation for gas  
539 chromatographic separation and measurement. *Org. Geochem.* 41, 580–585.  
540 <https://doi.org/10.1016/J.ORGGEOCHEM.2010.02.009>

541 Eivazi, F., Tabatabai, M.A., 1990. Factors affecting glucosidase and galactosidase activities  
542 in soils. *Soil Biol. Biochem.* 22, 891–897. [https://doi.org/10.1016/0038-0717\(90\)90126-](https://doi.org/10.1016/0038-0717(90)90126-K)  
543 [K](https://doi.org/10.1016/0038-0717(90)90126-K)

544 Fierer, N., Bradford, M.A., Jackson, R.B., 2007. Toward an ecological classification of soil  
545 bacteria. *Ecology* 88, 1354–1364. <https://doi.org/10.1890/05-1839>

546 Filley, T.R., Nierop, K.G.J., Wang, Y., 2006. The contribution of polyhydroxyl aromatic  
547 compounds to tetramethylammonium hydroxide lignin-based proxies. *Org. Geochem.*  
548 37, 711–727. <https://doi.org/10.1016/J.ORGGEOCHEM.2006.01.005>

549 Fujisaki, K., Chevallier, T., Chapuis-Lardy, L., Albrecht, A., Razafimbelo, T., Masse, D.,  
550 Ndour, Y.B., Chotte, J.-L., 2018. Soil carbon stock changes in tropical croplands are  
551 mainly driven by carbon inputs: A synthesis. *Agric. Ecosyst. Environ.* 259, 147–158.  
552 <https://doi.org/10.1016/j.agee.2017.12.008>

553 Gavrichkova, O., Moscatelli, M.C., Kuzyakov, Y., Grego, S., Valentini, R., 2010. Influence  
554 of defoliation on CO<sub>2</sub> efflux from soil and microbial activity in a Mediterranean  
555 grassland. *Agric. Ecosyst. Environ.* 136, 87–96.  
556 <https://doi.org/10.1016/j.agee.2009.11.015>

557 Gougoulias, C., Clark, J.M., Shaw, L.J., 2014. The role of soil microbes in the global carbon  
558 cycle: tracking the below-ground microbial processing of plant-derived carbon for  
559 manipulating carbon dynamics in agricultural systems. *J. Sci. Food Agric.* 94, 2362–  
560 2371. <https://doi.org/10.1002/jsfa.6577>

561 Havstad, K.M., Peters, D.P.C., Skaggs, R., Brown, J., Bestelmeyer, B., Fredrickson, E.,  
562 Herrick, J., Wright, J., 2007. Ecological services to and from rangelands of the United  
563 States. *Ecol. Econ.* 64, 261–268. <https://doi.org/10.1016/J.ECOLECON.2007.08.005>



564 Hedges, J.I., Ertel, J.R., 1982. Characterization of lignin by gas capillary chromatography of  
565 cupric oxide oxidation products. *Anal. Chem.* 54, 174–178.  
566 <https://doi.org/10.1021/ac00239a007>

567 Heitkötter, J., Heinze, S., Marschner, B., 2017. Relevance of substrate quality and nutrients  
568 for microbial C-turnover in top- and subsoil of a Dystric Cambisol. *Geoderma* 302, 89–  
569 99. <https://doi.org/10.1016/j.geoderma.2017.04.029>

570 IPCC, 2018. Special Report. Global Warming of 1.5 °C. <https://www.ipcc.ch/sr15/>.

571 Jenkinson, D.S., Brookes, P.C., Powlson, D.S., 2004. Measuring soil microbial biomass. *Soil*  
572 *Biol. Biochem.* 36, 5–7. <https://doi.org/10.1016/J.SOILBIO.2003.10.002>

573 Joergensen, R.G., Emmerling, C., 2006. Methods for evaluating human impact on soil  
574 microorganisms based on their activity, biomass, and diversity in agricultural soils. *J.*  
575 *Plant Nutr. Soil Sci.* 169, 295–309. <https://doi.org/10.1002/jpln.200521941>

576 Kallenbach, C.M., Frey, S.D., Grandy, A.S., 2016. Direct evidence for microbial-derived soil  
577 organic matter formation and its ecophysiological controls. *Nat. Commun.* 7, 1–10.  
578 <https://doi.org/10.1038/ncomms13630>

579 Kemmitt, S.J., Wright, D., Goulding, K.W.T., Jones, D.L., 2006. pH regulation of carbon and  
580 nitrogen dynamics in two agricultural soils. *Soil Biol. Biochem.* 38, 898–911.  
581 <https://doi.org/10.1016/j.soilbio.2005.08.006>

582 Koch, O., Tscherko, D., Kandeler, E., 2007. Temperature sensitivity of microbial respiration,  
583 nitrogen mineralization, and potential soil enzyme activities in organic alpine soils.  
584 *Global Biogeochem. Cycles* 21. <https://doi.org/10.1029/2007GB002983>

585 Kögel-Knabner, I., 2002. The macromolecular organic composition of plant and microbial  
586 residues as inputs to soil organic matter. *Soil Biol. Biochem.* 34, 139–162.  
587 [https://doi.org/10.1016/S0038-0717\(01\)00158-4](https://doi.org/10.1016/S0038-0717(01)00158-4)

588 Kögel, I., Bochter, R., 1985. Characterization of lignin in forest humus layers by high-  
589 performance liquid chromatography of cupric oxide oxidation products. *Soil Biol.*  
590 *Biochem.* 17, 637–640. [https://doi.org/10.1016/0038-0717\(85\)90040-9](https://doi.org/10.1016/0038-0717(85)90040-9)

591 Liang, C., Schimel, J.P., Jastrow, J.D., 2017. The importance of anabolism in microbial  
592 control over soil carbon storage. *Nat. Microbiol.* 2, 17105.  
593 <https://doi.org/10.1038/nmicrobiol.2017.105>

594 Marchus, K.A., Blankinship, J.C., Schimel, J.P., 2018. Environmental controls on  
595 extracellular polysaccharide accumulation in a California grassland soil. *Soil Biol.*  
596 *Biochem.* <https://doi.org/10.1016/j.soilbio.2018.07.009>

597 Marinari, S., Masciandaro, G., Ceccanti, B., Grego, S., 2000. Influence of organic and

598 mineral fertilisers on soil biological and physical properties. *Bioresour. Technol.* 72, 9–  
599 17. [https://doi.org/10.1016/S0960-8524\(99\)00094-2](https://doi.org/10.1016/S0960-8524(99)00094-2)

600 Marx, M.-C., Kandeler, E., Wood, M., Wermbter, N., Jarvis, S.C., 2005. Exploring the  
601 enzymatic landscape: distribution and kinetics of hydrolytic enzymes in soil particle-size  
602 fractions. *Soil Biol. Biochem.* 37, 35–48.  
603 <https://doi.org/10.1016/J.SOILBIO.2004.05.024>

604 McSherry, M.E., Ritchie, M.E., 2013. Effects of grazing on grassland soil carbon: a global  
605 review. *Glob. Chang. Biol.* 19, 1347–1357. <https://doi.org/10.1111/gcb.12144>

606 Medina-Roldán, E., Bardgett, R.D., 2011. Plant and soil responses to defoliation: A  
607 comparative study of grass species with contrasting life history strategies. *Plant Soil*  
608 344, 377–388. <https://doi.org/10.1007/s11104-011-0756-4>

609 Ngo, P.T., Rumpel, C., Dignac, M.-F., Billou, D., Duc, T.T., Jouquet, P., 2011.  
610 Transformation of buffalo manure by composting or vermicomposting to rehabilitate  
611 degraded tropical soils. *Ecol. Eng.* 37, 269–276.  
612 <https://doi.org/10.1016/j.ecoleng.2010.11.011>

613 Panikov, N.S., Sizova, M. V., 1996. A kinetic method for estimating the biomass of microbial  
614 functional groups in soil. *J. Microbiol. Methods* 24, 219–230.  
615 [https://doi.org/10.1016/0167-7012\(95\)00074-7](https://doi.org/10.1016/0167-7012(95)00074-7)

616 Piñeiro, G., Paruelo, J.M., Oesterheld, M., Jobbágy, E.G., 2010. Pathways of Grazing Effects  
617 on Soil Organic Carbon and Nitrogen. *Rangel. Ecol. Manag.* 63, 109–119.  
618 <https://doi.org/10.2111/08-255.1>

619 Post, W.M., Kwon, K.C., 2000. Soil carbon sequestration and land-use change: processes and  
620 potential. *Glob. Chang. Biol.* 6, 317–327. <https://doi.org/10.1046/j.1365-2486.2000.00308.x>

622 Puche, N.J.B., Senapati, N., Klumpp, K., Fléchar, C.R., Kirschbaum, M.U.F., Chabbi, A.,  
623 2019. Modelling carbon and water fluxes of managed grasslands: comparing flux  
624 variability and net carbon budgets between grazed and mowed systems. *Agronomy*.

625 Razavi, B.S., Blagodatskaya, E., Kuzyakov, Y., 2015. Nonlinear temperature sensitivity of  
626 enzyme kinetics explains canceling effect—a case study on loamy haplic Luvisol. *Front.*  
627 *Microbiol.* 6. <https://doi.org/10.3389/fmicb.2015.01126>

628 Rumpel, C., Crème, A., Ngo, P., Velásquez, G., Mora, M., Chabbi, A., 2015. The impact of  
629 grassland management on biogeochemical cycles involving carbon, nitrogen and  
630 phosphorus. *J. soil Sci. plant Nutr.* 15(2), 353–371. <https://doi.org/10.4067/S0718-95162015005000034>

631

- 632 Rumpel, C., Dignac, M.-F., 2006. Gas chromatographic analysis of monosaccharides in a  
633 forest soil profile: Analysis by gas chromatography after trifluoroacetic acid hydrolysis  
634 and reduction–acetylation. *Soil Biol. Biochem.* 38, 1478–1481.  
635 <https://doi.org/10.1016/J.SOILBIO.2005.09.017>
- 636 Rumpel, C., Kögel-Knabner, I., 2011. Deep soil organic matter---a key but poorly understood  
637 component of terrestrial C cycle. *Plant Soil* 338, 143–158.  
638 <https://doi.org/10.1007/s11104-010-0391-5>
- 639 Rumpel, C., Lehmann, J., Chabbi, A., 2018. ‘4 per 1,000’ initiative will boost soil carbon for  
640 climate and food security. *Nat.* 2020 5537686.
- 641 Sanaullah, M., Chabbi, A., Lemaire, G., Charrier, X., Rumpel, C., 2010. How does plant leaf  
642 senescence of grassland species influence decomposition kinetics and litter compounds  
643 dynamics? *Nutr. Cycl. Agroecosystems* 88, 159–171. [https://doi.org/10.1007/s10705-](https://doi.org/10.1007/s10705-009-9323-2)  
644 [009-9323-2](https://doi.org/10.1007/s10705-009-9323-2)
- 645 Schädel, C., Blöchl, A., Richter, A., Hoch, G., 2010. Quantification and monosaccharide  
646 composition of hemicelluloses from different plant functional types. *Plant Physiol.*  
647 *Biochem.* 48, 1–8. <https://doi.org/10.1016/J.PLAPHY.2009.09.008>
- 648 Schimel, J.P., Schaeffer, S.M., 2012. Microbial control over carbon cycling in soil. *Front.*  
649 *Microbiol.* <https://doi.org/10.3389/fmicb.2012.00348>
- 650 Senapati, N., Chabbi, A., Gastal, F., Smith, P., Mascher, N., Loubet, B., Cellier, P., Naisse,  
651 C., 2014. Net carbon storage measured in a mowed and grazed temperate sown  
652 grassland shows potential for carbon sequestration under grazed system. *Carbon Manag.*  
653 5, 131–144. <https://doi.org/10.1080/17583004.2014.912863>
- 654 Senapati, N., Chabbi, A., Giotri, A.F., Yeluripati, J.B., Smith, P., 2016. Modelling nitrous  
655 oxide emissions from mown-grass and grain-cropping systems: Testing and sensitivity  
656 analysis of DailyDayCent using high frequency measurements. *Sci. Total Environ.* 572,  
657 955–977. <https://doi.org/10.1016/j.scitotenv.2016.07.226>
- 658 Smith, P., Martino, D., Cai, Z., Gwary, D., Janzen, H., Kumar, P., McCarl, B., Ogle, S.,  
659 O’Mara, F., Rice, C., Scholes, B., Sirotenko, O., Howden, M., McAllister, T., Pan, G.,  
660 Romanenkov, V., Schneider, U., Towprayoon, S., Wattenbach, M., Smith, J., 2008.  
661 Greenhouse gas mitigation in agriculture. *Philos. Trans. R. Soc. B Biol. Sci.* 363, 789–  
662 813. <https://doi.org/10.1098/rstb.2007.2184>
- 663 Soussana, J.-F., Loiseau, P., Vuichard, N., Ceschia, E., Balesdent, J., Chevallier, T.,  
664 Arrouays, D., 2006. Carbon cycling and sequestration opportunities in temperate  
665 grasslands. *Soil Use Manag.* 20, 219–230. <https://doi.org/10.1111/j.1475->

666 2743.2004.tb00362.x

667 Strickland, M.S., Rousk, J., 2010. Considering fungal: Bacterial dominance in soils -  
668 Methods, controls, and ecosystem implications. *Soil Biol. Biochem.* 42, 1385–1395.  
669 <https://doi.org/10.1016/j.soilbio.2010.05.007>

670 Tautges, N.E., Chiartas, J.L., Gaudin, A.C.M., O’Geen, A.T., Herrera, I., Scow, K.M., 2019.  
671 Deep soil inventories reveal that impacts of cover crops and compost on soil carbon  
672 sequestration differ in surface and subsurface soils. *Glob. Chang. Biol.* 25, 3753–3766.  
673 <https://doi.org/10.1111/gcb.14762>

674 Thevenot, M., Dignac, M.-F.F., Rumpel, C., 2010. Fate of lignins in soils: A review. *Soil*  
675 *Biol. Biochem.* 42, 1200–1211. <https://doi.org/10.1016/J.SOILBIO.2010.03.017>

676 Vance, E.D., Brookes, P.C., Jenkinson, D.S., 1987. An extraction method for measuring soil  
677 microbial biomass C. *Soil Biol. Biochem.* 19, 703–707. [https://doi.org/10.1016/0038-](https://doi.org/10.1016/0038-0717(87)90052-6)  
678 [0717\(87\)90052-6](https://doi.org/10.1016/0038-0717(87)90052-6)

679 Wang, G., Mayes, M.A., Gu, L., Schadt, C.W., 2014. Representation of dormant and active  
680 microbial dynamics for ecosystem modeling. *PLoS One* 9.  
681 <https://doi.org/10.1371/journal.pone.0089252>

682 Wilson, C.H., Strickland, M.S., Hutchings, J.A., Bianchi, T.S., Flory, S.L., 2018. Grazing  
683 enhances belowground carbon allocation, microbial biomass, and soil carbon in a  
684 subtropical grassland. *Glob. Chang. Biol.* 24, 2997–3009.  
685 <https://doi.org/10.1111/gcb.14070>

686 Wutzler, T., Blagodatsky, S.A., Blagodatskaya, E., Kuzyakov, Y., 2012. Soil microbial  
687 biomass and its activity estimated by kinetic respiration analysis – Statistical guidelines.  
688 *Soil Biol. Biochem.* 45, 102–112. <https://doi.org/10.1016/j.soilbio.2011.10.004>

689 Xu, S., Silveira, M.L., Inglett, K.S., Sollenberger, L.E., Gerber, S., 2017. Soil microbial  
690 community responses to long-term land use intensification in subtropical grazing lands.  
691 *Geoderma* 293, 73–81. <https://doi.org/10.1016/j.geoderma.2017.01.019>

692

693

694

695 **Tables**696 **Table 1.** General soil properties under two grassland management practices (grazing and

697 mowing) at in surface soil (0-10 cm) and subsurface soil (20-30 cm).

	Treatment	pH	SOC content mg g <sup>-1</sup>	N mg g <sup>-1</sup>	δ <sup>13</sup> C ‰	δ <sup>15</sup> N ‰	C:N ratio
<b>Surface soil</b>	<b>Grazing</b>	5.95±0.09	21.4±0.81	2.2±0.09	-27.4±0.06	4.9±0.13	9.6±0.05
	<b>Mowing</b>	5.51±0.08	14.6±0.51	1.5±0.05	-27.0±0.05	5.0±0.09	9.6±0.07
<b>Subsurface soil</b>	<b>Grazing</b>	5.99±0.12	11.8±0.62	1.3±0.06	-26.7±0.06	6.2±0.08	9.2±0.06
	<b>Mowing</b>	6.01±0.13	8.6±0.44	0.9±0.05	-26.3±0.10	6.4±0.10	9.1±0.07
<b>ANCOVA, F value (P values)</b>							
SOC stocks in 2005		0.15 (0.70)	5.31 (0.03)	6.26 (0.02)	6.86 (0.01)	1.90 (0.18)	1.33 (0.26)
Treatment		3.37 (0.08)	30.3 (<0.001)	28.7 (<0.001)	17.2 (0.002)	0.35 (0.56)	1.19 (0.28)
Depth		5.89 (0.02)	181.8 (<0.001)	153.8 (<0.001)	132.5 (<0.001)	157.9 (<0.001)	52.5 (<0.001)
Treatment×Depth		4.27 (0.04)	0.68 (0.41)	0.86 (0.36)	0.28 (0.06)	0.25 (0.62)	0.37 (0.55)

698 Values are shown as the average of ten replicates and ±SE. Significant differences between the treatments are  
699 indicated by capital case letters. Lower case letters show significant differences with depth (P < 0.05).

700

701 **Table 2.** Non-cellulosic polysaccharides (NCP) signature in soil under two grassland  
 702 management practices (grazing and mowing) at two depths (0-10 cm and 20-30 cm).

	Treatment	NCP content	NCP content per SOC	NCP monomers ratios		
		mg g <sup>-1</sup>	mg g <sup>-1</sup> SOC	C6/C5 <sup>1</sup>	DesoxyC6/C5 <sup>2</sup>	Man/Xyl <sup>3</sup>
<b>Surface soil</b>	<b>Grazing</b>	6.61±0.23	308.98±6.3	0.80±0.02	0.35±0.01	0.54±0.02
	<b>Mowing</b>	4.45±0.18	306.63±11.5	0.84±0.02	0.34±0.01	0.61±0.02
<b>Subsurface soil</b>	<b>Grazing</b>	3.09±0.15	263.39±6.4	1.03±0.02	0.43±0.01	0.87±0.03
	<b>Mowing</b>	2.50±0.11	292.41±10.5	1.01±0.02	0.46±0.01	0.91±0.03
<b>ANCOVA, F value (P values)</b>						
SOC stocks in 2005		0.43 (0.52)				
Treatment		36.6 (<0.001)	2.50 (0.12)	5.42 (0.03)	3.81 (0.06)	4.74 (0.04)
Depth		241.1	0.11 (0.74)	0.87 (0.36)	0.01 (0.91)	0.64 (0.43)
Treatment×Depth		(<0.001)	11.5 (0.002)	122.2 (<0.001)	102.8 (<0.001)	166.3 (<0.001)
		19.7 (<0.001)	3.17 (0.08)	3.14 (0.09)	3.18 (0.08)	0.52 (0.48)

703  
 704 <sup>1</sup>C6/C5 – the ratio of C6- to C5- sugar monomers, <sup>2</sup>DesoxyC6/C5 – the ratio of desoxy C6- to desoxy  
 705 C5- sugar monomers, <sup>3</sup>Man/Xyl - the ratio of mannose to xylose. These ratios indicate the origin of non-  
 706 cellulosic polysaccharides (microbial or plant).  
 707

708

709 **Table 3.** Lignin signature in soil under two grassland management practices (grazing and  
 710 mowing) in surface soil (0-10 cm) and subsurface soil (20-30 cm).

	Treatment	Lignin content	Lignin content per SOC	Lignin monomers ratios			
		mg g <sup>-1</sup>	mg g <sup>-1</sup> SOC	C/V	S/V	(Ac/Al) <sub>v</sub>	(Ac/Al) <sub>s</sub>
<b>Surface soil</b>	<b>Grazing</b>	0.35±0.01	16.31±0.64	0.45±0.03	1.34±0.02	0.53±0.02	0.46±0.01
	<b>Mowing</b>	0.26±0.01	17.86±0.43	0.45±0.03	1.24±0.02	0.65±0.02	0.57±0.02
<b>Subsurface soil</b>	<b>Grazing</b>	0.19±0.02	16.22±0.57	0.37±0.02	1.33±0.03	0.66±0.01	0.54±0.02
	<b>Mowing</b>	0.16±0.01	18.86±0.89	0.37±0.03	1.30±0.02	0.63±0.03	0.56±0.02
<b>ANCOVA, F value (P values)</b>							
SOC stocks in 2005		0.05 (0.82)	10.9 (0.002)	1.59 (0.22)	1.99 (0.17)	0.08 (0.78)	18.6 (<0.001)
Treatment		15.3 (<0.001)	1.14 (0.29)	0.64 (0.43)	7.91 (0.01)	2.83 (0.10)	2.82 (0.10)
Depth		96.3 (<0.001)	0.62 (0.44)	8.58 (0.006)	0.95 (0.34)	9.13 (0.005)	13.1
Treatment×Depth		4.83 (0.04)	0.88 (0.36)	0.015 (0.90)	2.19 (0.15)	16.3 (<0.001)	<0.001


711 C/V – the ratio of cinnamyl phenols to syringyl phenols; S/V - the ratio of syringyl phenols to vanillyl  
 712 phenols; (Ac/Al)<sub>v</sub> – acid to aldehyde ratio of vanillyl phenols; (Ac/Al)<sub>s</sub> – acid to aldehyde ratio of syringyl  
 713 phenols. These ratios are indicators of lignin degradation state in soil.  
 714

715

716

717

**Figure captions**



718

719

**Figure 1.** (A) Soil microbial respiration (SMR), (B) soil microbial respiration (SMR)

720

per soil organic carbon (SOC), (C) microbial biomass carbon (MBC) per soil organic carbon

721

(SOC) and (D) metabolic quotient ( $qCO_2$ ) in soil under two grassland management practices

722

(grazing and mowing) in surface soil (0-10 cm) and subsurface soil (20-30 cm). Significant

723


differences between the treatments are indicated by \*, \*\* and \*\*\*, representing probability at

724

the 5, 1, and 0.1% levels, respectively.

725





726

727 **Figure 2.** (A) Microbial C:N ratio, (B) the percentage of active microbial biomass


728 (AMB) and (C) specific microbial growth rate ( $\mu$ ) in soil under two grassland management

729 practices (grazing and mowing) in surface soil (0-10 cm) and subsurface soil (20-30 cm).

730 Significant differences between the treatments are indicated by \*, \*\* and \*\*\*, representing

731 probability at the 5, 1, and 0.1% levels, respectively.

732



733

734

**Figure 3.** Boxplot of enzyme activity per unit of microbial biomass C (MBC) for nine

735

enzymes under two grassland management practices (grazing and mowing) in surface soil (0-

736


10 cm) and subsurface soil (20-30 cm). Significant differences between the treatments are

737

indicated by \*, \*\* and \*\*\*, representing probability at the 5, 1, and 0.1% levels, respectively.

738

739



740

741

**Figure 4.** Principal component analysis (PCA) for soil under grazing and mowing in

742

surface soil (0-10 cm) and in subsurface soil (20-30 cm). Only variables with quality of

743

representation ( $\cos^2$ ) higher than 0.75 was shown on PCA plot.