



HAL
open science

Management of grasslands by mowing versus grazing – impacts on soil organic matter quality and microbial functioning

Aliia Gilmullina, Cornelia Rumpel, Evgenia Blagodatskaya, Abad Chabbi

► To cite this version:

Aliia Gilmullina, Cornelia Rumpel, Evgenia Blagodatskaya, Abad Chabbi. Management of grasslands by mowing versus grazing – impacts on soil organic matter quality and microbial functioning. *Applied Soil Ecology*, 2020, 156, pp.1-9. 10.1016/j.apsoil.2020.103701 . hal-03140069

HAL Id: hal-03140069

<https://hal.inrae.fr/hal-03140069v1>

Submitted on 22 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Management of grasslands by mowing versus grazing – impacts on soil organic matter**
2 **quality and microbial functioning**

3 Gilmullina Aliia^{1,2}, Rumpel Cornelia³, Blagodatskaya Evgenia^{4,5}, Chabbi Abad^{1,2*}

4 ¹ UMR P3F, INRA, Lusignan, France

5 ² UMR ECOSYS, INRA, Thiverval-Grignon, France,

6 ³ Institute of Ecology and Environmental Sciences- Paris (iEES-Paris) UMR CNRS, INRA,

7 UPMC, Thiverval-Grignon, France

8 ⁴ Agro-Technology Institute, RUDN University, Moscow, Russia

9 ⁵ Dept. of Soil Ecology, Helmholtz Centre for Environmental Research – UFZ, Halle (Saale),

10 Germany

11 * - corresponding author, abad.chabbi@inra.fr

12

13

14 **Keywords**

15 Grassland, grazing, mowing, non-cellulosic polysaccharides, lignin, microbial activity

16

17

18 **Highlights**

19 • Grazing and mowing have contrasting effects on soil biogeochemical properties

20 • Grazing promoted more efficient microbial functioning

21 • Mowing leads to more degraded lignin than grazing

22 • Only microbial properties were sensitive to treatment effects in subsurface soil

23

24 **Abstract**

25 Although 30% of the European surface area is covered with grasslands, little is known
26 about the effect of their management on soil quality and biogeochemical cycling. Here, we
27 analysed soil from an experimental site in Western France, which had been under either
28 grazing or mowing regime for 13 years. We aimed to assess the effect of the two
29 management practices on the biogeochemical functioning of soil system. To this end we
30 compared soil organic matter (SOM) composition and microbial properties at two soil depths.
31 We analysed for elemental, lignin and non-cellulosic polysaccharide content and composition
32 and for microbial biomass, soil microbial respiration and enzyme activities. Our results
33 showed higher soil organic carbon (SOC) and nitrogen contents in the surface soil under
34 grazing as compared to mowing. Soil biogeochemical properties differed between grazing
35 and mowing treatments. In particular, soil under grazing showed lower lignin and higher
36 microbial biomass. Despite the similar non-cellulosic polysaccharide content under both
37 treatments, microbial community under mowing was characterised by higher enzyme
38 production per microbial biomass, leading to more degraded SOM in the mowing system as
39 compared to grazing. We conclude that grazing and mowing regimes impact differently the
40 biogeochemical soil functioning. Higher and more diverse carbon input under grazing
41 compared to mowing may lead to enhanced substrate availability and thus more efficient
42 microbial functioning, which could favour SOC sequestration through formation of microbial
43 products.

44

45

46

47 **1. Introduction**

48 Dangerous climate change can only be avoided if we succeed to remove CO₂ from the
49 atmosphere with negative emission technologies (IPCC, 2018). Soil organic carbon (SOC)
50 sequestration is a nature-based negative emission technology, which may be achievable at
51 scale through the introduction of sustainable management practices with permanent soil cover
52 (Rumpel et al., 2018). Permanent grasslands, which in Europe, occupy about 30% of the
53 agricultural area (Ec.europa.eu, 2018), are responsible for many ecosystem services including
54 forage for animal production and SOC storage (Havstad et al., 2007; Rumpel et al., 2015).
55 Biogeochemical cycling in grassland soils can be influenced by a variety of management
56 practices (Rumpel et al., 2015). The impact of these management practices on processes
57 impacting soil biogeochemical cycling via soil-plant interactions are poorly understood
58 (Dignac et al., 2017). These interactions result in contrasting effects of grassland
59 management on SOC storage potential (Post and Kwon, 2000; Rumpel et al., 2015; Smith et
60 al., 2008).

61 Grazing and mowing are the most frequently used grassland management practices.
62 Both practices lead to defoliation (removal of plant aboveground tissue). Defoliation alters
63 root exudation and C allocation in plants but the direction of these changes was found to be
64 contrasting (Bazot et al., 2005; Gavrichkova et al., 2010; Medina-Roldán and Bardgett,
65 2011), related to different climatic and pedological conditions (Pineiro *et al.*, 2010).

66 Defoliation under grazing management is caused by herbivores during several days
67 (Senapati et al., 2014). This process plays an important role in terms of carbon and nutrient
68 return (Soussana et al., 2006), because about 50-70% of the ingested biomass is returned to
69 soil in the form of excreta. In mowing systems, plant biomass is removed in a day with up to
70 20% of all cut biomass remaining as green litter in form of harvesting losses (Sanaullah et al.,

71 2010). In order to compensate for nutrient exportation during defoliation events, mineral
72 fertilisers are applied in mowing systems.

73 Due to the different types of biomass returned in the two systems, the quality of
74 biomass input also varies. Mowing systems receive only plant residues while input in grazing
75 systems comprises additionally animal depositions. Dung and urine inputs are characterised
76 by lower C:N ratio, higher amount of easily available compounds (Dungait et al., 2009) and
77 relatively stable compounds, such as crude proteins and fats (Dungait et al., 2005; Ngo et al.,
78 2011). Moreover, in grazing systems, there is a return of senescent brown litter, which
79 contains less N and less soluble compounds compared to the green litter returned as
80 harvesting losses in mowing systems (Sanaullah et al., 2010).

81 These differences may affect belowground processes (Wilson et al., 2018), SOC
82 formation and storage (Cotrufo et al., 2015; Rumpel et al., 2015) through their effect on the
83 soil microbial biomass and its activity (Liang et al., 2017). We therefore hypothesised that the
84 two management systems may lead to contrasting soil microbial functioning and affect
85 differently biogeochemical cycling. The effect of management has been analysed up to now
86 mainly in the first few centimetres of soil, although it has been shown that management can
87 affect SOC stored down to 2 m depth (Tautges et al., 2019). We thus hypothesised that
88 grassland management affects SOC below the first centimetres.

89 We focused on an experimental site with grazing and mowing as two contrasting
90 management practices under similar soil and climatic conditions. We aimed to evaluate the
91 differences in biogeochemical cycling in soil under the two different management practices at
92 two depths. To this end we analysed C and N contents, molecular signatures of
93 polysaccharide and lignin monomers. These variables were compared to the functioning of
94 the soil microbial communities, assessed by the analyses of soil microbial respiration, growth
95 kinetic parameters and activity of 9 enzymes as well as microbial biomass C and N.

96

97 2. Materials and methods

98 2.1. Site description and soil sampling

99 The field experiment is located in Lusignan (southwest of France, 46°25'12,91"N;
100 0°07'29,35"E) at the national long-term experimental observatory SOERE ACBB
101 (Agroecosystems, Biogeochemical Cycles and Biodiversity). The mean annual temperature
102 and precipitation for the period 2006–2010 were 11.2°C and 773 mm, (Senapati et al., 2014).
103 The landscape is flat. The soil is classified as a Dystric Cambisol with loamy texture (Chabbi
104 et al., 2009).

105 The current study is focused on two permanent sown grasslands (each of about 3 ha in
106 size), which were established in 2005 by sowing a mixture of three plant species (*Lolium*
107 *perenne*, *Festuca arundinacea*, *Dactylis glomerata* L.) in both treatments. In the grazing
108 system, legume *Trifolium repens* was included in the species mixture but covered only 5% of
109 grazed paddock in 2017. The mown grassland was cut four times per year with biomass
110 exported. To replace the exported nutrients, nitrogen (N) fertilizer was applied at rates
111 between 170 and 380 kg N ha⁻¹ year⁻¹ (Puche et al., 2019). Grazing in the grazed paddock
112 took place from March to December with 50 days per year using 15 to 20 livestock units per
113 hectare. Grazed grasslands received less nitrogen fertilization (60-150 kg N ha⁻¹ year⁻¹, Puche
114 et al., 2019) because nitrogen losses were additionally returned by dung and urine and
115 through the presence of the leguminous species. In order to compare the treatments at similar
116 N status, fertilizer application rates were adjusted to maintain the Nitrogen Nutrition Index
117 between 0.9 and 1.0 for both treatments, close to non-limiting nitrogen nutrition to near
118 maximum plant production (Senapati et al., 2016). Moreover, both sites were limed regularly
119 in order to neutralize acid pH.

120 Due to the large land requirements (3 ha for plots with cows), it was not possible to
121 establish and maintain a completely replicated field experiment including grazing treatment
122 for several decades. Limitations to generalization of the treatment effects due to the absence
123 of replication of the experiments were limited by choosing homogenous flat areas in close
124 proximity with similar land use history, climate, and soil type. Moreover, we carried out
125 baseline measurements, in form of geostatistical evaluation of the soils SOC and N contents
126 and included initial SOC stocks as a co-variate. These data showed that both plots were
127 significantly different in initial SOC and N contents (n=28). The SOC contents on mowing
128 plots varied between 9.9 and 13.7 mg g⁻¹ (average 12.0 ± 1.0 mg g⁻¹), while under grazing it
129 was between 11.9 and 19.1 mg g⁻¹ (average 14.8 ± 1.5 mg g⁻¹). N contents varied between 1.0
130 and 1.4 mg g⁻¹ (average 1.2 ± 0.1 mg g⁻¹) under mowing, while under grazing the values
131 ranged between 1.2 and 1.9 mg g⁻¹ (average 1.5 ± 0.1 mg g⁻¹). These previous analyses
132 indicated on average non-significant differences in SOC stock changes between grazing and
133 mowing after nine years of treatment (Crème et al., 2020). The study also showed
134 partitioning of the field into different zones with SOC gain and loss (Crème et al., 2020; Fig
135 S1, Supplementary materials).

136

137 Five replicated soil samples were taken from each of the two zones, giving a total of 10
138 replicated field samples per plot. Sampling took place in November 2017, 2 weeks and 5
139 months after the last grazing and mowing events, accordingly. The shortest distance between
140 samples was 25 m. Soil samples were collected with a mechanical auger (5cm Ø, 30cm) at
141 two depths: 0-10 cm (surface soil) and 20-30 cm (subsurface soil) giving in total 40 samples.
142 All samples were sieved through a 2-mm mesh. Thereafter, half of the samples was air-dried
143 and ground for measurements of physicochemical analysis and the other half was stored at
144 4°C before microbial analyses. Because of dry field conditions prior to measurements of

145 microbiological analysis, soil samples were moistened by distilled water to adjust 50% of
146 WHC and pre-incubated at 22 °C for 7 days.

147 2.2. *Soil general properties*

148 Soil pH (H₂O) was measured in a soil:water suspension (1:2.5 weight/volume). Soil
149 organic carbon (SOC), nitrogen (N) and stable isotopes (¹³C and ¹⁵N) contents were measured
150 using a CHN auto-analyser (Flash EA, Thermo Electron Corporation, Bremen, Germany)
151 coupled with an isotope ratio mass spectrometer. The isotopic ratios were calculated relative
152 to the Pee Dee Belemnite Standard (PDB) for C and relative to atmospheric N₂ for nitrogen.

153

154 2.3. *Soil chemical properties*

155 Lignin was analysed by the alkaline cupric oxide (CuO) oxidation method (Hedges and
156 Ertel, 1982; Kögel and Bochter, 1985). Briefly, oxidation was carried out under alkaline
157 conditions (2M NaOH) at 172 °C for 4 hours using 500 mg of air-dried soil, 250 mg of CuO,
158 50 mg of ammonium ferrous hexahydrate and 50 mg of glucose. After cooling, samples were
159 acidified with 5 M HCl and left overnight for humic acid precipitation. Removal of humic
160 acids was conducted through centrifugation (10 min at 10000 rpm) and followed by
161 extraction of phenolic oxidation products with C18 reversed phase columns. The phenols
162 were derivatized with BSTFA and quantified as trimethylsilyl derivatives by gas
163 chromatography with a HP gas chromatograph (HP GC 6890) equipped with a flame
164 ionization detector and a SGE BPX-5 column (50 m length, 0.25 mm inner diameter, 0.32 µm
165 coating). Samples were injected in split mode (1:10). The GC oven temperature was
166 programmed at 100 °C for 2 min, then increased from 100 to 172 °C at a heating rate of 8 °C
167 min⁻¹, from 172 to 184 °C at 4 °C min⁻¹, and from 184 to 300 °C at a rate of 10°C min⁻¹.

168 The internal standard ethylvanillin was added before the purification step to quantify
169 lignin recovery and the quantification standard phenylacetic acid was added before GC
170 analyses.

171 The total lignin content (mg g^{-1} dry soil) of the sample was determined as the sum of
172 phenolic oxidation products: vanillyl (V), syringyl (S) and p-coumaryl (C) in their acid (Ac),
173 aldehyde (Al) and ketone forms. Lignin content was also expressed as lignin content per SOC
174 (mg g^{-1} SOC). Lignin decomposition was assessed by the ratios of S, C to V and (Ac/Al)
175 ratios of V and S, which generally indicate decomposition state (Thevenot et al., 2010).

176 Non-cellulosic polysaccharides of plant and microbial origin (Kögel-Knabner, 2002)
177 were determined by gas chromatography after trifluoroacetic acid (TFA) hydrolysis and
178 reduction-acetylation using a method introduced by Rumpel and Dignac (2006) and modified
179 by Eder et al. (2010). The analysis was performed using 700 mg of soil samples. Briefly,
180 hydrolysis of non-cellulose polysaccharides was carried out at 105°C for 4 h with 10 ml of 4
181 M TFA. Thereafter, Myo-inositol was added as quantification standard to account for the
182 losses during the purification procedure. Removal of soil was performed by filtration through
183 glass fibre filters (Whatman GF/C $0.45\ \mu\text{m}$). Afterwards, TFA was evaporated using a
184 centrifugal Evaporator EZ-2 ENVI at 35°C for 4 hours and dry samples were left overnight in
185 the freezer. Thereafter, dry samples were dissolved in 0.5 ml of H_2O followed by the addition
186 of 0.9 EDTA in order to avoid co-precipitation of organic material with metal oxides and
187 hydroxides (Eder et al., 2010). One mL sodium borohydride (NaBH_4) in dimethylsulfoxide
188 ($20\ \text{g L}^{-1}$) was added for reduction of polysaccharide monomers into alditol forms and kept
189 at 40°C for 1.5 hours. Then, acetylation was conducted by addition of 0.2 mL acetic acid, 2
190 mL of acetic anhydride and 0.2 mL Methylimidazole. Acetylated alditols were extracted by 1
191 ml of dichloromethane and quantified with a HP GC 6890 gas chromatograph equipped with
192 a flame ionization detector. Separation was achieved with a 60 m fused silica capillary

193 column (SGE BPX 70, 0.32 mm internal diameter, 0.25 mm film thickness) under the
194 following temperature program: 170 to 250 °C at 8 °C.min⁻¹, followed by 12 min at 250 °C
195 (isothermal). Helium was used as the carrier gas at a flow rate of 1.0 mL min⁻¹. The injector
196 was kept at 250 °C and the detector at 260 °C. The non-cellulosic polysaccharides content of
197 soil samples was determined as the sum of monosaccharides: C5 (pentoses: xylose, ribose
198 and arabinose), C6 (hexoses: glucose, galactose and mannose), and desoxyC6
199 (desoxyhexoses: fucose and rhamnose) (Kögel-Knabner, 2002). A higher C6/C5 ratio
200 generally indicates higher contribution of microbial sugars.

201 2.4. Soil microbial properties

202 Microbial biomass C (MBC) and nitrogen (MBN) were determined by the chloroform
203 fumigation-extraction method (Vance et al., 1987). Dissolved organic C and N in fumigated
204 and non-fumigated soil samples were extracted in 0.05 M K₂SO₄ and were measured using a
205 multi C/N analyzer (multi C/N analyser 2100S, Analytic Jena). MBC and MBN were
206 calculated with a conversion factor of 0.45 (Jenkinson et al., 2004). For measuring soil
207 microbial respiration (SMR) a half gram of soil sample was placed in 2 ml Eppendorf tubes.
208 The CO₂ efflux was trapped in 3 ml of 0.1 M NaOH and determined by conductometry. The
209 metabolic quotient (qCO₂), reflecting decomposition activity (Anderson, 2003; Anderson and
210 Domsch, 1993), was calculated as soil microbial respiration expressed per gram of microbial
211 biomass carbon: $qCO_2 = SMR/MBC$ (μg CO₂-C g⁻¹ MBC h⁻¹).

212 We used microbial growth kinetics technique as an approach to estimate microbial
213 biomass activity state (Blagodatskaya and Kuzyakov, 2013). This approach is based on soil
214 respiratory response to unlimited nutrient amendments (Panikov and Sizova, 1996). For this
215 purpose, soil samples were treated with a solution (0.1 ml per g of dw soil) containing per g
216 soil: 10 mg glucose, 1.9 mg (NH₄)₂SO₄, 3.8 mg MgSO₄*7H₂O, 0.11 mg K₂HPO₄ and 1.68
217 mg KH₂PO₄ for surface soil samples and 10 mg glucose, 1.9 mg (NH₄)₂SO₄, 3.8 mg

218 MgSO₄*7H₂O, 0.53 mg K₂HPO₄ and 1.35 mg KH₂PO₄ for subsurface soil samples. The
219 amount of mineral salts was preliminary selected in order to avoid soil pH change of more
220 than 0.1 units after addition. For active microbial biomass (AMB) and specific growth rate
221 calculation, the results of substrate induced respiration rate were fitted with a model proposed
222 by Panikov and Sizova (Panikov and Sizova, 1996; Wutzler et al., 2012):

$$223 \quad CO_2(t) = A + B * \exp(\mu * t) \quad (1)$$

224 In order to estimate catabolic (decomposition) activity in regards to specific substrates
225 in soil, we measured extracellular enzyme activity using the fluorometric technique (Koch et
226 al., 2007; Marx et al., 2005; Razavi et al., 2015). Nine types of fluorogenic substrates based
227 on 4-methylumbelliferone (MUF) and 7-amino-4-methylcoumarin (AMC) were used: (1)
228 MUF- α -D-glucopyranoside for α -glucosidase, (2) MUF- β -D-glucopyranoside for β -
229 glucosidase, (3) MUF- β -D-galactopyranoside for β -galactosidase, (4) MUF- β -D-
230 xylopyranoside for β -xylosidase, (5) MUF- β -D-cellobioside for β -cellobiohydrolase, (6)
231 MUF-N-acetyl- β -D-glucosamide for chitinase, (7) Leucine-AMC for leucine aminopeptidase,
232 (8) MUF-heptanoate for lipase and (9) MUF-phosphate for phosphatase. Saturation
233 concentrations of fluorogenic substrates were determined in preliminary experiments and
234 comprised 20 μ mol g⁻¹ soil for all enzymes except lipase with 60 μ mol g⁻¹ soil. Briefly, a
235 water extract of soil (1:10) was homogenised by low-energy sonication (40 J s⁻¹ output
236 energy) for 60 s. Thereafter 50 ml of the soil suspension were added to 150 ml of each
237 substrate solution in a 96-well microplate. Fluorescence was measured at an excitation
238 wavelength of 355 nm and an emission wavelength of 460 nm (Victor3 1420-050 Multilabel
239 Counter, PerkinElmer, USA).

240

241 2.5. *Statistical analysis*

242 All results are presented as arithmetic means with standard error. The statistical
243 analyses were conducted by using R (Studio Version 1.1.447). We used analyses of
244 covariance (ANCOVA) to test treatment effect, depth effect and their interactions using
245 chemical and microbial variables with initial SOC stock as a covariate. The initial SOC
246 stocks data was obtained from exactly the same sampling points based on the geostatistical
247 evaluation before the beginning of the experiment. This procedure allowed us to account for
248 the lack of field replication by taking into account the original difference between the grazed
249 and mowed plots. In order to obtain better understanding of treatment and depth effects, non-
250 transformed data (except C and N contents) were subjected to Principal Component Analysis
251 (PCA) and the results were also tested by ANCOVA with initial SOC stock as a covariate.
252 The equations were fitted by non-linear regression, using Model Maker-3 software (SB
253 technology Ltd.).

254

255 **3. Results**

256 3.1. *Soil properties*

257 Soil physicochemical properties are presented in Table 1. The pH was not controlled by
258 initial SOC stock ($P=0.70$). Lower pH was found for both treatments in surface soil compared
259 to subsurface soil, although, the lowest pH value was recorded in surface soils under mowing
260 treatment. SOC and N contents were nearly twice as high in the surface soil compared to the
261 subsurface soil under both treatments. Even if SOC and N contents were dependant on initial
262 SOC stock ($P=0.03$ and 0.02 , respectively), there were still significant effects of depth
263 ($P<0.001$) and treatment ($P<0.001$) after correction by using it as covariate. C:N ratio differed
264 only between soil depths ($P<0.001$) showing slightly higher C:N ratios in surface soils as

265 compared to subsurface soils. $\delta^{13}\text{C}$ followed the same pattern as SOC content and the highest
266 enrichment was recorded for the surface soil of the grazing treatment (depth effect $P < 0.001$
267 and treatment effect $P = 0.002$). The $\delta^{15}\text{N}$ did not differ between the treatments and was
268 enriched in surface soils compared to subsurface soils.

269

270 3.2. Specific SOM compounds

271 Non-cellulosic polysaccharide (NCP) content was not affected by initial SOC stock
272 ($P = 0.52$) and there was treatment \times depth interaction (Table 2, $P < 0.001$). Grazing resulted in
273 higher NCP content in both depths compared to mowing. The NCP content per SOC (mg g^{-1}
274 soil C) was affected only by depth ($P = 0.002$). Concerning the NCP monomers ratio, C6/C5
275 and Man/Xyl ratios were controlled by initial SOC stock ($P = 0.03$ and 0.04 , respectively),
276 consequently, after ANCOVA application the treatment effect was vanished while depth
277 effect remained significant (Table 2, $P < 0.001$). All NCP monomers ratios were higher in
278 subsurface soil compared to surface soil under both treatments.

279 Lignin content was not affected by initial SOC stock correction ($P = 0.82$), so the effects
280 of depth ($P < 0.001$), treatment ($P < 0.001$) and their interactions ($P = 0.04$) remained significant
281 (Table 3). Lignin content was higher in surface soils than in subsurface soils and was higher
282 under grazing compared to mowing as well. Correcting for initial SOC stock caused the
283 elimination of all effects on lignin content per SOC content. The C/V ratio was affected only
284 by depth ($P = 0.006$) showing higher values in surface soils than in subsurface soils. The S/V
285 ratio was greater under grazing treatment than under mowing treatment at both depths even
286 after correction by initial SOC stock (Table 3, $P = 0.01$). Based on the presence of treatment \times
287 depth interaction $(\text{Ac/Al})_V$ and $(\text{Ac/Al})_S$ ratios were lower in the surface soil of grazing
288 treatment as compared to mowing treatment ($P < 0.001$). In contrast to surface soils, treatments
289 did not show any effects on these lignin ratios in subsurface soils.

290

291 3.3. Soil microbial properties

292 The soil microbial respiration (SMR) ranged between 0.2 and 0.7 $\mu\text{g CO}_2 \text{-C g}^{-1} \text{ h}^{-1}$
293 with highest values in the surface soil under grazing treatment (Fig. 1A). After correcting for
294 initial SOC stock, treatment \times depth interaction effect on SMR was significant (Table S1,
295 Supplementary materials, $P < 0.001$). Soil microbial respiration per SOC was around 33%
296 higher in the surface soil under grazing as compared to mowing (Fig. 1B). In contrast, it was
297 greater in the subsurface soil under mowing than under grazing treatment. Including initial
298 SOC stock as covariate resulted only in significant effect of treatment \times depth interaction on
299 soil microbial respiration per SOC (Table S1, Supplementary materials, $P = 0.004$).

300 MBC per SOC was highest in the surface soil under grazing (20 $\mu\text{g C mg}^{-1} \text{ SOC}$, Fig.
301 1C). Mowing treatment resulted in two times lower MBC per SOC in the surface soil
302 compared to grazing treatment. After correction for initial SOC stock, treatment ($P < 0.001$)
303 and their interaction ($P < 0.001$) showed significant effects on $q\text{CO}_2$. Mowing treatment
304 resulted in higher $q\text{CO}_2$ at both depths as compared to grazing treatment (Fig. 1D, $P = 0.02$).

305 Microbial C:N ratio ranged between 4.9 and 6.4. It was affected by treatments in all
306 depths showing higher values under mowing (Fig. 2A). After taking into account initial SOC
307 stock, the treatment effect was still significant (Table S1, Supplementary materials, $P < 0.001$).
308 Active microbial biomass was also higher under mowing at both depths compared to grazing
309 treatment (Fig. 2B, $P = 0.02$). The highest specific microbial growth rate (Fig. 2C) was
310 recorded in subsurface soils without difference between treatments. But in surface soils, the
311 specific microbial growth rate was higher under grazing than under mowing (Fig. 2C).
312 However, ANCOVA with initial SOC stock as covariate decreased the significance treatment
313 effects ($P = 0.05$) on specific microbial growth rate but increased the depth effect Table S1,
314 Supplementary materials, $P < 0.001$).

315 Treatment effect on absolute enzyme activities is presented only for leucine
316 aminopeptidase in surface soil and chitinase and phosphatase in subsurface soil (Fig S2,
317 Supplementary materials). When the initial SOC stock was used as covariate, treatment
318 differences between enzyme activities per MBC were observed for all enzymes (except
319 leucine aminopeptidase) in surface soil. Soil under mowing treatment showed 2-2.5 times
320 higher enzyme activity per MBC under mowing compared to soil under grazing (Fig. 3). The
321 differences between treatments were more pronounced in surface soil for activities of
322 chitinase, β -galactosidase, β -glucosidase and phosphatase (Table S1, Supplementary
323 materials).

324

325 3.4. *Principal component analysis*

326 Principal component analysis based on SOC normalised data of all soil properties
327 showed that the first two factors explained 54.4% of the variation (Fig. 4). The first
328 component (Dim1) was related to microbial functioning, as it was strongly associated with
329 the soil microbial properties MBC and MBN per SOC in negative direction. The positive
330 direction was related to the lipase activity per MBC. The second component (Dim2) was
331 explained by variables related to polysaccharides. It was positively correlated with enzymes
332 participating in polysaccharide degradation and negatively with polysaccharide ratios. The
333 clustering of samples allowed to separate surface soil and subsurface soil samples along both
334 axes, while surface soil samples were additionally separated by treatments along the first axes
335 (Fig. 4). Subsurface soil samples were differentiated from surface soil by high neutral
336 polysaccharide monomer ratio, low enzymes activities per MBC, MBC and MBN per SOC.
337 Treatments in surface soil were separated by C- and N-cycle enzyme activity and MBC and
338 MBN per SOC. We also applied ANCOVA with initial SOC stock as a covariate on new
339 PCA coordinates which resulted in significant effects of treatment, depth and their

340 interaction. Treatment effect was more pronounced on Dim1, while Dim2 was more affected
341 by depth.

342

343 **4. Discussion**

344 *4.1. Effect of grazing and mowing on chemical properties of surface soil*

345 Since the primary factor of SOM formation is organic matter input (Fujisaki et al.,
346 2018; Kögel-Knabner, 2002), higher SOC and N contents in the surface soil under grazing
347 system might be explained by greater C input compared to mowing systems. This was shown
348 through ecosystem flux measurements at these plots (Senapati et al., 2014). Moreover, dung
349 return comprising about 50-80% of plant biomass could also favour higher SOC and N
350 content under grazing (Soussana et al., 2006). Even if mowing leads to some biomass input in
351 the form of plant material lost during grass removal (Sanaullah et al., 2010), the amount is
352 not enough to reach a similar input level than under grazing. Additionally, the lower pH
353 under mowing could contribute to indirect losses of SOC via changing C cycle and microbial
354 functioning (Kemmitt et al., 2006). Consequently, our results suggest that temperate loamy
355 soil under grazing is more prone to higher SOC contents when compared to mowing.

356 With regards to the biogeochemical composition of SOC, we did not find any
357 differences in non-cellulosic polysaccharide concentrations. These results are in agreement
358 with other studies showing that the soils' polysaccharide content is more or less stable and
359 even plant removal does not have a strong effect on the total polysaccharide concentrations
360 (Marchus et al., 2018). Soil lignin content, in contrast, was lower under grazing than mowing.
361 As lignin is a biomarker for plant-derived organic matter and more difficult to decompose,
362 because it requires a specific enzyme system (Buswell et al., 1987; Thevenot et al., 2010),
363 lower exportation of plant biomass and lignin input via dung deposition in soil under grazing
364 would suggest the opposite trend. However, dung contains only small amounts of lignin

365 (Dungait et al., 2005), which is relatively instable being degraded during one year (Dungait et
366 al., 2008). All lignin parameters (except the C/V ratio) suggested that lignin was less
367 degraded in the grazing than the mowing system. More acid pH in fertilised mowing systems
368 could have favoured the activity of lignin-degrading fungi (Couto et al., 2006). In mowing
369 systems microbial activity is fuelled exclusively by plant litter, whereas in grazing systems
370 organic matter input is supplied also by animal depositions. We hypothesise that this could
371 lead to contrasting quantitative lignin inputs, but could also impact its decomposition. Our
372 data show that lignin degradation in the mowing system is slower and less complete than in
373 the grazing system, leading to accumulation of partially degraded lignin molecules (Filley et
374 al., 2006). Therefore, lignin in the mowing system was characterised by a higher state of
375 degradation and at the same time its contribution to SOC was higher as compared to the
376 grazing system.

377 4.2. *Effect of grazing treatment on biological properties of surface soil*

378 Higher maturity and sustainability of the grazing system was shown by higher MBC per
379 SOC together with a lower qCO_2 (Anderson and Domsch, 2010). Higher qCO_2 in the mowing
380 system indicates that the microbial communities were less efficient and respired more C to
381 maintain metabolic activity as compared to those under grazing (Anderson, 2003).
382 Microorganisms are the main SOM decomposers leading to release of greenhouse gases and
383 nutrients in natural as well as in managed soils (Bardgett et al., 2008; Gougoulias et al.,
384 2014). This is particularly relevant for grazed pastures. Higher soil microbial respiration and
385 microbial CO_2 –C per unit SOC (soil microbial respiration per SOC) in the grazing system
386 was probably related to dung input with a huge amount of easily available compounds (Chu
387 et al., 2007; Marinari et al., 2000).

388 Contrary to our expectations, absolute enzyme activity did not differ among the
389 treatments, even after normalisation by SOC. A treatment effect was only observed after

390 normalisation by MBC, which expresses microbial activity in terms of enzyme production.
391 The enzymatic activities per MBC were higher in the mowing system as compared to the
392 grazing one, indicating that microorganisms in mowed soil produced enzymes more actively
393 than those under grazing. Microbial communities in the mowing system stayed active and
394 were investing in enzyme production probably to adapt to less decomposable organic
395 materials with higher lignin contents (see above). This maintenance of active state requires a
396 lot of energy, consequently, it could change C-cycling rates and decomposition of SOM
397 (Schimel and Schaeffer, 2012; Wang et al., 2014).

398 Microbial communities in the mowed soil are probably characterised by a higher
399 contribution of fungi than those of the grazed soil because we recorded a higher C:N ratio of
400 the microbial biomass (Joergensen and Emmerling, 2006) and more acid pH. Lower specific
401 growth rates in the mowing system may indicate relative domination of K-strategists in the
402 microbial community, which are more adapted to nutrient poor conditions (Strickland and
403 Rousk, 2010; Xu et al., 2017) and the decomposition of specific substances, such as plant
404 material containing high amounts of biopolymers (Fontaine et al., 2003). As illustrated by
405 lower enzyme activity per MBC, microorganisms in the grazing system invested less energy
406 for the degradation of complex compounds than those of the mowing system, most probably
407 because of higher availability of easily decomposable substrates. These conditions favour r-
408 strategists (Fierer et al., 2007; Xu et al., 2017) and thus stimulate microbial activity, as shown
409 by higher MBC per SOC and higher soil microbial respiration under the grazing as compared
410 to mowing system. As a consequence, the biogeochemical soil functioning under the two
411 management practices is quite different. This may affect significantly SOM formation, which
412 is favoured in systems with intensive microbial processing of C input (Kallenbach et al.,
413 2016; Liang et al., 2017) thus corroborating the high SOC contents observed under grazing.

414 4.3. *Less pronounced treatment effects in subsurface soil*

415 Treatment effects on soil properties were less pronounced in subsurface soil compared
416 to surface soil. Enhanced leaching and activity of soil fauna (Bohlen et al., 2004; Rumpel and
417 Kögel-Knabner, 2011) promote nutrient transport to subsurface soil under grazing which
418 resulted in higher SOC and N contents in subsurface soil under grazing than the one under
419 mowing. Treatment effects in the subsurface soil were neither observed for non-cellulosic
420 polysaccharide content and origin nor for lignin content or its degradation status. Since
421 lignins are typical indicators of plant input (Kögel-Knabner, 2002), this could indicate that
422 grazing and mowing have only small effects on plant rooting behaviour at lower depths.

423 On the other hand, the treatment effects on MBC and MBN was also observable in
424 subsurface soil. Soil microbial respiration did not differ between the treatments but microbial
425 CO_2 –C per SOC and $q\text{CO}_2$ were higher in the subsurface soil under mowing indicating that
426 the microbial communities used C inefficiently, similarly to surface soil. Higher
427 galactosidase activity in the subsurface soil of the mowing treatment is related to higher
428 contribution of galactose monomers in grass roots compared to grass leaves (Schädel et al.,
429 2010). As lipase is hydrolysing triglycerides, higher lipase activity in the subsurface soil
430 indicates accumulation of lipid compounds at depth, which probably serve as C source for
431 microorganisms under C-limiting conditions (Heitkötter et al., 2017).

432 The absence of treatment separation for the subsurface soils on the PCA plot might
433 indicate that in deeper soil probably more time is required to make treatment effects
434 observable. It was interesting to note that chemical properties related to SOM composition
435 were not sensitive to treatment effects in the subsurface soil, whereas microbial properties
436 were. This is in agreement with other studies, which showed that microbial properties are
437 most sensitive to changes introduced by management activities (Allison and Martiny, 2008;
438 Bending et al., 2004).

439

440 **5. Conclusions**

441 In this study we investigated the effect of grazing and mowing treatments on soil
442 biogeochemical and microbial properties. Our data indicated significant differences in the soil
443 organic matter composition as well as microbial functioning of the two treatments. Both plots
444 were also characterized by contrasting SOC contents and pH values. The grazing system was
445 characterized by (1) more efficient microbial community and (2) less decomposed organic
446 matter as compared to the mowing system. We conclude that the harvesting regime by
447 grazing or mowing affects the biogeochemical functioning of grassland soils. Even though
448 both systems are favorable to SOC storage, grazing might be preferable to mowing because it
449 leads to better substrate quality and more efficient microbial functioning. Although SOM
450 changes were only evident in surface soil, microbial properties suggest that these processes
451 are also occurring in subsurface soil.

452

453 **6. Conflicts of interests**

454 We state that there is no conflict of interests.

455

456 **7. Acknowledgements**

457 The research leading to these results has received funding principally from the New
458 Zealand Government to support the objectives of the Livestock Research Group of the Global
459 Research Alliance on Agricultural Greenhouse Gases - Grants SOW12-GPLER-LCR-PM
460 (Proposal ID 16949-15 LCR), the ANR (ANR-11-INBS-0001), the CNRS-INSU and the
461 INRA/Region Nouvelle-Aquitaine. We are grateful to Xavier Charrier, Jean-François
462 Bouhiron, Jerome Chargelegue, Marie-Laure Decau for help with site access, organization,
463 and field sampling and to Karin Schmidt and Valerie Pouteau for laboratory assistance. Two

464 anonymous reviewers are acknowledged for their valuable comments on an earlier version of
465 the manuscript.

466

467 **8. References**

- 468 Abdalla, M., Hastings, A., Chadwick, D.R., Jones, D.L., Evans, C.D., Jones, M.B., Rees,
469 R.M., Smith, P., 2018. Critical review of the impacts of grazing intensity on soil organic
470 carbon storage and other soil quality indicators in extensively managed grasslands.
471 *Agric. Ecosyst. Environ.* 253, 62–81. <https://doi.org/10.1016/j.agee.2017.10.023>
- 472 Allison, S.D., Martiny, J.B.H., 2008. In the Light of Evolution, In the Light of Evolution.
473 National Academies Press, Washington, D.C. <https://doi.org/10.17226/12501>
- 474 Anderson, T.-H., 2003. Microbial eco-physiological indicators to asses soil quality. *Agric.*
475 *Ecosyst. Environ.* 98, 285–293. [https://doi.org/10.1016/S0167-8809\(03\)00088-4](https://doi.org/10.1016/S0167-8809(03)00088-4)
- 476 Anderson, T.-H., Domsch, K.H., 2010. Soil microbial biomass: The eco-physiological
477 approach. *Soil Biol. Biochem.* 42, 2039–2043.
478 <https://doi.org/10.1016/j.soilbio.2010.06.026>
- 479 Anderson, T.-H., Domsch, K.H., 1993. The metabolic quotient for CO₂ (qCO₂) as a specific
480 activity parameter to assess the effects of environmental conditions, such as ph, on the
481 microbial biomass of forest soils, *Soil Biology and Biochemistry*. Pergamon.
482 [https://doi.org/10.1016/0038-0717\(93\)90140-7](https://doi.org/10.1016/0038-0717(93)90140-7)
- 483 Bardgett, R.D., Freeman, C., Ostle, N.J., 2008. Microbial contributions to climate change
484 through carbon cycle feedbacks. *ISME J.* 2, 805–814.
485 <https://doi.org/10.1038/ismej.2008.58>
- 486 Bazot, S., Mikola, J., Nguyen, C., Robin, C., 2005. Defoliation-induced changes in carbon
487 allocation and root soluble carbon concentration in field-grown *Lolium perenne* plants:
488 Do they affect carbon availability, microbes and animal trophic groups in soil? *Funct.*
489 *Ecol.* 19, 886–896. <https://doi.org/10.1111/j.1365-2435.2005.01037.x>
- 490 Bending, G.D., Turner, M.K., Rayns, F., Marx, M.-C., Wood, M., 2004. Microbial and
491 biochemical soil quality indicators and their potential for differentiating areas under
492 contrasting agricultural management regimes. *Soil Biol. Biochem.* 36, 1785–1792.
493 <https://doi.org/10.1016/J.SOILBIO.2004.04.035>
- 494 Blagodatskaya, E., Kuzyakov, Y., 2013. Active microorganisms in soil: Critical review of
495 estimation criteria and approaches. *Soil Biol. Biochem.*

496 <https://doi.org/10.1016/j.soilbio.2013.08.024>

497 Bohlen, P.J., Pelletier, D.M., Groffman, P.M., Fahey, T.J., Fisk, M.C., 2004. Influence of
498 Earthworm Invasion on Redistribution and Retention of Soil Carbon and Nitrogen in
499 Northern Temperate Forests. *Ecosystems* 7, 13–27. [https://doi.org/10.1007/s10021-003-](https://doi.org/10.1007/s10021-003-0127-y)
500 [0127-y](https://doi.org/10.1007/s10021-003-0127-y)

501 Buswell, J.A., Odier, E., Kirk, T.K., 1987. Lignin Biodegradation. *Crit. Rev. Biotechnol.* 6,
502 1–60. <https://doi.org/10.3109/07388558709086984>

503 Chabbi, A., Kögel-Knabner, I., Rumpel, C., 2009. Stabilised carbon in subsoil horizons is
504 located in spatially distinct parts of the soil profile. *Soil Biol. Biochem.* 41, 256–261.
505 <https://doi.org/10.1016/j.soilbio.2008.10.033>

506 Chu, H., Lin, X., Fujii, T., Morimoto, S., Yagi, K., Hu, J., Zhang, J., 2007. Soil microbial
507 biomass, dehydrogenase activity, bacterial community structure in response to long-term
508 fertilizer management. *Soil Biol. Biochem.* 39, 2971–2976.
509 <https://doi.org/10.1016/J.SOILBIO.2007.05.031>

510 Cotrufo, M.F., Soong, J.L., Horton, A.J., Campbell, E.E., Haddix, M.L., Wall, D.H., Parton,
511 W.J., 2015. Formation of soil organic matter via biochemical and physical pathways of
512 litter mass loss. *Nat. Geosci.* 8, 776–779. <https://doi.org/10.1038/ngeo2520>

513 Couto, S.R., Moldes, D., Sanromán, M.A., 2006. Optimum stability conditions of pH and
514 temperature for ligninase and manganese-dependent peroxidase from *Phanerochaete*
515 *chrysosporium*. Application to in vitro decolorization of Poly R-478 by MnP. *World J.*
516 *Microbiol. Biotechnol.* 22, 607–612. <https://doi.org/10.1007/s11274-005-9078-0>

517 Dignac, M.-F., Derrien, D., Barré, P., Barot, S., Cécillon, L., Chenu, C., Chevallier, T.,
518 Freschet, G.T., Garnier, P., Guenet, B., Hedde, M., Klumpp, K., Lashermes, G., Maron,
519 P.-A., Nunan, N., Roumet, C., Basile-Doelsch, I., 2017. Increasing soil carbon storage:
520 mechanisms, effects of agricultural practices and proxies. A review. *Agron. Sustain.*
521 *Dev.* 37, 14. <https://doi.org/10.1007/s13593-017-0421-2>

522 Dungait, J.A.J., Bol, R., Bull, I.D., Evershed, R.P., 2009. Tracking the fate of dung-derived
523 carbohydrates in a temperate grassland soil using compound-specific stable isotope
524 analysis. *Org. Geochem.* 40, 1210–1218.
525 <https://doi.org/10.1016/j.orggeochem.2009.08.001>

526 Dungait, J.A.J., Bol, R., Evershed, R.P., 2005. Quantification of dung carbon incorporation in
527 a temperate grassland soil following spring application using bulk stable carbon isotope
528 determinations. *Isotopes Environ. Health Stud.* 41, 3–11.
529 <https://doi.org/10.1080/10256010500053516>

530 Dungait, J.A.J., Stear, N.A., van Dongen, B.E., Bol, R., Evershed, R.P., 2008. Off-line
531 pyrolysis and compound-specific stable carbon isotope analysis of lignin moieties: a
532 new method for determining the fate of lignin residues in soil. *Rapid Commun. Mass*
533 *Spectrom.* 22, 1631–1639. <https://doi.org/10.1002/rcm.3454>

534 Ec.europa.eu, 2018. Share of main land types in utilised agricultural area (UAA) by NUTS 2
535 regions [WWW Document]. URL <https://ec.europa.eu/eurostat/data/database> (accessed
536 1.23.19).

537 Eder, E., Spielvogel, S., Kölbl, A., Albert, G., Kögel-Knabner, I., 2010. Analysis of
538 hydrolysable neutral sugars in mineral soils: Improvement of alditol acetylation for gas
539 chromatographic separation and measurement. *Org. Geochem.* 41, 580–585.
540 <https://doi.org/10.1016/J.ORGGEOCHEM.2010.02.009>

541 Eivazi, F., Tabatabai, M.A., 1990. Factors affecting glucosidase and galactosidase activities
542 in soils. *Soil Biol. Biochem.* 22, 891–897. [https://doi.org/10.1016/0038-0717\(90\)90126-](https://doi.org/10.1016/0038-0717(90)90126-K)
543 [K](https://doi.org/10.1016/0038-0717(90)90126-K)

544 Fierer, N., Bradford, M.A., Jackson, R.B., 2007. Toward an ecological classification of soil
545 bacteria. *Ecology* 88, 1354–1364. <https://doi.org/10.1890/05-1839>

546 Filley, T.R., Nierop, K.G.J., Wang, Y., 2006. The contribution of polyhydroxyl aromatic
547 compounds to tetramethylammonium hydroxide lignin-based proxies. *Org. Geochem.*
548 37, 711–727. <https://doi.org/10.1016/J.ORGGEOCHEM.2006.01.005>

549 Fujisaki, K., Chevallier, T., Chapuis-Lardy, L., Albrecht, A., Razafimbelo, T., Masse, D.,
550 Ndour, Y.B., Chotte, J.-L., 2018. Soil carbon stock changes in tropical croplands are
551 mainly driven by carbon inputs: A synthesis. *Agric. Ecosyst. Environ.* 259, 147–158.
552 <https://doi.org/10.1016/j.agee.2017.12.008>

553 Gavrichkova, O., Moscatelli, M.C., Kuzyakov, Y., Grego, S., Valentini, R., 2010. Influence
554 of defoliation on CO₂ efflux from soil and microbial activity in a Mediterranean
555 grassland. *Agric. Ecosyst. Environ.* 136, 87–96.
556 <https://doi.org/10.1016/j.agee.2009.11.015>

557 Gougoulias, C., Clark, J.M., Shaw, L.J., 2014. The role of soil microbes in the global carbon
558 cycle: tracking the below-ground microbial processing of plant-derived carbon for
559 manipulating carbon dynamics in agricultural systems. *J. Sci. Food Agric.* 94, 2362–
560 2371. <https://doi.org/10.1002/jsfa.6577>

561 Havstad, K.M., Peters, D.P.C., Skaggs, R., Brown, J., Bestelmeyer, B., Fredrickson, E.,
562 Herrick, J., Wright, J., 2007. Ecological services to and from rangelands of the United
563 States. *Ecol. Econ.* 64, 261–268. <https://doi.org/10.1016/J.ECOLECON.2007.08.005>

564 Hedges, J.I., Ertel, J.R., 1982. Characterization of lignin by gas capillary chromatography of
565 cupric oxide oxidation products. *Anal. Chem.* 54, 174–178.
566 <https://doi.org/10.1021/ac00239a007>

567 Heitkötter, J., Heinze, S., Marschner, B., 2017. Relevance of substrate quality and nutrients
568 for microbial C-turnover in top- and subsoil of a Dystric Cambisol. *Geoderma* 302, 89–
569 99. <https://doi.org/10.1016/j.geoderma.2017.04.029>

570 IPCC, 2018. Special Report. Global Warming of 1.5 °C. <https://www.ipcc.ch/sr15/>.

571 Jenkinson, D.S., Brookes, P.C., Powlson, D.S., 2004. Measuring soil microbial biomass. *Soil*
572 *Biol. Biochem.* 36, 5–7. <https://doi.org/10.1016/J.SOILBIO.2003.10.002>

573 Joergensen, R.G., Emmerling, C., 2006. Methods for evaluating human impact on soil
574 microorganisms based on their activity, biomass, and diversity in agricultural soils. *J.*
575 *Plant Nutr. Soil Sci.* 169, 295–309. <https://doi.org/10.1002/jpln.200521941>

576 Kallenbach, C.M., Frey, S.D., Grandy, A.S., 2016. Direct evidence for microbial-derived soil
577 organic matter formation and its ecophysiological controls. *Nat. Commun.* 7, 1–10.
578 <https://doi.org/10.1038/ncomms13630>

579 Kemmitt, S.J., Wright, D., Goulding, K.W.T., Jones, D.L., 2006. pH regulation of carbon and
580 nitrogen dynamics in two agricultural soils. *Soil Biol. Biochem.* 38, 898–911.
581 <https://doi.org/10.1016/j.soilbio.2005.08.006>

582 Koch, O., Tscherko, D., Kandeler, E., 2007. Temperature sensitivity of microbial respiration,
583 nitrogen mineralization, and potential soil enzyme activities in organic alpine soils.
584 *Global Biogeochem. Cycles* 21. <https://doi.org/10.1029/2007GB002983>

585 Kögel-Knabner, I., 2002. The macromolecular organic composition of plant and microbial
586 residues as inputs to soil organic matter. *Soil Biol. Biochem.* 34, 139–162.
587 [https://doi.org/10.1016/S0038-0717\(01\)00158-4](https://doi.org/10.1016/S0038-0717(01)00158-4)

588 Kögel, I., Bochter, R., 1985. Characterization of lignin in forest humus layers by high-
589 performance liquid chromatography of cupric oxide oxidation products. *Soil Biol.*
590 *Biochem.* 17, 637–640. [https://doi.org/10.1016/0038-0717\(85\)90040-9](https://doi.org/10.1016/0038-0717(85)90040-9)

591 Liang, C., Schimel, J.P., Jastrow, J.D., 2017. The importance of anabolism in microbial
592 control over soil carbon storage. *Nat. Microbiol.* 2, 17105.
593 <https://doi.org/10.1038/nmicrobiol.2017.105>

594 Marchus, K.A., Blankinship, J.C., Schimel, J.P., 2018. Environmental controls on
595 extracellular polysaccharide accumulation in a California grassland soil. *Soil Biol.*
596 *Biochem.* <https://doi.org/10.1016/j.soilbio.2018.07.009>

597 Marinari, S., Masciandaro, G., Ceccanti, B., Grego, S., 2000. Influence of organic and

598 mineral fertilisers on soil biological and physical properties. *Bioresour. Technol.* 72, 9–
599 17. [https://doi.org/10.1016/S0960-8524\(99\)00094-2](https://doi.org/10.1016/S0960-8524(99)00094-2)

600 Marx, M.-C., Kandeler, E., Wood, M., Wermbter, N., Jarvis, S.C., 2005. Exploring the
601 enzymatic landscape: distribution and kinetics of hydrolytic enzymes in soil particle-size
602 fractions. *Soil Biol. Biochem.* 37, 35–48.
603 <https://doi.org/10.1016/J.SOILBIO.2004.05.024>

604 McSherry, M.E., Ritchie, M.E., 2013. Effects of grazing on grassland soil carbon: a global
605 review. *Glob. Chang. Biol.* 19, 1347–1357. <https://doi.org/10.1111/gcb.12144>

606 Medina-Roldán, E., Bardgett, R.D., 2011. Plant and soil responses to defoliation: A
607 comparative study of grass species with contrasting life history strategies. *Plant Soil*
608 344, 377–388. <https://doi.org/10.1007/s11104-011-0756-4>

609 Ngo, P.T., Rumpel, C., Dignac, M.-F., Billou, D., Duc, T.T., Jouquet, P., 2011.
610 Transformation of buffalo manure by composting or vermicomposting to rehabilitate
611 degraded tropical soils. *Ecol. Eng.* 37, 269–276.
612 <https://doi.org/10.1016/j.ecoleng.2010.11.011>

613 Panikov, N.S., Sizova, M. V., 1996. A kinetic method for estimating the biomass of microbial
614 functional groups in soil. *J. Microbiol. Methods* 24, 219–230.
615 [https://doi.org/10.1016/0167-7012\(95\)00074-7](https://doi.org/10.1016/0167-7012(95)00074-7)

616 Piñeiro, G., Paruelo, J.M., Oesterheld, M., Jobbágy, E.G., 2010. Pathways of Grazing Effects
617 on Soil Organic Carbon and Nitrogen. *Rangel. Ecol. Manag.* 63, 109–119.
618 <https://doi.org/10.2111/08-255.1>

619 Post, W.M., Kwon, K.C., 2000. Soil carbon sequestration and land-use change: processes and
620 potential. *Glob. Chang. Biol.* 6, 317–327. <https://doi.org/10.1046/j.1365-2486.2000.00308.x>

622 Puche, N.J.B., Senapati, N., Klumpp, K., Fléchar, C.R., Kirschbaum, M.U.F., Chabbi, A.,
623 2019. Modelling carbon and water fluxes of managed grasslands: comparing flux
624 variability and net carbon budgets between grazed and mowed systems. *Agronomy*.

625 Razavi, B.S., Blagodatskaya, E., Kuzyakov, Y., 2015. Nonlinear temperature sensitivity of
626 enzyme kinetics explains canceling effect—a case study on loamy haplic Luvisol. *Front.*
627 *Microbiol.* 6. <https://doi.org/10.3389/fmicb.2015.01126>

628 Rumpel, C., Crème, A., Ngo, P., Velásquez, G., Mora, M., Chabbi, A., 2015. The impact of
629 grassland management on biogeochemical cycles involving carbon, nitrogen and
630 phosphorus. *J. soil Sci. plant Nutr.* 15(2), 353–371. <https://doi.org/10.4067/S0718-95162015005000034>

631

- 632 Rumpel, C., Dignac, M.-F., 2006. Gas chromatographic analysis of monosaccharides in a
633 forest soil profile: Analysis by gas chromatography after trifluoroacetic acid hydrolysis
634 and reduction–acetylation. *Soil Biol. Biochem.* 38, 1478–1481.
635 <https://doi.org/10.1016/J.SOILBIO.2005.09.017>
- 636 Rumpel, C., Kögel-Knabner, I., 2011. Deep soil organic matter---a key but poorly understood
637 component of terrestrial C cycle. *Plant Soil* 338, 143–158.
638 <https://doi.org/10.1007/s11104-010-0391-5>
- 639 Rumpel, C., Lehmann, J., Chabbi, A., 2018. ‘4 per 1,000’ initiative will boost soil carbon for
640 climate and food security. *Nat.* 2020 5537686.
- 641 Sanaullah, M., Chabbi, A., Lemaire, G., Charrier, X., Rumpel, C., 2010. How does plant leaf
642 senescence of grassland species influence decomposition kinetics and litter compounds
643 dynamics? *Nutr. Cycl. Agroecosystems* 88, 159–171. [https://doi.org/10.1007/s10705-](https://doi.org/10.1007/s10705-009-9323-2)
644 [009-9323-2](https://doi.org/10.1007/s10705-009-9323-2)
- 645 Schädel, C., Blöchl, A., Richter, A., Hoch, G., 2010. Quantification and monosaccharide
646 composition of hemicelluloses from different plant functional types. *Plant Physiol.*
647 *Biochem.* 48, 1–8. <https://doi.org/10.1016/J.PLAPHY.2009.09.008>
- 648 Schimel, J.P., Schaeffer, S.M., 2012. Microbial control over carbon cycling in soil. *Front.*
649 *Microbiol.* <https://doi.org/10.3389/fmicb.2012.00348>
- 650 Senapati, N., Chabbi, A., Gastal, F., Smith, P., Mascher, N., Loubet, B., Cellier, P., Naisse,
651 C., 2014. Net carbon storage measured in a mowed and grazed temperate sown
652 grassland shows potential for carbon sequestration under grazed system. *Carbon Manag.*
653 5, 131–144. <https://doi.org/10.1080/17583004.2014.912863>
- 654 Senapati, N., Chabbi, A., Giotri, A.F., Yeluripati, J.B., Smith, P., 2016. Modelling nitrous
655 oxide emissions from mown-grass and grain-cropping systems: Testing and sensitivity
656 analysis of DailyDayCent using high frequency measurements. *Sci. Total Environ.* 572,
657 955–977. <https://doi.org/10.1016/j.scitotenv.2016.07.226>
- 658 Smith, P., Martino, D., Cai, Z., Gwary, D., Janzen, H., Kumar, P., McCarl, B., Ogle, S.,
659 O’Mara, F., Rice, C., Scholes, B., Sirotenko, O., Howden, M., McAllister, T., Pan, G.,
660 Romanenkov, V., Schneider, U., Towprayoon, S., Wattenbach, M., Smith, J., 2008.
661 Greenhouse gas mitigation in agriculture. *Philos. Trans. R. Soc. B Biol. Sci.* 363, 789–
662 813. <https://doi.org/10.1098/rstb.2007.2184>
- 663 Soussana, J.-F., Loiseau, P., Vuichard, N., Ceschia, E., Balesdent, J., Chevallier, T.,
664 Arrouays, D., 2006. Carbon cycling and sequestration opportunities in temperate
665 grasslands. *Soil Use Manag.* 20, 219–230. <https://doi.org/10.1111/j.1475->

666 2743.2004.tb00362.x

667 Strickland, M.S., Rousk, J., 2010. Considering fungal: Bacterial dominance in soils -
668 Methods, controls, and ecosystem implications. *Soil Biol. Biochem.* 42, 1385–1395.
669 <https://doi.org/10.1016/j.soilbio.2010.05.007>

670 Tautges, N.E., Chiartas, J.L., Gaudin, A.C.M., O’Geen, A.T., Herrera, I., Scow, K.M., 2019.
671 Deep soil inventories reveal that impacts of cover crops and compost on soil carbon
672 sequestration differ in surface and subsurface soils. *Glob. Chang. Biol.* 25, 3753–3766.
673 <https://doi.org/10.1111/gcb.14762>

674 Thevenot, M., Dignac, M.-F.F., Rumpel, C., 2010. Fate of lignins in soils: A review. *Soil*
675 *Biol. Biochem.* 42, 1200–1211. <https://doi.org/10.1016/J.SOILBIO.2010.03.017>

676 Vance, E.D., Brookes, P.C., Jenkinson, D.S., 1987. An extraction method for measuring soil
677 microbial biomass C. *Soil Biol. Biochem.* 19, 703–707. [https://doi.org/10.1016/0038-](https://doi.org/10.1016/0038-0717(87)90052-6)
678 [0717\(87\)90052-6](https://doi.org/10.1016/0038-0717(87)90052-6)

679 Wang, G., Mayes, M.A., Gu, L., Schadt, C.W., 2014. Representation of dormant and active
680 microbial dynamics for ecosystem modeling. *PLoS One* 9.
681 <https://doi.org/10.1371/journal.pone.0089252>

682 Wilson, C.H., Strickland, M.S., Hutchings, J.A., Bianchi, T.S., Flory, S.L., 2018. Grazing
683 enhances belowground carbon allocation, microbial biomass, and soil carbon in a
684 subtropical grassland. *Glob. Chang. Biol.* 24, 2997–3009.
685 <https://doi.org/10.1111/gcb.14070>

686 Wutzler, T., Blagodatsky, S.A., Blagodatskaya, E., Kuzyakov, Y., 2012. Soil microbial
687 biomass and its activity estimated by kinetic respiration analysis – Statistical guidelines.
688 *Soil Biol. Biochem.* 45, 102–112. <https://doi.org/10.1016/j.soilbio.2011.10.004>

689 Xu, S., Silveira, M.L., Inglett, K.S., Sollenberger, L.E., Gerber, S., 2017. Soil microbial
690 community responses to long-term land use intensification in subtropical grazing lands.
691 *Geoderma* 293, 73–81. <https://doi.org/10.1016/j.geoderma.2017.01.019>

692

693

694

695 **Tables**

696 **Table 1.** General soil properties under two grassland management practices (grazing and
 697 mowing) at in surface soil (0-10 cm) and subsurface soil (20-30 cm).

	Treatment	pH	SOC content mg g ⁻¹	N mg g ⁻¹	δ ¹³ C ‰	δ ¹⁵ N ‰	C:N ratio
Surface soil	Grazing	5.95±0.09	21.4±0.81	2.2±0.09	-27.4±0.06	4.9±0.13	9.6±0.05
	Mowing	5.51±0.08	14.6±0.51	1.5±0.05	-27.0±0.05	5.0±0.09	9.6±0.07
Subsurface soil	Grazing	5.99±0.12	11.8±0.62	1.3±0.06	-26.7±0.06	6.2±0.08	9.2±0.06
	Mowing	6.01±0.13	8.6±0.44	0.9±0.05	-26.3±0.10	6.4±0.10	9.1±0.07
ANCOVA, F value (P values)							
SOC stocks in 2005		0.15 (0.70)	5.31 (0.03)	6.26 (0.02)	6.86 (0.01)	1.90 (0.18)	1.33 (0.26)
Treatment		3.37 (0.08)	30.3 (<0.001)	28.7 (<0.001)	17.2 (0.002)	0.35 (0.56)	1.19 (0.28)
Depth		5.89 (0.02)	181.8 (<0.001)	153.8 (<0.001)	132.5 (<0.001)	157.9 (<0.001)	52.5 (<0.001)
Treatment×Depth		4.27 (0.04)	0.68 (0.41)	0.86 (0.36)	0.28 (0.06)	0.25 (0.62)	0.37 (0.55)

698 Values are shown as the average of ten replicates and ±SE. Significant differences between the treatments are
 699 indicated by capital case letters. Lower case letters show significant differences with depth (P < 0.05).

700

701 **Table 2.** Non-cellulosic polysaccharides (NCP) signature in soil under two grassland
 702 management practices (grazing and mowing) at two depths (0-10 cm and 20-30 cm).

	Treatment	NCP content	NCP content per SOC	NCP monomers ratios		
		mg g ⁻¹	mg g ⁻¹ SOC	C6/C5 ¹	DesoxyC6/C5 ²	Man/Xyl ³
Surface soil	Grazing	6.61±0.23	308.98±6.3	0.80±0.02	0.35±0.01	0.54±0.02
	Mowing	4.45±0.18	306.63±11.5	0.84±0.02	0.34±0.01	0.61±0.02
Subsurface soil	Grazing	3.09±0.15	263.39±6.4	1.03±0.02	0.43±0.01	0.87±0.03
	Mowing	2.50±0.11	292.41±10.5	1.01±0.02	0.46±0.01	0.91±0.03
ANCOVA, F value (P values)		0.43 (0.52)				
SOC stocks in 2005		36.6 (<0.001)	2.50 (0.12)	5.42 (0.03)	3.81 (0.06)	4.74 (0.04)
Treatment		241.1	0.11 (0.74)	0.87 (0.36)	0.01 (0.91)	0.64 (0.43)
Depth		(<0.001)	11.5 (0.002)	122.2 (<0.001)	102.8 (<0.001)	166.3 (<0.001)
Treatment×Depth		19.7 (<0.001)	3.17 (0.08)	3.14 (0.09)	3.18 (0.08)	0.52 (0.48)

703
 704 ¹C6/C5 – the ratio of C6- to C5- sugar monomers, ²DesoxyC6/C5 – the ratio of desoxy C6- to desoxy
 705 C5- sugar monomers, ³Man/Xyl - the ratio of mannose to xylose. These ratios indicate the origin of non-
 706 cellulosic polysaccharides (microbial or plant).
 707

708

709 **Table 3.** Lignin signature in soil under two grassland management practices (grazing and
 710 mowing) in surface soil (0-10 cm) and subsurface soil (20-30 cm).

	Treatment	Lignin content	Lignin content per SOC	Lignin monomers ratios			
		mg g ⁻¹	mg g ⁻¹ SOC	C/V	S/V	(Ac/Al) _v	(Ac/Al) _s
Surface soil	Grazing	0.35±0.01	16.31±0.64	0.45±0.03	1.34±0.02	0.53±0.02	0.46±0.01
	Mowing	0.26±0.01	17.86±0.43	0.45±0.03	1.24±0.02	0.65±0.02	0.57±0.02
Subsurface soil	Grazing	0.19±0.02	16.22±0.57	0.37±0.02	1.33±0.03	0.66±0.01	0.54±0.02
	Mowing	0.16±0.01	18.86±0.89	0.37±0.03	1.30±0.02	0.63±0.03	0.56±0.02
ANCOVA, F value (P values)							
SOC stocks in 2005		0.05 (0.82)	10.9 (0.002)	1.59 (0.22)	1.99 (0.17)	0.08 (0.78)	18.6 (<0.001)
Treatment		15.3 (<0.001)	1.14 (0.29)	0.64 (0.43)	7.91 (0.01)	2.83 (0.10)	2.82 (0.10)
Depth		96.3 (<0.001)	0.62 (0.44)	8.58 (0.006)	0.95 (0.34)	9.13 (0.005)	13.1
Treatment×Depth		4.83 (0.04)	0.88 (0.36)	0.015 (0.90)	2.19 (0.15)	16.3 (<0.001)	<0.001


711 C/V – the ratio of cinnamyl phenols to syringyl phenols; S/V - the ratio of syringyl phenols to vanillyl
 712 phenols; (Ac/Al)_v – acid to aldehyde ratio of vanillyl phenols; (Ac/Al)_s – acid to aldehyde ratio of syringyl
 713 phenols. These ratios are indicators of lignin degradation state in soil.
 714

715

716

717

Figure captions



718

719

Figure 1. (A) Soil microbial respiration (SMR), (B) soil microbial respiration (SMR)

720

per soil organic carbon (SOC), (C) microbial biomass carbon (MBC) per soil organic carbon

721

(SOC) and (D) metabolic quotient ($q\text{CO}_2$) in soil under two grassland management practices

722

(grazing and mowing) in surface soil (0-10 cm) and subsurface soil (20-30 cm). Significant


723

differences between the treatments are indicated by *, ** and ***, representing probability at

724

the 5, 1, and 0.1% levels, respectively.

725



726

727 **Figure 2.** (A) Microbial C:N ratio, (B) the percentage of active microbial biomass


728 (AMB) and (C) specific microbial growth rate (μ) in soil under two grassland management

729 practices (grazing and mowing) in surface soil (0-10 cm) and subsurface soil (20-30 cm).

730 Significant differences between the treatments are indicated by *, ** and ***, representing

731 probability at the 5, 1, and 0.1% levels, respectively.

732



733

734

735


736

737

738

739

Figure 3. Boxplot of enzyme activity per unit of microbial biomass C (MBC) for nine enzymes under two grassland management practices (grazing and mowing) in surface soil (0-10 cm) and subsurface soil (20-30 cm). Significant differences between the treatments are indicated by *, ** and ***, representing probability at the 5, 1, and 0.1% levels, respectively.



740

741

Figure 4. Principal component analysis (PCA) for soil under grazing and mowing in

742

surface soil (0-10 cm) and in subsurface soil (20-30 cm). Only variables with quality of

743

representation (\cos^2) higher than 0.75 was shown on PCA plot.