



HAL
open science

Diet during pregnancy: Influence of social characteristics and migration in the ELFE cohort

Manik Kadawathagedara, Namanjeet Ahluwalia, Marie-noelle Dufourg, Anne Forhan, Marie Aline Charles, Sandrine Lioret, Blandine Lauzon-guillain

► To cite this version:

Manik Kadawathagedara, Namanjeet Ahluwalia, Marie-noelle Dufourg, Anne Forhan, Marie Aline Charles, et al.. Diet during pregnancy: Influence of social characteristics and migration in the ELFE cohort. *Maternal and Child Nutrition*, 2021, pp.e13140. 10.1111/mcn.13140 . hal-03144495

HAL Id: hal-03144495

<https://hal.inrae.fr/hal-03144495>

Submitted on 31 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Diet during pregnancy: Influence of social characteristics and migration in the ELFE cohort

Manik Kadawathagedara¹ | Namanjeet Ahluwalia² | Marie-Noelle Dufourg³ |
Anne Forhan¹ | Marie Aline Charles^{1,3}  | Sandrine Lioret¹  |
Blandine de Lauzon-Guillain¹ 

¹Université de Paris, CRESS, INSERM, INRAE, Paris, France

²UREN, Faculty of Medicine, University of Paris 13, Paris, France

³INED, INSERM, Joint Unit Elfe, Paris, France

Correspondence

Sandrine Lioret, CRESS Eq 6 EAROH, Hôpital Paul Brousse, Batiment Leriche, 16 av. Paul Vaillant Couturier, 94807 Villejuif Cedex, France.

Email: sandrine.lioret@inserm.fr

Funding information

Agence Nationale de la Recherche, Grant/Award Number: ANR-19-CE36-0006; SF-DOHAD/Blédina; French Institute for Demographic Studies (INED)

Abstract

Better adherence to dietary guidelines during pregnancy is supposed to result in healthier perinatal outcomes. We aim to characterize the diets of pregnant women by hypothesis-driven and exploratory approaches and describe potential social determinants. Analyses included 12 048 mothers from the French nationwide ELFE birth cohort. Dietary intake over the last three months of the pregnancy was assessed by a food frequency questionnaire. Two hypothesis-driven scores (the Diet Quality score, based on benchmarks derived from the National Health and Nutrition Program Guidelines, and the PANDiet score, based on nutrient intake) were calculated. Exploratory dietary patterns were also identified by principal component analysis. Multiple linear regressions were used to assess associations of maternal social characteristics with dietary patterns, accounting for the possible effect modification by their migration status. Five dietary patterns were identified: the Western, Balanced, Bread and toppings, Processed products, and Milk and breakfast cereals. Younger maternal age, single motherhood, unemployment and the presence of older children in the household were related to a suboptimal diet during pregnancy. The less acculturated the women were, the healthier and less processed their diets were, independent of their socio-economic position. Several social determinants of the quality of women's diets were however moderated by their migration status. These findings shed light on the relations between indicators of social vulnerability, such as single motherhood and unemployment, and poorer diet quality. Given the reduced diet quality that accompanies the acculturation process, it is of paramount importance to identify the specific factors or obstacles that affect migrant women in maintaining their diet quality advantage over the majority population.

KEYWORDS

acculturation, birth cohort, food frequency questionnaire, maternal diet, migrant, pregnancy

Sandrine Lioret and Blandine de Lauzon-Guillain contributed equally to this work.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. Maternal & Child Nutrition published by John Wiley & Sons Ltd.

1 | INTRODUCTION

Maternal diet is the major prenatal source of nutrients and can thus influence embryonic, placental, and fetal growth (Blumfield, Hure, Macdonald-Wicks, Smith, & Collins, 2013). Accordingly, maternal diet during pregnancy appears to be crucial for developing fetuses and their later-life health status (Abu-Saad & Fraser, 2010; McMillen et al., 2008). In particular, depletion of the fetal environment during pregnancy can induce a fetal adaptive response of the body, that prioritizes the growth of vital organs such as the brain over the development of others and leads to metabolic alterations (Hales & Barker, 2001). These adaptations are assumed to be especially harmful when followed by overnutrition of the newborn in the postnatal period. Beyond insufficient or excessive energy intake (Blumfield, Hure, Macdonald-Wicks, Smith, & Collins, 2012), the current problem in high-income countries is that of unbalanced diet and suboptimal micronutrient intake (Blumfield et al., 2013), especially in disadvantaged populations (Bertin et al., 2016; Lioret et al., 2012; Roberts, Cavill, Hancock, & Rutter, 2013).

Among the latter, cultural and ethnic minority groups, defined as migrants or population subgroups of migrant or indigenous background (Osei-Kwasi et al., 2016), are reported to have a higher prevalence of diet-related noncommunicable diseases, along with a suboptimal diet, compared with the native-born population (Faskunger, Eriksson, Johansson, Sundquist, & Sundquist, 2009; Patel et al., 2006). Beyond socio-economic position, various sociocultural factors have been shown to have a greater influence on dietary behaviours in cultural minority groups, including acculturation, a multi-dimensional process in which individuals adopt characteristic ways of life from the culture they are now living in that differ from the primary learning of their cultural background (D'Souza, Jayaweera, & Pickett, 2016; Osei-Kwasi et al., 2016). In their reviews, D'Souza and colleagues (D'Souza et al., 2016) noted that traditional pregnancy diets vary from society to society, and Urquia and colleagues (Urquia et al., 2010) noted that the association between foreign-born status and birth outcomes also varies and depends on both migrant subgroup and destination country.

Initiated in 2001, the French National Nutrition and Health Program (PNNS) aims to improve the health status of the population by acting on one of its major determinants: nutrition (S. Hercberg, Chat-Yung, & Chauliac, 2008). This programme includes guidelines on dietary intake and physical activity for the general population, along with guidelines for specific populations, such as pregnant women. Although several national surveys have monitored dietary intake and adherence to dietary guidelines in the French adult population (Bertin et al., 2016; Castetbon et al., 2009; Dubuisson et al., 2010; Volatier, 2000), these studies included too few pregnant women to assess their adherence to these recommendations. Assessing dietary patterns of pregnant women with specifically designed studies is an important public health objective, to make it possible to identify the groups at greatest risk so that they can receive more tailored information and interventions.

Key messages

- Beyond the well-known association of maternal education level and age with diet quality, these findings underline the relations of some indicators of social vulnerability, such as single motherhood and unemployment, with poorer diet quality during pregnancy.
- The less acculturated pregnant women were, the healthier and less processed their diets were, regardless of their socioeconomic position.
- Several demographic and socioeconomic determinants of the quality of women's diets were moderated by their migration status.

Two main classes of methods are used to derive dietary patterns: first, the production of diet quality scores or indices based on a priori benchmarks from nutritional guidelines or on nutrient intake, and second, the use of exploratory statistical methods, such as principal component analysis and cluster analysis (Kant, 2004). The latter a posteriori data reduction techniques are commonly used in nutritional epidemiology to summarize the variation in food intakes into a small number of patterns and have shown good reproducibility across studies (Newby & Tucker, 2004).

The objective of this study was to characterize the diet of pregnant women living in France, by using both hypothesis-driven (a priori) and exploratory or data-driven (a posteriori) approaches, and then to assess its association with demographic and socio-economic characteristics, while its moderation by migration status is considered.

2 | MATERIAL AND METHODS

2.1 | Study population

This analysis is based on data from the ELFE (Etude Longitudinale Française depuis l'Enfance) study, a multidisciplinary nationwide birth cohort including 18 329 children born in 2011 in a random sample of 349 maternity units in mainland France (Charles et al., 2019). Enrolment began in April 2011 and took place during 25 selected days, over four waves of 4–8 days each, each wave covering a different season. Infants met the inclusion criteria if they were singletons or twins born after 33 weeks of gestation to mothers aged 18 years or older who were not planning to move outside of metropolitan France in the next 3 years.

Overall, 51% of eligible parents agreed to their child's participation. Data were collected in standardized interviews conducted by trained interviewers and by self-administered questionnaires. Efforts

were made at inclusion to reach women who were not totally fluent in French: in particular, the information letter and consent form were also edited in the three languages most often spoken in France (after French), i.e. Arabic, Turkish and English. The interviews were also conducted in Soninké, Bambara and Wolof.

2.2 | Maternal diet during pregnancy

2.2.1 | Data collection

At the hospital before discharge, the mothers enrolled in the study completed a self-administered food frequency questionnaire (FFQ) that described their dietary intake over the last three months of the pregnancy. Consumption of 125 food items was collected, including 12 nonalcoholic and 4 alcoholic beverages, with frequency assessed by a 7-item scale from 'never' to 'more than once a day'. Usual portion sizes for different food types were estimated with photos indicating five possible sizes, derived from the SU.VI.MAX portion book (S. Hercberg, Deheeger, & Preziosi, 2002) for 75 commonly eaten food items; and for the 50 remaining food items, the midportion in the SU.VI.MAX portion book was automatically assigned. After the transformation of FFQ frequency categories into daily frequencies, daily intake was assessed, for each food item, by combining frequency of intake and portion size. Nutrient intakes were then calculated for all women by multiplying the daily intake of each food item by the nutritional values in the SU.VI.MAX nutrient composition database (S. Hercberg, 2006). The 125 food items were then classified into 48 food groups. Women for whom more than 10 FFQ items were missing were excluded. Conversely, when 10 or fewer items were missing, the missing values were imputed by the sample median. Women were deemed likely misreporters of their food intake, and thus excluded, when their estimated total energy intake was less than 933 kcal/day (3rd percentile) or exceeded 5072 kcal/day (97th percentile). Women were also asked to report their frequency of intake of industrially processed foods on a five-item scale ranging from 'never' to 'always': reduced-sugar and reduced-fat products, grouped together in the diet food group; prepacked foods; canned foods; and ready-prepared dishes.

The pregnant women's adherence to the French nutritional guidelines (ANSES, 2016) was assessed with two complementary a priori approaches. The first, the Diet Quality score, was based on consumption of the main food groups, and the second, the PANDiet score, on a range of nutrient intakes. Calculation of the Diet Quality score therefore used 17 quantitative benchmarks (Table S1) adapted from the PNNS-GS (National Health and Nutrition Program Guideline Score) (Estaquio et al., 2009). The methodology applied the principle of the percentage of each guideline followed by women, consistent with the approach developed in the MoBa cohort (von Ruesten et al., 2014). The overall score, which can range from 0 to 17, is the sum of the scores obtained on each item: the higher the score, the better the adherence to the national guidelines. Details of this score have previously been published (Kadawathagedara et al., 2017). At

the same time, the PANDiet score, previously developed and validated among French pregnant women (Bianchi, Mariotti, Verger, & Huneau, 2016), was calculated. It aims to measure the overall quality of a woman's diet by combining the probability of adequate intake for different nutrients, based on recommendations for women in the third trimester of pregnancy. The PANDiet score ranges from 0 to 100, with a higher score again indicating better nutritional quality.

2.2.2 | FFQ validation study

A separate study sought to validate the FFQ. It enrolled participants from maternity clinics during their appointment with their obstetrician in major cities across France. Of the 128 women informed about the study by their physician and asked to participate, 105 consented. Among them, 67 (age: 31.4 ± 4.8 years) could give the study dietician information for at least one 24-h recall over the telephone (out of three initially planned within the last three months of pregnancy); and 62 women completed the FFQ assessing foods consumed during the last trimester. Participants received the study FFQ by mail a month before the planned due date and a telephone reminder to complete the FFQ within a week after their baby's birth. Participants with an incomplete FFQ ($n = 4$) and those for whom the estimated energy intake by FFQ was considered unrealistic ($n = 2$; i.e. 600 Kcal for one subject and 6000 Kcal for the other subject) were excluded from the analysis. The analysis covered the data from the 56 women who provided at least one 24-h recall and the FFQ were analysed. The methodology of this validation study is described in greater detail in the Supporting Information, as are its results (Table S2).

2.3 | Demographic and socio-economic characteristics

As the demographic and socio-economic characteristics collected at the two-month telephone interview were more detailed than at the maternity ward interview, we prioritized the data collected at 2 months when they conflicted. Demographic and socio-economic characteristics studied were as follows: mother's age at delivery (18–25, 25–29, 30–34 or ≥ 35 years), older children in the household (none, 1 or more), education level (up to lower secondary, upper secondary, intermediate, 3-year university degree, at least 5-year university degree), work status [employed, unemployed, not in the labour force (e.g. housewife, student, disabled, retired)], single motherhood (yes/no) and household income per consumption unit ($\leq \text{€}750$, $\text{€}751\text{--}1111$, $\text{€}1112\text{--}1500$, $\text{€}1501\text{--}1944$, $\text{€}1945\text{--}2500$, or $> \text{€}2500$ /month, that is, $\leq \$815$, $\$816\text{--}1207$, $\$1208\text{--}1630$, $\$1631\text{--}2113$, $\$2114\text{--}2717$, or $> \$2717$ /month), city size (rural area, $< 50\,000$ inhabitants, $50\,000\text{--}199\,999$ inhabitants, $200\,000\text{--}1\,999\,999$ inhabitants, Paris metropolitan area). Her migration status was defined in three categories: 'majority population', which included women born with a

French nationality to two French parents (in or outside of France); 'descendants of migrants' including women born in France with at least one non-French parent; and 'migrants' including women not born in France, and without French citizenship at birth.

2.4 | Sample selection

The study excluded infants whose parents withdrew consent during the first year ($n = 57$) (Figure S1). In twin births, random selection of one twin ($n = 287$ exclusions) made it possible to count each mother of twins only once in the analysis. The analyses excluded women who did not complete the FFQ ($n = 2,013$), or with more than 10 missing values ($n = 658$), with implausible energy intake ($n = 910$) or with missing data for use of industrially processed products during pregnancy ($n = 433$); this left 13 881 women for dietary pattern identification.

The complete case analysis, which excluded women with at least one missing value for demographic and socioeconomic characteristics, had a sample of 12 048 women.

Compared with the women included in these main analyses, the excluded women were younger (30.3 ± 5.6 years vs. 31.1 ± 4.7 years, $P < 0.0001$) and more likely to be single (9.8% vs. 3.1%, $P < 0.0001$) or to have been born abroad (19.5% vs. 7.1%, $P < 0.0001$), and less likely to have a Master's degree (17.2% vs. 21.1%, $P < 0.0001$) or to have been employed during pregnancy (56.6% vs. 76.4%, $P < 0.0001$).

2.5 | Statistical analyses

2.5.1 | Exploratory dietary patterns

Exploratory dietary patterns were derived by principal component analysis (PCA, FACTOR procedure; SAS software (Version 9.4, SAS Institute, Cary, NC, USA)) of the 46 standardized food groups consumed by more than half the sample. The number of factors to retain was determined with the diagram of eigenvalues (the Scree plot) and based on the interpretability of the factors (Cattell, 1966; Kline, 1994). A standardized score was calculated for each individual and pattern to quantify the individual's adherence to the pattern.

The association between each of the dietary scores, whether derived by hypothesis-driven or exploratory approaches, were analysed with Pearson's correlations. Recommendations for interpreting these correlation coefficients are: low 0.1–0.3, moderate 0.3–0.5 and high >0.5 (Cohen, 1988).

2.5.2 | Associations with demographic and socio-economic characteristics

Associations of demographic and socio-economic characteristics with maternal hypothesis-driven and exploratory dietary patterns

(outcomes) were tested by multiple linear regressions, which included all the demographic and socio-economic characteristics and were adjusted for variables related to the study design (maternity unit size and season of enrolment). Possible interactions (i.e. effect modifications) between migration status and all demographic and socio-economic characteristics were tested for their relations with the various dietary patterns.

2.5.3 | Sensitivity analyses

To take into account the inclusion procedure and biases related to nonconsent, the data were weighted (Juillard, 2015) in sensitivity analyses. Weighting also included calibration on margins from the state register's statistical data and the 2010 French National Perinatal study (Blondel, Lelong, Kermarrec, Goffinet, & National Coordination Group of the National Perinatal Surveys, 2012) on the following variables: age, region, marital status, migration status, level of education and primiparity. This weighting was calculated for the subsample that completed the FFQ.

In addition to the main analyses of the complete cases, analyses with multiple imputations dealt with missing data for demographic and socio-economic characteristics ($n = 13\ 618$). Missingness of each variable was less than 11% (10.6% for household income; 9.5% for maternal education level; 6.9% for maternal migration status; 6.9% for maternal employment; 6.4% for single parenthood; 6.4% for parity; and 0% for all other variables of interest). On the assumption that data were missing at random, five independent datasets were generated with the fully conditional specification method {MI procedure, FCS statement, NIMPUTE option; SAS software [Version 9.4, SAS Institute, Cary, NC, USA]} and pooled effect estimates were then calculated {MIANALYSE procedure SAS software}. The imputation models included all variables of interest after they were ranked in ascending order of missing data. Categorical variables were imputed with multinomial models, ordinal or binary variables with logistic regressions, and continuous variables with linear regressions.

Statistical analysis was performed with SAS software [SAS software ['SAS (computer program)', 2013] Version 9.4, SAS Institute, Cary, NC, USA]. The significance level was set at 0.05.

2.6 | Ethical considerations

Participating mothers provided written consent for themselves and for their child. Fathers signed the consent form for the child's participation when present at enrolment or were informed about their right to oppose it. The Advisory Committee for Treatment of Health Research Information (Comité Consultatif sur le Traitement des Informations pour la Recherche en Santé), the National Data Protection Authority (Commission Nationale Informatique et Libertés) and the National Statistics Council approved the ELFE study.

3 | RESULTS

3.1 | Identification of exploratory (PCA) dietary patterns in the ELFE study

Table 1 describes the sample's characteristics.

Five dietary patterns were identified that together explained 28% of the total variance. The first pattern was characterized by high positive factor loadings for French fries, fast foods, potatoes, mixed dishes, cakes and pastries, sauce (e.g. mayonnaise or ketchup), sweet snacks, red meat, potato chips and similar products, processed meat, candies and breaded fish (Table 2). It is labelled the 'Western' pattern. The second pattern was characterized by high positive factor loadings for cooked vegetables, raw vegetables, fish, fruit, whole-grain bread, legumes, yogurt and cottage cheese, and dried fruit and negative coefficients for sweetened beverages. It is referred to as the 'Balanced' pattern. The third pattern, with high coefficients for honey/jam, chocolate, butter and, to a lesser extent, cheese and bread, is named the 'Bread and toppings' pattern. The fourth pattern is labelled the 'Processed products' pattern in view of its high coefficients for prepacked foods, ready-prepared dishes, canned foods and diet foods. Finally, the fifth pattern was characterized by high positive coefficients for milk, cocoa powder, and breakfast cereals and negative coefficients for coffee. It is called the 'Milk and breakfast cereals' pattern.

The correlation between the Diet Quality and the PANDiet scores was moderate (0.43, $P < 0.0001$). The correlations of both the Diet Quality and the PANDiet scores with the exploratory Balanced

pattern score were respectively high (0.65, $P < 0.0001$) and moderate (0.42, $P < 0.0001$). The correlations of the remaining four exploratory dietary patterns with each of the Diet quality and the PANDiet scores were low, i.e. respectively -0.03 and 0.20 for the Western pattern ($P > 0.05$ and $P < 0.0001$); 0.01 and -0.05 for the Bread and toppings pattern ($P > 0.05$ and $P < 0.0001$); -0.03 and -0.14 for the Processed products pattern (both $P < 0.0001$); and 0.21 and 0.24 for the Milk and breakfast cereals pattern (both $P < 0.0001$).

3.2 | Demographic or socio-economic characteristics and maternal diet

Table S3 presents the bivariate associations of the maternal dietary patterns with demographic or socio-economic characteristics.

Moderating effects of maternal migration status

There was no interaction between migration status and any of the variables of interest for the Western, Bread and toppings and Milk and breakfast cereals dietary patterns. Conversely, a moderating effect of the variable 'migration status' was observed on the whole in the three healthier dietary patterns (i.e. Diet quality score, PANDiet score and Balanced) for the following factors: maternal education (P value for interactions ranging from <0.0001 to 0.046), maternal occupation (P value for interactions ranging from 0.0029 to 0.066), and household income (P value for interactions ranging from <0.0001 to 0.033). Interaction effects were also observed between migration status and both household income (P value = 0.09) and older children in the household (P value = 0.03) in their association with the Processed dietary pattern. Consequently, analyses involving the Western, Bread and toppings, and Milk and breakfast cereals dietary patterns used the entire sample, whereas the associations between demographic or socioeconomic characteristics and either of the three healthier or the Processed dietary patterns were assessed separately according to maternal migration status.

Western dietary pattern

Maternal age, maternal education level and household income were inversely related to this dietary pattern, while older children in the household, unemployment during pregnancy and single motherhood were positively related to it (Table 3). Maternal migration status was not associated with this dietary pattern.

Bread and toppings dietary pattern

Maternal age, maternal education level, older children in the household and household income were positively associated with the Bread and toppings pattern (Table 3). Maternal occupation and single motherhood were not associated with it. Migrant women and, to a lesser extent, descendants of migrants, had lower scores on the Bread and topping pattern than the majority population did.

Milk and breakfast cereals dietary pattern

Maternal age and older children in the household were negatively related to this dietary pattern (Table 3). Pregnant women of higher maternal education level scored higher on it, and migrant women higher than the majority population. No association was observed with household income, maternal occupation or single motherhood.

TABLE 1 Sample characteristics ($n = 12\,048$): The ELFE study, 2011

	% (n) ^a
Maternal age at delivery (years), mean (SD)	31.1 (4.7)
Maternal Master's degree: yes	21.1%
Maternal migration status	
Migrant	7.1%
Descendant of migrant	9.9%
Majority population	83.0%
Single motherhood	3.1%
First child	44.1%
Maternal work status during pregnancy	
Employed	76.4%
Unemployed	11.1%
Not in the labour force	12.5%
Household income per consumption unit (€/month) mean (SD) ^b	1686 (930)
Diet quality score ^c , mean (SD)	12.8 (1.2)
PANDiet score ^d , mean (SD)	55.7 (9.1)

^aNumbers are in % unless otherwise indicated.

^bThis corresponds to \$1832 (1011).

^cThe diet quality score could range from 0 to 17.

^dThe PANDiet score could range from 0 to 100.

TABLE 2 Factor loadings for dietary patterns during pregnancy, derived from principal component analysis ($n = 13\,881$): The ELFE study, 2011

	Pattern 1	Pattern 2	Pattern 3	Pattern 4	Pattern 5
French fries	0.54	-0.11	-0.30	-0.11	-0.21
Fast foods	0.51	-0.10	-0.07	0.12	-0.03
Potatoes	0.51	0.14	-0.30	-0.11	-0.16
Mixed dishes	0.50	0.06	-0.30	0.03	0.05
Cakes and pastries	0.49	-0.22	0.24	-0.10	-0.01
Sauce	0.44	-0.21	0.02	0.00	-0.14
Biscuits and chocolate or cereal bars	0.44	-0.24	0.35	-0.07	0.03
Red meat	0.40	0.16	-0.25	-0.02	-0.08
Chips and savoury biscuits	0.40	-0.18	0.09	0.04	-0.16
Processed meat	0.39	-0.04	0.06	0.17	-0.18
Candy	0.38	-0.27	0.25	-0.08	-0.07
Breaded fish	0.38	0.05	-0.24	-0.06	0.04
Dairy dessert	0.34	-0.14	0.23	-0.08	0.13
Other starchy foods	0.31	0.21	-0.23	0.00	0.00
Cream	0.29	0.05	0.21	0.02	-0.08
Chocolate spread	0.27	-0.26	0.25	-0.13	0.06
Ice cream	0.25	-0.08	0.05	-0.06	-0.08
Poultry	0.24	0.23	-0.19	0.06	-0.02
Shellfish	0.19	0.18	-0.13	0.00	0.04
Fruit juice	0.19	-0.08	0.15	-0.11	0.06
Cooked vegetables	0.21	0.60	-0.18	-0.14	0.06
Raw vegetables	0.10	0.56	0.05	0.01	-0.06
Fish	0.19	0.47	-0.11	0.00	0.06
Fruit	0.14	0.47	0.11	-0.17	0.10
Wholegrain bread	-0.07	0.44	0.10	0.06	0.10
Legumes	0.14	0.41	-0.20	-0.15	0.09
Yogurt and cottage cheese	0.04	0.37	0.23	0.11	0.17
Dried fruit	0.01	0.37	0.13	-0.16	0.11
Tea	-0.06	0.32	0.18	0.11	-0.13
Eggs	0.28	0.32	-0.13	-0.02	0.02
Oil, margarine	0.16	0.30	0.24	0.11	-0.16
Water	0.00	0.26	0.11	0.01	0.13
Sweetened beverages	0.34	-0.36	0.00	-0.02	-0.15
Honey/jam	0.01	0.28	0.49	-0.07	0.02
Chocolate	0.15	0.08	0.48	-0.04	0.11
Butter	0.12	0.19	0.44	0.05	-0.08
Cheese	0.20	0.31	0.32	0.10	-0.04
White bread	0.13	0.11	0.30	0.05	-0.05
Prepacked foods	0.17	-0.08	0.05	0.62	0.34
Ready-prepared dishes	0.19	-0.11	-0.08	0.59	0.34
Canned foods	0.20	-0.04	-0.05	0.52	0.32
Diet foods	0.06	0.11	-0.11	0.32	0.08
Milk	0.15	-0.08	-0.07	-0.33	0.56
Cocoa powder	0.19	-0.20	0.09	-0.33	0.53
Breakfast cereals	0.14	-0.09	-0.04	-0.16	0.38
Coffee	0.07	0.08	0.09	0.23	-0.40
% explained variance	7.9	6.8	4.7	3.7	3.5
Proposed label	Western	Balanced	Bread and toppings	Processed products	Milk and breakfast cereals

TABLE 3 Multivariate associations between demographic and socio-economic characteristics and maternal diet for the whole sample ($n = 12\,048$): The ELFE study, 2011

	PCA pattern 1: Western	PCA pattern 3: Bread and toppings	PCA pattern 5: Milk and breakfast
Maternal age at deliver			
<25 years	0.22 [0.15; 0.29] ^a	-0.08 [-0.15; -0.01]	0.08 [0.01; 0.16]
25–29 years	0 [ref]	0 [ref]	0 [ref]
30–34 years	-0.06 [-0.10; -0.02]	0.13 [0.09; 0.17]	-0.12 [-0.17; -0.08]
35 years or more	-0.18 [-0.23; -0.13]	0.15 [0.10; 0.20]	-0.22 [-0.28; -0.17]
Maternal education level			
Up to lower secondary	0.31 [0.20; 0.41]	-0.37 [-0.48; -0.26]	-0.20 [-0.31; -0.08]
Upper secondary	0.22 [0.17; 0.28]	-0.25 [-0.30; -0.19]	-0.17 [-0.23; -0.12]
Intermediate	0.09 [0.04; 0.14]	-0.14 [-0.19; -0.09]	-0.06 [-0.11; 0.00]
3-year university degree	0.06 [0.01; 0.11]	-0.02 [-0.08; 0.03]	0.02 [-0.03; 0.08]
At least 5-year university degree	0 [ref]	0 [ref]	0 [ref]
Maternal migration status			
Migrant	-0.01 [-0.07; 0.06]	-0.58 [-0.65; -0.51]	0.24 [0.17; 0.31]
Descendant of migrant	0.03 [-0.03; 0.09]	-0.25 [-0.31; -0.19]	0.03 [-0.03; 0.09]
Majority population	0 [ref]	0 [ref]	0 [ref]
Single motherhood			
No	0 [ref]	0 [ref]	0 [ref]
Yes	0.11 [0.01; 0.21]	-0.07 [-0.17; 0.03]	-0.01 [-0.11; 0.10]
Older children in household			
ELFE child is the first child	0 [ref]	0 [ref]	0 [ref]
At least one older child	0.18 [0.14; 0.22]	0.09 [0.05; 0.13]	-0.17 [-0.21; -0.13]
Maternal work status during pregnancy			
Employed	0 [ref]	0 [ref]	0 [ref]
Unemployed	0.09 [0.03; 0.14]	0.03 [-0.02; 0.09]	0.05 [-0.01; 0.11]
Not in the labour force	0.05 [-0.01; 0.11]	-0.03 [-0.09; 0.02]	-0.02 [-0.08; 0.04]
Household income			
≤€750/month ^b	0.18 [0.11; 0.26]	-0.26 [-0.34; -0.19]	0.01 [-0.07; 0.09]
€751–1111/month ^c	0.08 [0.02; 0.14]	-0.14 [-0.19; -0.08]	-0.03 [-0.10; 0.03]
€1112–1500/month ^d	0 [ref]	0 [ref]	0 [ref]
€1501–1944/month ^e	-0.05 [-0.09; 0.00]	0.05 [0.01; 0.10]	0.01 [-0.04; 0.06]
€1945–2500/month ^f	-0.09 [-0.14; -0.03]	0.11 [0.06; 0.17]	0.06 [0.00; 0.12]
>€2500/month ^g	-0.12 [-0.19; -0.05]	0.10 [0.03; 0.17]	-0.02 [-0.09; 0.06]

^aValues are estimates [95% CI] from multivariable linear regressions, also adjusted for region of residence, maternity unit size and enrolment wave; the global significance was indicated with the following labels: *** for $p < 0.001$, ** for $p < 0.01$ and * for $p < 0.05$.

^bCorresponding to ≤\$815/month.

^cCorresponding to \$816–1207/month.

^dCorresponding to \$1208–1630/month.

^eCorresponding to \$1631–2113/month.

^fCorresponding to \$2114–2717/month.

^gCorresponding to >\$2717/month.

Diet quality score, PANDiet score, and balanced dietary pattern

Although positive associations between maternal education and healthy dietary patterns were observed among all women, effect sizes were stronger in the majority population (Figure 1). Household income

was positively associated with these patterns only in the majority population. Higher adherence to the PANDiet and Balanced patterns were observed in women not in the labour force only in descendants of migrants; no association was observed between maternal




(a) *Migrants*(b) *Descendants of migrants*(c) *Majority of population*

FIGURE 1 Multivariate associations between demographic or socio-economic characteristics and maternal diet by maternal migration status: The ELFE study, 2011. Figure 1a presents the multivariate associations among migrants ($n = 853$), Figure 1b those among descendants of migrants ($n = 1195$), and Figure 1c those among the majority population ($n = 10\,000$). All multivariable linear regressions presented in this figure were adjusted for region of residence, maternity unit size and recruitment wave. ^aCorresponding to $\leq \$815$ /month; ^bcorresponding to $\$816$ – 1207 /month; ^ccorresponding to $\$1208$ – 1630 /month; ^dcorresponding to $\$1631$ – 2113 /month; ^ecorresponding to $\$2114$ – 2717 /month; ^fcorresponding to $> \$2717$ /month

occupation and healthy dietary patterns in either migrants or the majority population. Age was positively related to healthy dietary patterns in all women, regardless of their migration status, and single motherhood, negatively; there was no clear relation to the presence of older children in the household.

Processed dietary pattern

None of maternal age, maternal education level and maternal occupation was associated with this dietary pattern (Figure 1). Household income was positively associated with it in the majority population and in descendants of migrants, but not in migrants. The presence of older children was associated with greater adherence to this pattern in descendants of migrants only. Single motherhood was associated with a higher score on this pattern, regardless of migration status.

The analyses based on weighted data or on multiple imputations of missing data produced very similar findings (data not shown).

4 | DISCUSSION

This study is the first to provide insights into the demographic and socio-economic determinants of pregnant women's diet in France, while taking the possible moderating effects of migration status into account. Five distinct dietary patterns were identified and labelled as Western, Balanced, Bread and toppings, Processed products, and Milk and breakfast cereals. Overall, older age and foreign cultural background were related to better quality diets during pregnancy, whether quality was assessed by hypothesis-driven or exploratory approaches. The reverse was true for single motherhood, unemployment, and older children in the household, all of which were associated with sub-optimal diets. Maternal education and household income were negatively associated with the Western dietary pattern in the overall population, while their association with the healthier dietary patterns was moderated by the mother's migration status: their traditional positive association with a healthy diet was essentially confirmed in the majority population, and to a lesser extent in mothers from different cultural backgrounds. This acculturation process was also suggested for the Processed dietary pattern, which had higher scores in households with higher incomes, in the majority population and to a lesser extent in descendants of migrants, but not in migrant mothers.

Comparison of the maternal and household characteristics associated with maternal diet in a US study (Bodnar & Siega-Riz, 2002) showed that a high score on the *Diet Quality Index for Pregnancy* was positively associated with women's age, education, and income. Similar relations between women's characteristics and diet quality during pregnancy have been reported in Norway (Hillesund, Bere, Haugen, & Overby, 2014) and Spain (Rodriguez-Bernal et al., 2010). The Norwegian study demonstrated that adherence to Nordic or Norwegian dietary guidelines increased slightly with age, education and exercise frequency. Likewise, in the ELFE study, age at delivery and education were positively associated with healthy patterns (i.e. the Diet Quality and the PANDiet scores and the balanced dietary pattern), and negatively with the Western dietary pattern. Interestingly,

even after adjustment for education and employment, family income remained negatively associated with the Western dietary pattern and positively associated with the Bread and toppings pattern. These findings are notably similar to those observed among women from the general French population for the relations of age and education level to balanced and healthier dietary patterns (Bertin et al., 2016).

Our findings however further showed that the relation of maternal education to the healthier patterns was strongest in the majority population, lower in descendants of migrants and still lower among migrants. This moderating effect of migration status was also evident for another socio-economic indicator – household income. On the one hand, it was positively associated with healthy patterns only in the majority population; on the other hand, it was also positively related to the Processed pattern in this population and, to a less extent, in descendants of migrants, but again, not among migrants. Overall, the less acculturated pregnant women were, the healthier and less processed their diet, regardless of their socio-economic position. The so-called 'healthy migrant effect', which results from a selective process where migrants are on average healthier than, on the one hand, those who stayed behind in the country of origin, and, on the other hand, the majority population born in the destination country (Kennedy, Kidd, McDonald, & Biddle, 2014), might in part explain these findings. It has also been reported that among migrant pregnant women and mothers, cultural factors play a more important role than traditional socio-economic indicators (such as education and income) (Kumar, Holmboe-Ottesen, Lien, & Wandel, 2004; Osei-Kwasi et al., 2016) and result in stronger adherence to their own, apparently healthier, social and cultural dietary traditions. Dietary behaviours are influenced by a variety of factors, such as cultural identity and desire to maintain traditional food identity (which presumably does not include processed foods), religious beliefs and prescriptions, social networks, social norms, and the social role of foods, beliefs and perceptions of healthy food (D'Souza et al., 2016; Osei-Kwasi et al., 2016). Similarly, previous findings about breastfeeding have underlined that migrant women are more likely to begin by continuing habits from their country of origin and then progressively become acculturated (Kersuzan, Tichit, & Thierry, 2018). It should be noted that the Bread and topping dietary pattern seems to reflect both the traditional bread and jam/honey French breakfast and the bread/cheese third course of traditional French meals, which might explain in part migrant women's lesser adherence to it. Importantly, acculturation, approximated in the current study by the migration status variable, was associated with poorer diet quality (as measured by any of the Diet quality score, PANDiet score, and Balanced and Processed dietary patterns), along with socio-economic inequalities that emerged in the descendants of migrants. It has been suggested that migration is often accompanied by disruption of social ties and norms, a reduction in socio-economic position, material hardships, literacy challenges, increased stress and the experience of discrimination – all potential obstacles to maintaining traditional healthy dietary patterns in groups of foreign origin across generations (Allen et al., 2014).

Consistent with other studies (Bodnar & Siega-Riz, 2002; Hillesund et al., 2014; Rodriguez-Bernal et al., 2010), single

motherhood was related, independently of migration status and socioeconomic position, to lower adherence to guidelines, along with higher scores on the Western and Processed patterns. Similar associations were also observed with unemployment and older children in the household. In a context of higher social vulnerability, obstacles to the availability and accessibility of healthy food, along with time constraints, may partly explain these associations (Dlugonski & Motl, 2013; Mills et al., 2017; Pollard, Kirk, & Cade, 2002) and should inform nutritional interventions targeted at more at-risk families.

The *Diet Quality* score was constructed to consider adherence to guidelines for all of the core food groups. These groups, however, contributed in different ways to the various dietary patterns derived from PCA: the Western pattern was mainly characterized by meat and fish, and the Balanced pattern by fruit and vegetables. High factor loadings were observed for dairy products in the Balanced pattern (yogurt), the Bread and toppings pattern (cheese), and the Milk and breakfast cereals pattern (milk). These exploratory dietary patterns therefore encompass different dimensions of diet quality.

Only few studies have combined hypothesis-driven and exploratory approaches to characterize maternal diet. Principal component analysis is a standard technique used in population studies to explore dietary patterns (Bertin et al., 2016; Lioret et al., 2012; Northstone, 2012; Northstone, Emmett, & Rogers, 2008), which takes the potential effect of interactions within combinations of foods and nutrients into account (Newby & Tucker, 2004). While the hypothesis-driven approach promotes international comparisons and enables the exploration of global adherence to nutritional guidelines, the exploratory approach, which identifies specific patterns within a country and various dimensions of diet quality, may enable the provision of more tailored food-based recommendations. It should be noted that the two complementary hypothesis-driven quality scores used in the current study (one derived from food group consumption and the other one based on nutrient intake) were moderately correlated ($r = 0.43$). Both scores were also strongly correlated with the Balanced dietary pattern. These correlations support the external validity of the hypothesis-driven and exploratory scores.

The ELFE-FFQ provided reasonably good estimates of food and nutrient intake compared with repeated 24-h recalls (Table S2); correlation coefficients are similar to those reported in other validation studies among pregnant women (Cheng et al., 2008; Erkkola et al., 2001; Mikkelsen, Osler, & Olsen, 2006; Mouratidou, Ford, & Fraser, 2006; Pinto et al., 2010; Robinson, Godfrey, Osmond, Cox, & Barker, 1996), although a US study noted higher correlations (Brown et al., 1996). There is no general consensus on a 'satisfactory' level of correlation. In most validation studies, correlation coefficients between methods are considered poor when <0.30 ; fair when between 0.3 to 0.49; and good when >0.50 (Hankin, Wilkens, Kolonel, & Yoshizawa, 1991). Adjustment for energy and correction for random intraindividual variation tended to increase the magnitude of the correlation coefficients (Brantsaeter, Haugen, Alexander, & Meltzer, 2008; Brown et al., 1996; Erkkola et al., 2001;

Mouratidou et al., 2006). Correlation coefficients are commonly employed and reported in validation studies, but the level of agreement assessed by cross-classification into quartiles or quintiles of their intake is considered more appropriate to evaluate the ability of an FFQ to rank individuals into low or high intakes relative to the reference method (Garrow, 1995). In the ELFE study, only a very small percentage of the women was grossly misclassified (classified into opposing quintiles in FFQ vs. 24-h recalls). It should be noted that a review of 90 studies of energy intake during pregnancy in high-income countries found an average intake of 2340 Kcal in the third trimester, very close to the FFQ estimates reported here (Blumfield et al., 2012). The mean PANDiet score was also very close to the value previously estimated in a population of pregnant women or in any case of childbearing age (Bianchi et al., 2016). It is however recognized that the FFQ was completed after delivery and collected dietary intake retrospectively over the last three months of pregnancy, which does not necessarily reflect the diet across all the pregnancy period.

The ELFE cohort is a nationwide study of births in 2011 in metropolitan France (excluding very premature babies). The main analyses were conducted on the complete case sample. It must also be acknowledged that, as it is often the case in cohorts, women not participating in the cohort or those excluded from the analyses were more strongly socially disadvantaged and more likely to have been born abroad, which is a potential limitation in terms of generalizability of the findings. We also acknowledge that whereas the main documents were edited in the three languages most often spoken in France (after French), this was not done for the FFQ. Even if women could be helped to complete the questionnaire, this limitation probably led to a higher probability of missing dietary information among women with lower literacy in French. Nonetheless, addressing missing data for demographic and socioeconomic characteristics by the multiple imputation method, or running the analysis using weighting to account for selection bias, produced results consistent with those of the main analyses, thus suggesting that selection bias has limited impact on our findings. Furthermore, dietary patterns derived from PCA in the current study were overall consistent with literature based on cross-sectional studies, for which selection bias is supposedly smaller (Bertin et al., 2016). These findings support the robustness of the results. Two specific limitations about the analysis of the migrant effect must be acknowledged. The first is the selection of migrant women by their ability to read French (which was necessary for them to be able to complete the FFQ); this may have resulted in an overrepresentation of educated women in the migrant group. However, enough variability remained to observe an association with education level for some of the dietary indices. Second, the FFQ may have underrepresented specific foods consumed by migrants. The principal strengths of the ELFE study also include the large sample and the wide range of socio-economic, cultural and demographic variables collected, which made it possible to disentangle the relation between the various components of social background and diet at the same time that the moderating effect of maternal migration status is taken into account.

5 | CONCLUSION

The present study used both hypothesis-driven and exploratory approaches to characterize the diet of pregnant women more comprehensively. In addition to two scores indicating higher adherence to dietary (Diet quality score) and nutritional (PANDiet score) guidelines, five distinct data-driven dietary patterns were identified and labelled as Western, Balanced, Bread and toppings, Processed products, and Milk and breakfast cereals. These findings from the large, nationwide ELFE birth cohort suggest that younger maternal age, single motherhood, and unemployment – all indicators of greater social vulnerability – as well as the presence of older children in the household are determinants of a suboptimal diet during pregnancy. The consideration of migration status makes it possible to show that the less acculturated pregnant women were, the healthier and less processed their diets were, independent of socio-economic position. Several demographic and socio-economic determinants of the quality of women's diets were moderated by their migration status. These original findings should inform the development of future nutritional interventions during pregnancy. Such interventions would be more efficient and likely more effective if directed at more socially disadvantaged populations. Given the reduction in diet quality that appears to accompany the acculturation process, it is of paramount importance to identify the specific factors or obstacles that affect migrant women in maintaining their diet quality advantage over the majority population and thereby prevent the emergence of diet-related health inequalities for themselves and their offspring.

ACKNOWLEDGMENTS

We thank the scientific coordinators (B. Geay, H. Léridon, C. Bois, J. L. Lanoé, X. Thierry and C. Zaros), IT and data managers, statisticians (M. Cheminat, C. Ricourt, A. Candea and S. de Visme), administrative and family communication staff, and study technicians (C. Guevel, M. Zoubiri, L. G. L. Gravier, I. Milan and R. Popa) of the ELFE coordination team as well as the families that gave their time for the study. We would like also to thank the French Institute for Demographic Studies (INED) researchers for the sociodemographic variables created as part of the ANR-funded Veniromond project. We thank Jo Ann Cahn for her help in preparing the manuscript. Permission from all persons named in the acknowledgements has been obtained. This study was funded by two grants from the French National Research Agency (ANR-12-DSSA-0001, within the framework “Social determinants of health”; and ANR-19-CE36-0006). Manik Kadawathagedara received a grant from SF-DOHAD/Blédina. The Elfe survey is a joint project between the French Institute for Demographic Studies (INED) and the National Institute of Health and Medical Research (INSERM), in partnership with the French blood transfusion service (Etablissement français du sang, EFS), Santé publique France, the National Institute for Statistics and Economic Studies (INSEE), the Direction générale de la santé (DGS, part of the Ministry of Health and Social Affairs), the Direction générale de la prévention des risques (DGPR, Ministry for the Environment), the Direction de la recherche, des études, de l'évaluation et des statistiques (DREES, Ministry of

Health and Social Affairs), the Département des études, de la prospective et des statistiques (DEPS, Ministry of Culture), and the Caisse nationale des allocations familiales (CNAF), with the support of the Ministry of Higher Education and Research and the Institut national de la jeunesse et de l'éducation populaire (INJEP). Via the RECONAI platform, it receives a government grant managed by the National Research Agency under the “Investissements d'avenir” programme (ANR-11-EQPX-0038). The funders had no role in the study design, data collection and analysis, decision to publish or preparation of the manuscript.

DATA AVAILABILITY STATEMENT

The data underlying the findings cannot be made freely available for ethical and legal restrictions imposed, because this study includes a substantial number of variables that, together, could be used to re-identify the participants based on a few key characteristics and then be used to have access to other personal data. Therefore, the French ethics authority strictly forbids making these data freely available. However, they can be obtained upon request from the ELFE principal investigator. Readers may contact marie-aline.charles@inserm.fr to request the data.

CONFLICTS OF INTEREST

The other authors had no conflicts of interest relevant to this article to disclose.

CONTRIBUTIONS

MK conceptualized and designed the study, conducted statistical analyses, drafted the initial manuscript, and approved the final manuscript as submitted; MND and AF designed the data collection instruments, supervised data collection, critically reviewed the manuscript, and approved the final manuscript as submitted; NA, MAC, SL and BLG conceptualized and designed the study, contributed to data analysis and interpretation of the study, reviewed and revised the manuscript, and approved the final manuscript as submitted.

ORCID

Marie Aline Charles  <https://orcid.org/0000-0003-4025-4390>

Sandrine Lioret  <https://orcid.org/0000-0002-2483-7820>

Blandine de Lauzon-Guillain  <https://orcid.org/0000-0001-5887-8842>

REFERENCES

- Abu-Saad, K., & Fraser, D. (2010). Maternal nutrition and birth outcomes. *Epidemiologic Reviews*, 32(1), 5–25. <https://doi.org/10.1093/epirev/mxq001>
- Allen, J. D., Caspi, C., Yang, M., Leyva, B., Stoddard, A. M., Tamers, S., ... Sorensen, G. C. (2014). Pathways between acculturation and health behaviors among residents of low-income housing: The mediating role of social and contextual factors. *Social Science & Medicine*, 123, 26–36. <https://doi.org/10.1016/j.socscimed.2014.10.034>
- ANSES. (2016). Avis de l'ANSES relatif à l'actualisation des repères du PNNS: Révision des références nutritionnelles en vitamines et minéraux pour la population générale adulte.

- Bertin, M., Touvier, M., Dubuisson, C., Dufour, A., Havard, S., Lafay, L., ... Lioret, S. (2016). Dietary patterns of French adults: Associations with demographic, socio-economic and behavioural factors. *Journal of Human Nutrition and Dietetics*, 29(2), 241–254. <https://doi.org/10.1111/jhn.12315>
- Bianchi, C. M., Mariotti, F., Verger, E. O., & Huneau, J. F. (2016). Pregnancy requires major changes in the quality of the diet for nutritional adequacy: Simulations in the French and the United States populations. *PLoS ONE*, 11(3), e0149858. <https://doi.org/10.1371/journal.pone.0149858>
- Blondel, B., Lelong, N., Kermarrec, M., Goffinet, F., & National Coordination Group of the National Perinatal Surveys. (2012). Trends in perinatal health in France from 1995 to 2010. Results from the French National Perinatal Surveys. *Journal de Gynecologie, Obstetrique et Biologie de la Reproduction*, 41(4), e1–e15. <https://doi.org/10.1016/j.jgyn.2012.04.014>
- Blumfield, M. L., Hure, A. J., Macdonald-Wicks, L., Smith, R., & Collins, C. E. (2012). Systematic review and meta-analysis of energy and macronutrient intakes during pregnancy in developed countries. *Nutrition Reviews*, 70(6), 322–336. <https://doi.org/10.1111/j.1753-4887.2012.00481.x>
- Blumfield, M. L., Hure, A. J., Macdonald-Wicks, L., Smith, R., & Collins, C. E. (2013). A systematic review and meta-analysis of micronutrient intakes during pregnancy in developed countries. *Nutrition Reviews*, 71(2), 118–132. <https://doi.org/10.1111/nure.12003>
- Bodnar, L. M., & Siega-Riz, A. M. (2002). A diet quality index for pregnancy detects variation in diet and differences by sociodemographic factors. *Public Health Nutrition*, 5(6), 801–809. <https://doi.org/10.1079/PHN2002348>
- Brantsaeter, A. L., Haugen, M., Alexander, J., & Meltzer, H. M. (2008). Validity of a new food frequency questionnaire for pregnant women in the Norwegian Mother and Child Cohort Study (MoBa). *Maternal & Child Nutrition*, 4(1), 28–43. <https://doi.org/10.1111/j.1740-8709.2007.00103.x>
- Brown, J. E., Buzzard, I. M., Jacobs, D. R., Hannan, P. J., Kushi, L. H., Barosso, G. M., & Schmid, L. A. (1996). A food frequency questionnaire can detect pregnancy-related changes in diet. *Journal of the American Dietetic Association*, 96(3), 262–266. [https://doi.org/10.1016/s0002-8223\(96\)00078-8](https://doi.org/10.1016/s0002-8223(96)00078-8)
- Castetbon, K., Vernay, M., Malon, A., Salanave, B., Deschamps, V., Roudier, C., ... Hercberg, S. (2009). Dietary intake, physical activity and nutritional status in adults: The French nutrition and health survey (ENNS, 2006–2007). *The British Journal of Nutrition*, 102(5), 733–743. <https://doi.org/10.1017/S0007114509274745>
- Cattell, R. B. (1966). The Scree test for the number of factors. *Multivariate Behav Res*, 1(2), 245–276. https://doi.org/10.1207/s15327906mbr0102_10
- Charles, M. A., Thierry, X., Lanoe, J. L., Bois, C., Dufour, M. N., Popa, R., ... Geay, B. (2019). Cohort profile: The French national cohort of children ELFE: Birth to 5 years. *International Journal of Epidemiology*, 49, 368–369j. <https://doi.org/10.1093/ije/dyz227>
- Cheng, Y., Yan, H., Dibley, M., Shen, Y., Li, Q., & Zeng, L. (2008). Validity and reproducibility of a semi-quantitative food frequency questionnaire for use among pregnant women in rural China. *Asia Pacific Journal of Clinical Nutrition*, 17, 166–177.
- Cohen, J. (1988). *Statistical power analysis for the behavioural science*. New York: Hillsdale NJ, Erlbaum L.
- Dlugonski, D., & Motl, R. W. (2013). Marital status and motherhood: Implications for physical activity. *Women & Health*, 53(2), 203–215. <https://doi.org/10.1080/03630242.2013.767304>
- D'Souza, L., Jayaweera, H., & Pickett, K. E. (2016). Pregnancy diets, migration, and birth outcomes. *Health Care for Women International*, 37(9), 964–978. <https://doi.org/10.1080/07399332.2015.1102268>
- Dubuisson, C., Lioret, S., Touvier, M., Dufour, A., Calamassi-Tran, G., Volatier, J. L., & Lafay, L. (2010). Trends in food and nutritional intakes of French adults from 1999 to 2007: Results from the INCA surveys. *The British Journal of Nutrition*, 103(7), 1035–1048. <https://doi.org/10.1017/S0007114509992625>
- Erkkola, M., Karppinen, M., Javanainen, J., Rasanen, L., Knip, M., & Virtanen, S. M. (2001). Validity and reproducibility of a food frequency questionnaire for pregnant Finnish women. *American Journal of Epidemiology*, 154(5), 466–476. <https://doi.org/10.1093/aje/154.5.466>
- Estaquio, C., Kesse-Guyot, E., Deschamps, V., Bertrais, S., Dauchet, L., Galan, P., ... Castetbon, K. (2009). Adherence to the French Programme National Nutrition Sante Guideline score is associated with better nutrient intake and nutritional status. *Journal of the American Dietetic Association*, 109(6), 1031–1041. <https://doi.org/10.1016/j.jada.2009.03.012>
- Faskunger, J., Eriksson, U., Johansson, S. E., Sundquist, K., & Sundquist, J. (2009). Risk of obesity in immigrants compared with Swedes in two deprived neighbourhoods. *BMC Public Health*, 9, 304. <https://doi.org/10.1186/1471-2458-9-304>
- Garrow, J. S. (1995). Validation of methods for estimating habitual diet: Proposed guidelines. *European Journal of Clinical Nutrition*, 49(4), 231–232.
- Hales, C. N., & Barker, D. J. (2001). The thrifty phenotype hypothesis. *British Medical Bulletin*, 60, 5–20. <https://doi.org/10.1093/bmb/60.1.5>
- Hankin, J. H., Wilkens, L. R., Kolonel, L. N., & Yoshizawa, C. N. (1991). Validation of a quantitative diet history method in Hawaii. *American Journal of Epidemiology*, 133(6), 616–628. <https://doi.org/10.1093/oxfordjournals.aje.a115934>
- Hercberg, S. (2006). *Table de composition des aliments SU.VI.MAX*, Paris: Economica.
- Hercberg, S., Chat-Yung, S., & Chauliac, M. (2008). The French National Nutrition and Health Program: 2001–2006–2010. *International Journal of Public Health*, 53(2), 68–77. <https://doi.org/10.1007/s00038-008-7016-2>
- Hercberg, S., Deheeger, M., & Preziosi, P. (2002). *SU.VI.MAX. Portions alimentaires: Manuel photos pour l'estimation des quantités*. Paris: Editions Polytechnica.
- Hillesund, E. R., Bere, E., Haugen, M., & Overby, N. C. (2014). Development of a New Nordic Diet score and its association with gestational weight gain and fetal growth—A study performed in the Norwegian Mother and Child Cohort Study (MoBa). *Public Health Nutrition*, 17(9), 1909–1918. <https://doi.org/10.1017/S1368890014000421>
- Juillard, H. (2015). Weighting of Elfe survey data at time 0. pandora.vjf.inserm.fr/public/.
- Kadawathagedara, M., Kersuzan, C., Wagner, S., Tichit, C., Gojard, S., Charles, M. A., ... de Lauzon-Guillain, B. (2017). Adéquation des consommations alimentaires des femmes enceintes de l'étude ELFE aux recommandations du Programme national nutrition santé. *Cahiers de Nutrition et de Diététique*, 52(2), 78–88. <https://doi.org/10.1016/j.cnd.2016.12.001>
- Kant, A. K. (2004). Dietary patterns and health outcomes. *Journal of the American Dietetic Association*, 104(4), 615–635. <https://doi.org/10.1016/j.jada.2004.01.010>
- Kennedy, S., Kidd, M. P., McDonald, J. T., & Biddle, N. (2014). The healthy immigrant effect: Patterns and evidence from four countries. *Journal of International Migration and Integration*, 16(2), 317–332. <https://doi.org/10.1007/s12134-014-0340-x>
- Kersuzan, C., Tichit, C., & Thierry, X. (2018). Breastfeeding practices among immigrants and natives in France, from the ELFE cohort. *Population*, 73(3), 545–566.
- Kline, P. (1994). *An easy guide to factor analysis*. London: Routledge.
- Kumar, B. N., Holmboe-Ottesen, G., Lien, N., & Wandel, M. (2004). Ethnic differences in body mass index and associated factors of adolescents from minorities in Oslo, Norway: A cross-sectional study. *Public Health Nutrition*, 7(8), 999–1008. <https://doi.org/10.1079/phn2004644>

- Lioret, S., McNaughton, S. A., Crawford, D., Spence, A. C., Hesketh, K., & Campbell, K. J. (2012). Parents' dietary patterns are significantly correlated: Findings from the Melbourne Infant Feeding Activity and Nutrition Trial Program. *The British Journal of Nutrition*, 108(3), 518–526. <https://doi.org/10.1017/S0007114511005757>
- McMillen, I. C., MacLaughlin, S. M., Muhlhausler, B. S., Gentili, S., Duffield, J. L., & Morrison, J. L. (2008). Developmental origins of adult health and disease: The role of periconceptual and foetal nutrition. *Basic & Clinical Pharmacology & Toxicology*, 102(2), 82–89. <https://doi.org/10.1111/j.1742-7843.2007.00188.x>
- Mikkelsen, T. B., Osler, M., & Olsen, S. F. (2006). Validity of protein, retinol, folic acid and n-3 fatty acid intakes estimated from the food-frequency questionnaire used in the Danish National Birth Cohort. *Public Health Nutrition*, 9(6), 771–778. <https://doi.org/10.1079/phn2005883>
- Mills, S., White, M., Brown, H., Wrieden, W., Kwasnicka, D., Halligan, J., ... Adams, J. (2017). Health and social determinants and outcomes of home cooking: A systematic review of observational studies. *Appetite*, 111, 116–134. <https://doi.org/10.1016/j.appet.2016.12.022>
- Mouratidou, T., Ford, F., & Fraser, R. B. (2006). Validation of a food-frequency questionnaire for use in pregnancy. *Public Health Nutrition*, 9(4), 515–522. <https://doi.org/10.1079/phn2005876>
- Newby, P. K., & Tucker, K. L. (2004). Empirically derived eating patterns using factor or cluster analysis: A review. *Nutrition Reviews*, 62(5), 177–203. <https://doi.org/10.1301/nr.2004.may.177-203>
- Northstone, K. (2012). Dietary patterns: The importance of sex differences. *The British Journal of Nutrition*, 108(3), 393–394. <https://doi.org/10.1017/S0007114511006337>
- Northstone, K., Emmett, P. M., & Rogers, I. (2008). Dietary patterns in pregnancy and associations with nutrient intakes. *The British Journal of Nutrition*, 99(2), 406–415. <https://doi.org/10.1017/S0007114507803977>
- Osei-Kwasi, H. A., Nicolaou, M., Powell, K., Terragni, L., Maes, L., Stronks, K., ... Holdsworth, M. (2016). Systematic mapping review of the factors influencing dietary behaviour in ethnic minority groups living in Europe: A DEDIPAC study. *International Journal of Behavioral Nutrition and Physical Activity*, 13, 85. <https://doi.org/10.1186/s12966-016-0412-8>
- Patel, J. V., Vyas, A., Cruickshank, J. K., Prabhakaran, D., Hughes, E., Reddy, K. S., ... Durrington, P. N. (2006). Impact of migration on coronary heart disease risk factors: Comparison of Gujaratis in Britain and their contemporaries in villages of origin in India. *Atherosclerosis*, 185(2), 297–306. <https://doi.org/10.1016/j.atherosclerosis.2005.06.005>
- Pinto, E., Severo, M., Correia, S., dos Santos Silva, I., Lopes, C., & Barros, H. (2010). Validity and reproducibility of a semi-quantitative food frequency questionnaire for use among Portuguese pregnant women. *Maternal & Child Nutrition*, 6(2), 105–119. <https://doi.org/10.1111/j.1740-8709.2009.00199.x>
- Pollard, J., Kirk, S. F., & Cade, J. E. (2002). Factors affecting food choice in relation to fruit and vegetable intake: A review. *Nutrition Research Reviews*, 15(2), 373–387. <https://doi.org/10.1079/NRR200244>
- Roberts, K., Cavill, N., Hancock, C., & Rutter, H. (2013). *Social and economic inequalities in diet and physical activity*. London: Public Health England.
- Robinson, S., Godfrey, K., Osmond, C., Cox, V., & Barker, D. (1996). Evaluation of a food frequency questionnaire used to assess nutrient intakes in pregnant women. *European Journal of Clinical Nutrition*, 50(5), 302–308.
- Rodriguez-Bernal, C. L., Rebagliato, M., Iniguez, C., Vioque, J., Navarrete-Munoz, E. M., Murcia, M., ... Ballester, F. (2010). Diet quality in early pregnancy and its effects on fetal growth outcomes: The Infancia y Medio Ambiente (Childhood and Environment) Mother and Child Cohort Study in Spain. *The American Journal of Clinical Nutrition*, 91(6), 1659–1666. <https://doi.org/10.3945/ajcn.2009.28866>
- SAS [computer program] (Version 9.4). (2013). Cary, NC: SAS Institute Inc.
- Urquia, M. L., Glazier, R. H., Blondel, B., Zeitlin, J., Gissler, M., Macfarlane, A., ... Gagnon, A. J. (2010). International migration and adverse birth outcomes: Role of ethnicity, region of origin and destination. *Journal of Epidemiology and Community Health*, 64(3), 243–251. <https://doi.org/10.1136/jech.2008.083535>
- Volatier, J.-L. (2000). *Enquête INCA individuelle et nationale Sur les consommations alimentaires. (individual and National Food Consumption INCA survey)*. Paris: Tec&Doc.
- Von Ruesten, A., Brantsaeter, A. L., Haugen, M., Meltzer, H. M., Mehlig, K., Winkvist, A., & Lissner, L. (2014). Adherence of pregnant women to Nordic dietary guidelines in relation to postpartum weight retention: Results from the Norwegian Mother and Child Cohort Study. *BMC Public Health*, 14, 75. <https://doi.org/10.1186/1471-2458-14-75>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Kadawathagedara M, Ahluwalia N, Dufourg M-N, et al. Diet during pregnancy: Influence of social characteristics and migration in the ELFE cohort. *Matern Child Nutr*. 2021:e13140. <https://doi.org/10.1111/mcn.13140>