



HAL
open science

The kisspeptin system in domestic animals: what we know and what we still need to understand of its role in reproduction

Massimiliano Beltramo, Vincent Robert, C. Decourt

► **To cite this version:**

Massimiliano Beltramo, Vincent Robert, C. Decourt. The kisspeptin system in domestic animals: what we know and what we still need to understand of its role in reproduction. *Domestic Animal Endocrinology*, 2020, 73, pp.1-11. 10.1016/j.domaniend.2020.106466 . hal-03150449

HAL Id: hal-03150449

<https://hal.inrae.fr/hal-03150449>

Submitted on 14 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 The Kisspeptin system in domestic animals: what we know and what we still need to understand of
2 its role in reproduction.

3

4 Massimiliano Beltramo¹, Vincent Robert¹, Caroline Decourt^{1,a}

5

6

7 ¹INRAe (CNRS, UMR7247, Université de Tours, IFCE) UMR85 Physiologie de la Reproduction et des
8 Comportements F-37380 Nouzilly, France¹

9

10 **Corresponding Authors :**

11 Massimiliano BELTRAMO

12 UMR85, Physiologie de la Reproduction et des Comportements, INRAe.

13 37380 Nouzilly, France

14 E-mail Massimiliano.Beltramo@inrae.fr

15 Tel 00 33 2 47 42 73 60

16 Fax 00 33 2 47 42 77 43

17

18

19 **Declaration of interest:** MB is inventor of the patent containing the kisspeptin analog C6.

20

21 **CRedit Author Statement: Massimiliano Beltramo:** Conceptualization, Writing Original Draft,
22 Investigation, Funding acquisition. **Vincent Robert:** Investigation, Writing Review and Editing.

23 **Caroline Decourt:** Conceptualization, Investigation, Writing Review and Editing.

24

^aPresent address: Centre for Neuroendocrinology and Department of Anatomy, University of Otago School of Biomedical Sciences | PO Box 913 | Dunedin 9054 University of Otago, NZ.

25

26 **Abstract**

27 The discovery of the kisspeptin (Kp) system stirred a burst of research in the field of reproductive
28 neuroendocrinology. In the last 15 years the organization and activity of the system including its
29 neuroanatomical structure, its major physiological functions, and its main pharmacological properties
30 were outlined. To this endeavor, the use of genetic tools to delete and to restore Kp system
31 functionality in a specific tissue was essential. At present, there is no question as to the key role of
32 the Kp system in mammalian reproduction. However, easily applicable genetic manipulations are
33 unavailable for domestic animals. Hence, many essential details on the physiological mechanisms
34 underlying its action on domestic animals require further investigation. The potentially different
35 effects of the various Kp isoforms, the precise anatomical localization of the Kp receptor, and the
36 respective role played by the two main populations of Kp cells in different species are only few of the
37 questions that remain unanswered and that will be illustrated in this review. Furthermore, the
38 application of synthetic pharmacological tools to manipulate the Kp system is still in its infancy but
39 has produced some interesting results suggesting the possibility to develop new methods to manage
40 reproduction in domestic animals. In spite of a decade and a half of intense research effort, much
41 work is still required to achieve a comprehensive understanding of the influence of the Kp system on
42 reproduction. Furthermore, Kp system ramifications in other physiological functions are emerging
43 and open new research perspectives.

44 **Key words:** Kisspeptin, reproduction, ovulation, GnRH, LH, pharmacology, agonist.

45

46

47 **1. Introduction**

48 The discovery that reproduction in mammals is mastered by a brain neuropeptide, kisspeptin (Kp), is
49 the most recent breakthrough in reproductive neuroendocrinology. Our knowledge of the Kp system
50 has grown exponentially and several excellent reviews on the subject are available [\[1-4\]](#). To elucidate
51 Kp system functions, most studies used rodents or primates, but investigations in domestic animals
52 are less copious. Nevertheless, these studies are extremely interesting from a comparative point of
53 view and capable of opening new opportunities to manage livestock reproduction. In the following
54 sections we will briefly delineate the biological action of Kp and point to a few outstanding questions
55 that we deem particularly interesting. Namely, we will discuss the anatomical distribution of the Kp
56 system, its main physiological effects in reproduction and some pharmacological aspects crucial for
57 the development of applications to manage domestic animal reproduction. Many other important
58 and exciting features of the Kp system still require a substantial research effort. To gain a broader
59 perspective on these aspects the reader could refer to other reviews on domestic animals [\[5-8\]](#).

60

61

62 **2. The Kp system**

63 A role for Kp in reproduction was first discovered in human patients showing loss of function
64 mutations of the Kp receptor, *KISS1R* (also known as GPR54). These mutations lead to
65 hypogonadotropic hypogonadism, lack of puberty and infertility [\[9, 10\]](#). Further studies showing that
66 GnRH neurons express *Kiss1r* [\[11\]](#), and the fact that Kp administration triggers GnRH release [\[12, 13\]](#)
67 demonstrated that Kp stimulates the reproductive axis by a direct action on GnRH neurons. Kp also
68 has a role in modulating the timing of puberty onset as shown in rodents, where repeated
69 administration of Kp to prepubertal female rats advanced vaginal opening [\[14\]](#). Two major
70 populations of Kp neurons in the hypothalamus have been identified: one in the preoptic area (POA)

71 region and the other in the arcuate nucleus (ARC). Neurons of both populations bear estrogen
72 receptor alpha, progesterone receptors and/or androgen receptors [15-18]. These finding and others
73 [19] imply that Kp neurons are the target of sexual steroid feedback. Kp neurons in the ARC colocalize
74 with neurokinin B and dynorphin and are referred to as KNDy neurons from the initials of the
75 neurotransmitters they produce [20]. Localization and pharmacological studies have shown that
76 KNDy neurons express the neurokinin B receptor NK3R [21] and the K-opioid receptor that is the
77 cognate dynorphin receptor [22]. These results were instrumental in formulating the hypothesis that
78 by an autocrine mechanism neurokinin B would initiate Kp release from KNDy neurons and
79 dynorphin would terminate it [23]. The concept that KNDy neurons are the GnRH pulse generator is
80 now widely accepted [24].

81

82 **3. *KISS1* and *KISS1R* genes**

83 In vertebrates, three different genes encoding for Kp and four genes encoding for its receptor have
84 been identified. In mammals, with the exception of the monotremata (e.g. platypus), only one gene
85 for the Kp (*KISS1*) and one for its receptor (*KISS1R*) are present [25].
86 *KISS1R* is a G protein-coupled receptor encoded by a highly conserved sequence. In humans *KISS1R*
87 loss or gain of function mutations result in infertility [9, 10] and precocious puberty [26] respectively.
88 Study in transgenic mice further support the importance of *Kiss1R* in puberty onset and fertility [27].
89 The existence of a loss of function mutation in a species or breed of domestic animals would be
90 unexpected due to the importance of reproductive capacities in these animals and the resulting loss
91 with such mutation. However, the identification of polymorphisms in the *KISS1* or *KISS1R* and their
92 correlation with reproductive traits is an interesting topic for investigation. In goat breeds the
93 presence of specific *KISS1* and/or *KISS1R* single nucleotide polymorphisms (SNPs) was correlated with
94 variations in litter size and age at puberty [28, 29] as well as with an increase in circulating levels of
95 estradiol and progesterone [30]. In *Kiss1R* haplo-insufficient mice, the litter size is reduced compared

96 to wild type mice [31]. This would concur with a role of the Kp system to define the litter size.
97 Nevertheless, a more thorough characterization is required to establish firmly if the reported SNPs
98 would be responsible or not for the modifications observed in these reproductive traits. A better
99 understanding of the functional correlate of *KISS1R* SNPs may reveal new criteria for the genetic
100 selection of domestic animals.

101 Processing of the Kp precursor leads to several endogenous Kp isoforms. The shortest active
102 sequence, 10 amino acids (Kp10), corresponds to the C-terminal part of the precursor. This sequence
103 is extremely well conserved in mammals with only few variants identified. The remaining of the
104 sequence is significantly less conserved. Three other Kp isoforms have been described, two short
105 ones (Kp13 and Kp14, the number indicates the length of the amino acid sequence), and a longer one
106 that varies among species studied (i.e. Kp54 in humans, Kp53 in cattle, in the sheep, and in the goat,
107 and Kp52 in the mouse). Preliminary data obtained in the rat and in the sheep showed the existence
108 of an additional isoform, Kp16, and suggested that Kp16 and Kp13 are actually the most abundant
109 isoforms [32]. Unfortunately, a complete account of these results was never published. The presence
110 of different isoforms raises a major question: do the different isoforms serve different physiological
111 functions? This question remains at present unanswered. The vast majority of the knowledge gained
112 on the Kp system was obtained by applying the Kp10 and to a much lesser extent the Kp54. The other
113 isoforms have received very little attention. To the best of our knowledge, *in vitro* data are available
114 only for the hKp13, showing binding affinity and potency in a calcium mobilization assay similar to
115 that of hKp10 [33].

116
117 The small variations observed in the Kp10 sequence may have physiological consequences, but these
118 are not easily predictable based on the sequence's structure. Interestingly the human (h) and ovine
119 Kp10 (hKp10 and oKp10) differ in the C-terminal: such that the last amino acid is a phenylalanine in
120 the human sequence and a tyrosine in the sheep (figure 1A). This amino acid is part of the
121 pharmacophore region, the region bearing the biological activity [34]. A modification of this region

122 could influence molecule's activity. However, when tested *in vitro* in a calcium mobilization assay (for
123 detail on the assay see [35]) hKp10 and oKp10 have superimposable profiles (figure 1B).
124 Furthermore, when injected into the ewe both peptides produced an increase of LH and FSH plasma
125 levels that were identical in amplitude and duration [36]. Studies in goats using intravenous
126 administration of hKp10 produced similar results [37, 38]. Surprisingly, in Hereford cross-bred
127 heifers, a comparison of the effect of hKp10 and cattle Kp10, that is structurally identical to oKp10
128 (figure 1A), on LH plasma concentration showed that hKp10 elicited a more pronounced effect than
129 cattle Kp10. In addition, repeated injections of hKp10 over a 2-hour period triggered ovulation, but
130 this was not the case when cattle Kp10 was used [39].

131 The equine Kp10 (eKp10) also has a single amino acid difference compared to the oKp10, in position
132 2 an arginine is substituted by a valine. This position is outside the Kp10 pharmacophore [34] and,
133 therefore, this modification should bear no consequence on the activity. At odds with this prediction,
134 when tested in a calcium mobilization assay on HEK293 cells transfected with the human *KISS1R*, the
135 eKp10 was actually less potent than oKp10 and hKp10 (figure 1B). Consistent with this result when
136 injected iv in the ewe (15 nmol/ewe, N=5 per group) the eKp10 is less potent *in vivo* than the oKp10
137 (figure 1C and D). The effect of eKp10 was extensively studied in the horse, where it was able to
138 stimulate the release of gonadotropins. However, in contrast to what found with oKp10 in the sheep
139 [36], it was unable to trigger or advance ovulation [40] and significantly disrupted normal sexual
140 receptivity in the estrous mare [41]. At present, it is difficult to draw any conclusion from this
141 scattered data. Nonetheless, they indicate that a single amino acid modification could result in a
142 significant change of Kps' capacity to activate the receptor of a different species. Considering that
143 often the hKp10 was used to study the physiology of the Kp system in other mammals, these data
144 point to the importance of using the appropriate endogenous ligand for the species under evaluation
145 in order to obtain meaningful physiological data.

146

147 **4. Anatomical localization of the Kp system**

148 Initial studies by RT-PCR reported the expression of *KISS1R* gene in several tissues including the brain,
149 the pituitary gland, the placenta and the pancreas [42-44]. Subsequent research focusing on the
150 brain showed its presence in the hypothalamus [45]. However, only a handful of studies describing a
151 more precise brain localization of *Kiss1r* are available. This information was obtained by applying *in*
152 *situ* hybridization in the rat [46, 47] and in the rhesus monkey [48] or by using a β -gal reporter gene
153 in the mouse [11]. Receptor expression was detected in several brain regions including the olfactory
154 bulb, the hypothalamus, the septum, the hippocampus, the thalamus, and the brainstem. The
155 distribution pattern appears significantly discordant between species suggesting either a different
156 sensitivity of the methods or a species-specific dependent expression of the receptor. This last
157 possibility deserves further investigation because it might have important implications to understand
158 the physiological role of the Kp system in different species.

159 On the other hand, double labelling studies in rats, monkeys and ewes consistently reported the
160 expression of *KISS1R* gene in GnRH neurons [47-49]. These data confirm that a direct effect of Kp on
161 GnRH neurons is a common feature.

162 Detection of the KISS1R protein has proven a difficult task. Most of the available antibodies are
163 insufficiently characterized lacking for example specificity validation performed in *Kiss1r* knockout
164 animals. Evaluation of the antibody using western blotting often results in multiple bands and
165 inconsistent molecular weights. Furthermore, use of antibodies raised against the rodent or human
166 receptor to detected KISS1R in other species without performing stringent controls in the target
167 species is common, leading to arguable results. This represents an obstacle to our understanding of
168 Kp action. The development of new tools, either dependable antibodies against the receptor or
169 tagged molecules allowing its detection and localization, would be an important advance in the field.
170 An attempt in this direction has been done by the creation of synthetic Kp-based probes labelled
171 with a fluorescent tag [50]. These probes were characterized successfully in *in vitro* systems, but
172 were not assessed for receptor localization on tissue. We have recently described the synthesis of a

173 tagged Kp-derived molecule capable of covalently binding KISS1R *in vitro* [51]. Further development
174 of this molecule to improve its stability is ongoing and might become a valuable tool for receptor
175 localization studies.

176 The expression of the *KISS1* gene and that of the related protein has been investigated in several
177 species. In mammals, two main neuronal populations sited in the diencephalon have been described.
178 One, located in the ARC, has been recognized in rodents [52-54], in primates [48, 55-57], in the sheep
179 [15, 58], in the goat [59, 60], in the horse [61], in the pig [62], in cattle [63], in the dromedary camel
180 [64], and in the cat [65]. The presence of the second one, located in the POA, has been reported in
181 rodents [52-54], in the sheep [15, 58], in the goat [59], in cattle [63], in the dromedary camel [64],
182 and in the cat [65], but data are contradictory in the horse [61, 66] and in primates [56, 57]. The
183 precise localization of this second population also varies with species: in rodents it is adjacent to the
184 ventricle whereas it is more lateral in other mammals [53].

185 Discrepancy on the localization of Kp in the horse could arise from the use of different antibodies. In
186 one study, Kp-like immunoreactivity was found not only in the POA but also in the dorso-medial
187 nucleus [66]. In this study an antibody from Phoenix Pharmaceutical was used, which have shown to
188 have cross reactivity with other members of the RFamide family [67] and NPVF (the precursor mRNA
189 for RFamide-related peptide 3) has been detected in the dorsomedial paraventricular hypothalamic
190 nuclei of the mare [68]. This antibody gave similar results in the sheep [69], but there is no evidence
191 of the presence of mRNA encoding for Kp in the dorso-medial region of the sheep hypothalamus
192 [58]. There is a single report of this antibody labelling GnRH neurons in the sheep [69], but these
193 results could not be reproduced using an antisera specific for Kp [53]. It is likely that this antibody
194 cross reacts with other neuropeptides of the RFamide family, such as NPVF, showing a sequence
195 similar to that of Kp. Nevertheless, the existence of a Kp POA population cannot be ruled out based
196 on antibody cross-reactivity alone because NPVF was not detected in the POA [68].

197 In addition, the Kp POA population is sexually dimorphic with females having a larger number of cells.
198 Indeed, several reports indicate an important role of sex steroids, in particular estrogens (Es), in
199 regulating the size of this population [1]. High levels of E, similar to that observed during the pre-
200 ovulatory period, would increase the number of neurons expressing Kp. In contrast, ovariectomy,
201 ablating ovarian E production, would decrease it to an almost undetectable level. The study reporting
202 the absence of Kp in the POA of the mare was performed on brain dissected between 2 and 4 hours
203 after ovulation [61]. At this stage of the ovulatory cycle E concentration is rapidly declining and is
204 lowered by 2/3 compared to its maximal level [70], leading to an E negative feedback. It is possible
205 that localization studies performed at a moment of the cycle when circulating E is high (i.e. the pre-
206 ovulatory phase) may confirm the existence of a neuronal population expressing Kp in the POA of the
207 mare.

208 The question about the existence of this neuronal population is not trivial. In the current view, the Kp
209 ARC population would be mainly involved in controlling the pulsatile release of GnRH and submitted
210 to the E negative feedback. Optogenetic studies in rodents [71-73] and combined pharmacological
211 and electrophysiological analyses in goats [74] converge in favor of this hypothesis. On the other
212 hand, the POA population would relay the E positive feedback triggering the increase of GnRH
213 secretion that leads to the pre-ovulatory surge [75]. However, in the sheep data are less clear-cut.
214 Some data suggest a rodent-like mechanism showing an activation, concomitant with the LH surge,
215 of POA neurons, but not of ARC neurons [76]. Others support a combined contribution of the two
216 populations in generating the ovine pre-ovulatory surge [77, 78]. Finally, an E implant in the
217 mediobasal hypothalamus induces a surge [79], and cells expressing *KISS1R* mRNA in the caudal ARC
218 were more abundant in the pre-ovulatory period than during the other phase of the cycle. These
219 data suggest a prominent role for the ARC Kp population in pre-ovulatory surge induction [80].

220 In the ewe the caudal portion of the Kp neurons of the ARC are less sensitive to E negative feedback
221 during the breeding season [81]. The abundance of the estrogen receptor α in the Kp neurons of the
222 ARC of pre-pubertal ewe was compared in ovariectomized (OVX) and OVX and E (OVX+E) replaced

223 animals. The OVX+E showed a decrease of estrogen receptor α in the rostral part on the nucleus and
224 a tendency to a decrease in the medial part and no effect on the caudal portion. Similar results were
225 obtained for the number of neurons expressing Kp [82]. The possibility that the ARC Kp neurons of
226 the ewe (and perhaps other species) would convey both positive and negative estrogen feedback to
227 GnRH neurons is an interesting hypothesis that remains to be further substantiated. If the absence of
228 a POA Kp population in the mare is confirmed, this species would be an interesting research model to
229 test this hypothesis.

230 In the ewe, a combined regulatory effect of sexual steroids and photoperiod has also been observed.
231 In OVX+E ewes, a lower level of Kp was observed in the mediobasal hypothalamus and in POA during
232 the non-breeding season compared to the breeding season [19] and switching from long to short
233 photoperiod induced an increase of Kp expressing neurons in POA and ARC [83].

234

235 A third population of Kp neurons has been detected in the medial amygdala of rodents [52, 84], a
236 region shaping social, emotional as well as sexual behaviors. Kp expression in these neurons is
237 sexually dimorphic, with males showing a larger population. Gonadectomy decreased *Kiss1*
238 expression in both sexes whereas treatment with either testosterone or E increased it [84].

239 Application of anterograde and retrograde tracers showed the existence of reciprocal connections
240 between Kp neurons located in the amygdala and the accessory olfactory bulb. A direct connection of
241 amygdala Kp neurons with GnRH neurons of the POA was also observed. Therefore, Kp neurons in
242 the medial amygdala could be part of the pathways relaying information controlling sexual behaviors
243 related to odor cues [85]. In small ruminants the so-called male effect, applied to resume the
244 ovulatory cycle during the seasonal anoestrus, is mediated mainly by odor cues. In the goat, input
245 from the medial nucleus of the amygdala to Kp neurons of the ARC has been proposed to play a role
246 in the male effect [86]. However, in domestic animals the role of Kp neurons located in the central
247 amygdala in processing and relaying odor cues remains unexplored. Actually, even the existence of

248 such a population in domestic species has not been demonstrated yet. This interesting research area
249 warrants further investigation.

250

251 The presence of the Kp system in the pituitary gland and an *in vitro* direct effect of Kp administration
252 on LH and FSH secretion and gene expression was reported in several species [42-44, 87-91] (see
253 [92, 93] for review on the subject). Even though available results suggest a potential direct
254 stimulation of gonadotrophs by Kp, nevertheless effects were mainly of small amplitude. The most
255 compelling evidence has suggested a minor, if any, role of Kp in stimulating the pituitary gland. In
256 particular, rescue of Kp signaling in GnRH neurons in *Kiss1r* KO mice completely reinstates a
257 reproductive phenotype [94, 95]. In hypothalamo-pituitary disconnected ewes there was no
258 temporal correlation between Kp levels recorded in the portal blood and LH pulsatile pattern
259 measured in the jugular vein [88]. Conversely, Kp peripheral administration is highly correlated with
260 GnRH release and its effect on gonadotropin secretion is blocked by a GnRH receptor antagonist [48,
261 96-98].

262

263 Expression and localization data indicate the presence of Kp and its receptor also in the reproductive
264 organs. In the female of various species, both Kp and its receptor have been localized in the ovary, in
265 the uterus and in the placenta [93]. Immunoreactivity was observed in granulosa cells, cumulus cells,
266 theca cells, stromal cells, trophoblast giant cells, etc. Localization differs between species and
267 sometime contrasting results in the same species were reported [99]. Contradictory findings could be
268 explained partially by the use of tissue obtained at different times of the estrus cycle under the
269 influence of a different hormonal milieu. Nevertheless, this variability is puzzling and methodological
270 biases are a concern. Some immunohistochemistry results suffer from poor characterization of the
271 antibodies that make them unreliable. Caution should be taken also when considering expression
272 data obtained by RT-PCR. They are sometime undependable due to an excessively high number of
273 amplification cycles leading to the generation of false positives.

274 Despite these notes of caution, some convincing results for a local role of Kp on the female
275 reproductive apparatus of rodents are available. In *Kiss1r* null mice ovary and uterus have a reduced
276 weight and no ovulation occurs [9]. In these mice, a treatment combining GnRH and gonadotrophins
277 triggers ovulation. However, null mice release a significantly reduced number of oocytes than WT
278 suggesting that *Kiss1r* in the ovary is required for a full recovery of ovarian activity [31]. In line with
279 this result *Kiss1r* haplo-insufficient mice display a syndrome resembling premature ovarian failure
280 with decline in ovulatory rate and reduction of pre-antral follicles [31]. The potential implication of
281 Kp in this phenomenon was discovered by studying the neurotrophin BDNF (brain-derived
282 neurotrophic factor) and its receptor NTRK2 (neurotrophic tyrosine kinase receptor type 2). In mice a
283 specific deletion of NTRK2 in oocyte results in oocyte death and early adulthood infertility similar to
284 that observed in *Kiss1r* haplo-insufficient mice. Experimental data suggest that Kp would act in
285 concert with BDNF to mediate the effect of gonadotrophins on NTRK2 and on ovarian function [100].
286 Clarifying the physiological action of Kp in the ovary of domestic animals would be particularly
287 interesting and relevant to optimize ovulation rate. Expression of the Kp system in the uterus has
288 been associated with embryo implantation [101]. In *Kiss1^{-/-}* mice application of E and gonadotropins
289 coupled to a superovulation protocol induces normal ovulation, egg fertilization and initial embryo
290 development; however implantation fails. When *Kiss1^{-/-}* embryos are transferred to a WT female
291 they implant successfully, indicating a maternal defect in the *Kiss^{-/-}* females. *Kiss1r* expression in the
292 uterus is greater in *Kiss^{-/-}* females and the distribution within the subluminal epithelium is different
293 from the WT females. Treatment with Kp10 failed to rescue embryos' implantation in *Kiss1^{-/-}* mice
294 [101]. In the uterus of *Kiss1^{-/-}* mice endometrial glands are poorly developed [101, 102] evoking the
295 possibility that this defect is the consequence of the lack of a Kp local action during uterus
296 development. Results showing that E treatment of *Kiss1^{-/-}* mice or selective restoration of Kp
297 signaling in GnRH neurons of *Kiss1^{-/-}* mice only partially restored endometrial gland formation
298 support this hypothesis [95]. Furthermore, the expression of the cytokine leukemia inhibitory factor,

299 essential for embryo implantation, is reduced in *Kiss1*^{-/-} uterine gland. Administration of leukemia
300 inhibitory factor to hormone-primed *Kiss1*^{-/-} partially rescued implantation [101].

301 Similarly to rodents, endometrial gland formation in sheep and pigs takes place during the postnatal
302 period, pointing to the relevance of these results for the physiology of domestic animals.

303 The presence and physiological significance of the Kp system in the male reproductive apparatus has
304 received less attention. In the mouse, the most convincing data indicate that the expression of the
305 genes for Kp and *Kiss1r* in the gonad is restricted to haploid spermatids. However, no protein for Kp
306 was detected, suggesting a translational repression, and Kp administration had no effect on
307 testosterone release from an immortalized Leydig cell line [103]. Conversely, in Leydig cells from goat
308 testes *Kiss1* and *Kiss1r* genes expression was increased by gonadotropins treatment. Incubation of
309 Leydig cells with the Kp antagonist Kp234 reduced testosterone secretion implying a potential role of
310 Kp in androgen production [104]. The conflicting results obtained in these studies could be due to a
311 physiological difference between species or due to the use of a cell line versus cells from primary
312 culture. A drawback of the goat study is the lack of evidence about a positive effect (i.e. stimulation
313 of testosterone production by Kp administration). Of note, Kp234 has been applied to counteract the
314 effect on Kp in various experimental systems [105]. However, in the dog it was unable to block the
315 effect of Kp both *in vitro* and *in vivo* [106]. Furthermore, it has been reported that p234 and p356,
316 the next generation analog of p234, exhibit weak agonist activity at *Kiss1r* in mice [107] and *in vitro*
317 [106]. These findings raise doubts that the observed effect would be mediated by the Kp system. The
318 fact that GnRH-specific rescue of *Kiss1r* null mice was sufficient to restore a normal fertile phenotype
319 and testes were not different from that of WT [94] support a minor role, if any, of Kp in modulating
320 testicular functions. Nevertheless, further studies on the subject would be welcome to clarify this
321 issue.

322

323

324 5. Pharmacological modulation of the kisspeptin system

325 A thorough understanding of Kp system functions requires the capacity to pharmacologically
326 modulate its action. Modulation of the Kp system was initially explored by injecting the endogenous
327 Kps. Of the various isoforms, Kp10 has been the most broadly used because it is easy to synthesize
328 and highly efficacious. Except in humans, where the effects of hKp54 have been quite well studied
329 (see [108] for a review on the subject), few studies have evaluated the effect of the longer isoforms.
330 The effect of mKp52 was tested by icv injection in the mouse, showing that at femtomolar
331 concentration it was able to increase the plasma concentration of LH and FSH [52]. In the cow,
332 intravenous administration of cattle Kp53 triggered an increase of LH, but not FSH, plasma
333 concentration and ovulation in 1 out of 4 cows [109]. In females of several species (rodents, ewe,
334 goat, cow, pig, horse, and dog) a single injection of Kp10 triggers a rapid, albeit short-lasting, increase
335 of LH and/or FSH [36, 37, 40, 45, 52, 54, 97, 110-113]. In the ewe during the non-breeding season a
336 perfusion of Kp10 lasting 24 hours or longer induced the ovulation in about 75% of the treated
337 animals [36, 114]. In prepubertal ewes, hourly administration of Kp10 for 24 hours induced an
338 increase of LH and ovulation [115]. These results corroborated the idea that the Kp system would be
339 an interesting target to modulate the female reproductive cycle. A major drawback in using Kp10 is
340 its short half-life due to rapid degradation and excretion. This is a relevant obstacle when prolonged
341 activation of the system is required, as in the case of ovulation induction or for the treatment of
342 chronic reproductive pathologies in humans (e.g. hypothalamic amenorrhea).

343 A possible way to overcome this problem is the use of the longer isoforms eliciting a protracted
344 activation of the system. In the male rat rKp52 produced an increase of LH of larger amplitude and
345 longer duration never matched, even at high doses, by rKp10 or hKp10 [116]. Consistently,
346 experiments performed in the male mouse indicated an increase of LH of longer duration and larger
347 amplitude after hKp54 compared to hKp10 injection [117]. At odds with these results, however, a
348 study on human males reported no significant difference in the effect of hKp10 and hKp54 [118]. The
349 hKp54 isoform was also used in women but a prolonged action was only obtained after perfusion for

350 several hours [119]. Interestingly, in women administration of hKp54 increased plasma LH and FSH
351 level regardless to the phase of the ovulatory cycle. Conversely, hKp10 is effective on the pre-
352 ovulatory phase but not in the follicular phase [120]. In ruminants Kp10 injection during the luteal
353 phase triggered an increase of LH [37, 121, 122] even though, at least in the ewe, it was lower than
354 that induced during the mid and late follicular phase [122, 123].

355 Regardless to the different effects observed between sexes, isoforms, or different phases of the
356 ovulatory cycle the overall duration of the stimulation after a single injection remains relatively short.
357 To tackle this problem synthetic analogs based on Kp10 structure were designed to reduce
358 degradation and increase their circulating half-life. Their potential use to manage livestock
359 reproduction has been recently reviewed [7, 124]. The effects of these analogs (TAK683 and C6) have
360 been extensively characterized in small ruminants. TAK683 was tested under different conditions in
361 the goat. It consistently increased LH plasma level, with an efficacy depending on the phase of the
362 ovulatory cycle [125-128], and induced ovulations. However, no data are available on the fertility of
363 these ovulations [126]. The other analog, C6, induced fertile ovulations in small ruminants (sheep
364 and goats) when injected as a substitute of PMSG (pregnant mare serum gonadotropin) after a
365 progestogen priming [129, 130]. The treatment was efficacious both in the breeding as well as in the
366 non-breeding season. These results are proof of the concept that the Kp system is a suitable target
367 for ovulation synchronization and induction in small ruminants.

368 C6 was recently evaluated in steroid-primed pre-pubertal gilts where initial results showed that it
369 evokes an LH surge and ovulation [131]. While non data are available on the effect of Kp analogs in
370 the cow, data obtained with Kp10 in pre-pubertal and adult cows demonstrate an increase of plasma
371 LH levels [112, 121, 132-135]. Furthermore, in Hereford cross-bred heifers under low plasma
372 progesterone concentrations repeated administration of hKp10 induced ovulation [39]. Interestingly,
373 a difference in the capacity of cattle Kp10 to trigger LH increase was observed in different cattle
374 species with Holstein cows (*Bos taurus*) being more responsive than Gyr cows (*Bos indicus*) [112]. The
375 potential impact of this difference on the capacity to trigger ovulation remains to be explored.

376

377 There is only a handful of studies on the effect of Kp and its analogs on male domestic animals.

378 In adult and pre-pubertal male goats an intravenous injection of hKp10 stimulated the release of LH

379 and testosterone [38]. Similar results were also obtained in adult rams by intramuscular injection of

380 the Kp analog C6 [124]. In pre-pubertal Japanese bull calves a bolus injection of hKp10 increased both

381 LH and FSH with a temporal profile similar to that observed in other species. Interestingly the effect

382 was more pronounced in males compared to females of the same age [133]. This result is reminiscent

383 of data obtained with the C6 in the mouse showing a longer lasting effect on LH plasma

384 concentration in males compared to females [129].

385 hKp10 and C6 were recently tested in pre-pubertal Holstein bull calves using a treatment consisting

386 of a daily intramuscular injection for 4 days. hKp10 had no effect on LH and FSH concentration on

387 both the first and last day of treatment [136]. This is not completely surprising because of the low

388 dose used (20 nmol/calf) and the route of administration. Experiments comparing intravenous and

389 intramuscular injection of hKp10 in pre-pubertal bull calves applying a higher dose (about 550

390 nmol/calf) showed a significantly reduced effect of intramuscular compared to intravenous injection

391 [133]. On the other hand, the first injection of C6 in bull calves increased LH plasma concentration for

392 several hours but the increase after the fourth injection was limited if any. Interestingly, C6 had no

393 immediate effect on circulating FSH, but at the fourth injection FSH level were lower compared to

394 control [136].

395 Further investigations will be required to evaluate if repeated administration of Kp or its analogs may

396 be a valuable approach to stimulate male sexual behaviors during the non-breeding season in small

397 ruminants and to advance the onset of puberty in bulls.

398

399 **6. Conclusions**

400

401 In mammals an overall similar scenario has emerged showing that the Kp system plays a key role in
402 stimulating the hypothalamo-pituitary-gonad axis. Yet significant species-specific differences exist
403 and their potential implications in shaping the peculiar reproductive features of various domestic
404 species is still unclear. To fully appreciate the many facets of these species-specific differences and
405 their physiological reverberations in domestic animal reproduction, several aspects need to be
406 further investigated. In particular differences in receptor and peptide localization, the physiological
407 relevance of the different Kp isoforms, the creation of new tools for receptor localization, the
408 implication of the Kp system in mediating sexual behavior, etc. are all exciting under-investigated
409 areas of research.

410 Outside reproduction, the involvement of the Kp system in other physiological functions ranging
411 from control of metabolism to motivation to stress responses has been proposed. These functions
412 are indirectly involved in reproduction and may complement the more direct effect of the Kp system
413 on the reproductive axis. However, they have been only partially characterized in primates and
414 rodents and remain almost unexplored in domestic animals. To conclude we could say that while
415 much has been learned, more remains to be discovered on the physiological ramifications of the Kp
416 system in reproduction control.

417

418 **7. Acknowledgements.** Funding: This work was supported by the French National Research
419 Agency (ANR, Kiss project, ANR-15-CE20-0015-01), but had not role in study design; in the collection,
420 analysis and interpretation of data; in the writing of the report; and in the decision to submit the
421 article for publication. We thank the professor Greg Anderson for revising the article.

422

423 **8. Bibliography**

424 [1] L. Pinilla, E. Aguilar, C. Dieguez, R.P. Millar, M. Tena-Sempere, Kisspeptins and reproduction:
425 physiological roles and regulatory mechanisms, *Physiol Rev*, 92 (2012) 1235-1316.

426 [2] A.N. Comninou, W.S. Dhillon, Emerging Roles of Kisspeptin in Sexual and Emotional Brain
427 Processing, *Neuroendocrinology*, 106 (2018) 195-202.

428 [3] I. Franceschini, E. Desroziers, Development and Aging of the Kisspeptin-GPR54 System in the
429 Mammalian Brain: What are the Impacts on Female Reproductive Function?, *Front Endocrinol*
430 (Lausanne), 4 (2013) 22.

431 [4] S.H. Yeo, W.H. Colledge, The Role of Kiss1 Neurons As Integrators of Endocrine, Metabolic, and
432 Environmental Factors in the Hypothalamic-Pituitary-Gonadal Axis, *Front Endocrinol (Lausanne)*, 9
433 (2018) 188.

434 [5] H. Okamura, T. Yamamura, Y. Wakabayashi, Kisspeptin as a master player in the central control of
435 reproduction in mammals: an overview of kisspeptin research in domestic animals, *Anim Sci J*, 84
436 (2013) 369-381.

437 [6] J.A. Daniel, C.D. Foradori, B.K. Whitlock, J.L. Sartin, Reproduction and beyond, kisspeptin in
438 ruminants, *J Anim Sci Biotechnol*, 6 (2015) 23.

439 [7] C.A. Lents, Review: kisspeptin and reproduction in the pig, *Animal*, (2019) 1-14.

440 [8] C.J. Scott, J.L. Rose, A.J. Gunn, B.M. McGrath, Kisspeptin and the regulation of the reproductive
441 axis in domestic animals, *J. Endocrinol.*, 240 (2019) R1-R16.

442 [9] S.B. Seminara, S. Messenger, E.E. Chatzidaki, R.R. Thresher, J.S. Acierno, Jr., J.K. Shagoury, Y. Bo-
443 Abbas, W. Kuohung, K.M. Schwinof, A.G. Hendrick, D. Zahn, J. Dixon, U.B. Kaiser, S.A. Slaugenhaupt,
444 J.F. Gusella, S. O'Rahilly, M.B. Carlton, W.F. Crowley, Jr., S.A. Aparicio, W.H. Colledge, The GPR54
445 gene as a regulator of puberty, *N Engl J Med*, 349 (2003) 1614-1627.

446 [10] N. de Roux, E. Genin, J.C. Carel, F. Matsuda, J.L. Chaussain, E. Milgrom, Hypogonadotropic
447 hypogonadism due to loss of function of the KiSS1-derived peptide receptor GPR54, *Proc Natl Acad
448 Sci U S A*, 100 (2003) 10972-10976.

449 [11] A.E. Herbison, X. de Tassigny, J. Doran, W.H. Colledge, Distribution and postnatal development
450 of Gpr54 gene expression in mouse brain and gonadotropin-releasing hormone neurons,
451 *Endocrinology*, 151 (2010) 312-321.

452 [12] S. Messenger, E.E. Chatzidaki, D. Ma, A.G. Hendrick, D. Zahn, J. Dixon, R.R. Thresher, I. Malinge, D.
453 Lomet, M.B. Carlton, W.H. Colledge, A. Caraty, S.A. Aparicio, Kisspeptin directly stimulates
454 gonadotropin-releasing hormone release via G protein-coupled receptor 54, *Proc Natl Acad Sci U S A*,
455 102 (2005) 1761-1766.

456 [13] A. Caraty, D. Lomet, S. Me, D. Guillaume, M. Beltramo, N. Evans, GnRH release into the
457 hypophyseal portal blood of the ewe mirrors both pulsatile and continuous intravenous infusion of
458 kisspeptin: an insight into kisspeptin's mechanism of action, *J Neuroendocrinol*, 25 (2013) 537-546.

459 [14] V.M. Navarro, R. Fernandez-Fernandez, J.M. Castellano, J. Roa, A. Mayen, M.L. Barreiro, F.
460 Gaytan, E. Aguilar, L. Pinilla, C. Dieguez, M. Tena-Sempere, Advanced vaginal opening and precocious
461 activation of the reproductive axis by KiSS-1 peptide, the endogenous ligand of GPR54, *J Physiol*, 561
462 (2004) 379-386.

463 [15] I. Franceschini, D. Lomet, M. Cateau, G. Delsol, Y. Tillet, A. Caraty, Kisspeptin immunoreactive
464 cells of the ovine preoptic area and arcuate nucleus co-express estrogen receptor alpha, *Neurosci
465 Lett*, 401 (2006) 225-230.

466 [16] J. Clarkson, X. d'Anglemont de Tassigny, A.S. Moreno, W.H. Colledge, A.E. Herbison, Kisspeptin-
467 GPR54 signaling is essential for preovulatory gonadotropin-releasing hormone neuron activation and
468 the luteinizing hormone surge, *J Neurosci*, 28 (2008) 8691-8697.

469 [17] M.N. Lehman, C.M. Merkley, L.M. Coolen, R.L. Goodman, Anatomy of the kisspeptin neural
470 network in mammals, *Brain Res.*, 1364 (2010) 90-102.

471 [18] K. Iwata, Y. Kunimura, K. Matsumoto, H. Ozawa, Effect of androgen on Kiss1 expression and
472 luteinizing hormone release in female rats, *J Endocrinol*, 233 (2017) 281-292.

473 [19] C. Urias-Castro, J.A. Arreguin-Arevalo, C. Magee, E.R. Weber, T.M. Nett, Hypothalamic
474 concentrations of kisspeptin and gonadotropin-releasing hormone during the breeding season and
475 non-breeding season in ewes, *Am J Reprod Immunol*, 82 (2019) e13146.

476 [20] G. Cheng, L.M. Coolen, V. Padmanabhan, R.L. Goodman, M.N. Lehman, The
477 kisspeptin/neurokinin B/dynorphin (KNDy) cell population of the arcuate nucleus: sex differences and
478 effects of prenatal testosterone in sheep, *Endocrinology*, 151 (2010) 301-311.

479 [21] M. Amstalden, L.M. Coolen, A.M. Hemmerle, H.J. Billings, J.M. Connors, R.L. Goodman, M.N.
480 Lehman, Neurokinin 3 receptor immunoreactivity in the septal region, preoptic area and
481 hypothalamus of the female sheep: colocalisation in neurokinin B cells of the arcuate nucleus but not
482 in gonadotrophin-releasing hormone neurones, *J Neuroendocrinol*, 22 (2010) 1-12.

483 [22] Y. Wakabayashi, T. Nakada, K. Murata, S. Ohkura, K. Mogi, V.M. Navarro, D.K. Clifton, Y. Mori, H.
484 Tsukamura, K. Maeda, R.A. Steiner, H. Okamura, Neurokinin B and dynorphin A in kisspeptin neurons
485 of the arcuate nucleus participate in generation of periodic oscillation of neural activity driving
486 pulsatile gonadotropin-releasing hormone secretion in the goat, *The Journal of neuroscience*, 30
487 (2010) 3124-3132.

488 [23] R.L. Goodman, L.M. Coolen, M.N. Lehman, A role for neurokinin B in pulsatile GnRH secretion in
489 the ewe, *Neuroendocrinology*, 99 (2014) 18-32.

490 [24] T.M. Plant, The neurobiological mechanism underlying hypothalamic GnRH pulse generation: the
491 role of kisspeptin neurons in the arcuate nucleus, *F1000Res*, 8 (2019).

492 [25] J. Pasquier, A.G. Lafont, K. Rousseau, B. Querat, P. Chemineau, S. Dufour, Looking for the bird
493 Kiss: evolutionary scenario in sauropsids, *BMC Evol Biol*, 14 (2014) 30.

494 [26] M.G. Teles, S.D. Bianco, V.N. Brito, E.B. Trarbach, W. Kuohung, S. Xu, S.B. Seminara, B.B.
495 Mendonca, U.B. Kaiser, A.C. Latronico, A GPR54-activating mutation in a patient with central
496 precocious puberty, *N Engl J Med*, 358 (2008) 709-715.

497 [27] W.H. Colledge, Transgenic mouse models to study Gpr54/kisspeptin physiology, *Peptides*, 30
498 (2009) 34-41.

499 [28] G.L. Cao, M.X. Chu, L. Fang, R. Di, T. Feng, N. Li, Analysis on DNA sequence of KiSS-1 gene and its
500 association with litter size in goats, *Mol Biol Rep*, 37 (2010) 3921-3929.

501 [29] X. An, T. Ma, J. Hou, F. Fang, P. Han, Y. Yan, H. Zhao, Y. Song, J. Wang, B. Cao, Association
502 analysis between variants in KISS1 gene and litter size in goats, *BMC Genet.*, 14 (2013) 63.

503 [30] M.S. El-Tarabany, A.W. Zagloul, A.A. El-Tarabany, A. Awad, Association analysis of polymorphism
504 in KISS1 gene with reproductive traits in goats, *Anim Reprod Sci*, 180 (2017) 92-99.

505 [31] F. Gaytan, D. Garcia-Galiano, M.D. Dorfman, M. Manfredi-Lozano, J.M. Castellano, G.A. Dissen,
506 S.R. Ojeda, M. Tena-Sempere, Kisspeptin receptor haplo-insufficiency causes premature ovarian
507 failure despite preserved gonadotropin secretion, *Endocrinology*, 155 (2014) 3088-3097.

508 [32] A. Caraty, J. Leprince, L. D., B. Lefranc, A. Bourmaud, M.E. Sebert, M. Beltramo, I. Franceschini, H.
509 Vaudry, Kisspeptin isoforms in the hypothalamus : what new, 2nd World Conference Kisspeptin
510 Signaling in the Brain, (2012).

511 [33] R. Quillet, S. Ayachi, F. Bihel, K. Elhabazi, B. Ilien, F. Simonin, RF-amide neuropeptides and their
512 receptors in Mammals: Pharmacological properties, drug development and main physiological
513 functions, *Pharmacol Ther*, 160 (2016) 84-132.

514 [34] E. Gutierrez-Pascual, J. Leprince, A.J. Martinez-Fuentes, I. Segalas-Milazzo, R. Pineda, J. Roa, M.
515 Duran-Prado, L. Guilhaudis, E. Desperrois, A. Lebreton, L. Pinilla, M.C. Tonon, M.M. Malagon, H.
516 Vaudry, M. Tena-Sempere, J.P. Castano, In vivo and in vitro structure-activity relationships and
517 structural conformation of Kisspeptin-10-related peptides, *Mol. Pharmacol.*, 76 (2009) 58-67.

518 [35] M. Beltramo, V. Robert, M. Galibert, J.B. Madinier, P. Marceau, H. Dardente, C. Decourt, N. De
519 Roux, D. Lomet, A.F. Delmas, A. Caraty, V. Aucagne, Rational design of triazololipopeptides analogs of
520 kisspeptin inducing a long-lasting increase of gonadotropins, *J. Med. Chem.*, 58 (2015) 3459-3470.

521 [36] A. Caraty, J.T. Smith, D. Lomet, S. Ben Said, A. Morrissey, J. Cognie, B. Doughton, G. Baril, C.
522 Briant, I.J. Clarke, Kisspeptin synchronizes preovulatory surges in cyclical ewes and causes ovulation
523 in seasonally acyclic ewes, *Endocrinology*, 148 (2007) 5258-5267.

524 [37] T. Hashizume, H. Saito, T. Sawada, T. Yaegashi, A.A. Ezzat, K. Sawai, T. Yamashita, Characteristics
525 of stimulation of gonadotropin secretion by kisspeptin-10 in female goats, *Anim Reprod Sci*, 118
526 (2010) 37-41.

527 [38] H. Saito, T. Sawada, T. Yaegashi, Y. Goto, J. Jin, K. Sawai, T. Hashizume, Kisspeptin-10 stimulates
528 the release of luteinizing hormone and testosterone in pre- and post-pubertal male goats, *Anim Sci J*,
529 83 (2012) 487-492.

530 [39] C.E.P. Leonardi, F.C.F. Dias, G.P. Adams, E.R. Araujo, J. Singh, Kisspeptin induces ovulation in
531 heifers under low plasma progesterone concentrations, *Theriogenology*, 141 (2019) 26-34.

532 [40] C. Decourt, A. Caraty, C. Briant, D. Guillaume, D. Lomet, D. Chesneau, L. Lardic, G. Duchamp, F.
533 Reigner, P. Monget, L. Dufourny, M. Beltramo, H. Dardente, Acute injection and chronic perfusion of

534 kisspeptin elicit gonadotropins release but fail to trigger ovulation in the mare, *Biol Reprod*, 90 (2014)
535 1-12.

536 [41] C. Magee, J.E. Bruemmer, T.M. Nett, E.L. Squires, C.M. Clay, Kisspeptide in the estrous mare: is it
537 an appropriate ovulation-inducing agent?, *Theriogenology*, 78 (2012) 1987-1996.

538 [42] M. Kotani, M. Detheux, A. Vandenbogaerde, D. Communi, J.M. Vanderwinden, E. Le Poul, S.
539 Brezillon, R. Tyldesley, N. Suarez-Huerta, F. Vandeput, C. Blanpain, S.N. Schiffmann, G. Vassart, M.
540 Parmentier, The metastasis suppressor gene KiSS-1 encodes kisspeptins, the natural ligands of the
541 orphan G protein-coupled receptor GPR54, *J Biol Chem*, 276 (2001) 34631-34636.

542 [43] S. Funes, J.A. Hedrick, G. Vassileva, L. Markowitz, S. Abbondanzo, A. Golovko, S. Yang, F.J.
543 Monsma, E.L. Gustafson, The KiSS-1 receptor GPR54 is essential for the development of the murine
544 reproductive system, *Biochem Biophys Res Commun*, 312 (2003) 1357-1363.

545 [44] A.I. Muir, L. Chamberlain, N.A. Elshourbagy, D. Michalovich, D.J. Moore, A. Calamari, P.G.
546 Szekeres, H.M. Sarau, J.K. Chambers, P. Murdock, K. Steplewski, U. Shabon, J.E. Miller, S.E.
547 Middleton, J.G. Darker, C.G. Larminie, S. Wilson, D.J. Bergsma, P. Emson, R. Faull, K.L. Philpott, D.C.
548 Harrison, AXOR12, a novel human G protein-coupled receptor, activated by the peptide KiSS-1, *J Biol*
549 *Chem*, 276 (2001) 28969-28975.

550 [45] V.M. Navarro, J.M. Castellano, R. Fernandez-Fernandez, M.L. Barreiro, J. Roa, J.E. Sanchez-
551 Criado, E. Aguilar, C. Dieguez, L. Pinilla, M. Tena-Sempere, Developmental and hormonally regulated
552 messenger ribonucleic acid expression of KiSS-1 and its putative receptor, GPR54, in rat
553 hypothalamus and potent luteinizing hormone-releasing activity of KiSS-1 peptide, *Endocrinology*,
554 145 (2004) 4565-4574.

555 [46] D.K. Lee, T. Nguyen, G.P. O'Neill, R. Cheng, Y. Liu, A.D. Howard, N. Coulombe, C.P. Tan, A.T. Tang-
556 Nguyen, S.R. George, B.F. O'Dowd, Discovery of a receptor related to the galanin receptors, *FEBS*
557 *Lett*, 446 (1999) 103-107.

558 [47] S. Higo, S. Honda, N. Iijima, H. Ozawa, Mapping of Kisspeptin Receptor mRNA in the Whole Rat
559 Brain and its Co-Localisation with Oxytocin in the Paraventricular Nucleus, *J Neuroendocrinol*, 28
560 (2016).

561 [48] M. Shahab, C. Mastronardi, S.B. Seminara, W.F. Crowley, S.R. Ojeda, T.M. Plant, Increased
562 hypothalamic GPR54 signaling: a potential mechanism for initiation of puberty in primates, *Proc Natl*
563 *Acad Sci U S A*, 102 (2005) 2129-2134.

564 [49] Q. Li, A. Roa, I.J. Clarke, J.T. Smith, Seasonal variation in the gonadotropin-releasing hormone
565 response to kisspeptin in sheep: possible kisspeptin regulation of the kisspeptin receptor,
566 *Neuroendocrinology*, 96 (2012) 212-221.

567 [50] M. Kaneda, R. Misu, H. Ohno, A. Hirasawa, N. Ieda, Y. Uenoyama, H. Tsukamura, K. Maeda, S.
568 Oishi, N. Fujii, Design and synthesis of fluorescent probes for GPR54, *Bioorg Med Chem*, 22 (2014)
569 3325-3330.

570 [51] W. Vannecke, C. Ampe, M. Van Troys, M. Beltramo, A. Madder, Cross-Linking Furan-Modified
571 Kisspeptin-10 to the KISS Receptor, *ACS Chem Biol*, 12 (2017) 2191-2200.

572 [52] M.L. Gottsch, M.J. Cunningham, J.T. Smith, S.M. Popa, B.V. Acohido, W.F. Crowley, S. Seminara,
573 D.K. Clifton, R.A. Steiner, A role for kisspeptins in the regulation of gonadotropin secretion in the
574 mouse, *Endocrinology*, 145 (2004) 4073-4077.

575 [53] J. Clarkson, X. d'Anglemont de Tassigny, W.H. Colledge, A. Caraty, A.E. Herbison, Distribution of
576 kisspeptin neurones in the adult female mouse brain, *J Neuroendocrinol*, 21 (2009) 673-682.

577 [54] T.J. Greives, A.O. Mason, M.A. Scotti, J. Levine, E.D. Ketterson, L.J. Kriegsfeld, G.E. Demas,
578 Environmental control of kisspeptin: implications for seasonal reproduction, *Endocrinology*, 148
579 (2007) 1158-1166.

580 [55] S. Ramaswamy, K.A. Guerriero, R.B. Gibbs, T.M. Plant, Structural interactions between kisspeptin
581 and GnRH neurons in the mediobasal hypothalamus of the male rhesus monkey (*Macaca mulatta*) as
582 revealed by double immunofluorescence and confocal microscopy, *Endocrinology*, 149 (2008) 4387-
583 4395.

584 [56] M. Shibata, R.L. Friedman, S. Ramaswamy, T.M. Plant, Evidence that down regulation of
585 hypothalamic KiSS-1 expression is involved in the negative feedback action of testosterone to

586 regulate luteinising hormone secretion in the adult male rhesus monkey (*Macaca mulatta*), *J*
587 *Neuroendocrinol*, 19 (2007) 432-438.

588 [57] A.M. Rometo, S.J. Krajewski, M.L. Voytko, N.E. Rance, Hypertrophy and increased kisspeptin
589 gene expression in the hypothalamic infundibular nucleus of postmenopausal women and
590 ovariectomized monkeys, *The Journal of clinical endocrinology and metabolism*, 92 (2007) 2744-
591 2750.

592 [58] J.T. Smith, C.M. Clay, A. Caraty, I.J. Clarke, KiSS-1 messenger ribonucleic acid expression in the
593 hypothalamus of the ewe is regulated by sex steroids and season, *Endocrinology*, 148 (2007) 1150-
594 1157.

595 [59] S. Matsuyama, S. Ohkura, K. Mogi, Y. Wakabayashi, Y. Mori, H. Tsukamura, K. Maeda, M.
596 Ichikawa, H. Okamura, Morphological evidence for direct interaction between kisspeptin and
597 gonadotropin-releasing hormone neurons at the median eminence of the male goat: an
598 immunoelectron microscopic study, *Neuroendocrinology*, 94 (2011) 323-332.

599 [60] S. Ohkura, K. Takase, S. Matsuyama, K. Mogi, T. Ichimaru, Y. Wakabayashi, Y. Uenoyama, Y. Mori,
600 R.A. Steiner, H. Tsukamura, K.I. Maeda, H. Okamura, Gonadotrophin-releasing hormone pulse
601 generator activity in the hypothalamus of the goat, *J Neuroendocrinol*, 21 (2009) 813-821.

602 [61] C. Decourt, Y. Tillet, A. Caraty, I. Franceschini, C. Briant, Kisspeptin immunoreactive neurons in
603 the equine hypothalamus Interactions with GnRH neuronal system, *J Chem Neuroanat*, 36 (2008)
604 131-137.

605 [62] J. Tomikawa, T. Homma, S. Tajima, T. Shibata, Y. Inamoto, K. Takase, N. Inoue, S. Ohkura, Y.
606 Uenoyama, K. Maeda, H. Tsukamura, Molecular characterization and estrogen regulation of
607 hypothalamic KISS1 gene in the pig, *Biol Reprod*, 82 313-319.

608 [63] R.C. Cardoso, B.R. Alves, S.M. Sharpton, G.L. Williams, M. Amstalden, Nutritional Programming
609 of Accelerated Puberty in Heifers: Involvement of Pro-Opiomelanocortin Neurones in the Arcuate
610 Nucleus, *J Neuroendocrinol*, 27 (2015) 647-657.

611 [64] H. Ainani, N. El Bousmaki, V.J. Poirel, M.R. Achaaban, M. Ouassat, M. Piro, P. Klosen, V.
612 Simonneaux, K. El Allali, The dromedary camel displays annual variation in hypothalamic kisspeptin
613 and Arg-Phe-amide-related peptide-3 according to sex, season, and breeding activity, *J Comp Neurol*,
614 (2019).

615 [65] O. Amelkina, P. Tanyapanyachon, A. Thongphakdee, K. Chatdarong, Identification of feline Kiss1
616 and distribution of immunoreactive kisspeptin in the hypothalamus of the domestic cat, *J Reprod*
617 *Dev*, 65 (2019) 335-343.

618 [66] C. Magee, C.D. Foradori, J.E. Bruemmer, J.A. Arreguin-Arevalo, P.M. McCue, R.J. Handa, E.L.
619 Squires, C.M. Clay, Biological and anatomical evidence for kisspeptin regulation of the hypothalamic-
620 pituitary-gonadal axis of estrous horse mares, *Endocrinology*, 150 (2009) 2813-2821.

621 [67] F.G. Revel, M. Saboureau, P. Pevet, V. Simonneaux, J.D. Mikkelsen, RFamide-related peptide
622 gene is a melatonin-driven photoperiodic gene, *Endocrinology*, 149 (2008) 902-912.

623 [68] J.F. Thorson, L.D. Prezotto, R.C. Cardoso, S.M. Sharpton, J.F. Edwards, T.H. Welsh, Jr., P.K. Riggs,
624 A. Caraty, M. Amstalden, G.L. Williams, Hypothalamic distribution, adenohipophyseal receptor
625 expression, and ligand functionality of RFamide-related peptide 3 in the mare during the breeding
626 and nonbreeding seasons, *Biol. Reprod.*, 90 (2014) 28.

627 [69] S. Pompolo, A. Pereira, K.M. Estrada, I.J. Clarke, Colocalization of kisspeptin and gonadotropin-
628 releasing hormone in the ovine brain, *Endocrinology*, 147 (2006) 804-810.

629 [70] O.J. Ginther, M.D. Utt, D.R. Bergfelt, M.A. Beg, Controlling interrelationships of progesterone/LH
630 and estradiol/LH in mares, *Anim Reprod Sci*, 95 (2006) 144-150.

631 [71] S.Y. Han, T. McLennan, K. Czielesky, A.E. Herbison, Selective optogenetic activation of arcuate
632 kisspeptin neurons generates pulsatile luteinizing hormone secretion, *Proc Natl Acad Sci U S A*, 112
633 (2015) 13109-13114.

634 [72] J. Clarkson, S.Y. Han, R. Piet, T. McLennan, G.M. Kane, J. Ng, R.W. Porteous, J.S. Kim, W.H.
635 Colledge, K.J. Iremonger, A.E. Herbison, Definition of the hypothalamic GnRH pulse generator in
636 mice, *Proc Natl Acad Sci U S A*, 114 (2017) E10216-E10223.

637 [73] H.J. McQuillan, S.Y. Han, I. Cheong, A.E. Herbison, GnRH Pulse Generator Activity Across the
638 Estrous Cycle of Female Mice, *Endocrinology*, 160 (2019) 1480-1491.

639 [74] T. Yamamura, Y. Wakabayashi, K. Sakamoto, H. Matsui, M. Kusaka, T. Tanaka, S. Ohkura, H.
640 Okamura, The Effects of Chronic Subcutaneous Administration of an Investigational Kisspeptin
641 Analog, TAK-683, on Gonadotropin-Releasing Hormone Pulse Generator Activity in Goats,
642 *Neuroendocrinology*, 100 (2014) 250-264.

643 [75] F. Matsuda, K. Nakatsukasa, Y. Suetomi, Y. Naniwa, D. Ito, N. Inoue, Y. Wakabayashi, H.
644 Okamura, K.I. Maeda, Y. Uenoyama, H. Tsukamura, S. Ohkura, The Luteinising Hormone Surge-
645 Generating System is Functional in Male Goats as in Females: Involvement of Kisspeptin Neurones in
646 the Medial Preoptic Area, *J. Neuroendocrinol.*, 27 (2015) 57-65.

647 [76] G.E. Hoffman, W.W. Le, I. Franceschini, A. Caraty, J.P. Advis, Expression of fos and in vivo median
648 eminence release of LHRH identifies an active role for preoptic area kisspeptin neurons in
649 synchronized surges of LH and LHRH in the ewe, *Endocrinology*, 152 (2011) 214-222.

650 [77] C.M. Merkle, K.L. Porter, L.M. Coolen, S.M. Hileman, H.J. Billings, S. Drews, R.L. Goodman, M.N.
651 Lehman, KNDy (kisspeptin/neurokinin B/dynorphin) neurons are activated during both pulsatile and
652 surge secretion of LH in the ewe, *Endocrinology*, 153 (2012) 5406-5414.

653 [78] J.T. Smith, Q. Li, A. Pereira, I.J. Clarke, Kisspeptin neurons in the ovine arcuate nucleus and
654 preoptic area are involved in the preovulatory luteinizing hormone surge, *Endocrinology*, 150 (2009)
655 5530-5538.

656 [79] A. Caraty, C. Fabre-Nys, B. Delaleu, A. Locatelli, G. Bruneau, F.J. Karsch, A. Herbison, Evidence
657 that the mediobasal hypothalamus is the primary site of action of estradiol in inducing the
658 preovulatory gonadotropin releasing hormone surge in the ewe, *Endocrinology*, 139 (1998) 1752-
659 1760.

660 [80] K.M. Estrada, C.M. Clay, S. Pompolo, J.T. Smith, I.J. Clarke, Elevated KiSS-1 expression in the
661 arcuate nucleus prior to the cyclic preovulatory gonadotrophin-releasing hormone/lutenising
662 hormone surge in the ewe suggests a stimulatory role for kisspeptin in oestrogen-positive feedback,
663 *J. Neuroendocrinol.*, 18 (2006) 806-809.

664 [81] J.T. Smith, L.M. Coolen, L.J. Kriegsfeld, I.P. Sari, M.R. Jaafarzadehshirazi, M. Maltby, K. Bateman,
665 R.L. Goodman, A.J. Tilbrook, T. Ubuka, G.E. Bentley, I.J. Clarke, M.N. Lehman, Variation in kisspeptin
666 and RFamide-related peptide (RFRP) expression and terminal connections to gonadotropin-releasing
667 hormone neurons in the brain: a novel medium for seasonal breeding in the sheep, *Endocrinology*,
668 149 (2008) 5770-5782.

669 [82] M.N. Bedenbaugh, M. D'Oliveira, R.C. Cardoso, S.M. Hileman, G.L. Williams, M. Amstalden,
670 Pubertal Escape From Estradiol Negative Feedback in Ewe Lambs Is Not Accounted for by Decreased
671 ESR1 mRNA or Protein in Kisspeptin Neurons, *Endocrinology*, 159 (2018) 426-438.

672 [83] S. Chalivoix, A. Bagnolini, A. Caraty, J. Cognie, B. Malpoux, L. Dufourny, Effects of photoperiod on
673 kisspeptin neuronal populations of the ewe diencephalon in connection with reproductive function,
674 *J. Neuroendocrinol.*, 22 (2010) 110-118.

675 [84] J. Kim, S.J. Semaan, D.K. Clifton, R.A. Steiner, S. Dhamija, A.S. Kauffman, Regulation of Kiss1
676 expression by sex steroids in the amygdala of the rat and mouse, *Endocrinology*, 152 (2011) 2020-
677 2030.

678 [85] R. Pineda, F. Plaisier, R.P. Millar, M. Ludwig, Amygdala Kisspeptin Neurons: Putative Mediators of
679 Olfactory Control of the Gonadotropic Axis, *Neuroendocrinology*, 104 (2017) 223-238.

680 [86] K. Sakamoto, Y. Wakabayashi, T. Yamamura, T. Tanaka, Y. Takeuchi, Y. Mori, H. Okamura, A
681 population of kisspeptin/neurokinin B neurons in the arcuate nucleus may be the central target of
682 the male effect phenomenon in goats, *PLoS One*, 8 (2013) e81017.

683 [87] M. Bellingham, P.A. Fowler, M.R. Amezcaga, S.M. Rhind, C. Cotinot, B. Mandon-Pepin, R.M.
684 Sharpe, N.P. Evans, Exposure to a complex cocktail of environmental endocrine-disrupting
685 compounds disturbs the kisspeptin/GPR54 system in ovine hypothalamus and pituitary gland,
686 *Environ. Health Perspect.*, 117 (2009) 1556-1562.

687 [88] J.T. Smith, A. Rao, A. Pereira, A. Caraty, R.P. Millar, I.J. Clarke, Kisspeptin is present in ovine
688 hypophysial portal blood but does not increase during the preovulatory luteinizing hormone surge:

689 evidence that gonadotropes are not direct targets of kisspeptin in vivo, *Endocrinology*, 149 (2008)
690 1951-1959.

691 [89] N. Richard, G. Galmiche, S. Corvaisier, A. Caraty, M.L. Kottler, KiSS-1 and GPR54 genes are co-
692 expressed in rat gonadotrophs and differentially regulated in vivo by oestradiol and gonadotrophin-
693 releasing hormone, *J Neuroendocrinol*, 20 (2008) 381-393.

694 [90] V.M. Navarro, J.M. Castellano, R. Fernandez-Fernandez, S. Tovar, J. Roa, A. Mayen, R. Nogueiras,
695 M.J. Vazquez, M.L. Barreiro, P. Magni, E. Aguilar, C. Dieguez, L. Pinilla, M. Tena-Sempere,
696 Characterization of the potent luteinizing hormone-releasing activity of KiSS-1 peptide, the natural
697 ligand of GPR54, *Endocrinology*, 146 (2005) 156-163.

698 [91] S. Suzuki, H. Kadokawa, T. Hashizume, Direct kisspeptin-10 stimulation on luteinizing hormone
699 secretion from bovine and porcine anterior pituitary cells, *Anim Reprod Sci*, 103 (2008) 360-365.

700 [92] M.D. Gahete, M.C. Vazquez-Borrego, A.J. Martinez-Fuentes, M. Tena-Sempere, J.P. Castano,
701 R.M. Luque, Role of the Kiss1/Kiss1r system in the regulation of pituitary cell function, *Mol Cell*
702 *Endocrinol*, 438 (2016) 100-106.

703 [93] M. Bhattacharya, A.V. Babwah, Kisspeptin: beyond the brain, *Endocrinology*, 156 (2015) 1218-
704 1227.

705 [94] M. Kirilov, J. Clarkson, X. Liu, J. Roa, P. Campos, R. Porteous, G. Schutz, A.E. Herbison,
706 Dependence of fertility on kisspeptin-Gpr54 signaling at the GnRH neuron, *Nat Commun*, 4 (2013)
707 2492.

708 [95] S. Leon, A. Barroso, M.J. Vazquez, D. Garcia-Galiano, M. Manfredi-Lozano, F. Ruiz-Pino, V. Heras,
709 A. Romero-Ruiz, J. Roa, G. Schutz, M. Kirilov, F. Gaytan, L. Pinilla, M. Tena-Sempere, Direct Actions of
710 Kisspeptins on GnRH Neurons Permit Attainment of Fertility but are Insufficient to Fully Preserve
711 Gonadotropic Axis Activity, *Scientific reports*, 6 (2016) 19206.

712 [96] M.S. Irwig, G.S. Fraley, J.T. Smith, B.V. Acohido, S.M. Popa, M.J. Cunningham, M.L. Gottsch, D.K.
713 Clifton, R.A. Steiner, Kisspeptin activation of gonadotropin releasing hormone neurons and regulation
714 of KiSS-1 mRNA in the male rat, *Neuroendocrinology*, 80 (2004) 264-272.

715 [97] H. Matsui, Y. Takatsu, S. Kumano, H. Matsumoto, T. Ohtaki, Peripheral administration of
716 metastin induces marked gonadotropin release and ovulation in the rat, *Biochem Biophys Res*
717 *Commun*, 320 (2004) 383-388.

718 [98] V.M. Navarro, J.M. Castellano, R. Fernandez-Fernandez, S. Tovar, J. Roa, A. Mayen, M.L. Barreiro,
719 F.F. Casanueva, E. Aguilar, C. Dieguez, L. Pinilla, M. Tena-Sempere, Effects of KiSS-1 peptide, the
720 natural ligand of GPR54, on follicle-stimulating hormone secretion in the rat, *Endocrinology*, 146
721 (2005) 1689-1697.

722 [99] K.L. Hu, H. Zhao, H.M. Chang, Y. Yu, J. Qiao, Kisspeptin/Kisspeptin Receptor System in the Ovary,
723 *Front Endocrinol (Lausanne)*, 8 (2017) 365.

724 [100] M.D. Dorfman, C. Garcia-Rudaz, Z. Alderman, B. Kerr, A. Lomniczi, G.A. Dissen, J.M. Castellano,
725 D. Garcia-Galiano, F. Gaytan, B. Xu, M. Tena-Sempere, S.R. Ojeda, Loss of Ntrk2/Kiss1r signaling in
726 oocytes causes premature ovarian failure, *Endocrinology*, 155 (2014) 3098-3111.

727 [101] M. Calder, Y.M. Chan, R. Raj, M. Pampillo, A. Elbert, M. Noonan, C. Gillio-Meina, C. Caligioni,
728 N.G. Berube, M. Bhattacharya, A.J. Watson, S.B. Seminara, A.V. Babwah, Implantation failure in
729 female Kiss1^{-/-} mice is independent of their hypogonadic state and can be partially rescued by
730 leukemia inhibitory factor, *Endocrinology*, 155 (2014) 3065-3078.

731 [102] X. d'Anglemont de Tassigny, L.A. Fagg, J.P. Dixon, K. Day, H.G. Leitch, A.G. Hendrick, D. Zahn, I.
732 Franceschini, A. Caraty, M.B. Carlton, S.A. Aparicio, W.H. Colledge, Hypogonadotropic hypogonadism
733 in mice lacking a functional Kiss1 gene, *Proc Natl Acad Sci U S A*, 104 (2007) 10714-10719.

734 [103] H. Mei, J. Doran, V. Kyle, S.H. Yeo, W.H. Colledge, Does Kisspeptin Signaling have a Role in the
735 Testes?, *Front Endocrinol (Lausanne)*, 4 (2013) 198.

736 [104] H. Samir, K. Nagaoka, G. Watanabe, Effect of kisspeptin antagonist on goat in vitro Leydig cell
737 steroidogenesis, *Theriogenology*, 121 (2018) 134-140.

738 [105] A.K. Roseweir, R.P. Millar, Kisspeptin Antagonists, in : Kauffman A. S. and Smith J. T., eds
739 *Kisspeptin Signaling in Reproductive Biology*. New York, NY: Springer, 784 (2013) 159-186.

740 [106] C.H.J. Albers-Wolthers, J. de Gier, M. Walen, P.J.S. van Kooten, C.B. Lambalk, P.A.J. Leegwater,
741 B.A.J. Roelen, A.C. Schaefers-Okkens, V. Rutten, R.P.M. Millar, H.S. Kooistra, In vitro and in vivo
742 effects of kisspeptin antagonists p234, p271, p354, and p356 on GPR54 activation, *PLoS One*, 12
743 (2017) e0179156.

744 [107] X. Liu, R. Porteous, X. d'Anglemont de Tassigny, W.H. Colledge, R. Millar, S.L. Petersen, A.E.
745 Herbison, Frequency-dependent recruitment of fast amino acid and slow neuropeptide
746 neurotransmitter release controls gonadotropin-releasing hormone neuron excitability, *The Journal*
747 *of neuroscience*, 31 (2011) 2421-2430.

748 [108] S.A. Clarke, W.S. Dhillon, Kisspeptin across the human lifespan:evidence from animal studies and
749 beyond, *J Endocrinol*, 229 (2016) R83-98.

750 [109] Y. Naniwa, K. Nakatsukasa, S. Setsuda, S. Oishi, N. Fujii, F. Matsuda, Y. Uenoyama, H.
751 Tsukamura, K. Maeda, S. Ohkura, Effects of full-length kisspeptin administration on follicular
752 development in Japanese Black beef cows, *J Reprod Dev*, 59 (2013) 588-594.

753 [110] K.H. Albers-Wolthers, J. de Gier, H.S. Kooistra, V.P. Rutten, P.J. van Kooten, J.J. de Graaf, P.A.
754 Leegwater, R.P. Millar, A.C. Schaefers-Okkens, Identification of a novel kisspeptin with high
755 gonadotrophin stimulatory activity in the dog, *Neuroendocrinology*, 99 (2014) 178-189.

756 [111] H. Kadokawa, M. Matsui, K. Hayashi, N. Matsunaga, C. Kawashima, T. Shimizu, K. Kida, A.
757 Miyamoto, Peripheral administration of kisspeptin-10 increases plasma concentrations of GH as well
758 as LH in prepubertal Holstein heifers, *J Endocrinol*, 196 (2008) 331-334.

759 [112] G.G. Macedo, R.D. Mingoti, E.O.S. Batista, B.M. Monteiro, L.M. Vieira, R.V. Barletta, M.C.
760 Wiltbank, G.P. Nogueira, F.P. Renno, J.R. Maio, P.S. Baruselli, Profile of LH release in response to
761 intramuscular treatment with kisspeptin in *Bos indicus* and *Bos taurus* prepubertal heifers,
762 *Theriogenology*, 125 (2019) 64-70.

763 [113] C.A. Lents, N.L. Heidorn, C.R. Barb, J.J. Ford, Central and peripheral administration of kisspeptin
764 activates gonadotropin but not somatotropin secretion in prepubertal gilts, *Reproduction*, 135 (2008)
765 879-887.

766 [114] M.E. Sebert, D. Lomet, S.B. Said, P. Monget, C. Briant, R.J. Scaramuzzi, A. Caraty, Insights into
767 the mechanism by which kisspeptin stimulates a preovulatory LH surge and ovulation in seasonally
768 acyclic ewes: potential role of estradiol, *Domest Anim Endocrinol*, 38 (2010) 289-298.

769 [115] J.S. Redmond, G.G. Macedo, I.C. Velez, A. Caraty, G.L. Williams, M. Amstalden, Kisspeptin
770 activates the hypothalamic-adenohypophyseal-gonadal axis in prepubertal ewe lambs, *Reproduction*,
771 141 (2011) 541-548.

772 [116] V. Pheng, Y. Uenoyama, T. Homma, Y. Inamoto, K. Takase, K. Yoshizawa-Kumagaye, S. Isaka,
773 T.X. Watanabe, S. Ohkura, J. Tomikawa, K. Maeda, H. Tsukamura, Potencies of centrally- or
774 peripherally-injected full-length kisspeptin or its C-terminal decapeptide on LH release in intact male
775 rats, *J Reprod Dev*, 55 (2009) 378-382.

776 [117] X. d'Anglemont de Tassigny, C.N. Jayasena, K.G. Murphy, W.S. Dhillon, W.H. Colledge,
777 Mechanistic insights into the more potent effect of KP-54 compared to KP-10 in vivo, *PLoS One*, 12
778 (2017) e0176821.

779 [118] C.N. Jayasena, A. Abbara, S. Narayanaswamy, A.N. Comninou, R. Ratnasabapathy, P. Bassett,
780 J.T. Mogford, Z. Malik, J. Calley, M.A. Ghatei, S.R. Bloom, W.S. Dhillon, Direct comparison of the effects
781 of intravenous kisspeptin-10, kisspeptin-54 and GnRH on gonadotrophin secretion in healthy men,
782 *Hum Reprod*, 30 (2015) 1934-1941.

783 [119] C.N. Jayasena, A. Abbara, J.D. Veldhuis, A.N. Comninou, R. Ratnasabapathy, A. De Silva, G.M.
784 Nijher, Z. Ganiyu-Dada, A. Mehta, C. Todd, M.A. Ghatei, S.R. Bloom, W.S. Dhillon, Increasing LH
785 pulsatility in women with hypothalamic amenorrhoea using intravenous infusion of Kisspeptin-54,
786 *The Journal of clinical endocrinology and metabolism*, 99 (2014) E953-961.

787 [120] C.N. Jayasena, G.M. Nijher, A.N. Comninou, A. Abbara, A. Januszewski, M.L. Vaal, L.
788 Sriskandarajah, K.G. Murphy, Z. Farzad, M.A. Ghatei, S.R. Bloom, W.S. Dhillon, The effects of
789 kisspeptin-10 on reproductive hormone release show sexual dimorphism in humans, *The Journal of*
790 *clinical endocrinology and metabolism*, 96 (2011) E1963-1972.

791 [121] C.E.P. Leonardi, F.C.F. Dias, G.P. Adams, J. Singh, Effect of Kisspeptin-10 on plasma luteinizing
792 hormone concentrations and follicular dynamics during the luteal phase in cattle, *Theriogenology*,
793 119 (2018) 268-274.

794 [122] J.T. Smith, S.N. Saleh, I.J. Clarke, Seasonal and cyclical change in the luteinizing hormone
795 response to kisspeptin in the ewe, *Neuroendocrinology*, 90 (2009) 283-291.

796 [123] J.T. Smith, Q. Li, K.S. Yap, M. Shahab, A.K. Roseweir, R.P. Millar, I.J. Clarke, Kisspeptin is
797 essential for the full preovulatory LH surge and stimulates GnRH release from the isolated ovine
798 median eminence, *Endocrinology*, 152 (2011) 1001-1012.

799 [124] M. Beltramo, C. Decourt, Towards new strategies to manage livestock reproduction using
800 kisspeptin analogs, *Theriogenology*, 112 (2018) 2-10.

801 [125] Y. Goto, N. Endo, K. Nagai, S. Ohkura, Y. Wakabayashi, A. Tanaka, H. Matsui, M. Kusaka, H.
802 Okamura, T. Tanaka, Ovarian and hormonal responses to follicular phase administration of
803 investigational metastin/kisspeptin analog, TAK-683, in goats, *Reproduction in domestic animals*, 49
804 (2014) 338-342.

805 [126] N. Endo, C. Tamesaki, S. Ohkura, Y. Wakabayashi, H. Matsui, A. Tanaka, T. Watanabe, H.
806 Okamura, T. Tanaka, Differential changes in luteinizing hormone secretion after administration of the
807 investigational metastin/kisspeptin analog TAK-683 in goats, *Anim Reprod Sci*, 159 (2015) 87-93.

808 [127] L.P. Rahayu, M.E. Behiry, N. Endo, T. Tanaka, Effect of investigational kisspeptin/metastin
809 analog, TAK-683, on luteinizing hormone secretion at different stages of the luteal phase in goats, *J
810 Reprod Dev*, 63 (2017) 221-226.

811 [128] N. Kanai, N. Endo, S. Ohkura, Y. Wakabayashi, H. Matsui, H. Matsumoto, K. Ishikawa, A. Tanaka,
812 T. Watanabe, H. Okamura, T. Tanaka, An administration of TAK-683 at a minimally effective dose for
813 luteinizing hormone stimulation under the absence of the ovary induces luteinizing hormone surge in
814 ovary-intact goats, *J Reprod Dev*, 63 (2017) 305-310.

815 [129] C. Decourt, V. Robert, K. Anger, M. Galibert, J.B. Madinier, X. Liu, H. Dardente, D. Lomet, A.F.
816 Delmas, A. Caraty, A.E. Herbison, G.M. Anderson, V. Aucagne, M. Beltramo, A synthetic kisspeptin
817 analog that triggers ovulation and advances puberty, *Scientific reports*, 6 (2016) 26908.

818 [130] C. Decourt, V. Robert, D. Lomet, K. Anger, M. Georgelin, K. Poissenot, M.T. Pellicer-Rubio, V.
819 Aucagne, M. Beltramo, The kisspeptin analog C6 is a possible alternative to PMSG (pregnant mare
820 serum gonadotropin) for triggering synchronized and fertile ovulations in the Alpine goat, *PLoS One*,
821 14 (2019) e0214424.

822 [131] C. Ralph, R. Kirkwood, M. Beltramo, V. Aucagne, A. Tilbrook, A single injectin of compound 6
823 evoked an LH surge and ovulation in 18 week old gilts, *Proceedings of the Annual Meeting of the
824 Endocrine Society*, (2018) 8.

825 [132] V. Pottapenjara, S.R. Rajanala, C. Reddy, A. Gangineni, K. Avula, S.K. Bejjanki, S. Sathagopam, S.
826 Kesharwani, S. Velmurugan, Kisspeptin Modulates Luteinizing Hormone Release and Ovarian
827 Follicular Dynamics in Pre-pubertal and Adult Murrah Buffaloes, *Front Vet Sci*, 5 (2018) 149.

828 [133] A. Ezzat Ahmed, H. Saito, T. Sawada, T. Yaegashi, T. Yamashita, T. Hirata, K. Sawai, T.
829 Hashizume, Characteristics of the stimulatory effect of kisspeptin-10 on the secretion of luteinizing
830 hormone, follicle-stimulating hormone and growth hormone in prepubertal male and female cattle, *J
831 Reprod Dev*, 55 (2009) 650-654.

832 [134] B.K. Whitlock, J.A. Daniel, R.R. Wilborn, S.P. Rodning, H.S. Maxwell, B.P. Steele, J.L. Sartin,
833 Interaction of estrogen and progesterone on kisspeptin-10-stimulated luteinizing hormone and
834 growth hormone in ovariectomized cows, *Neuroendocrinology*, 88 (2008) 212-215.

835 [135] H. Kadokawa, S. Suzuki, T. Hashizume, Kisspeptin-10 stimulates the secretion of growth
836 hormone and prolactin directly from cultured bovine anterior pituitary cells, *Anim Reprod Sci*, 105
837 (2008) 404-408.

838 [136] P.A. Parker, E.A. Coffman, K.G. Pohler, J.A. Daniel, V. Aucagne, M. Beltramo, B.K. Whitlock,
839 Acute and subacute effects of a synthetic kisspeptin analog, C6, on serum concentrations of
840 luteinizing hormone, follicle stimulating hormone, and testosterone in prepubertal bull calves,
841 *Theriogenology*, 130 (2019) 111-119.

842
843

844 **Figure Caption**


845 **Figure 1.** (A) Kp10 sequences of representative species. In red are indicated the amino acid variants
846 in the sequence. (B) Concentration activity response of human, ovine, and equine Kp10 on a calcium
847 mobilization assay on HEK293 transfected with the human *KISS1R*. Human and ovine Kp10 have very
848 similar potency and efficacy, whereas equine Kp10 is about 10 times less potent. (C & D) Effect of the
849 injection of ovine or equine Kp10 (15 nmol/animal) in the ewe. Blood samples (2 mL) were collected
850 from the jugular vein every 15 minutes from 30 minutes before the injection of the drug to 45
851 minutes after, and then every 30 minutes until experiment end. LH was measured with an RIA
852 methods as previously described [129]. Both molecules increase the LH plasma concentration.
853 However, the maximal amplitude (C) and the total amount (D) of LH released are significantly less
854 after the injection of the equine Kp10. The black arrow in C indicates the time of molecule injection.
855

Figure 1

A


Homo sapiens	YNWNSFGLRF
Pan troglodytes	YNWNSFGLRF
Macaca mulatta	YNWNSFGLRF
Capra hircus	YNWNSFGLRY
Ovis aries	YNWNSFGLRY
Bos taurus	YNWNSFGLRY
Sus scrofa	YNWNSFGLRY
Equus caballus	YRWNSFGLRY
Mus musculus	YNWNSFGLRY
Felis catus	YNWNSFGLRY
Canis lupus familiaris	YNWNVFGLRY

B



	oKP10	hKp10	eKP10
EC50	1.209e-010	1.406e-010	2.746e-009

C



D

