



HAL
open science

Selective breeding for better farmed fish: how to be more efficient ?

Marc Vandeputte

► **To cite this version:**

Marc Vandeputte. Selective breeding for better farmed fish: how to be more efficient ?. AQUAFARM, Jan 2017, Pordenone, Italy. hal-03155356

HAL Id: hal-03155356

<https://hal.inrae.fr/hal-03155356>

Submitted on 1 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




Selective breeding for better farmed fish: how to be more efficient ?


Marc Vandeputte

GABI, INRA, AgroParisTech, Université Paris-Saclay Jouy en Josas, France
Ifremer, Palavas les Flots, France

AQUAFARM congress, Pordenone, Italy, 26 January 2017




Aquaculture: a growing challenge



- The increase in consumption is and will be met by aquaculture
➔ pressing need to increase competitiveness and sustainability

Selective breeding: a major productivity enhancer

➤ Simultaneous comparison of 1957 and 2001 broiler strains



➤ Body weight
810g → 3950g

➤ Carcass Yield
60.8% → 74.4%

➤ Breast Yield
11.4% → 21.3%

All were fed the same feed




Havenstein, 2006, Lohmann Information 41: 30-37

Early fish breeding programs

- With pedigree: Norway salmon (1972)




- Without pedigree: French PROSPER (1986)



- One major goal: faster growth

Increasing growth: it works well!



High potential of fish because:

- Often starts from wild (domestication)
- High selection intensities possible
- Short generations (tropical species)

Aquaculture breeding in Europe (2016)


12 species
54 programs
>31 with pedigree

Sources: Chavanne et al., 2016
Aquacult. Int. 24: 1287-1305





Traits selected



Chavanne et al., 2016, *Aquacult. Int.* 24:1287-1307

FISHBOOST
The next level of aquaculture breeding

Growth remains the #1 trait



But is it growth that we really need to improve?

Review

The importance of selective breeding in aquaculture to meet future demands for animal protein: A review

Trygve Gjedrem ^{a,b}, Nick Robinson ^a, Morten Rye ^{b,*}

^a Nofima, PO Box 5010, N-1432 Ås, Norway

^b AkvaGen Genetics Center, N-6900 Sandnessjøen, Norway

ARTICLE INFO

Article history:
 Received 21 July 2011
 Received in revised form 3 April 2012
 Accepted 9 April 2012
 Available online 16 April 2012

Keywords:
 Aquaculture
 Potential
 Selective breeding
 Animal protein

ABSTRACT

Aquaculture is the fastest growing food production industry, and the vast majority of aquaculture products are derived from Asia. The quantity of aquaculture products directly consumed is now greater than that resulting from conventional fisheries. The nutritional value of aquatic products compares favourably with meat from farm animals because they are rich in micronutrients and contain high levels of healthy omega-3 fatty acids. Compared with farm animals, fish are more efficient converters of energy and protein. If the aquaculture sector continues to expand at its current rate, production will reach 132 million tonnes of fish and shellfish and 43 million tonnes of seaweed in 2020. Future potential for marine aquaculture production can be estimated based on the length of coastline, and for freshwater aquaculture from available land area in different countries. The average marine production in 2005 was 103 tonnes per km coastline, varying from 0 to 1721 (China). Freshwater aquaculture production in 2005 averaged 0.17 tonnes/ha, varying from 0 to close to 6 tonnes per ha (Bangladesh), also indicating potential to dramatically increase freshwater aquaculture output. Simple estimations indicate potential for a 20-fold increase in world aquaculture production. Limits imposed by the availability of feed resources would be lessened by growing more herbivorous species and by using more of genetically improved stocks. Aquaculture generally trails far behind plant and farm animal industries in utilizing selective breeding as a tool to improve the biological efficiency of production. It is estimated that at present less than 10% of aquaculture production is based on genetically improved stocks, despite the fact that annual genetic gains reported for aquatic species are substantially higher than that of farm animals. **With an average genetic gain in growth rate of 12.5% per generation, production may be dramatically increased if genetically improved animals are used.** Importantly, animals selected for faster growth have also been shown to have improved feed conversion and higher survival, implying that increased use of selectively bred stocks leads to better utilization of limited resources such as feed, labour, water, and available land and sea areas.

© 2012 Elsevier B.V. All rights reserved.


Contents

1. Introduction	118
2. Status of aquaculture production	118
3. Fish and shellfish for human consumption: comparison with meat from farm animals	119
4. Efficiency of fish production	119
5. Potential of aquaculture production	119
5.1. Growth in aquaculture production	119

With an average genetic gain in growth rate of 12.5% per generation, production may be dramatically increased if genetically improved animals are used





The problem...



The solution?



Economic value of TGC et FCR



500t/yr, 1.35€/kg

TGC


- 8.33
- 8.93
- 9.53
- 10.11
- 11.94

- Variable traits
 - FCR: 0.81 ($\Delta=0.06/\text{generation}$)
 - TGC: 8.33 ($\Delta=+0.6/\text{generation}$)
- Limiting factors
 - Density 230 kg/m³
 - N-NH₃ : 40 kg/day

- FCR is always profitable
- TGC is profitable only if density is the limiting factor
- Extension to other systems show similar results

Besson et al. 2014, J Anim. Sci. 92:5394-5405.

Changing perspective: towards efficiency



How can we improve those traits by selective breeding ?
 Will faster growing fish also be better for efficiency ?

Fillet yield: indirect selection

➤ Carp

- Head size



+fillet **-fillet** $r_A = -0.86$

Selection for smaller head:
+1.4% fillet/gen. (avg 31,2%)




Kocour et al., 2007. Aquaculture 270: 43-50



ongoing work:




➤ Trout




Haffray et al., 2012. Aquaculture 368: 145-152

Selection for disease resistance



Selection of survivors

Sib selection



Indirect (QTL / genomic)





Fig. 1. Survival by family in a WSSV challenge study of CAMACO stocks.

2-18% gain/generation in multitrait
Gjedrem, 2015, JMSE 3: 146-153:



Moen et al, 2015, Genetics


Cuellar-Angel et al, 2012 Aquaculture 368: 36-39
IPN: Okamoto et al 1993 Aquaculture 117 71-76:

Selecting for feed efficiency ?


- **Highly profitable but not selected for**
- **Individual feed intake on long periods not measurable**
 - ➔ **candidate FE not measurable**
- **Indirect criteria:**
 - **FE of individually housed fish**
 - **X-ray point FI measurements**
 - **Weight loss at fasting**
 - **Growth ??**



Weight loss at fasting



Grima et al, 2010a. *Aquaculture* 300: 50-58




Grima et al, 2010b. *Aquaculture* 302: 169-174

Validation by experimental selection in sea bass ongoing




Does selection for faster growth imply better FE ?

Atlantic salmon




Thodesen et al 1999
 Aquaculture 180: 237-246

Japanese flounder




Ogata et al 2002
 Aquaculture 211: 183-193

Brown trout



Sanchez et al 2004
 J.Anim.Sci. 82: 2865-2875



Nile tilapia in RAS



Hans Komen, ABGC, WUR

➔ Growth may contribute but we need more data on FE genetics

Selecting « vegetarian » rainbow trout





V-gain=
 $(S-T)/T$

M-gain=
 $(S-T)/T$

Le Boucher et al., 2012, PLoS ONE 7: e44898


Efficiency also reduces environmental impact



Besson et al., 2016, J. Cleaner Prod. 116: 100-109

What are the limits to selection for efficiency ?

- Ever faster growing fish is (probably) possible
- But an immortal fillet block growing without feed will not be possible...
- What are the limits ?
- Will efficient fish be less robust ?



Nestor et al. 1996 Poultry Sci 75: 1180-1191
28 gen: 400g → 5500g (♂ 16 wks BW)



Conclusion

- **Efficiency is both economically and environmentally beneficial**
- **Major traits: feed efficiency, processing yields, disease resistance, adaptation to alternative diets**
- **Some (little ?) progress may be expected by correlated response to selection for growth**
- **Specific research needed on phenotyping, genetic and genomic bases of efficiency traits**

- **A major key for sustained growth of aquaculture production in a finite world**