

Selective breeding to reduce the ecological footprint of aquaculture

Marc Vandeputte, Mathieu Besson, François Allal, Florence Phocas, Edwige

Quillet

▶ To cite this version:

Marc Vandeputte, Mathieu Besson, François Allal, Florence Phocas, Edwige Quillet. Selective breeding to reduce the ecological footprint of aquaculture. Aquaculture Europe 2017, Oct 2017, Dubrovnik, Croatia. . hal-03156701

HAL Id: hal-03156701 https://hal.inrae.fr/hal-03156701

Submitted on 2 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SELECTIVE BREEDING TO REDUCE THE ECOLOGICAL FOOTPRINT OF AQUACULTURE



M. Vandeputte, M. Besson, F. Allal, F. Phocas, E. Quillet



The big question

Fish farming has to grow to meet increased demand for fish

Selective breeding is an efficient method that has mostly been used

to increase production in quantity (growth rate)

How to use this potential to improve production efficiency and decrease global and local environmental impact ?

Our vision



Producing 1 kg of sea bass fillet requires 5.8 kg of compound feed All efficiency traits have a genetic basis that can be selected Could sea bass reach the efficiency of rainbow trout ?

1.9 kg 🗘 1.7 kg 🦾 1.6 kg 🏳 1.0 kg Trout

5.8 kg 🗇 2.9 kg 🦾 2.6 kg 🖓 1.0 kg 🖉 Sea bass

Challenges to be met

Develop efficient phenotyping methods and/or indirect predictors of efficiency traits

Evaluate how genomic selection can help use less (more precise) phenotypes

Ultimately, co-adapt fish strains and novel, agro-ecological production systems





This research has received funding from the European Union's FP7 and Horizon 2020 research and innovation programmes. This output reflects the views only of the author(s), and the European Union cannot be held responsible fpor any use which may be made of the information contained therein.

