



HAL
open science

Poplar Propolis Ethanolic Extract Reduces Body Weight Gain and Glucose Metabolism Disruption in High-Fat Diet-Fed Mice

Nicolas Cardinault, Franck Tourniaire, Julien Astier, Charlène Couturier, Estelle Perrin, Julie Dalifard, Eva Seipelt, Lourdes Mounien, Claire Letullier, Lauriane Bonnet, et al.

► To cite this version:

Nicolas Cardinault, Franck Tourniaire, Julien Astier, Charlène Couturier, Estelle Perrin, et al.. Poplar Propolis Ethanolic Extract Reduces Body Weight Gain and Glucose Metabolism Disruption in High-Fat Diet-Fed Mice. *Molecular Nutrition and Food Research*, 2020, 64 (18), pp.2000275. 10.1002/mnfr.202000275 . hal-03157023

HAL Id: hal-03157023

<https://hal.inrae.fr/hal-03157023v1>

Submitted on 14 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Poplar propolis ethanolic extract reduces body weight gain and glucose metabolism**
2 **disruption in high-fat diet-fed mice.**

3

4 Nicolas Cardinault^{3*}, Franck Tourniaire^{1,2*}, Julien Astier¹, Charlène Couturier¹, Estelle Perrin¹,
5 Julie Dalifard¹, Eva Seipelt¹, Lourdes Mounien¹, Claire Letullier³, Lauriane Bonnet¹, Esma
6 Karkeni¹, Naïma Delbah⁴, Stéphane Georgé⁴, Jean-François Landrier^{1,2}

7

8 ¹ Aix-Marseille Université, C2VN, INRA, INSERM, 13000, Marseille, France

9 ² CriBioM, Criblage Biologique Marseille, Faculté de médecine de la Timone, Marseille, France.

10 ³ Pollenergie, La Grabère, 47450 Saint Hilaire de Lusignan, France.

11 ⁴ CTCPA, 449 Avenue Clément Ader, 84911 Avignon.

12 * These authors contributed equally to this work.

13

14 Running title: Propolis improves obesity and associated disorders

15

16 *Corresponding author and person to whom reprint requests should be addressed:*

17 Jean-François Landrier, C2VN ; UMR 1260 INRA/1263 INSERM/Université d'Aix-Marseille,
18 27 Bd Jean Moulin, 13385 Marseille cedex 05, France. Phone: +33 4 91 29 42 75; E-mail: [jean-](mailto:jean-francois.landrier@univ-amu.fr)
19 [francois.landrier@univ-amu.fr](mailto:jean-francois.landrier@univ-amu.fr)

20

21 Keywords : adipose tissue, insulin resistance, NRF2, obesity, propolis.

22

23 **Abstract**

24 **Scope:** Current evidence supports the beneficial effect of polyphenols on the management of
25 obesity and associated comorbidities. This is the case for propolis, a polyphenol-rich substance
26 produced by bees. The aim of the present study was to evaluate the effect of a poplar propolis
27 ethanolic extract (PPEE) on obesity and glucose homeostasis, and to unveil its putative
28 molecular mechanisms of action.

29 **Methods and results:** Male high-fat (HF) diet-fed mice were administered PPEE for 12 weeks.
30 PPEE supplementation reduced the HF-mediated adiposity index, adipocyte hypertrophy and
31 body weight gain. It also improved HOMA-IR and fasting glucose levels. Gene expression
32 profiling of adipose tissue (AT) showed an induction of mRNA related to lipid catabolism and
33 mitochondrial biogenesis and inhibition of mRNA coding for inflammatory markers.
34 Interestingly, several Nrf2-target genes were induced in AT following administration of PPEE.
35 The ability of PPEE to induce the expression of Nrf2-target genes was studied in adipocytes.
36 PPEE was found to transactivate the Nrf2 response element and the Nrf2 DNA-binding,
37 suggesting that part of the effect of PPEE could be mediated by Nrf2.

38 **Conclusion:** PPEE supplementation may represent an interesting preventive strategy to tackle
39 the onset of obesity and associated metabolic disorders.

40

41

42

43 **Introduction**

44

45 Obesity is a major metabolic risk factor for type 2 diabetes mellitus (T2DM) and a public health
46 problem throughout the world. In T2DM, insulin signaling is progressively lost, leading to an
47 increase in pancreatic insulin production and pancreas exhaustion at the latest stages of the
48 disease [1]. T2DM represents 90-95% of diabetes mellitus cases worldwide and s a multifactorial
49 disease in which obesity plays a central role [2]. Indeed, obesity generates low-grade
50 inflammation, which is related to the activation of inflammatory pathways (JNK and NF- κ B) in
51 adipose tissue (AT) [3] and increased production of cytokines such as interleukin-6 (IL-6),
52 interleukin-1beta (IL-1 β), tumor necrosis factor alpha (TNF- α), and chemokines [4] (e.g.
53 monocyte chemotactic protein 1 (MCP-1)) and other mediators of the immune response [3],
54 including microRNA [5-7]. When AT lipid storage capacities are exhausted, ectopic fat depots
55 develop in other tissues, including insulin sensitive tissues (e.g. liver, skeletal muscle). Such
56 ectopic fat accumulation, together with inflammation, interferes with insulin signaling and is
57 central in the development and persistence of insulin resistance[8], observed in T2DM. So far,
58 drug-based therapeutic approaches to tackle obesity and associated disorders remain limited and
59 have either deleterious side effects or poor efficiency. Therefore, alternative “natural” treatments
60 are urgently needed.

61

62 Propolis is a complex natural resinous substance from buds and exudates of selected plant
63 sources, collected by bees, mixed with wax, and transformed in the presence of bee enzymes [9].

64 Propolis has been used as a folk medicine in many countries since ancient times and has been
65 reported to possess various pharmacological effects such as immunomodulatory, antitumoral,
66 antimicrobial, antiviral, antifungal, anti-inflammatory and antioxidant properties [10]. These

67 effects are mostly attributed to the nature and amount of phenolic compounds contained in
68 propolis. The chemical composition of propolis is highly dependent upon its plant source(s) and
69 varies with the site of collection [9].

70 Interestingly, several clinical trials have been performed to test the effect of propolis
71 supplementation in T2DM patients [11]. These recently reviewed and meta-analyzed studies
72 suggest that propolis supplementation decreases fasting blood glucose and Hb1Ac levels [11]and
73 indicate that propolis is able to delay or slow down the progression of the disease, and acts by
74 decreasing oxidative stress and inflammatory processes associated with T2DM.

75

76 Preclinical studies have been performed in animal models of obesity and/or diabetes, to
77 investigate the phenotypical impact of propolis supplementation in the diet [12-17]. The
78 molecular mechanisms underlying the observed effects in terms of obesity management and
79 improvement of glucose homeostasis, however, still remain unclear. Thus, the aim of the present
80 study was to evaluate the effect of a standardized poplar propolis ethanolic extract (PPEE) on
81 obesity glucose homeostasis and the physiology of the adipose tissue in high-fat diet-fed mice.
82 The putative molecular mechanisms involved in these effects was also investigated.

83

84 **Materials and methods**

85

86 **Propolis source and extraction**

87 Poplar propolis was collected in the southern regions of France using dietary propolis traps
88 according to special specifications to minimize contamination. It was stored at -20 °C prior to
89 extraction. The latter was performed according to a previously published method [18]. Briefly,
90 the propolis sample was cut into small pieces and soaked into 75% ethanol for 8 days. Following
91 filtration, the filtrate was concentrated under reduced pressure using a rotary evaporator, to
92 afford a poplar propolis ethanolic extract (PPEE). The dry residue was ground to a fine powder
93 by adding food excipients and kindly provided by Pollenergie (Agen, France). The PPEE powder
94 used in this study fully complies with European regulations on food supplements and those more
95 specific of propolis.

96

97 **Total polyphenol content** -The total polyphenol content was measured according to the method
98 described by Popova *et al.*. The results were expressed in mg/100 g of propolis powder as
99 pinocembrin/galangin equivalent for poplar propolis. Methanolic solutions of
100 pinocembrin/galangin (2/1) were used for calibration purposes [19]. The phenolic profile was
101 determined by HPLC as previously reported. [20]. Briefly, a sample of 1 g of PPEE powder was
102 dissolved in 30 mL of ethanol (70%) in a 50 mL Falcon tube then heated for 2h at 70 °C.
103 Following centrifugation, the alcoholic phase was recovered and the extraction was repeated
104 twice. The three extracts were combined and adjusted to 100 mL with 70% ethanol in a
105 volumetric flask. PP solution was finally diluted 1 to 50 times in methanol / water (1:1).
106 Separation was achieved by HPLC using a Symmetry (250 x 4.6 mm, C₁₈) column (Waters, Saint

107 Quentin en Yvelines, France). A linear gradient of acetonitrile and 0.2% aqueous formic acid
108 (v/v) was used as the mobile phase with a flow rate of 1.2 mL/min and DAD data were acquired
109 in the 200–450 nm range. Polyphenolic compounds were identified by comparing their retention
110 time and UV characteristics to authentic standards. The total polyphenol content and the detailed
111 composition of polyphenols present in PPEE are reported on Table 1.

112

113 **Animal Experiments** - Six-week-old male C57BL/6J mice were obtained from Janvier Labs (Le
114 Genest Saint Isle, France) and fed *ad libitum* with control food (chow diet A04, Safe-diets,
115 France) and with full access to drinking water for a 1-week acclimatization period. The animals
116 were maintained at 22 °C under a 12-h light/dark cycle and a 20 % relative humidity level. The
117 mice (3 or 4 per cage) were randomly assigned (10 per group) into one of three experimental
118 groups depending on their diet, i.e. standard (chow diet A04 from Safe diets), high fat (HF: 45%
119 energy from lipids, Test Diet ref. 58V8) or HF-supplemented with PPEE (20 mg/mouse/day,
120 corresponding to 4.5 mg of total polyphenols/mouse/day). The pellets were replaced every 3
121 days. The weight gain was measured once a week and dietary intake was recorded every two
122 weeks. After 12 weeks, the mice were fasted overnight and blood was collected by cardiac
123 puncture using a general anesthetic (sevoflurane, Baxter, France). The plasma was isolated by
124 centrifugation at 3000 rpm for 15 min at 4 °C and was stored at -80 °C until further analysis.
125 Animals were euthanized by cervical dislocation under general anesthesia, and their tissues
126 (liver and various white AT deposits) were collected, weighted and stored at -80 °C until further
127 analysis. This protocol was approved by the local Ethics committee and the French Minister of
128 Research and Education (n° 01549.04).

129

130 **Dosage information** – The PPEE has been mixed in the food and delivered to mice at the dosage
131 of 20 mg / mouse / day, which corresponding to 4.5 mg of total polyphenols / mouse / day, for 12
132 weeks. This dosage was chosen to be relevant in human, as it correspond to a dose of 12 mg of
133 total polyphenol / kg of body weight, taking into account the conversion factor for dose
134 translation between mice and human [21].

135

136 **Oral glucose and insulin tolerance tests** - Glucose and insulin tolerance tests were performed
137 two weeks before animals were euthanized. For the glucose tolerance test, the mice were fasted
138 for 16 h and then gavaged with glucose (2 g/kg). For the insulin tolerance test, the mice were
139 fasted for 5 h and then injected *i.p.* with insulin (0.5 U/kg). Blood samples were taken from the
140 tail tips at specific time intervals to measure glucose levels (Accu-Check glucometer, Roche).

141

142 **Biochemical analyses** – The concentration of glucose in plasma was evaluated using glucose
143 RTU (bioMerieux, Craponne, France). Triglycerides and free fatty acids (FFA) were measured
144 using a colorimetric test (RANDOX, Cruclin, Co. Antrim, United Kingdom). Insulin was
145 measured using an enzyme-linked immuno-sorbent assay ELISA (ALPCO Diagnostics, New
146 Hampshire, United States). Leptin and adiponectin were quantified by ELISA (R&D Systems
147 quantikine ELISA). β -hydroxybutyrate concentration was measured using a colorimetric test
148 according to the manufacturer's procedure (BEN srl, Milano, Italy). The HOMA-IR index was
149 calculated according to the following formula: fasting insulin (microU/L) x fasting glucose
150 (nmol/L)/22.5.

151

152 **Adipose tissue histology** - Epididymal AT samples were fixed in 10% buffered formalin,
153 embedded in paraffin and sliced to prepare 5 μm tissue sections that were stained with
154 hematoxylin and eosin (H&E) as previously reported [22]. The images were captured by a light
155 microscope (EZAD, Leica, Germany). The adipocyte area (μm^2) were determined using the
156 Image J software.

157
158 **Cell culture** - 3T3-L1 preadipocytes (ATCC, Manassas, VA) were seeded in 3.5-cm diameter
159 dishes at a density of 15×10^4 cells/well. Cells were grown in DMEM supplemented with 10%
160 FBS at 37 °C in a 5% CO₂ humidified atmosphere as previously reported [23]. To induce
161 differentiation, 2-day post-confluent 3T3-L1 preadipocytes (day 0) were stimulated for 48 h with
162 0.5 mM isobutylmethylxanthine, 0.25 μM dexamethasone, and 1 $\mu\text{g}/\text{mL}$ insulin in DMEM
163 supplemented with 10 % FBS. The cells were maintained in DMEM supplemented with 10%
164 FBS and 1 $\mu\text{g}/\text{mL}$ insulin. To examine the effect of PPEE, 3T3-L1 adipocytes received either
165 PPEE (3 - 30 $\mu\text{g}/\text{mL}$) or ethanol only (0.1%). All treatments were performed on day eight. The
166 data obtained are the mean of three independent experiments, each performed in triplicate.

167
168 **RNA interference** - The 3T3-L1 differentiated cells were seeded in 6-well plates and transfected
169 with either targeted siRNA (against Nrf2) or a non-targeting siRNA according to the
170 manufacturer's instructions (Dharmacon, Inc. Lafayette, CO) and as previously reported [24, 25].
171 Briefly, the cells were transfected overnight using a mixture of 100 nM siRNA and 2 μL of
172 DharmaFECT reagent per well. The media were subsequently replaced with fresh media with or
173 without PPEE and the plates were re-incubated for 24 h.

174

175 **Reporter assays** - The reporter plasmid containing the gene for firefly luciferase under the
176 control of the wild-type or mutated antioxidant response element (ARE) [26] was transfected
177 into 3T3-L1 preadipocyte cells, together with the pGL4 plasmid (1 µg) coding for the renilla
178 luciferase. The transfection was performed using Lipofectamine LTX (0.9 µL) (Thermo, Saint-
179 Aubin, France) and the Plus Reagent (1 µL) as previously reported [27]. After overnight
180 incubation the medium was replaced by DMEM supplemented with 10% FBS and PPEE (30
181 µg/mL) . After 24 h, the cells were lyzed and assayed for luciferase activity using a Dual-Glo
182 luciferase assay system (Promega, Madison, WI) with firefly luciferase activity normalized to
183 renilla luciferase. The transfection experiments were performed in triplicate and repeated three
184 times independently.

185

186 **Nfr2 transAM assay** - 3T3-L1 cells were incubated for 6 h with or without PPEE (30 µg/mL).
187 Nuclei were prepared as recommended and used in the TransAM Nrf2 assay according to the
188 manufacturer's protocol (Active Motif, La Hulpe, Belgium)

189

190 **RNA extraction and real-time PCR** - Total RNA was extracted from the liver, kidney and
191 epididymal AT or from cells using TRIzol reagent (Thermo, Courtaboeuf, France). Total RNA (1
192 µg) was used to synthesize cDNAs using random primers and Moloney murine leukemia virus
193 reverse transcriptase (Thermo, Courtaboeuf, France). Real-time quantitative PCR analyses were
194 performed using the Mx3005P Real-Time PCR System (Stratagene, La Jolla, USA) as previously
195 described [28]. For each condition, the expression was quantified in duplicate, and 18S rRNA
196 was used as the endogenous control in the comparative cycle threshold (CT) method [29].

197

198 **Statistical analysis** - All data are expressed as the mean \pm SEM. Significant differences between
199 the control and treated groups were determined using ANOVA, followed by the Tukey-Kramer
200 post hoc test using the Statview software (SAS Institute, Cary, NC). $P < 0.05$ was considered
201 statistically significant.

202

203 **Results**

204

205 *Poplar propolis ethanolic extract supplementation protects against diet-induced obesity*

206 The results showed that PPEE supplementation (20 mg/mouse/day, equivalent to 4.5 mg of total
207 polyphenols/mouse/day) significantly limited weight gain ($p<0.0001$) caused by a HF diet (Fig
208 1A and 1B). The energy intake was higher in the HF diet fed group compared to the control
209 group, but no significant difference ($p<0.002$) was observed in the PPEE supplemented group
210 compared to the HF diet fed group (Fig 1C). The limitation of body weight gain in PPEE
211 supplemented mice was associated with a reduction of absolute and relative fat mass in
212 epididymal, retroperitoneal or inguinal fat pads (Fig 1D and 1E, $p<0.05$). Consequently, the
213 adiposity index (sum of epididymal, inguinal and retroperitoneal AT mass relative to total body
214 mass) in PPEE supplemented mice was reduced, compared to HF diet fed mice (Fig 1F,
215 $p<0.001$). Interestingly no significant difference of adiposity index was observed between PPEE
216 supplemented mice and control fed mice after 12 weeks of supplementation (Fig 1F).
217 Triglycerides, free fatty acids and alanine transaminase (ALAT) were neither affected by the HF
218 diet nor the EEPP supplementation ($p>0.05$). Adiponectin was not affected by PPEE
219 supplementation compared to the HF diet fed group, but was significantly reduced compared to
220 the control group ($p<0.05$). The levels of leptin and β -hydroxybutyrate were increased in the HF
221 diet fed group compared to the control group and were reduced in the PPEE supplemented group
222 ($p<0.05$) (Table 2).

223

224 *Poplar propolis ethanolic extract supplementation improves glucose homeostasis*

225 The effect of PPEE supplementation on glucose homeostasis was evaluated using the oral
226 glucose tolerance test (OGTT) and the insulin tolerance test (ITT). PPEE supplementation
227 improved ($p<0.001$), glucose tolerance in OGTT (Fig 2A) as highlighted by area under the curve
228 (AUC) of the glycemic response (Fig 2B) which was reduced compared to what is observed in
229 the group of HF fed animals. Additionally, PPEE supplementation reduced blood glucose levels,
230 compared to HF fed mice (Fig 2C and 2D). In fasted conditions, insulinemia and glycaemia that
231 were induced by the HF diet, were normalized in PPEE supplemented mice and were not
232 significantly different from control mice (Fig 2E, $p<0.05$). Similarly, the HOMA-IR index was
233 induced ($p<0.005$) by the HF diet and corrected by PPEE supplementation (Fig 2F,

234

235 *Poplar propolis ethanolic extract supplementation changes adipocyte cellularity and modulates*
236 *gene expression in adipose tissue*

237 To evaluate the effect of PPEE supplementation on the phenotype of AT, epididymal AT was
238 used for histological analysis. A HF diet promoted adipocyte hypertrophy with a significant
239 increase in the adipocyte area compared to the control group. Interestingly, this effect was
240 significantly prevented ($p<0.05$) by PPEE supplementation (Fig 3A). Furthermore, the effect of
241 PPEE on inflammation in AT was evaluated by measuring the expression of different markers of
242 inflammation including cytokines (Tnfa) and chemokines (Chemokine C-C motif ligand 5 (Ccl5)
243 and Ccl2). As expected, the HF diet induced a strong increase in the mRNA level of these
244 inflammatory markers compared to the control group., PPEE supplementation, on the other hand,
245 caused a significant decrease ($p<0.05$) of those levels (Fig 3B).

246 To understand the modification of cellularity and fat mass increase, the expression of genes
247 encoding for transcription factors involved in energy metabolism, fatty acid oxidation and

248 mitochondrial biogenesis were studied. PPEE supplementation stimulated ($p<0.05$) the
249 expression of genes promoting fatty acid oxidation, such as the peroxisome proliferator activated
250 receptor α (Ppara), the carnitine palmitoyl-transferase 1a (Cpt1a), the medium- and long-chain
251 acyl-CoA dehydrogenases (Mcad and Lcad), and adipose triglyceride lipase (Atgl), compared to
252 HF diet fed mice (Fig 3C). PPEE supplementation also induced ($p<0.05$) genes related to white
253 AT browning, including genes encoding for the uncoupling protein 1 (Ucp1), the transcription
254 factors peroxisome proliferator-activated receptor gamma coactivator 1-alpha (Ppargc1a), PR
255 domain containing 16 (Prdm16) and cell death-inducing DNA fragmentation factor α -like
256 effector A (Cidea) (Fig 3D) as well as the mitochondrial DNA relative content (Fig 3E).

257

258 *Poplar propolis ethanolic extract supplementation induces the Nrf2 pathway in adipose tissue*
259 *and adipocytes*

260 PPEE contains several polyphenols considered as ligands of the natural nuclear factor (erythroid-
261 derived 2)-like 2 (Nrf2) [30], and Nrf2 agonists are known to induce a similar metabolic
262 phenotype [31] the activation of this signaling pathway was investigated. PPEE supplementation
263 induced the gene expression of the glutamate-cysteine ligase catalytic subunit (Gclc), glutamate-
264 cysteine ligase modifier subunit (Gclm), NADPH: quinone oxidoreductase 1 (Nqo1) and heme
265 oxygenase 1 (Hmox1) in the adipose tissue of mice ($p<0.05$) (Fig 4A). No effect on Nrf2 gene
266 expression was observed. PPEE (30 $\mu\text{g/mL}$) significantly ($p<0.05$) induced the expression of
267 Nrf2, Gclc, Gclm and Nqo1 in 3T3-L1 cells (Fig 4B). The ability of PPEE to transactivate a Nrf2
268 response element (ARE) was confirmed using transient transfection. PPEE increased luciferase
269 activity in 3T3-L1 cells transfected with the wildtype ARE (ARE wt), whereas no induction was
270 observed with the mutated ARE (ARE mut) (Fig 4C, $p<0.05$). Furthermore, the increased ability

271 of Nrf2 to bind to its DNA binding domain in the presence of PPEE was confirmed in 3T3-L1
272 cells (TransAM Nrf2 assay; Fig 4D, $p<0.05$). The involvement of Nrf2 in the PPEE-mediated
273 induction of Nrf2, Gclc, Gclm and Nqo1 was confirmed by RNA interference. This PPEE-
274 mediated induction was blunted or abolished in 3T3-L1 cells transfected with siRNA against Nrf2
275 compared to control siRNA (Fig. 4E, 4F, 4G and 4H, $p<0.05$).

276

277

278 **Discussion**

279

280 In the present study, we reported a beneficial role of PPEE supplementation on weight gain
281 limitation and glucose homeostasis improvement in mice fed a HF diet. In addition, we provided
282 evidences of a direct effect of poplar propolis supplementation on AT biology, strongly
283 suspected to be related to Nfr2 signaling activation in adipocytes.

284 We observed that PPEE supplementation reduced weight gain in mice fed a HF diet. This seems
285 to be mainly related to the limitation of fat mass accumulation as highlighted by a reduced
286 adiposity index. It is noteworthy that these morphological changes were not caused by a reduced
287 energy intake. Interestingly, both subcutaneous (inguinal) and visceral (epididymal and
288 retroperitoneal) adipose pads masses were reduced by PPEE supplementation. The plasma levels
289 of leptin, which is known to be correlated to fat mass, s [32] also decreased. Similar results have
290 been previously observed with Brazilian propolis supplementation, which reduced adiposity and
291 limited the development of perirenal and mesenteric fat depots in high fat-fed rats. It was
292 postulated that the anti-obesity effect of propolis correlated to PPAR γ inhibition in AT, and
293 reduced triglycerides absorption (after a fat loading test [12]). In mice subjected to a high fat diet,
294 Brazilian propolis also limited body weight gain and visceral AT mass [14], or both epididymal
295 and subcutaneous fat masses. No modification of energy expenditure was observed under
296 propolis supplementation, but an increase of fecal lipids was observed, suggesting that Brazilian
297 propolis reduced intestinal fat absorption [13]. The molecular mechanism(s) underlying the effect
298 of propolis on weight gain limitation remains elusive.

299 We studied the cellularity of adipocytes and found that PPEE supplementation could normalize
300 the mean adipocyte area compared to control mice. To explain this effect, we quantified gene

301 expression linked to lipid metabolism. We observed that PPEE supplementation induced the
302 expression of genes coding for proteins involved in fatty acid oxidative metabolism (Ppara α ,
303 Cpt1, Mcad, Lcad). This suggests that PPEE drove a global fatty acid oxidative program.
304 Previous studies have reported the induction of PPAR α , a master regulator of fatty acid
305 oxidation, in the liver of rats supplemented with Brazilian propolis [12]. We also observed a
306 strong induction of genes linked to mitochondria biogenesis and function, such as Ucp1,
307 Ppargc1 α , Prdm16 and Cidea, and an increase in DNA mitochondrial content with PPEE
308 supplementation. The stimulation of mitochondrial biogenesis and fatty acid oxidative capacity
309 in white adipocytes, when linked to increased energy expenditure in such cells through increased
310 energy uncoupling, *via* Ucp1 expression and/or waste (e.g., futile cycles), is a potential novel
311 target for the control of obesity and related comorbidities [33], and could explain the PPEE-
312 induced anti-obesity effect. We, and others, have already proposed a similar mechanism to
313 explain the anti-obesity effect of other molecules,(e.g. *trans*-retinoic acid) [34].

314 In this study, we also observed a beneficial effect of PPEE on glucose homeostasis. Indeed,
315 PPEE supplementation in our diet induced obesity (DIO) mice model improved fasted glycemia
316 and insulinemia and the HOMA-IR index. An the glucose tolerance and sensitivity to insulin
317 were also improved, but were probably overestimated since doses were not adjusted to total
318 mass [35]. Such observations agree with a recent meta-analysis reporting that propolis
319 administrated to T2DM patients decreased fasting blood glucose and Hb1Ac levels[11], and with
320 several preclinical studies on the effect of propolis on glucose homeostasis. These studies
321 showed that propolis partly restored glucose tolerance and insulin sensitivity in ob/ob mice [15].
322 The anti-diabetic effect of propolis has also been described in streptozotocin /high fat diet fed

323 rodents [16, 17, 36], where it was found to stop systemic and pancreatic β -cells inflammation
324 and oxidative stress [17, 37].

325 AT plays a major role in the insulin resistance genesis in the context of obesity, notably through
326 the establishment of a low-grade inflammatory status [3]. The effect of propolis on glucose
327 homeostasis could be due, at least in part, to the impact of PPEE on the inflammation present in
328 AT. When investigating the expression of genes linked to inflammation, we observed that a HF
329 fed diet induced pro-inflammatory cytokines. This was reversed by PPEE supplementation,
330 suggesting that PPEE reduced the inflammatory process in AT. It is presently unclear if such
331 effect is a cell-autonomous effect in adipocytes or if it is related to the reduction of fat mass
332 accumulation, which is associated with an indirect anti-inflammatory effect. This will require
333 further investigations.

334 To further identify the molecular mechanism of action of PPEE we focused on Nrf2. Several
335 compounds found in propolis have been demonstrated to induce the Nrf2 signaling pathway [30],
336 including CAPE [38]. We observed that a large set of Nrf2-target genes were induced in white
337 AT following PPEE supplementation, suggesting that it mediated a strong antioxidant effect in
338 white AT. In addition, we demonstrated a cell-autonomous effect of PPEE supplementation in
339 adipocytes in our in vitro studies. We stimulated the induction of the canonical Nrf2 target genes
340 and demonstrated the specific involvement of Nrf2 using RNA interference. We demonstrated
341 that PPEE transactivated a wild type ARE but not a mutated ARE, and that PPEE induced Nrf2
342 DNA-binding. This demonstrated that PPEE was able to activate the Nrf2 signaling pathway in
343 adipocytes.

344 Nrf2 belongs to the Cap'n collar type transcription factor family (for review [39-41]). In the
345 inactive state, Nrf2 is bound to the protein Keap 1, itself associated with the cytoskeleton, which

346 prevents Nrf2 accumulation in the nucleus. The dissociation of Nrf2 and Keap 1 enables nuclear
347 translocation of Nrf2, which binds to ARE after heterodimerisation with the transcription factor
348 Maf. This type of response element is present in many genes encoding antioxidant enzymes:
349 Nqo1, Gclc, Gclm, Hmox-1 and phase II enzymes. In general, molecules capable of inducing
350 Nrf2 will induce the antioxidant system and detoxification system for xenobiotics and other
351 potentially deleterious molecules. The role of Nrf2 in the control of AT metabolism and
352 adipogenesis has also been extensively studied but remains far from clear [42, 43]. Mice lacking
353 Nrf2 have been found to display either a lean phenotype [44, 45] or no protection against obesity
354 [46, 47]. Nevertheless, if the role of Nrf2 invalidation remains complex, the effect of Nrf2
355 agonists are clearer and more consensual. Treatment of mice with CDDO-IM fed a HF diet
356 prevented body weight gain possibly through a Nrf2-dependant modulation of hepatic lipid
357 metabolism [48]. Similarly, oltipraz, a well-known Nrf2 inducer [49], as well as glycyrrhizin
358 [50]., blunted the HF-mediated obesity and insulin resistance in mice. Recently, glucoraphanin
359 administration was also found to prevent obesity, insulin resistance and NAFLD in HF fed mice
360 via the Nrf2 pathway [31]. The mechanisms by which Nrf2 activators restore energy homeostasis
361 in DIO mice models remains elusive. Such beneficial effect of Nrf2 inducers could be related to
362 an enhancement of mitochondrial biogenesis and function as reported in several cells types
363 including cardiomyocytes [51] and hippocampus neurons [52]. In addition, Nrf2 activation is
364 well-known to reduce oxidative stress and inflammation, which interfere with energy
365 homeostasis and insulin sensitivity [53, 54]. Nrf2 activation could also improve metabolic
366 disorders induced by a HF diet. Finally, the effect of Nrf2 activation could be related to the
367 induction of lipid catabolism by PPEE in AT, since Nrf2 signaling has already been associated to
368 an induction of lipid oxidation in the liver [55]. All these metabolic pathways are strongly

369 interconnected and involved in mediating the Nrf2 effects elicited by PPEE. Further work is
370 required to elucidate their relative contribution.

371

372 In conclusion, our results indicate that EEPP protects against HF-mediated obesity by limiting
373 weight gain and regulating glucose homeostasis This is likely to be achieved via interfering with
374 the mitochondrial biogenesis,uncoupling activity, enhancing of lipid oxidation, and limiting
375 oxidative stress and inflammation

376

377

378

379 **Author contributions**

380 Study design (NC, FT, JFL), data collection (FT, CC, JA, EP, JD, ES, LM, CL, LB, EK, ND), data
381 interpretation (NC, FT, LM, SG, JFL), manuscript preparation (NC, JFL) and funds collection (NC JFL).

382 All the authors have read and approved the final version of this manuscript.

383

384 **Acknowledgements**

385 This work has been partly funded by INRAE, INSERM and Aix Marseille University and la
386 Région Nouvelle-Aquitaine. We thank Dr. Véronique Seidel for English language editing.

387

388

389 **Conflict of interest statement**

390 NC and CL are employed by Pollenergie. Other authors have nothing to disclose.

391

392

393 **References**

394

- 395 [1] Zheng, Y., Ley, S. H., Hu, F. B., Global aetiology and epidemiology of type 2 diabetes mellitus and its
396 complications. *Nature Reviews Endocrinology* 2017, *14*, 88-98.
- 397 [2] Riaz, H., Khan, M. S., Siddiqi, T. J., Usman, M. S., Shah, N., Goyal, A., Khan, S. S., Mookadam, F.,
398 Krasuski, R. A., Ahmed, H., Association Between Obesity and Cardiovascular Outcomes: A Systematic
399 Review and Meta-analysis of Mendelian Randomization Studies. *JAMA Netw Open* 2018, *1*, e183788.
- 400 [3] Gregor, M. F., Hotamisligil, G. S., Inflammatory mechanisms in obesity. *Annu Rev Immunol* 2011, *29*,
401 415-445.
- 402 [4] Tourniaire, F., Romier-Crouzet, B., Lee, J. H., Marcotorchino, J., Gouranton, E., Salles, J., Malezet, C.,
403 Astier, J., Darmon, P., Blouin, E., Walrand, S., Ye, J., Landrier, J. F., Chemokine Expression in Inflamed
404 Adipose Tissue Is Mainly Mediated by NF-kappaB. *PLoS One* 2013, *8*, e66515.
- 405 [5] Karkeni, E., Astier, J., Tourniaire, F., El Abed, M., Romier, B., Gouranton, E., Wan, L., Borel, P., Salles,
406 J., Walrand, S., Ye, J., Landrier, J. F., Obesity-associated Inflammation Induces microRNA-155 Expression
407 in Adipocytes and Adipose Tissue: Outcome on Adipocyte Function. *J Clin Endocrinol Metab* 2016, *101*,
408 1615-1626.
- 409 [6] Ying, W., Riopel, M., Bandyopadhyay, G., Dong, Y., Birmingham, A., Seo, J. B., Ofrecio, J. M., Wollam,
410 J., Hernandez-Carretero, A., Fu, W., Li, P., Olefsky, J. M., Adipose Tissue Macrophage-Derived Exosomal
411 miRNAs Can Modulate In Vivo and In Vitro Insulin Sensitivity. *Cell* 2017, *171*, 372-384 e312.
- 412 [7] Landrier, J.-F., Derghal, A., Mounien, L., MicroRNAs in Obesity and Related Metabolic Disorders. *Cells*
413 2019, *8*, 859.
- 414 [8] Olefsky, J. M., Glass, C. K., Macrophages, inflammation, and insulin resistance. *Annu Rev Physiol*
415 2010, *72*, 219-246.
- 416 [9] Sforcin, J. M., Biological Properties and Therapeutic Applications of Propolis. *Phytother Res* 2016, *30*,
417 894-905.
- 418 [10] Pasupuleti, V. R., Sammugam, L., Ramesh, N., Gan, S. H., Honey, Propolis, and Royal Jelly: A
419 Comprehensive Review of Their Biological Actions and Health Benefits. *Oxid Med Cell Longev* 2017,
420 2017, 1259510.
- 421 [11] Karimian, J., Hadi, A., Pourmasoumi, M., Najafgholizadeh, A., Ghavami, A., The efficacy of propolis
422 on markers of glycemic control in adults with type 2 diabetes mellitus: A systematic review and meta-
423 analysis. *Phytother Res* 2019, *33*, 1616-1626.
- 424 [12] Ichi, I., Hori, H., Takashima, Y., Adachi, N., Kataoka, R., Okihara, K., Hashimoto, K., Kojo, S., The
425 beneficial effect of propolis on fat accumulation and lipid metabolism in rats fed a high-fat diet. *J Food*
426 *Sci* 2009, *74*, H127-131.
- 427 [13] Sakai, T., Ohhata, M., Fujii, M., Oda, S., Kusaka, Y., Matsumoto, M., Nakamoto, A., Taki, T.,
428 Nakamoto, M., Shuto, E., Brazilian Green Propolis Promotes Weight Loss and Reduces Fat Accumulation
429 in C57BL/6 Mice Fed A High-Fat Diet. *Biol Pharm Bull* 2017, *40*, 391-395.
- 430 [14] Koya-Miyata, S., Arai, N., Mizote, A., Taniguchi, Y., Ushio, S., Iwaki, K., Fukuda, S., Propolis prevents
431 diet-induced hyperlipidemia and mitigates weight gain in diet-induced obesity in mice. *Biol Pharm Bull*
432 2009, *32*, 2022-2028.
- 433 [15] Kitamura, H., Naoe, Y., Kimura, S., Miyamoto, T., Okamoto, S., Toda, C., Shimamoto, Y., Iwanaga, T.,
434 Miyoshi, I., Beneficial effects of Brazilian propolis on type 2 diabetes in ob/ob mice: Possible
435 involvement of immune cells in mesenteric adipose tissue. *Adipocyte* 2013, *2*, 227-236.

436 [16] Peng, S., Wei, P., Lu, Q., Liu, R., Ding, Y., Zhang, J., Beneficial Effects of Poplar Buds on
437 Hyperglycemia, Dyslipidemia, Oxidative Stress, and Inflammation in Streptozotocin-Induced Type-2
438 Diabetes. *J Immunol Res* 2018, 2018, 7245956.

439 [17] Chen, L. H., Chien, Y. W., Chang, M. L., Hou, C. C., Chan, C. H., Tang, H. W., Huang, H. Y., Taiwanese
440 Green Propolis Ethanol Extract Delays the Progression of Type 2 Diabetes Mellitus in Rats Treated with
441 Streptozotocin/High-Fat Diet. *Nutrients* 2018, 10.

442 [18] Kosalec, I., Bakmaz, M., Pepeljnjak, S., Vladimir-Knezevic, S., Quantitative analysis of the flavonoids
443 in raw propolis from northern Croatia. *Acta Pharm* 2004, 54, 65-72.

444 [19] Popova, M., Bankova, V., Butovska, D., Petkov, V., Nikolova-Damyanova, B., Sabatini, A. G.,
445 Marcazzan, G. L., Bogdanov, S., Validated methods for the quantification of biologically active
446 constituents of poplar-type propolis. *Phytochem Anal* 2004, 15, 235-240.

447 [20] Gardana, C., Scaglianti, M., Pietta, P., Simonetti, P., Analysis of the polyphenolic fraction of propolis
448 from different sources by liquid chromatography-tandem mass spectrometry. *J Pharm Biomed Anal*
449 2007, 45, 390-399.

450 [21] Reagan-Shaw, S., Nihal, M., Ahmad, N., Dose translation from animal to human studies revisited.
451 *FASEB J* 2008, 22, 659-661.

452 [22] Fenni, S., Hammou, H., Astier, J., Bonnet, L., Karkeni, E., Couturier, C., Tourniaire, F., Landrier, J. F.,
453 Lycopene and tomato powder supplementation similarly inhibit high-fat diet induced obesity,
454 inflammatory response, and associated metabolic disorders. *Mol Nutr Food Res* 2017.

455 [23] Karkeni, E., Marcotorchino, J., Tourniaire, F., Astier, J., Peiretti, F., Darmon, P., Landrier, J. F.,
456 Vitamin D limits chemokine expression in adipocytes and macrophage migration in vitro and in male
457 mice. *Endocrinology* 2015, 156, 1782-1793.

458 [24] Karkeni, E., Bonnet, L., Astier, J., Couturier, C., Dalifard, J., Tourniaire, F., Landrier, J. F., All-trans-
459 retinoic acid represses chemokine expression in adipocytes and adipose tissue by inhibiting NF-kappaB
460 signaling. *J Nutr Biochem* 2017, 42, 101-107.

461 [25] Bonnet, L., Karkeni, E., Couturier, C., Astier, J., Dalifard, J., Defoort, C., Svilar, L., Martin, J. C.,
462 Tourniaire, F., Landrier, J. F., Gene Expression Pattern in Response to Cholecalciferol Supplementation
463 Highlights Cubilin as a Major Protein of 25(OH)D Uptake in Adipocytes and Male Mice White Adipose
464 Tissue. *Endocrinology* 2018, 159, 957-966.

465 [26] Liu, Y., Kern, J. T., Walker, J. R., Johnson, J. A., Schultz, P. G., Luesch, H., A genomic screen for
466 activators of the antioxidant response element. *Proc Natl Acad Sci U S A* 2007, 104, 5205-5210.

467 [27] Fenni, S., Astier, J., Bonnet, L., Karkeni, E., Gouranton, E., Mounien, L., Couturier, C., Tourniaire, F.,
468 Bohm, V., Hammou, H., Landrier, J. F., (all-E)- and (5Z)-Lycopene Display Similar Biological Effects on
469 Adipocytes. *Mol Nutr Food Res* 2019, 63, e1800788.

470 [28] Bonnet, L., Hachemi, M. A., Karkeni, E., Couturier, C., Astier, J., Defoort, C., Svilar, L., Martin, J. C.,
471 Tourniaire, F., Landrier, J. F., Diet induced obesity modifies vitamin D metabolism and adipose tissue
472 storage in mice. *J Steroid Biochem Mol Biol* 2019, 185, 39-46.

473 [29] Livak, K. J., Schmittgen, T. D., Analysis of relative gene expression data using real-time quantitative
474 PCR and the 2(-Delta Delta C(T)) Method. *Methods* 2001, 25, 402-408.

475 [30] Matzinger, M., Fischhuber, K., Heiss, E. H., Activation of Nrf2 signaling by natural products-can it
476 alleviate diabetes? *Biotechnol Adv* 2018, 36, 1738-1767.

477 [31] Nagata, N., Xu, L., Kohno, S., Ushida, Y., Aoki, Y., Umeda, R., Fuke, N., Zhuge, F., Ni, Y., Nagashimada,
478 M., Takahashi, C., Suganuma, H., Kaneko, S., Ota, T., Glucoraphanin Ameliorates Obesity and Insulin
479 Resistance Through Adipose Tissue Browning and Reduction of Metabolic Endotoxemia in Mice.
480 *Diabetes* 2017, 66, 1222-1236.

481 [32] Coppari, R., Bjorbaek, C., Leptin revisited: its mechanism of action and potential for treating
482 diabetes. *Nat Rev Drug Discov* 2012, 11, 692-708.

483 [33] Flachs, P., Rossmeisl, M., Kuda, O., Kopecky, J., Stimulation of mitochondrial oxidative capacity in
484 white fat independent of UCP1: a key to lean phenotype. *Biochim Biophys Acta* 2013, 1831, 986-1003.

485 [34] Tourniaire, F., Musinovic, H., Gouranton, E., Astier, J., Marcotorchino, J., Arreguin, A., Bernot, D.,
486 Palou, A., Bonet, M. L., Ribot, J., Landrier, J. F., All-trans retinoic acid induces oxidative phosphorylation
487 and mitochondria biogenesis in adipocytes. *J Lipid Res* 2016, 56, 1100-1109.

488 [35] McGuinness, O. P., Ayala, J. E., Laughlin, M. R., Wasserman, D. H., NIH experiment in centralized
489 mouse phenotyping: the Vanderbilt experience and recommendations for evaluating glucose
490 homeostasis in the mouse. *American Journal of Physiology-Endocrinology and Metabolism* 2009, 297,
491 E849-E855.

492 [36] Li, Y., Chen, M., Xuan, H., Hu, F., Effects of encapsulated propolis on blood glycemic control, lipid
493 metabolism, and insulin resistance in type 2 diabetes mellitus rats. *Evid Based Complement Alternat*
494 *Med* 2012, 2012, 981896.

495 [37] Nna, V. U., Abu Bakar, A. B., Md Lazin, M., Mohamed, M., Antioxidant, anti-inflammatory and
496 synergistic anti-hyperglycemic effects of Malaysian propolis and metformin in streptozotocin-induced
497 diabetic rats. *Food Chem Toxicol* 2018, 120, 305-320.

498 [38] Stahli, A., Maheen, C. U., Strauss, F. J., Eick, S., Sculean, A., Gruber, R., Caffeic acid phenethyl ester
499 protects against oxidative stress and dampens inflammation via heme oxygenase 1. *Int J Oral Sci* 2019,
500 11, 6.

501 [39] Nguyen, T., Yang, C. S., Pickett, C. B., The pathways and molecular mechanisms regulating Nrf2
502 activation in response to chemical stress. *Free Radic Biol Med* 2004, 37, 433-441.

503 [40] Owuor, E. D., Kong, A. N., Antioxidants and oxidants regulated signal transduction pathways.
504 *Biochem Pharmacol* 2002, 64, 765-770.

505 [41] Nair, S., Li, W., Kong, A. N., Natural dietary anti-cancer chemopreventive compounds: redox-
506 mediated differential signaling mechanisms in cytoprotection of normal cells versus cytotoxicity in
507 tumor cells. *Acta Pharmacol Sin* 2007, 28, 459-472.

508 [42] Vomhof-Dekrey, E. E., Picklo, M. J., Sr., The Nrf2-antioxidant response element pathway: a target for
509 regulating energy metabolism. *J Nutr Biochem* 2012, 23, 1201-1206.

510 [43] Seo, H. A., Lee, I. K., The role of Nrf2: adipocyte differentiation, obesity, and insulin resistance. *Oxid*
511 *Med Cell Longev* 2013, 2013, 184598.

512 [44] Pi, J., Leung, L., Xue, P., Wang, W., Hou, Y., Liu, D., Yehuda-Shnaidman, E., Lee, C., Lau, J., Kurtz, T.
513 W., Chan, J. Y., Deficiency in the nuclear factor E2-related factor-2 transcription factor results in
514 impaired adipogenesis and protects against diet-induced obesity. *J Biol Chem* 2010, 285, 9292-9300.

515 [45] Chartoumpakis, D. V., Ziros, P. G., Psyrogiannis, A. I., Papavassiliou, A. G., Kyriazopoulou, V. E.,
516 Sykiotis, G. P., Habeos, I. G., Nrf2 represses FGF21 during long-term high-fat diet-induced obesity in
517 mice. *Diabetes* 2011, 60, 2465-2473.

518 [46] Zhang, Y. K., Wu, K. C., Liu, J., Klaassen, C. D., Nrf2 deficiency improves glucose tolerance in mice fed
519 a high-fat diet. *Toxicol Appl Pharmacol* 2012, 264, 305-314.

520 [47] Collins, A. R., Gupte, A. A., Ji, R., Ramirez, M. R., Minze, L. J., Liu, J. Z., Arredondo, M., Ren, Y., Deng,
521 T., Wang, J., Lyon, C. J., Hsueh, W. A., Myeloid deletion of nuclear factor erythroid 2-related factor 2
522 increases atherosclerosis and liver injury. *Arterioscler Thromb Vasc Biol* 2012, 32, 2839-2846.

523 [48] Shin, S., Wakabayashi, J., Yates, M. S., Wakabayashi, N., Dolan, P. M., Aja, S., Liby, K. T., Sporn, M. B.,
524 Yamamoto, M., Kensler, T. W., Role of Nrf2 in prevention of high-fat diet-induced obesity by synthetic
525 triterpenoid CDDO-imidazolide. *Eur J Pharmacol* 2009, 620, 138-144.

526 [49] Yu, Z., Shao, W., Chiang, Y., Foltz, W., Zhang, Z., Ling, W., Fantus, I. G., Jin, T., Oltipraz upregulates
527 the nuclear factor (erythroid-derived 2)-like 2 [corrected](NRF2) antioxidant system and prevents insulin
528 resistance and obesity induced by a high-fat diet in C57BL/6J mice. *Diabetologia* 2011, 54, 922-934.

529 [50] Abo El-Magd, N. F., El-Mesery, M., El-Karef, A., El-Shishtawy, M. M., Glycyrrhizin ameliorates high
530 fat diet-induced obesity in rats by activating Nrf2 pathway. *Life Sci* 2018, 193, 159-170.

- 531 [51] Piantadosi, C. A., Carraway, M. S., Babiker, A., Suliman, H. B., Heme oxygenase-1 regulates cardiac
532 mitochondrial biogenesis via Nrf2-mediated transcriptional control of nuclear respiratory factor-1. *Circ*
533 *Res* 2008, *103*, 1232-1240.
- 534 [52] Hota, K. B., Hota, S. K., Chaurasia, O. P., Singh, S. B., Acetyl-L-carnitine-mediated neuroprotection
535 during hypoxia is attributed to ERK1/2-Nrf2-regulated mitochondrial biosynthesis. *Hippocampus* 2012,
536 *22*, 723-736.
- 537 [53] Furukawa, S., Fujita, T., Shimabukuro, M., Iwaki, M., Yamada, Y., Nakajima, Y., Nakayama, O.,
538 Makishima, M., Matsuda, M., Shimomura, I., Increased oxidative stress in obesity and its impact on
539 metabolic syndrome. *J Clin Invest* 2004, *114*, 1752-1761.
- 540 [54] Houstis, N., Rosen, E. D., Lander, E. S., Reactive oxygen species have a causal role in multiple forms
541 of insulin resistance. *Nature* 2006, *440*, 944-948.
- 542 [55] Tanaka, Y., Aleksunes, L. M., Yeager, R. L., Gyamfi, M. A., Esterly, N., Guo, G. L., Klaassen, C. D., NF-
543 E2-related factor 2 inhibits lipid accumulation and oxidative stress in mice fed a high-fat diet. *J*
544 *Pharmacol Exp Ther* 2008, *325*, 655-664.

545

546

547

548 **Figure legends**

549 **Fig 1. Poplar propolis ethanolic extract limits weight gain associated with diet-induced**
550 **obesity.** **A.** Body weight evolution was quantified for each mouse ($n = 10$ per group). **B.** Before
551 sacrifice, animal weight was recorded. **C.** Energy intake was quantified by measuring food intake
552 every two weeks for a period of 12 weeks. **D.** Absolute organ weights. **E.** Relative organ weights
553 (absolute organ weight / body weight). **F.** Adiposity index was calculated by dividing the sum of
554 adipose tissues weight by the body weight of animal. Values are presented as means \pm SEM.
555 Bars not sharing the same letter were significantly different in Tukey-Kramer post hoc test $p <$
556 $0,05$.

557
558 **Fig 2. Poplar propolis ethanolic extract improves glucose homeostasis.** **A.** Glucose levels
559 during the oral glucose tolerance test (OGTT; $n=10$ mice per group). **B.** Area under the curve
560 (AUC) for the glycemie response during OGTT. **C.** Glycemia evolution during the insulin
561 tolerance test (ITT; $n=10$ mice per group). **D.** Area under the curve for the glycemie response
562 during ITT. **E.** Fasted glycemia and insulinemia of mice ($n=10$ mice per group). **F.** HOMA-IR
563 mean values. Values are presented as means \pm SEM. Bars not sharing the same letter were
564 significantly different in Tukey-Kramer post hoc test $p < 0,05$.

565
566 **Fig 3. Poplar propolis ethanolic extract effect on adipocyte cellularity, gene expression and**
567 **DNA mitochondrial content in adipose tissue.** **A.** Representative histological images of
568 epididymal fat pads H&E stained, taken at 10X magnification. Adipocyte area determined using
569 Image J software. **B.** Relative expression of epididymal adipose tissue genes related to

570 inflammatory markers measured through qPCR and expressed relative to 18S ribosomal RNA.
571 **C.** Relative expression of epididymal adipose tissue genes related to lipid metabolism measured
572 through qPCR and expressed relative to 18S ribosomal RNA. **D.** Relative expression of
573 epididymal adipose tissue genes related to mitochondrial metabolism measured through qPCR
574 and expressed relative to 18S ribosomal RNA. **E.** Mitochondrial DNA quantification in
575 epididymal adipose tissue. Mitochondrial DNA to nuclear DNA ratio was determined by qPCR.
576 N=10 mice per each group, values are presented as means \pm SEM. Bars not sharing the same
577 letter were significantly different in Tukey-Kramer post hoc test $p < 0,05$.

578

579 **Fig 4. Poplar propolis ethanolic extract induces Nrf2 pathway in adipose tissue and**
580 **adipocytes.** **A.** Relative expression of epididymal adipose tissue Nrf2-target genes coding for
581 antioxidant enzymes measured through qPCR and expressed relative to 18S ribosomal RNA
582 (n=10 mice per group). **B.** 3T3-L1 adipocytes were incubated with growing concentrations of
583 PPEE for 24 h. Expression of Nrf2-target genes coding for antioxidant enzymes measured
584 through qPCR and expressed relative to 18S ribosomal RNA. **C.** 3T3-L1 adipocytes were
585 transiently transfected with plasmid containing wild-type or mutated antioxidant response
586 element (ARE) and cells were incubated for 24 h with 30 μ g/ml of PPEE. Luminescence was
587 monitored and standardized. **D.** 3T3-L1 adipocytes were incubated with PPEE (30 μ g/ml) for 24
588 h. DNA binding activity was monitored using transAM Nrf2 assay. **E to H.** 3T3-L1 adipocytes
589 were transiently transfected with non-targeted (NT) siRNA or siRNA directed against Nrf2 for
590 24h. cells were incubated for an additional 24h, and gene expression was quantified by qPCR
591 and expressed relative to 18S ribosomal RNA. Values are presented as means \pm SEM of 3

592 independent cultures per treatment condition. Bars not sharing the same letter were significantly
593 different in Tukey-Kramer post hoc test $p < 0,05$.

594

595

596

597

598