



HAL
open science

The effects of food quantity, light, and temperature on clearance rates in freshwater bivalves (Cyrenidae and Unionidae)

Simon Pouil, Amber Hills, Teresa Mathews

► **To cite this version:**

Simon Pouil, Amber Hills, Teresa Mathews. The effects of food quantity, light, and temperature on clearance rates in freshwater bivalves (Cyrenidae and Unionidae). *Hydrobiologia*, 2021, 848 (3), pp.675-689. 10.1007/s10750-020-04472-y . hal-03163618

HAL Id: hal-03163618

<https://hal.inrae.fr/hal-03163618>

Submitted on 10 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **The effects of food quantity, light, and temperature on clearance rates**
2 **in freshwater bivalves (Cyrenidae and Unionidae)**

3
4 Simon Pouil¹, Amber Hills¹, Teresa J. Mathews¹

5
6 ¹ Environmental Sciences Division, Oak Ridge National Laboratory, Oak Ridge, TN, 37830, USA

7
8 * Correspondence: Teresa J. Mathews
9 Aquatic Ecology Group
10 Environmental Sciences Division
11 Oak Ridge National Laboratory
12 Oak Ridge, TN, 37830, USA
13 E-mail: mathewstj@ornl.gov

14
15 **Acknowledgements**

16 This work was funded by the US Department of Energy's Oak Ridge Office of Environmental Management (ORO-
17 DOE) and URS CH2M Oak Ridge LLC (UCOR) and is a product of ORNL's Mercury Remediation Technology
18 Development Program. Authors acknowledge Dr. Dan Hua, Senior Scientist at TWRA's C-RAC, and her team who
19 reared and provided the juveniles *Lampsilis ovata* used in this study. Special thanks to Allison Fortner, Michael
20 Jones, Trent Jett, and Nikki Jones and Louise Stevenson (ORNL) for assistance with field collections of *Corbicula*
21 *fluminea* and/or their constructive comments on the manuscript.

22 This manuscript has been authored by UT-Battelle, LLC under Contract No. DE-AC05-00OR22725 with the
23 U.S. Department of Energy. The United States Government retains and the publisher, by accepting the article
24 for publication, acknowledges that the United States Government retains a non-exclusive, paid-up,
25 irrevocable, worldwide license to publish or reproduce the published form of this manuscript, or allow others
26 to do so, for United States Government purposes. The Department of Energy will provide public access to
27 these results of federally sponsored research in accordance with the DOE Public Access Plan
28 (<http://energy.gov/downloads/doe-public-access-plan>).

29 **Abstract**

30 Assessing the environmental variables that influence freshwater bivalve filtration activity is key to better understand
31 the functioning of stream ecosystems. In the present study, the effects of light ($19 \pm 2 \mu\text{mol m}^{-2} \text{s}^{-1}$ and $0 \pm 0 \mu\text{mol}$
32 $\text{m}^{-2} \text{s}^{-1}$) and temperature (10, 15, 20 and 25 °C) on the clearance rates of two bivalve species: the Asian clam
33 *Corbicula fluminea* (Cyrenidae) and pocketbook *Lampsilis ovata* (Unionidae) were assessed under controlled
34 laboratory conditions at two different concentrations of algal food (i.e. 10,000 cells mL^{-1} and 100,000 cells mL^{-1}).
35 Clearance rates for *C. fluminea* varied from 0 to 491 $\text{mL g}^{-1} \text{h}^{-1}$ while the observed range for *L. ovata* was 0 to 905
36 $\text{mL g}^{-1} \text{h}^{-1}$. We found that the relative contribution of the tested variables was species-dependent. While temperature
37 plays a major role in the clearance rates of *C. fluminea*, food concentration was the most significant variable in the
38 clearance rate of *L. ovata*. Our results confirm the complex interactions between abiotic and biotic factors in
39 freshwater bivalve filtration activity. Overall, clearance rates are highly variable especially in *C. fluminea* and
40 presumably regulated by other untested factors suggesting high plasticity in filtration.

41

42 **Keywords:** Asian clam, Environmental variables, Filtration, Freshwater mussel, Pocketbook

43 **Introduction**

44 Freshwater bivalves are ecosystem engineers that feed by filtering suspended particles, including phytoplankton,
45 zooplankton, bacteria and fine organic detritus (Nichols et al., 2005; Haag, 2012; Marescaux et al., 2016) from the
46 water column. Through their filtration activity, freshwater bivalves can significantly modulate the availability of
47 resources to other organisms. For example, by filtering suspended particles out of the water column and transferring
48 these resources to substrates (i.e. as feces or pseudofeces), they play an important role in benthic-pelagic coupling in
49 stream systems (Howard & Cuffey, 2006) and can significantly affect water quality (Pigneur et al., 2014).

50 In addition to affecting energy and nutrient recycling and storage, freshwater mussels can create structural habitat
51 and can affect food web structures. Thus, losses or overabundance of bivalves may lead to, among other things, a
52 disruption in trophic networks or a deeply modified habitat for other organisms (Marescaux et al., 2016; Vaughn,
53 2018). In addition, because bivalve filtration activity affects the dynamics of contaminants in streams, there are
54 growing considerations regarding the use of freshwater bivalves for contaminant mitigation or bioremediation in
55 lakes and industrial effluents (Bianchi et al., 2014; Rosa et al., 2014; Domingues et al., 2020). For all of the
56 aforementioned reasons, a whole field of research is dedicated to the assessment of filtration rates in bivalves.

57 The filtration rates of freshwater bivalves are affected by environmental parameters including both abiotic and biotic
58 factors. Among the abiotic factors, several studies have already shown that temperature plays an important role for
59 many species (Aldridge et al., 1995; Lim et al., 2005; Loayza-Muro & Elias-Letts, 2007) while other work
60 demonstrated effects of pH, water velocity and turbidity (Ackerman, 1999; Loayza-Muro & Elias-Letts, 2007;
61 Tuttle- Raycraft & Ackerman, 2019). In addition, recent evidence shows that light may affect filtration rates in the
62 Asian clam *Corbicula fluminea* (Müller, 1774) (Hills et al., 2020). Numerous studies have also shown that biotic
63 factors such as food availability and particle type (Sprung & Rose, 1988), as well as reproduction and growth cycle
64 (Hornbach et al., 1984; Viergutz et al., 2012) can influence filtration rates.

65 Although a considerable amount of data is available regarding filtration rates on widespread invasive species such as
66 *Corbicula* and *Dreissena* species (see Marescaux et al., 2016), limited attention is paid to Unionid mussels. Overall,
67 the filtering capacity of Unionidae is often reported to be lower than the filtration capacity of *Corbicula*. Indeed, in
68 Unionidae, individual filtration rates for various species usually range between 18 and 2,000 mL ind⁻¹ h⁻¹
69 (Tankersley & Dimock, 1993; Gatenby et al., 1996; Tankersley, 1996; Loayza-Muro & Elias-Letts, 2007; Gatenby
70 et al., 2013; Mistry & Ackerman, 2017, 2018; Tuttle-Raycraft & Ackerman, 2018, 2019) though filtration rates of up

71 to 10,000 mL ind⁻¹ h⁻¹ were reported for the mucket *Actinonaias ligamentina* (Lamarck, 1819) (>100 g, Baker &
72 Hornback, 2001) compared to filtration rates between 29 and 3,000 mL ind⁻¹ h⁻¹ for *Corbicula* species (see
73 Marescaux et al., 2016). Nevertheless, there are some discrepancies in the data available as shown by the wide
74 variation in measured filtration rates. Indeed, very few studies have directly compared the filtration rates of different
75 bivalves, especially for freshwater species, in the same study using the same methodology (Kryger & Riisgård,
76 1988; Marescaux et al., 2016), making it difficult to compare filtration rate values among species and assess how
77 environmental variables affect them.

78 In this study, we aimed to test the effects of light and temperature on the clearance rates (i.e. volume of water
79 cleared of suspended particles per unit of time (Riisgård, 2001) of two bivalve species: the Asian clam *C. fluminea*
80 (Cyrenidae) and the pocketbook mussel *Lampsilis ovata* (Say, 1817) (Unionidae) under controlled laboratory
81 conditions at two different concentrations of algal food: ‘low’ food (i.e. 10,000 cells mL⁻¹) and ‘high’ food (i.e.
82 100,000 cells mL⁻¹). Using collected data from two experiments, the relative contribution of each tested variable (i.e.
83 food, temperature and light) on the clearance rate of each species was assessed.

84

85 **Methods**

86 Origin and acclimation of bivalves

87 Two-year-old juvenile *L. ovata* (pocketbook) and adult *C. fluminea* (Asian clam) were chosen in order to compare
88 native to exotic species in the USA. *C. fluminea* were collected from Sewee Creek in Meigs county, TN, USA, while
89 *L. ovata* came from Tennessee Wildlife Resources Agency’s hatchery, Cumberland River Aquatic Center (TWRA’s
90 C-RAC). Bivalves were brought to the laboratory in a cooler with an air bubbler. No mortality was recorded during
91 transportation. The bivalves were kept in a 700-L tank supplied with flow-through water from First Creek on the
92 Oak Ridge Reservation in Oak Ridge, TN (water renewal: 50-150 L h⁻¹; ambient temperature: 15-18 °C; light/dark:
93 12h/12h), and acclimated to the laboratory for at least three months prior to the experiment. In addition to the supply
94 of food particles coming from the water inlet (i.e. ~1,000 particles mL⁻¹), bivalves were fed a daily diet of fresh
95 algae: *Chlamydomonas reinhardtii* (Dangeard, 1888) and *Navicula* sp. (approx. 7 x 10⁷ cells ind⁻¹ d⁻¹) using a
96 peristaltic pump.

97 Four weeks before the experiment, individuals of each species were randomly placed in four 40-L closed-system
98 aquaria (n = 8 per species) and acclimated to the targeted temperatures (10, 15, 20 and 25 °C). Each aquarium was

99 equipped with an airlift foam filter and a plastic tray containing a 1-cm layer of fine gravel (1-4 mm) to encourage
100 natural burrowing activity of the bivalves. During the beginning of the acclimation period, temperatures were
101 gradually adjusted ($\sim 1\text{ }^{\circ}\text{C d}^{-1}$) and then stabilized at the targeted temperatures for at least 20 days before starting the
102 experiments. Each aquarium was placed in a large water bath to aid temperature control of the small volume. During
103 the aquaria-acclimation period, bivalves were fed exclusively on *C. reinhardtii* (1.2×10^7 cells $\text{ind}^{-1} \text{d}^{-1}$) and daily
104 water renewals (i.e. 10-20%) were performed.

105 Temperature and light were measured three times a day in each aquarium using a hand-held thermometer (Oakton®
106 RTD thermometer) and a PAR meter (Quantum Flux® Apogée), respectively. Conductivity, pH, dissolved oxygen,
107 total dissolved solids (TDS) were monitored 5 times a week in each aquarium using a logging multiparameter meter
108 (HANNA® Hi 9829). Additional measurements were regularly taken in each aquarium for NH_3 and NO_2^- (HACH®
109 SL 1000). Values of water parameters are summarized in Table 1.

110

111 Algae culture

112 The unicellular green algae *C. reinhardtii* (5-6 μm in diameter) was selected for its availability and ease of
113 maintenance in laboratory conditions. This chlorophyte remains in suspension because it is flagellated, facilitating
114 the homogeneity of the particles requires to properly estimate clearance rates. Although this species has been shown
115 to be filtered by both clams and mussels (Boltovskoy et al., 1995; Gatenby et al., 1996), is not among the dominant
116 species in the diet of such bivalves and no nutritional analysis has been performed in the present study. *C.*
117 *reinhardtii* culture was grown in 10-L clear Nalgene carboys containing WC medium (Guillard, 1975) (temperature:
118 $23\text{-}24\text{ }^{\circ}\text{C}$; light intensity: $150\text{-}200\text{ }\mu\text{mol m}^{-2} \text{s}^{-1}$; light/dark: 12h/12h). Algal concentrations were evaluated by
119 imaging flow-cytometry (see “Sample analysis” section for details). For the entire experiment (i.e. acclimation and
120 clearance rates assessment), the algae used were harvested during their exponential growth phase (i.e. after 7 ± 2 d at
121 $1.4 \pm 0.5 \times 10^6$ cells mL^{-1} ; mean \pm SD).

122

123 Experimental procedures

124 To assess the effects of food concentration on clearance rates of *C. fluminea* and *L. ovata*, two experiments were
125 performed under the same temperature (10, 15, 20 and $25\text{ }^{\circ}\text{C}$) and light conditions ($19 \pm 2\text{ }\mu\text{mol m}^{-2} \text{s}^{-1}$ and 0 ± 0
126 $\mu\text{mol m}^{-2} \text{s}^{-1}$). In Experiment 1, bivalves were acclimated to temperature and food ration (1.2×10^7 cells $\text{ind}^{-1} \text{d}^{-1}$),

127 then clearance rates were assessed using *C. reinhardtii* (dry weight of 83 pg cell⁻¹, Pickhardt & Fisher, 2007) at a
128 concentration of 10,000 cells mL⁻¹ (i.e. 0.8 mg (dry weight) L⁻¹; ‘low food’ condition). In Experiment 2, after an
129 additional week of acclimation to a 10-fold increase of the food ration (12 x 10⁷ cells ind⁻¹ d⁻¹), clearance rates were
130 assessed at the second *C. reinhardtii* concentration (i.e. 100,000 cells mL⁻¹ or 8 mg (dry weight) L⁻¹; ‘high food’
131 condition). The detailed experimental schedule is described in Fig. 1.

132 In each experiment, eight individuals of each bivalve species were randomly selected for each experimental
133 temperature (*C. fluminea*: 2.06 ± 0.35 g wet wt, 16.9 ± 0.8 mm shell length; *L. ovata*: 0.54 ± 0.13 g wet wt, 16.4 ±
134 1.2 mm shell length). On the day of the experiment, each individual was weighed and placed in a food-grade PET
135 container filled with 200 mL of stream water. Temperature was kept constant in the containers at targeted
136 experimental treatments by using chilled or heated water baths. Three containers per temperature treatment were
137 used as controls with no bivalve. Slight air bubbling in each container kept water well-circulated. Bivalves were
138 acclimated to experimental containers for 4 h. The sample sizes (i.e. n = 8 per temperature for each species)
139 followed the recommendations from (Salerno et al., 2018).

140 At the start of the experiment, concentrated live *C. reinhardtii* cells were spiked into each plastic container to add
141 the targeted initial concentration (see above). Algae cells were concentrated by centrifugation (5000 RPM for 5 min)
142 and resuspension in Milli-Q water. The use of concentrated algae allowed a reduction of spike volume (i.e. ~500-
143 600 µL) and avoided significant changes in water volume to preserve physical and chemical parameters.

144 The cell concentration of the initial spike was checked immediately by analyzing a 3-mL sample from three
145 randomly selected containers in each temperature treatment using flow imaging cytometry (see “Sample analysis”
146 section). A 4-mL sample was taken from each plastic container five minutes after the algae spike and then a second
147 4-mL sample at the end of the 35-minute filtration period. Samples were placed in 5-mL snap-cap centrifuge tubes
148 containing 800 µL of 10% formalin solution, a common preservative for phytoplankton samples (Mukherjee et al.,
149 2014) and mixed immediately.

150 In a single day, each experiment was first performed in light conditions (i.e. at 2PM: 19 ± 2 µmol m⁻² s⁻¹). Each PET
151 container was cleaned and rinsed with dechlorinated tap water, refilled with stream water at the appropriate
152 temperature, and the same organisms were acclimated for another 4-h period. The same experiment was then
153 repeated in dark conditions (i.e. at 8PM: 0 ± 0 µmol m⁻² s⁻¹). This protocol, without photoperiod modification,
154 maintains the circadian rhythm in the studied bivalves throughout the experiments.

155

156 Sample analysis

157 Algal cell counts were performed by flow imaging cytometry (FlowCam® Benchtop B3 Model) following a
158 standard procedure within 30 days of sampling to limit the effect of preservative. The preserved samples were
159 analyzed in AutoImage mode in which particles of the flow sample are imaged and captured at a regular interval,
160 with no fluorescence measurements being taken. Therefore, every particle (phytoplankton, aggregates, inorganic,
161 and so on) ranging from 2 to 50 μm equivalent spherical diameter (ESD) was counted and imaged. For each sample,
162 a maximum of either 10,000 particles or 1 mL were analyzed. A 10X objective was used in the sample analysis, and
163 the instrument count-calibration was checked using beads of a known size (Zarauz & Irigoien, 2008). Invalid
164 pictures (i.e. bubbles, repeated images) were removed from the image database, through visual recognition. Each
165 sample was run in at least duplicate, but if the coefficient of variation was $\geq 20\%$ between measurements, a third
166 measurement was taken. In that case, the two results within the variation limits were used in the mean calculation; if
167 all three results were outside variation limits, then the mean was calculated from all three. Mean values were then
168 used for clearance rate calculations.

169

170 Computations and statistics

171 Clearance rates (expressed as $\text{mL g}^{-1} \text{h}^{-1}$) were calculated two ways, using whole-body wet weight and soft parts wet
172 weight, according to the following equation modified from Coughlan (1969) (see Mistry & Ackerman, 2018):

173
$$CR = \frac{V}{nt} \left(\ln \frac{C_i}{C_f} - \ln \frac{C'_i}{C'_f} \right)$$

174 Where V is the volume of water in the PET container (mL), n is the number of individuals per container ($n=1$), t is
175 the length of the experiment (h), C_i and C_f are initial and final cell concentrations respectively in the container with
176 bivalves while, and C'_i and C'_f are average initial and final cell concentrations respectively in the containers used as
177 controls (with no bivalve). Based on calculations, all the individuals whose valves were closed at the end of the
178 experiment and with $CR < 1 \text{ mL g}^{-1} \text{h}^{-1}$ were considered as non-filtering.

179 Temperature coefficient (Q_{10}) values were calculated using the following equation:

$$Q_{10} = \left(\frac{R_2}{R_1} \right)^{\frac{10}{t_2 - t_1}}$$

180 Where R_1 is the clearance rate at t_1 temperature, and R_2 is the clearance rate at t_2 temperature.
181 For each food concentration test (i.e. ‘low’ and ‘high’ food), the data were analyzed using a three-way mixed
182 Analysis of Variance (ANOVA) to examine the effect of light (i.e. ‘dark’ and ‘light’ within-subjects factor) and
183 temperature (10, 15, 20 and 25 °C) on the clearance rates of the two bivalve species (*C. fluminea* and *L. ovata*). All
184 two-way and three-way interactions were examined ($\alpha = 0.05$). The data were first checked for normality (Shapiro’s
185 test) and homoscedasticity (Levene’s test). T-tests were then used to identify pairwise differences for significant
186 factors in the model. P-values were then adjusted using the Bonferroni multiple testing correction method. All
187 statistical analyses were performed in R v. 3.6.3 (R Development Core Team, 2020), using the packages *lmerTest*,
188 *rstatix* and *tidyverse*.

189 The relative contribution of each variable (food, temperature, and light) on the clearance rates of each species was
190 assessed from the data of the two experiments through the *lmg* approach (Groemping, 2007) using the package
191 *relaimpo*. This package does not cover linear mixed models, so we accommodated clustered data to a design that
192 contains clusters by applying a generalized linear model (GLM) with the *svyglm* function from the package *survey*
193 with a linear link function and gaussian distribution. The bootstrapping approach was subsequently used to test
194 significant differences in the contribution of each variable in the model; this takes care of the dependence between
195 the data from the same individuals.

196

197 **Results**

198 **General**

199 Overall, across all temperature, light and food conditions, clearance rates for *C. fluminea* (whole-body wet weight:
200 2.10 ± 0.34 g, mean \pm SD; min-max: 1.37-2.80 g) ranged from 0 to 591 mL g⁻¹ h⁻¹ (median value: 181 mL g⁻¹ h⁻¹)
201 equiv. to 0 to 1243 mL ind⁻¹ h⁻¹ (median value: 385 mL ind⁻¹ h⁻¹). Clearance rates for *L. ovata* (whole-body wet
202 weight: 0.56 ± 0.13 g, mean \pm SD; min-max: 0.33-0.89 g) ranged from 0 to 905 mL g⁻¹ h⁻¹ (median value: 233 mL g⁻¹
203 h⁻¹) equiv. to 0 to 507 mL ind⁻¹ h⁻¹ (median value: 133 mL ind⁻¹ h⁻¹).

204 Results are noticeably different when the calculation of the clearance rates are based only on the soft-parts wet
205 weight, the organic fraction involved in filtration activity, which represents $9 \pm 2\%$ ($n = 10$) of the total body weight
206 in *C. fluminea* and $28 \pm 4\%$ in *L. ovata* ($n = 5$). In this case, the clearance rates of *C. fluminea* are systematically

207 higher than those calculated for *L. ovata* with values 0 to 6517 mL g soft parts⁻¹ h⁻¹ (median value: 1991 mL g soft
208 parts⁻¹ h⁻¹) against 0 to 3234 mL g soft parts⁻¹ h⁻¹ (median value: 834 mL g soft parts⁻¹ h⁻¹) in *L. ovata*.

209 In the following sections, results are presented by experiment (i.e. Experiment 1 performed at 10,000 cells mL⁻¹ and
210 Experiment 2 performed at 100,000 cells mL⁻¹). As indicated in “Computations and statistics” section, combined
211 results from the two experiments were used to estimate the relative contribution of food, temperature and light on
212 clearance rate.

213

214 Experiment 1

215 The first experiment was performed using a low concentration of food (i.e. 10,000 cells mL⁻¹). In this treatment,
216 combining all the temperature and light conditions, clearance rates ranged from 0 to 472 mL g⁻¹ h⁻¹ (median value:
217 186 mL g⁻¹ h⁻¹) for *C. fluminea* (n = 64) while median clearance rate of *L. ovata* (n = 64) was 2.5-fold higher (i.e.
218 455 mL g⁻¹ h⁻¹) and values were ranging from 0 to 905 mL g⁻¹ h⁻¹ (Fig. 2). Interestingly, the proportion of individuals
219 filtering during this experiment also differed according to species. While most of the *L. ovata* filtered regardless of
220 the temperature and light condition (88-100% of individuals filtering), the proportion of individuals filtering was
221 more variable for *C. fluminea*: in both light and dark conditions, 100% of the individuals filtered at 15 °C , but only
222 88% at 10 and 20 °C and 75% at 25 °C (Fig. 3).

223 A three-way mixed ANOVA was performed to evaluate the effects of temperature and light on the clearance rates of
224 the two bivalve species. The three-way interaction between temperature, light and species was not significant (F(3,
225 56) = 0.74, *P* = 0.535), but a statistically significant simple two-way interaction was found between temperature and
226 species, (F(3, 56) = 5.29, *P* = 0.003) indicating that effect of temperature is species dependent. Thus, while there
227 were significant effects of temperature on clearance rates of *L. ovata* in the dark (F(3, 28) = 7.10, *P* = 0.001), no
228 statistical difference was observed for *C. fluminea* maintained in the dark (F(3, 28) = 1.77, *P* = 0.176). For both
229 species, no significant effect of temperature was found in light condition. Pairwise comparisons tests revealed that
230 for *L. ovata* in the dark, clearance rates were significantly different between 10 °C and 25 °C (*P* < 0.001). There was
231 no significant difference for other temperatures (*P* > 0.05, Fig. 3). In low food conditions, Q₁₀ values were 1.56
232 (dark) and 1.08 (light) in *C. fluminea* compared to 1.25 to 1.73 in dark and light conditions respectively in *L. ovata*.

233

234 Experiment 2

235 The second experiment was performed after one week of acclimation to a ten-fold higher daily food ration to assess
236 the clearance rate at a high concentration of food (i.e. 100,000 cells mL⁻¹). When all temperature and light
237 conditions were combined, the clearance rates ranged from 0 to 591 mL g⁻¹ h⁻¹ (median value: 154 mL g⁻¹ h⁻¹) for *C.*
238 *fluminea* (n = 64) and 0 to 453 mL g⁻¹ h⁻¹ (median value: 62 mL g⁻¹ h⁻¹) for *L. ovata* (n = 64, Fig. 4). While median
239 values were similar at the two food concentrations in *C. fluminea*, median values for *L. ovata* clearance rate was
240 more than 7-fold higher at the 'high' food concentration than the one observed at the 'low' food concentration. The
241 proportion of individuals filtering also differed between the two experiments. In the 'high' food condition, in dark,
242 all the *C. fluminea* filtered at all temperatures. For the same species, filtration activity was highly dependent on
243 temperature and light condition (38% of individuals filtered at 10 °C vs. 100% at 15 and 25 °C). For *L. ovata*, in the
244 dark, all the individuals filtered at the extreme temperatures (i.e. 10 °C and 25 °C) while this proportion decreased to
245 88% and 63% at 20 °C and 15 °C, respectively. The proportion of *L. ovata* individuals filtering in light increased
246 with increasing temperature from 50% at 10 °C and 15 °C to 88% at 25 °C (Fig. 5).

247 As for the 'low' food concentration tested, a three-way mixed ANOVA was used to evaluate the effects of
248 temperature and light conditions on the clearance rates *C. fluminea* and *L. ovata* revealed that the interaction
249 between temperature, light and species was not significant (F(3, 56) = 1.71, *P* = 0.176, Table 2). In contrast, the two-
250 way interaction between temperature and species was significant (F(3, 56) = 3.29, *P* = 0.027). There were significant
251 effects of temperature on clearance rates of *C. fluminea* in the dark (F(3, 28) = 19.6, *P* < 0.001) and in the light (F(3,
252 28) = 4.83 *P* = 0.008). Significant effects were also found for *L. ovata* in the dark (F(3, 28) = 9.46, *P* < 0.001) and in
253 the light (F(3, 28) = 4.83, *P* = 0.017). In the dark condition, clearance rates gradually increased with increasing
254 temperatures for the two species (*P* < 0.01). In the light condition, clearance rates of the two species were only
255 significantly different between the extreme temperatures tested (i.e. 10 °C and 25 °C, *P* < 0.05, Fig. 4). In high food
256 conditions, Q₁₀ values differed only slightly with light in *C. fluminea* with values of 4.81 in the dark and 4.53 in the
257 light while a broader range was observed for *L. ovata* (i.e. Q₁₀ of 2.41 in dark and 3.32 in light condition).

258

259 Relative contribution of food, temperature and light on clearance rate

260 GLM models were implemented using the data from each experiment to assess the effect of explanatory variables
261 (i.e. food, temperature and light) on clearance rates (Table 3). The lmg approach was then used to estimate the
262 relative contribution of each variable. For *C. fluminea*, the GLM revealed that temperature (*P* < 0.001) and to a

263 lesser extent light ($P = 0.022$) played a significant role in clearance rates, but clearance rates were not affected by
264 the 10-fold difference in food concentration ($P = 0.87$). Interestingly, for *C. fluminea* only 16% of the variance
265 observed in clearance rates was explained by the model. Within the explained variance (16%), temperature had the
266 highest contribution (80%, CI: 64-98%) while light had moderate and food had virtually no effect on the clearance
267 rate (19%, CI: 2-28% and 0%, CI: 0-9%, respectively). For *L. ovata*, clearance rate was significantly affected by
268 food, temperature ($P < 0.001$) and to a lesser extent light ($P = 0.048$) with a high proportion of the observed variance
269 explained by these three variables (67%). Within the explained variance, food clearly has the highest contribution
270 (81%, CI: 72-89%) while temperature and light have moderate effects on the clearance rate (18%, CI: 11-22% and
271 2%, CI: 0-6%, respectively, Fig. 6).

272

273 **Discussion**

274 Our experimental approach addresses the critical need to examine the relative contribution of biotic and abiotic
275 factors in the clearance rates of freshwater bivalves. To the best of our knowledge, this is the first study to examine
276 the effects and relative contribution of light, temperature and food in the clearance rates of two freshwater bivalve
277 species: the invasive Asian clam *C. fluminea* (Cyrenidae) and the native pocketbook *L. ovata* (Unionidae).

278 Overall, the filtration rates observed in our study were mostly in the range of those previously reported. In their
279 review, Marescaux et al. (2016) reported clearance rate values ranging from 29 mL ind⁻¹ h⁻¹ (Boltovskoy et al.,
280 1995) to 3,252 mL ind⁻¹ h⁻¹ (Viergutz et al., 2012) in *C. fluminea*. This wide range suggests plasticity in the filtration
281 activity of *C. fluminea* in response to environmental changes. Less information is available in the literature for *L.*
282 *ovata*, but recent studies highlighted the high filtration rate capacity of other *Lampsilis* species (*L. fasciola*
283 (Rafinesque, 1820) and *L. siliquoidea* (Barnes, 1823), 250-2,000 mL ind⁻¹ h⁻¹) compared to other freshwater mussels
284 species such as *Villosa iris* (Lea, 1829) and *Ligumia nasuta* (Say, 1817) of similar size (18-400 mL ind⁻¹ h⁻¹ and
285 150-500 mL ind⁻¹ h⁻¹, respectively; Mistry & Ackerman, 2018; Tuttle-Raycraft & Ackerman, 2018, 2019). Although
286 we used smaller individuals of *L. ovata* than the studies performed on other *Lampsilis* species (14-19 mm shell
287 length in the present study vs. ~50-120 mm shell length for previous studies), we confirmed the high filtration
288 capacity of this species seen in previous studies.

289 While we found filtration rates of the two studied species are highly dependent on environmental conditions,
290 interestingly, when only the organic biomass of bivalves is considered in the calculation of the rate clearances, *C.*

291 *fluminea* systematically showed a higher filtration capacity than that of *L. ovata*. This observation is true regardless
292 of the temperature, light and food conditions that we tested. This result demonstrates the importance of taking into
293 account the organic mass (sometimes expressed in g of C, e.g. Marescaux et al., 2016) in the calculation of clearance
294 rates in bivalves. Bivalve clearance rates can vary in proportion to gill surface area (Galbraith et al., 2009). Payne &
295 Miller (1995) found that the allometric relationship between gill surface area (GA, mm²) and shell length (SL, mm)
296 for *C. fluminea* can be approximately described by a linear relationship based on a regression model: $GA = 63.2 +$
297 $0.78 \times SL^2$. Similarly, Galbraith et al. (2009) found in four species of Unionids that, on average, $GA = 2.21 \times SL$.
298 Using these equations, we estimated that gill surface areas were $285 \pm 20 \text{ mm}^2$ in *C. fluminea* used in the present
299 study and only $36 \pm 3 \text{ mm}^2$ in *L. ovata*. Although these estimates need to be confirmed by measurements, such
300 differences suggest that clearance rates performances of *C. fluminea* may be explained, at least partially, by a large
301 gill surface area compared to Unionids.

302 Effects of numerous factors on the clearances rates of *Corbicula* and Unionidae species have been studied including
303 abiotic factors such as temperature (Aldridge et al., 1995; Lim et al., 2005; Loayza-Muro & Elias-Letts, 2007), pH,
304 water velocity, and turbidity (Ackerman, 1999; Loayza-Muro & Elias-Letts, 2007; Tuttle- Raycraft & Ackerman,
305 2019), and biotic factors including food quality and quantity (Sprung & Rose, 1988), and life-history traits such as
306 reproduction and growth cycle (Hornbach et al., 1984; Viergutz et al., 2012). However, the interactions between the
307 different factors and their relative contribution in clearance rates remain largely unknown. In the present study, we
308 found that contribution of each tested variable is species dependent. While temperature was the main variable
309 affecting the clearance rates with virtually no change related to food concentration in *C. fluminea*, food
310 concentration plays a major role in clearance with a limited role of temperature in *L. ovata*. For both species, the
311 effects of light remain limited although changes in light conditions did affect the clearance rate in *C. fluminea*. Such
312 findings highlight the complex interactions between abiotic and biotic factors that potentially affect bivalve
313 clearance rates.

314 Among abiotic factors, temperature is key because bivalves are poikilothermic organisms, responding to temperature
315 changes through behavioral and physiological adaptations. For example, Rodland et al. (2008) found that
316 temperature (from 10 to 34 °C) affects frequency and duration of shell closure in two freshwater mussels: *Anodonta*
317 *cygnea* (Linnaeus, 1758) and *Margaritifera falcata* (Gould, 1850), with a duration of intervals of valve closure
318 decreasing in both species as temperature rises. These results are in accordance with Block et al. (2013) who found

319 that latency in valve opening and foot extension was highest at the lowest experimental temperature (10 °C)
320 compared to the other experimental treatments (i.e. 20 and 30 °C) in their study. Regarding the effects of
321 temperature on bivalve metabolism, Vidal et al. (2002) highlighted that detoxification mechanisms in *C. fluminea*
322 are affected by change in temperatures (10 and 20 °C). In this study, we found that temperature effects on clearance
323 rates were species-dependent and dependent on light and food conditions. This is especially true for *C. fluminea*:
324 while at low food concentration (i.e. 10,000 cells mL⁻¹), no effect of temperature was found, we found an increase in
325 clearance rates at increasing temperatures (from 10 to 25 °C) at the high food concentration (i.e. 100,000 cells mL⁻¹).
326 These results are in accordance with Lim et al. (2005) who found that clearance rates of *C. fluminea* remained close
327 to 0 at 5 °C, increased linearly with water temperature up to approx. 25 °C, and drastically fell at temperatures above
328 25 °C. This suggests that the range of temperatures we studied covers the thermal preference of this species. For *L.*
329 *ovata*, we found a similar pattern in response to temperature changes. Such results are consistent with the field study
330 of Vanderploeg et al. (1995) who found a positive non-linear relationship between filtration rates in *L. radiata*
331 (Gmelin, 1791) and temperature measured in their habitat (from 8 to 25 °C). In a recent laboratory study, Malish &
332 Woolnough (2019) found that clearance rates of *L. cardium* were higher in a trial performed in June (25.0-26.5 °C)
333 than during the same experiment performed in May (17.0-18.5 °C). In the present study, the effects of temperature
334 on the filtration activity of the studied species were highlighted through the calculations of temperature coefficient
335 (Q₁₀). Overall, the average Q₁₀ was 2.12 for *C. fluminea* compared to 1.56 for *L. ovata*. Nevertheless, Q₁₀ was
336 strongly variable depending on the experimental conditions (i.e. food availability and light intensity) with values
337 ranging from 1.08 to 4.81 for *C. fluminea* and from 1.25 to 3.31 for *L. ovata*. These results suggest that the intensity
338 of the effects of temperature is modulated by other environmental conditions.

339 For both species, the effects of temperature in the two food conditions were light-dependent. Effects of temperature
340 were stronger in the dark (i.e. $0 \pm 0 \mu\text{mol m}^{-2} \text{s}^{-1}$) than in the light (i.e. $19 \pm 2 \mu\text{mol m}^{-2} \text{s}^{-1}$). These results are in line
341 with field observations showing that seasonal differences in clearance rates of freshwater bivalves were not
342 necessarily only related to temperature changes but also to the combined effects with other factors (Viergutz et al.,
343 2012). Here we show that the effects of temperature on clearance rate differ between the two studied species and are
344 also closely related to light and to food availability.

345 There is limited information available regarding the effects of light on clearance rates in freshwater bivalves;
346 however, there are a few relevant studies. For example, Ortmann & Grieshaber (2003) used gaping activity in *C.*

347 *fluminea* to estimate filtration activity in the field and reported a circadian rhythm with valves being closed in the
348 morning but open in the afternoon. In addition, Duchini et al. (2015) found that *Limnoperna fortunei* (Dunker, 1857)
349 is more mobile in darkness than in light. Kobak & Nowacki (2007) found that *D. polymorpha* shows light avoidance
350 behavior. Hills et al. (2020) found higher clearance rates in the dark for *C. fluminea* than *Utterbackia imbecillis*
351 (Say, 1829). The present study confirmed differences observed in the effects of light in clearance rates between *C.*
352 *fluminea* and Unionidae species.

353 Food quantity is one of the most studied biotic factors in the assessment of filtration activity in freshwater bivalves
354 (e.g. Roper & Hickey, 1995; Lim et al., 2005; Gatenby et al., 2013). In the present study, we found that clearance
355 rates for *C. fluminea* are overall higher at the highest concentration of food tested. Interestingly, we also found that
356 the effects of temperature on clearance rates are most pronounced at the highest concentration of food tested. This
357 can be related to the reduced or even zero filtration observed whatever the temperature in a large proportion of
358 individuals exposed to a low concentration of food. Altogether these results suggest that *C. fluminea* is efficient in
359 regulating its filtering activity according to the available food resource and that the filtration activity is a trade-off
360 between the energy expended and that acquired by the ingestion of filtered particles (Ortmann & Grieshaber, 2003).
361 Interestingly, an opposite trend was observed in *L. ovata* with lower clearance rates measured at the highest food
362 concentration. To our knowledge, there is no similar study in the literature for this species. We assume that such
363 difference may be related to lower energy requirements of individuals of *L. ovata* (~ 0.5 g) compared to *C. fluminea*
364 whose total weight is 4-fold higher (~ 2 g). Thus, the energy requirements are likely to be covered more quickly for
365 *L. ovata* when the availability of food is high. Nevertheless, further investigations assessing the metabolism of these
366 species through a wide range of temperatures and food concentrations are needed to support this assumption.

367 In North American freshwater ecosystems, native Unionids coexist with exotic bivalve species such as *C. fluminea*
368 and *Dreissena polymorpha* (Pallas, 1771). The threatened status of Unionid mussels has been attributed, along with
369 other factors such as space and habitat availability, to food competition with exotic and invasive bivalve species
370 (e.g., Ferreira-Rodríguez et al., 2018; Strayer & Malcom, 2018). Although experimental evidence of this
371 competition between species is rare in the literature thus far, field investigations have highlighted the declines of
372 Unionid populations due to exploitative competition for food after invasion by *Corbicula* and *Dreissena* species
373 (Burlakova et al., 2014; Strayer & Malcom, 2007). Comparative studies on food preferences in Unionids and
374 *Corbicula* and *Dreissena* have demonstrated that these bivalves compete for the same food resources, and this may

375 result in the reduced availability of preferred and nutritious food items for Unionid mussels. In the present study, we
376 highlighted that *C. fluminea* has a great ability to modulate filtration rate depending on the environmental conditions
377 presumably to optimize food intake. These results are not surprising considering that *Corbicula* has a much faster
378 growth rate than Unionids, which have longer lifespans and later sexual maturation (Anthony et al., 2001), requiring
379 high energy.

380 In combination with food competition, our findings suggest that differential responses to changing environmental
381 conditions in filtration rate between *Corbicula fluminea* and native mussels may be key to understanding the decline
382 of Unionids populations after invasions by Asian clams. Nevertheless, climate change may strongly affect current
383 coexistence and competitive relationships between Unionids and *Corbicula* species. Indeed, we found that *C.*
384 *fluminea* filtration is strongly driven by temperature. Thus, it is crucial to take temperature into account when
385 discussing food competition, because the metabolic rate, and thus the requirement for food, generally increases with
386 rising temperature in bivalves (Vohmann et al., 2010). An increase of 5°C results in an approximately two-fold
387 increase in energy demand in *C. fluminea* (Ortmann & Grieshaber 2003). Increasing temperature under limited food
388 conditions can lead to a negative energy balance and thus to an enhanced starvation (Vohmann et al., 2010).
389 Compared to many other bivalves, *C. fluminea* is less able to compensate in energy losses at high temperatures by
390 decreasing its metabolic activity (McMahon 2002). Although less information is available regarding metabolism in
391 Unionids, such findings illustrate that climate change may also change filtration in bivalves and ultimately affect
392 competition between species.

393

394 **Conclusion**

395 In the present study, we confirmed through multi-parameter laboratory experiments performed in controlled
396 conditions that complex interactions between abiotic and biotic factors affect freshwater bivalve filtration activity.
397 Overall, clearance rates are highly variable in *C. fluminea* and presumably regulated by additional untested factors,
398 suggesting high plasticity in filtration of this species. Such findings are useful to understand field observations
399 where temporal variations of clearance rates in bivalves are often difficult to link to a unique environmental
400 variable. Furthermore, our study provides data for modeling bivalve filtration in stream systems highlighting that
401 filtration activity is not constant through time and potentially highly variable between individuals especially in *C.*
402 *fluminea*. Refining models by using such data may avoid filtration rate overestimation, particularly when clams

403 switch to pedal feeding at low phytoplankton concentrations (Marescaux et al., 2016). Such findings help to properly
404 evaluate the potential of such organisms to mitigate/bioremediate contamination of anthropogenic origins in such
405 fragile ecosystems.

406

407 **Authors Contributions**

408 S.P. and A.H. designed and performed the experiments and analyzed samples. S.P. performed statistical analyses,
409 interpreted the results and wrote the first draft of the paper. T.J.M. supervised the experiments. T.J.M. and A.H.
410 corrected and improved the manuscript.

411

412 **Data Availability Statement**

413 Data are available upon personal request.

414

415 **Conflict of Interest Statement**

416 Authors declare no conflict of interest.

417

418 **ORCID**

419 *Simon Pouil* <https://orcid.org/0000-0003-1531-0362>

420 *Amber Hills* <https://orcid.org/0000-0002-5249-3033>

421 *Teresa J. Mathews* <https://orcid.org/0000-0001-6780-1142>

422 **References**

- 423 Ackerman, J. D. 1999. Effect of velocity on the filter feeding of dreissenid mussels (*Dreissena polymorpha* and
424 *Dreissena bugensis*): Implications for trophic dynamics. Canadian Journal of Fisheries and Aquatic
425 Sciences 56: 1551-1561.
- 426 Aldridge, D. W., Payne, B. S., & A. C. Miller, 1995. Oxygen consumption, nitrogenous excretion, and filtration
427 rates of *Dreissena polymorpha* at acclimation temperatures between 20 and 32 °C. Canadian Journal of
428 Fisheries and Aquatic Sciences 52: 1761-1767.
- 429 Baker, S. M., & D. J. Hornbach, 2001. seasonal metabolism and biochemical composition of two unionid mussels,
430 *Actinonaias ligamentina* and *Amblema plicata*. Journal of Molluscan Studies 67: 407-416.
- 431 Bianchi, V. A., Castro, J. M., Rocchetta, I., Bieczynski, F., & C. M. Luquet, 2014. Health status and bioremediation
432 capacity of wild freshwater mussels (*Diplodon chilensis*) exposed to sewage water pollution in a glacial
433 Patagonian lake. Fish & Shellfish Immunology 37: 268-277.
- 434 Block, J. E., Gerald, G. W., & T. D. Levine, 2013. Temperature effects on burrowing behaviors and performance in
435 a freshwater mussel. *Journal of Freshwater Ecology*, 28(3), 375-384. DOI:
436 10.1080/02705060.2013.767218.
- 437 Boltovskoy, D., Izaguirre, I., & N. Correa, 1995. Feeding selectivity of *Corbicula fluminea* (Bivalvia) on natural
438 phytoplankton. Hydrobiologia 312: 171-182.
- 439 Burlakova, L. E., Tulumello, B. L., Karatayev, A. Y., Krebs, R. A., Schloesser, D. W., Paterson, W. L., Griffith, T.
440 A., Scott, M. W., Crail, T., & D. T. Zanatta, 2014. Competitive replacement of invasive congeners may
441 relax impact on native species: Interactions among zebra, quagga, and native Unionid mussels. PLOS One
442 9(12): e114926.
- 443 Coughlan, J. 1969. The estimation of filtering rate from clearance of suspensions. Marine Biology 2: 356-358.
- 444 Domingues, A., Rosa, I. C., Pinto da Costa, J., Rocha-Santos, T. A. P., Gonçalves, F. J. M., Pereira, R., & J. L.
445 Pereira, 2020. Potential of the bivalve *Corbicula fluminea* for the remediation of olive oil wastewaters.
446 Journal of Cleaner Production 252: 119773.
- 447 Duchini, D., Boltovskoy, D., & F. Sylvester, 2015. Detachment, displacement and reattachment activity in a
448 freshwater byssate mussel (*Limnoperna fortunei*): The effects of light, temperature and substratum
449 orientation. Biofouling 31: 599-611.

450 Ferreira-Rodríguez, N., Sousa, R., & I. Pardo, 2018. Negative effects of *Corbicula fluminea* over native freshwater
451 mussels. *Hydrobiologia* 810(1): 85-95.

452 Galbraith, H. S., Frazier, S. E., Allison, B., Vaughn, C.C., 2009. Comparison of gill surface morphology across a
453 guild of suspension-feeding unionid bivalves, *Journal of Molluscan Studies* 75(2): 103-107.

454 Gatenby, C. M., Kreeger, D., Patterson, M., Marini, M., & R. Neves, 2013. Clearance rates of *Villosa iris* (Bivalvia:
455 Unionidae) fed different rations of the alga *Neochloris oleoabundans*. *Freshwater Mollusk Biology and*
456 *Conservation* 16: 9-20.

457 Gatenby, C. M., Neves, R. J., & B. C. Parker, 1996. Influence of sediment and algal food on cultured juvenile
458 freshwater mussels. *Journal of the North American Benthological Society* 15: 597-609.

459 Groemping, U. 2007. Relative importance for linear regression in R: The package *relaimpo*. *Journal of Statistical*
460 *Software* 17: 1-27.

461 Guillard, R. R. 1975. Culture of phytoplankton for feeding marine invertebrates. In Chanley, M. H., Matoira, H. &
462 L. Walter (eds) *Culture of Marine nvertebrate Animals*. Springer, Boston: 29-60.

463 Haag, W. R. 2012. *North American Freshwater Mussels: Natural history, ecology, and conservation*. Cambridge
464 University Press, Cambridge, New York.

465 Hills, A., Pouil, S., Hua, D. & T. J. Mathews, 2020. Effects of light on the clearance rates of freshwater bivalves: the
466 exotic *Corbicula fluminea* and the native *Utterbackia imbecillis*. *Aquatic Ecology*(), 1-8.
467 <https://doi.org/10.1007/s10452-020-09793-7>

468 Hornbach, D. J., Way, C., Wissing, T. E., & A. J. Burky, 1984. Effects of particle concentration and season on the
469 filtration rates of the freshwater clam, *Sphaerium striatinum* Lamarck (Bivalvia: Pisidiidae). *Hydrobiologia*
470 108: 83-96.

471 Howard, J. K., & K. M. Cuffey, 2006. The functional role of native freshwater mussels in the fluvial benthic
472 environment. *Freshwater Biology* 51: 460-474.

473 Kobak, J., & P. Nowacki, 2007. Light-related behaviour of the zebra mussel (*Dreissena polymorpha*, Bivalvia).
474 *Fundamental and Applied Limnology* 169: 341-352.

475 Kryger, J., & H. U. Riisgård, 1988. Filtration rate capacities in 6 species of European freshwater bivalves. *Oecologia*
476 77: 34-38.

477 Lim, K.-H., Shin, H.-C., & J.-S. Yang, 2005. The influence of water temperature and food concentration on the
478 filtration rates of the Asiatic clam, *Corbicula fluminea*. The Korean Journal of Malacology 21: 19-24.

479 Loayza-Muro, R., & Elias-Letts, R. 2007. Responses of the mussel *Anodontites trapesialis* (Unionidae) to
480 environmental stressors: Effect of pH, temperature and metals on filtration rate. Environmental Pollution
481 149: 209-215.

482 Malish, M. C., & D. A. Woolnough, 2019. Varied physiological responses of *Amblema plicata* and *Lampsilis*
483 *cardium* exposed to rising temperatures. Freshwater Science 38: 842-855.

484 Marescaux, J., Falisse, E., Lorquet, J., Van Doninck, K., Beisel, J.-N., & J.-P. Descy, 2016. Assessing filtration rates
485 of exotic bivalves: Dependence on algae concentration and seasonal factors. Hydrobiologia 777: 67-78.

486 McMahon, R. F. 2002. Evolutionary and physiological adaptations of aquatic invasive animals: r selection versus
487 resistance. Canadian Journal of Fisheries and Aquatic Sciences 59: 1235-1244.

488 Mistry, R., & J. D. Ackerman 2017. Algal flux affects the clearance rates of recently metamorphosed freshwater
489 mussels. Aquatic Sciences 79: 139-148.

490 Mistry, R., & J. D. Ackerman 2018. Flow, flux, and feeding in freshwater mussels. Water Resources Research 54:
491 7619-7630.

492 Mukherjee, A., Das, S., Bhattacharya, T., De, M., Maiti, T., & T. Kumar De, 2014. Optimization of phytoplankton
493 preservative concentrations to reduce damage during long-term storage. Biopreservation and Biobanking
494 12: 139-147.

495 Nichols, S. J., Silverman, H., Dietz, T. H., Lynn, J. W., & D. L. Garling, 2005. Pathways of food uptake in native
496 (Unionidae) and introduced (Corbiculidae and Dreissenidae) freshwater bivalves. Journal of Great Lakes
497 Research 31: 87-96.

498 Ortmann, C., & M. K. Grieshaber, 2003. Energy metabolism and valve closure behaviour in the Asian clam
499 *Corbicula fluminea*. Journal of Experimental Biology 206: 4167-4178.

500 Payne, B. S., & A. Miller, 1995. Palp to gill area ratio of bivalves: A sensitive indicator of elevated suspended
501 solids. Regulated Rivers Research & Management 11: 193-200.

502 Pickhardt, P. C., & N. S. Fisher, 2007. Accumulation of inorganic and methylmercury by freshwater phytoplankton
503 in two contrasting water bodies. Environmental Science & Technology 41(1): 125-131.

504 Pigneur, L.-M., Falisse, E., Roland, K., Everbecq, E., Delière, J.-F., Smits, J. S., Van Doninck, K., & J.-P. Descy,
505 2014. Impact of invasive Asian clams, *Corbicula* spp., on a large river ecosystem. *Freshwater Biology* 59:
506 573-583.

507 R Development Core Team 2020. R: A Language and Environment for Statistical Computing. R Foundation for
508 Statistical Computing, Vienna.

509 Riisgård, H. U. 2001. On measurement of filtration rates in bivalves - the stony road to reliable data: Review and
510 interpretation. *Marine Ecology Progress Series* 211: 275-291.

511 Rodland, D. L., Schöne, B. R., Baier, S., Zhang, Z., Dreyer, W., & N. A. Page, 2008. Changes in gape frequency,
512 siphon activity and thermal response in the freshwater bivalves *Anodonta cygnea* and *Margaritifera*
513 *falcata*. *Journal of Molluscan Studies* 75: 51-57.

514 Roper, D. S., & C. W. Hickey, 1995. Effects of food and silt on filtration, respiration and condition of the freshwater
515 mussel *Hyridella menziesi* (Unionacea: Hyriidae): Implications for bioaccumulation. *Hydrobiologia* 312:
516 17-25.

517 Rosa, I. C., Costa, R., Goncalves, F., & J. L. Pereira, 2014. Bioremediation of metal-rich effluents: Could the
518 invasive bivalve *Corbicula fluminea* work as a biofilter? *Journal of Environmental Quality* 43: 1536-1545.

519 Salerno, J., Gillis, P. L., Bennett, C. J., Sibley, P. K., & R. S. Prosser, 2018. Investigation of clearance rate as an
520 endpoint in toxicity testing with freshwater mussels (Unionidae). *Ecotoxicology and Environmental Safety*
521 163, 165-171.

522 Sprung, M., & U. Rose, 1988. Influence of food size and food quantity on the feeding of the mussel *Dreissena*
523 *polymorpha*. *Oecologia* 77: 526-532.

524 Strayer, D. L., & H. M. Malcom, 2018. Long-term responses of native bivalves (Unionidae and Sphaeriidae) to a
525 *Dreissena* invasion. *Freshwater Science* 37(4): 697-711.

526 Strayer, D. L., & H. M. Malcom, 2007. Effects of zebra mussels (*Dreissena polymorpha*) on native bivalves: The
527 beginning of the end or the end of the beginning? *Journal of the North American Benthological Society* 26:
528 111-122.

529 Tankersley, R. A. 1996. Multipurpose gills: effect of larval brooding on the feeding physiology of freshwater
530 unionid mussels. *Invertebrate Biology* 115: 243-255.

531 Tankersley, R. A., & R. V. Dimock Jr, 1993. The effect of larval brooding on the filtration rate and particle-retention
532 efficiency of *Pyganodon cataracta* (Bivalvia: Unionidae). Canadian Journal of Zoology 71: 1934-1944.

533 Tuttle-Raycraft, S., & J. D. Ackerman, 2018. Does size matter? Particle size vs. quality in bivalve suspension
534 feeding. Freshwater Biology 63: 1560-1568.

535 Tuttle- Raycraft, S., & J. D. Ackerman, 2019. Living the high turbidity life: The effects of total suspended solids,
536 flow, and gill morphology on mussel feeding. Limnology and Oceanography 64: 2526-2537.

537 Vanderploeg, H. A., Liebig, J. R., & T. F. Nalepa, 1995. From picoplankton to microplankton: Temperature-driven
538 filtration by the unionid bivalve *Lampsilis radiata siliquoidea* in Lake St. Clair. Canadian Journal of
539 Fisheries and Aquatic Sciences 52: 63-74.

540 Vaughn, C. C. 2018. Ecosystem services provided by freshwater mussels. *Hydrobiologia* 810: 15-27.

541 Vidal, M. L., Basseres, A., & J. F. Narbonne, 2002. Influence of temperature, pH, oxygenation, water-type and
542 substrate on biomarker responses in the freshwater clam *Corbicula fluminea* (Muller). Comparative
543 Biochemistry and Physiology C-Toxicology & Pharmacology 132: 93-104.

544 Viegutz, C., Linn, C., & M. Weitere, 2012. Intra- and interannual variability surpasses direct temperature effects on
545 the clearance rates of the invasive clam *Corbicula fluminea*. Marine Biology 159: 2379-2387.

546 Vohmann, A., Borcherdig, J., Kureck, A., Bij de Vaate, A., Arndt, H., & M. Weitere, 2010. Strong body mass
547 decrease of the invasive clam *Corbicula fluminea* during summer. Biological Invasions 12(1): 53-64.

548 Zarauz, L., & X. Irigoien, 2008. Effects of Lugol's fixation on the size structure of natural nano-microplankton
549 samples, analyzed by means of an automatic counting method. Journal of Plankton Research 30: 1297-
550 1303.

551 Table 1. Summary of water quality parameters measured for the four temperature conditions during the experiment.

552 Values are means \pm SD. DO: dissolved oxygen, TDS: total dissolved solids.

| Parameters | Experimental treatment | | | |
|----------------------------------------------------|------------------------|-------------------|-------------------|-------------------|
| | 10°C | 15°C | 20°C | 25°C |
| Temperature (°C) | 10.3 \pm 0.1 | 14.9 \pm 0.2 | 20.3 \pm 0.2 | 24.9 \pm 0.2 |
| pH | 8.02 \pm 0.10 | 8.05 \pm 0.11 | 8.08 \pm 0.12 | 8.26 \pm 0.13 |
| DO saturation (%) | 92.2 \pm 2.3 | 91.9 \pm 2.4 | 91.8 \pm 2.5 | 92.4 \pm 2.0 |
| DO (mg L ⁻¹) | 9.82 \pm 0.42 | 8.96 \pm 0.28 | 8.03 \pm 0.25 | 7.49 \pm 0.27 |
| Conductivity A (μ S cm ⁻¹) | 162 \pm 16 | 164 \pm 18 | 173 \pm 19 | 178 \pm 24 |
| TDS (mg L ⁻¹) | 85 \pm 11 | 85 \pm 11 | 88 \pm 12 | 89 \pm 12 |
| NH ₃ -N (mg L ⁻¹) | 0.05 \pm 0.04 | 0.03 \pm 0.02 | 0.02 \pm 0.03 | 0.02 \pm 0.02 |
| NO ₂ ⁻ (mg L ⁻¹) | 0.014 \pm 0.019 | 0.019 \pm 0.015 | 0.005 \pm 0.004 | 0.004 \pm 0.008 |

553

554 Table 2. Three-way mixed analysis of variance (ANOVA) results. Effect of light (i.e. ‘dark’ and ‘light’ within-
 555 subjects factor) and temperature (10, 15, 20 and 25 °C) on the clearance rates of bivalves of two different species (*C.*
 556 *fluminea* and *L. ovata*) at two different concentrations of food: ‘low’ food (i.e. 10,000 cells mL⁻¹) and ‘high’ food
 557 (i.e. 100,000 cells mL⁻¹) experiments.

| Effect | Degrees of freedom | | Low food | | High food | |
|----------------------------------|--------------------|-------|----------|----------------|-----------|----------------|
| | numDF | denDF | F-value | p-value | F-value | p-value |
| Temperature | 3 | 56 | 6.36 | < 0.001 | 25.31 | < 0.001 |
| Species | 1 | 56 | 77.77 | < 0.001 | 20.81 | < 0.001 |
| Light | 1 | 56 | 0.01 | 0.913 | 37.54 | < 0.001 |
| Temperature x Species | 3 | 56 | 5.29 | 0.003 | 3.29 | 0.027 |
| Temperature x Light | 3 | 56 | 1.77 | 0.164 | 2.38 | 0.079 |
| Species x Light | 1 | 56 | 0.41 | 0.525 | 2.77 | 0.102 |
| Temperature x Species x Light | 3 | 56 | 0.74 | 0.535 | 1.71 | 0.176 |

558

559 Table 3. Generalized linear model (GLM) for *C. fluminea* and *L. ovata*. Effects of each explanatory variable (i.e.
 560 food, temperature and light) on bivalve clearance rates. Coefficient of determination R^2 was 0.16 for *C. fluminea* and
 561 0.67 for *L. ovata*.

| Predictor | <i>C. fluminea</i> | | <i>L. ovata</i> | |
|---------------|----------------------|-------------------|----------------------|-------------------|
| | Estimate (\pm SD) | p-value | Estimate (\pm SD) | p-value |
| Intercept | 41.22 (46.06) | 0.374 | -118.10 (35.01) | 0.001 |
| Temperature | 10.53 (2.62) | < 0.001 | 14.01 (2.07) | < 0.001 |
| Food (Low) | -4.41 (28.48) | 0.877 | 334.25 (25.14) | < 0.001 |
| Light (Light) | -58.03 (24.59) | 0.022 | -46.57 (20.81) | 0.029 |

562

563 Figure captions

564

565 **Fig. 1.** Diagram of the protocol used to test effects of temperatures (10, 15, 20 and 25°C) on the clearance rates of
566 *Corbicula fluminea* and *Lampsilis ovata*. Experiments were repeated the same day using the same individuals at two
567 light intensities (i.e. ‘dark’: $0 \pm 0 \mu\text{mol m}^{-2} \text{s}^{-1}$ and ‘light’: $19 \pm 2 \mu\text{mol m}^{-2} \text{s}^{-1}$) first at ‘low’ concentration of food
568 (i.e. 10,000 cells mL^{-1}) and then, one week later at ‘high’ concentration of food (i.e. 100,000 cells mL^{-1}) after
569 acclimation to a 10-fold higher food daily ration.

570

571 **Fig. 2.** Effects of temperatures (10, 15, 20 and 25°C) on the clearance rates, expressed in g of whole body wet
572 weight (A) and g of soft parts wet weight (B), of *Corbicula fluminea* and *Lampsilis ovata* exposed to two light
573 intensities (i.e. ‘dark’: $0 \pm 0 \mu\text{mol m}^{-2} \text{s}^{-1}$ and ‘light’: $19 \pm 2 \mu\text{mol m}^{-2} \text{s}^{-1}$) and low concentration of food (i.e. 10,000
574 cells mL^{-1}). Symbols denote significant differences. Statistical differences printed in panel A are identical for panel
575 B.

576

577 **Fig. 3.** Percent of individuals filtering during the ‘low food’ (i.e. 10,000 cells mL^{-1}) experiment for *Corbicula*
578 *fluminea* and *Lampsilis ovata* exposed to two light intensities (i.e. ‘dark’: $0 \pm 0 \mu\text{mol m}^{-2} \text{s}^{-1}$ and ‘light’: $19 \pm 2 \mu\text{mol}$
579 $\text{m}^{-2} \text{s}^{-1}$) and four temperatures (10, 15, 20 and 25 °C).

580


581 **Fig. 4.** Effects of temperature (10, 15, 20 and 25 °C) on the clearance rates, expressed in g of whole body wet
582 weight (A) and g of soft parts wet weight (B), of *Corbicula fluminea* and *Lampsilis ovata* exposed to two light
583 intensities (i.e. ‘dark’: $0 \pm 0 \mu\text{mol m}^{-2} \text{s}^{-1}$ and ‘light’: $19 \pm 2 \mu\text{mol m}^{-2} \text{s}^{-1}$) and high concentration of food (i.e.
584 100,000 cells mL^{-1}). Symbols denote significant differences. Statistical differences printed in panel A are identical
585 for panel B.

586


587 **Fig. 5.** Percent of individuals filtering during the ‘high’ food (i.e. 100,000 cells mL^{-1}) experiment for *Corbicula*
588 *fluminea* and *Lampsilis ovata* exposed to two light intensities (i.e. ‘dark’: $0 \pm 0 \mu\text{mol m}^{-2} \text{s}^{-1}$ and ‘light’: $19 \pm 2 \mu\text{mol}$
589 $\text{m}^{-2} \text{s}^{-1}$) and four temperatures (10, 15, 20 and 25 °C).

590


591 **Fig. 6.** Relative contribution of each explanatory variable (i.e. food, temperature and light) on the clearance rates of
592 each species: *Corbicula fluminea* and *Lampsilis ovata*. Values are means with 95% bootstrap confidence intervals.




593 Figure 1




594 Figure 2




595 Figure 3



596 Figure 4



597 Figure 5



598 Figure 6