



HAL
open science

Pathogenic Leptospira and water quality in African cities: A case study of Cotonou, Benin

Honoré Houéménou, Philippe Gauthier, Gualbert Houemenou, Daouda Mama, Abdoukarim Alassane, Akilou Socohou, Henri-Joël Dossou, Sylvestre Badou, Mathieu Picardeau, Sarah Tweed, et al.

► To cite this version:

Honoré Houéménou, Philippe Gauthier, Gualbert Houemenou, Daouda Mama, Abdoukarim Alassane, et al.. Pathogenic Leptospira and water quality in African cities: A case study of Cotonou, Benin. Science of the Total Environment, 2021, 774, pp.145541. 10.1016/j.scitotenv.2021.145541 . hal-03166929

HAL Id: hal-03166929

<https://hal.inrae.fr/hal-03166929>

Submitted on 9 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Pathogenic *Leptospira* and water quality in African cities: a case study of Cotonou, Benin

Authors: Houéménou Honoré^{1,2*}, Gauthier Philippe³, Houéménou Gualbert⁴, Mama Daouda², Alassane Abdoukarim², Socohou Akilou², Dossou Henri-Joël^{4,5}, Badou Sylvestre⁴, Picardeau Mathieu⁶, Tweed Sarah⁷, Leblanc Marc¹ and Dobigny Gauthier^{3,4}

Affiliations

¹ University of Avignon, Hydrogeology Laboratory, UMR EMMAH, Avignon, France

² INE, Laboratory of Applied Hydrology, University of Abomey-Calavi, 01 B.P. 4521 Cotonou, Benin

³ IRD, CBGP (UMR IRD-INRAE-Cirad-Montpellier SupAgro), MUSE, Montferrier-sur-Lez, France

⁴ Laboratory of Research in Applied Biology, University of Abomey-Calavi, EPAC, Cotonou, Benin

⁵ IGATE, Université d'Abomey-Calavi, Cotonou, Benin

⁶ Institut Pasteur de Paris, Laboratoire des Spirochètes, Paris, France

⁷ G-EAU (UMR IRD-IRSTEA-Cirad-Agro Paris Tech-Montpellier SupAgro), Montpellier, France

* **corresponding author:** h.houemenou@yahoo.com

1 1. INTRODUCTION

2 Leptospirosis is a bacterial disease caused by spirochetes of the genus *Leptospira*. Leptospire
3 infect and develop inside renal tubules of a variety of Vertebrates, essentially mammals, notably rodents,
4 which excrete the pathogen into the external environment. Humans are infected following contact with
5 contaminated water or humid soils. Symptoms range from null to very severe and death (reviewed in
6 Haake & Levett 2015). It is considered that one million cases occur annually in the world, among which
7 60,000 have fatal issues (Costa et al. 2015). Leptospirosis has been mainly associated with rice
8 agriculture and breeding as well as water recreational activities (Mwachui et al. 2015). Flooding
9 episodes may also trigger local leptospiral epidemics (e.g., Thaipadungpanit et al. 2013), with one out
10 of eight flood event-associated epidemics being due to leptospirosis (Cann et al. 2013). In addition, more
11 and more evidence suggest that the disease may occur in urban settings, especially socio-economically
12 disadvantaged, informal and poorly sanitized ones, such as slums (e.g., Ko et al. 1999; Cornwall et al.
13 2016), sometimes with higher waterborne leptospire concentrations compared to surrounding rural
14 areas (e.g. in Peru, Ganoza et al. 2006). As a consequence, ongoing urbanization together with
15 multiplying extreme climatic events are expected to increase the risk of human leptospirosis (Lau et al.
16 2010). This may be particularly true in Africa where the disease is present but yet remains poorly
17 documented (reviews in de Vries et al. 2014; Allan et al. 2015). In particular, we are aware of no study
18 focusing on soil- or waterborne leptospire in Africa (see Tab. 2 in the review by Bierque et al. 2020).

19 The so-called « Abidjan-Lagos corridor » (ALC) is a 700km-long conurbation that sprawls
20 along the West African Atlantic coast. It currently houses >25 million people who live within and in the
21 surroundings of large adjacent cities such as Lagos (11.8 million inhabitants), Porto-Novo (570,000
22 inhabitants), Cotonou (1.5 million inhabitants), Lomé (1.7 million inhabitants), Accra (4.4 million
23 inhabitants) and Abidjan (4.7 million inhabitant) (2015 data; OECD/Africapolis project,
24 www.africapolis.org, 2020). It is expected to reach ca. 34 million city dwellers by 2025 (UN Habitat,
25 2014). This very rapid and mostly uncontrolled urbanization translates into the creation and expansion
26 of vast socially disadvantaged areas where pollution, access to basic services (e.g., health care,
27 education, transport, sanitation, waste management) and acceptable housing conditions are rare, thus

28 raising important environmental and health issues. In addition, the ALC comprises many lakes,
29 mangroves and swamps which form a dense hydrographic network. Together with a subequatorial
30 climate, low altitude and flatness, this makes this West African coastal region extremely susceptible to
31 flooding events. For instance, 43% of Cotonou, Benin, is flooded one to two months a year either
32 following Lake Nokoué overflows or rain accumulation in shallows. Such episodes directly affect
33 200,000 inhabitants and indirectly impact many others through service interruption, resource
34 unavailability or degraded water quality (PCUG3C, 2010; Houéménou et al. 2019b). In addition, they
35 are often enhanced by poor drainage due to defective or crowded sanitation network as well as anarchic
36 land use (PCUG3C, 2010).

37 These conditions as well as the omnipresence of anthropophilous rodents (Houéménou et al.
38 2019a) elevate the risk of leptospiral contamination in the ALC (Dobigny et al. 2018). Accordingly, up
39 to 18.9% of commensal rodents were found pathogenic *Leptospira*-positive in cities from south Benin
40 (Houéménou et al. 2014, 2019a), suggesting that leptospires may massively circulate in the ALC urban
41 environment, though with strong prevalence variations in both space and time (Houéménou et al.,
42 2019a). Mean annual temperatures (ca. 27°C) and rainfalls (ca. 800-1,600mm) are highly favorable to
43 *Leptospira* survival. However, although waters and humid soils appear as the corner stone of human
44 contamination (Barragan et al. 2017; Bierque et al. 2020), little is known about the precise environmental
45 conditions that are compatible with pathogenic *Leptospira* outside of its mammalian host (see for
46 instance Chang et al. 1948; Gordon-Smith & Turner 1961; Khairani-Bejo et al. 2004; Wojcik-Fatla et
47 al. 2014; André-Fontaine et al. 2015; reviewed in Barragan et al. 2017 and Bierque et al. 2020),
48 especially in urban habitats where pollution may be very important (Lapworth et al. 2017). The present
49 study aims are: (i) to confirm the presence of pathogenic leptospires out of their hosts in Cotonou waters,
50 (ii) to determine the physico-characteristics of waters where it was found, and (iii) to compare these
51 physico-characteristics with the chemical spectrum of all waters within the city. Altogether, our
52 preliminary data allow us to explore for the first time the ability of leptospires to evolve in water within
53 an urban polluted habitat in Africa.

55 2. MATERIAL AND METHODS

56

57 2.1 Study area and sampling

58

59 Cotonou town is located in the coastal sandy plain of Benin between the Atlantic Ocean at the
60 south and Lake Nokoué at the north (Fig. 1). The littoral zone in the south of Benin is characterized by
61 a subequatorial climate. The average annual rainfall for Cotonou is 1,300 mm (Yabi and Afouda, 2012).
62 Seasonal variations during the year are marked by a large rainy season from mid-March to mid-July
63 followed by a small rainy season from mid-September to mid-November alternated, respectively, by a
64 small dry season from mid-July to mid-September and a large dry season from mid-November to mid-
65 March.

66 The study area involves 3 districts located in the core city of Cotonou: Ladji, Agla and Saint
67 Jean (Fig. 1a) that are distinct in the sources and duration of inundation and pond waters. Ladji is located
68 at the edge as well as upon the Lake Nokoué, thus displaying houses built both on hard ground and on
69 stilt pegs (Fig. 1b). Lacustrine waters are permanent, but flooding of zones adjacent to the lake only
70 occurs at the end of the rainy season (i.e., September and October) following the rise in lake levels.
71 Therefore, the area has both long-standing lake water and temporary pond waters. Permanent ponds are
72 also present due to dry season groundwater discharge (Houéménou et al. 2019b). Ladji is densely
73 populated, has essentially informal housing development and is a very poor area where even the most
74 basic public services are usually missing. In particular, waste management is almost inexistent, garbage
75 are omnipresent and are even often used as embankment material. Such an environment favors
76 proliferation of rodents, which are abundant and infest 60-100% of households depending on the season
77 (Dobigny et al. 2019).

78 Agla is a recent but rapidly expanding district within a vast lowland that is extensively flooded
79 early in the rainy season (i.e., starting from June) following rainfall accumulation. Similar to Ladji,
80 permanent ponds are also present in Agla due to the extensive low-lying areas. The permanent ponds

81 are formed by groundwater discharge in the dry months, whereas during the winter months the rainfall
82 inputs to the ponds reverse the hydraulic gradients, resulting in groundwater recharge ponds
83 (Houéménou et al. 2019b). The habitat in Agla includes hard-built houses as well as very precarious
84 cabins, with the poorest inhabitants usually gathering around or within these floodable low-lying areas
85 (Fig. 1b). These low-lying areas are also widely used as dumping sites for household waste (Fig. 1b).
86 Rodents are abundant in households, with infestation rates always $\geq 90\%$ (Dobigny et al. 2019).

87 Saint-Jean is an old and formal district that has large open sewers and many houses that are
88 solidly constructed. This district is not floodable *per se*, but large ponds may stand for several days after
89 heavy rains. Rodents are also abundant in ca. 90% of households (Dobigny et al. 2019).

90
91 In 2017 and 2018, the three districts of Cotonou (Agla, Ladjì and Saint Jean; Fig.1b) were
92 monitored for both water quality parameters (physical parameters, major ions and trace elements) and
93 leptospire presence. Water sampling was organized concomitantly (i.e. within the same week) in the
94 three districts during the dry season (March 2017 and February 2018) as well as at the beginning (June
95 2017 and June 2018) and at the end (October 2017) of the rainy season. In total, 193 water samples were
96 collected: 85 in Agla, 53 in Ladjì and 55 in Saint Jean. Eighty-three samples were tap waters and 61, 29,
97 17 and 3 samples were from groundwater wells, temporary ponds, permanent ponds and Lake Nokoué,
98 respectively. This corresponds to 51, 64, 55, 5 and 18 samples collected in March 2017, June 2017,
99 October 2017, February 2018 and June 2018, respectively. The detailed distribution of samples among
100 districts, periods and water types are provided in Table 1. The temporary ponds were differentiated from
101 the permanent ponds by noting locations where the waters were present in the wet season but not in the
102 dry season. Both temporary and permanent ponds were sampled in Ladjì (8 and 6, respectively) and
103 Agla (21 and 11, respectively), but not in St Jean where ponds only last a few days.

104

105 Cartography of each studied area was performed within the same month immediately after each
106 of the five seasonal sampling campaigns. The land use categories and the extent of surface waters

107 (diameter > 3m) were systematically delineated using Open Street Map Tracker v.0.6.11 (Tools Open
108 Street Map) on an Android smartphone. A Geographic Information System was produced under QGIS
109 v.2.14 (QGIS Development team 2016). Surface water superfices were computed using the
110 LandscapeMetric package (option level class) in RStudio v.3.6.1 (RStudio Team 2016).

111

112 **2.2 Screening of waterborne pathogenic *Leptospira***

113

114 **2.2.1 Test of a filtration-based method of water-borne *Leptospira* screening**

115 Published studies on water-borne leptospires usually rely on high-speed centrifugation methods
116 and freezing preservation of field samples (e.g., Ganoza et al. 2006; Munoz-Zanzi et al. 2014; Lall et al.
117 2016; Riedeger et al. 2016; Thibeaux et al. 2017; Sato et al. 2019). These approaches are poorly adapted
118 to field surveys, especially in developing countries where lab facilities are usually hardly available
119 and/or distantly located. In order to adapt protocols of *Leptospira* screening of waters to the latter
120 conditions, we developed a simple filtration-based method that can be used in the field in absence of
121 costly and high-tech materials. To do so, we first experimentally compared the percentage of bacteria
122 recovery from mineral water (EC=180 μ S/cm) using centrifugation on the one hand, and filtration on the
123 other hand. For centrifugation tests, lab water samples were inoculated with various amounts (from
124 36,800 to 3,680,000 cells per sample, each sample in sextuplets) of living culture-originating *Leptospira*
125 *interrogans* serovar Icterohaemorrhagiae strain Verdun (Spirochete Laboratory, Pasteur Institute of Paris,
126 National Reference Center for Leptospirosis and WHO Collaborative Center). Centrifugation consisted
127 in spinning 15mL of inoculated lab water samples at 4,000 rpm during 30 minutes. Water-borne DNA
128 was extracted immediately after centrifugation. Filtration was performed on 500mL of inoculated lab
129 water samples (representing 2,000 to 2,000,000 cells per sample, each sample in triplicates) that were
130 sequentially filtered on 0.45 μ m and 0.1 μ m nitrocellulose membranes. Filters were then stored in ethanol
131 96° for subsequent DNA extraction.

132 Total DNA was extracted from both centrifuged and filtered samples using the Biobasics EZ-
133 10 Spin column Genomic DNA Minipreps kit, and eluted in 50µL of elution buffer. The same amounts
134 of expected bacteria were extracted directly from 20µL of culture medium as a positive control.

135 Leptospire detection and quantification were performed following a probe-based qPCR method
136 that targets a 199bp-long fragment of the *LipL32* gene using a LightCycler® 480 (Roche Diagnostics)
137 in 96-well micro-titre plates with a 10µL final volume for each reaction (see Dobigny et al. 2015). All
138 qPCR investigations were conducted in duplicates. Negative and positive controls were systematically
139 included in each qPCR detection plates. Extraction negative controls consisted in lab water that was
140 filtered and extracted in the same conditions than the inoculated water samples, while negative qPCR
141 controls consisted in molecular grade water used for qPCR mix preparation steps. Positive controls
142 consisted in the qPCR reactions that were used to construct the standard curve (i.e. DNA directly
143 extracted from *Leptospira* culture).

144

145 ***2.2.2 Leptospire screening in field water samples***

146 For each of the 193 samples, one liter was collected during the morning. The 1 L bottles were
147 rinsed at least twice with the sample water before collection. Up to 500ml of water aliquots were filtered
148 for *Leptospira* screening within the same day, once in the lab. All field water samples were first filtered
149 using 5µm filter papers to retain the large suspended particles. Waters were then filtered using 0.45µm
150 filter papers. Clogging sometimes occurred, especially when filtering turbid waters. In such cases,
151 successive filters were used until 500mL in total had been filtered (e.g. filter 1 used to filter, say, 250mL;
152 filter 2 to filter 150mL; filter 3 to filter the last 100mL). Each filter was preserved in an independent
153 tube, thus allowing us to perform independent lysis. Digestion products were then pooled to go through
154 the same single DNA extraction Qiagen© column. DNA extraction and leptospirosis quantification
155 proceeded as described above, except that for each water sample both 5µm and 0.45µm filter papers
156 were pooled for DNA extraction.

157

158 **2.3 Physico-chemical analysis of waters**

159

160 A wide spectrum of physico-chemical parameters were measured for each water sample
161 investigated for *Leptospira*. These included the field parameters temperature, electrical conductivity
162 (EC), dissolved oxygen (DO) and pH, and laboratory analysed parameters stable isotopes, major ions
163 and trace elements. The methods and results of these hydrochemical data are detailed in Houéménou et
164 al. (2019b, 2020).

165 The comparison of physico-parameters of leptospire-positive and leptospire-negative waters was
166 performed using univariate plots. Principal Component Analysis (PCA) was used under XLSTAT
167 (Addinsoft 2019) to explore in a multivariate manner the chemical characteristics of water samples
168 where pathogenic leptospires were found compared to those with no leptospires. In addition, using the
169 same parameters as for the PCA, a hierarchical cluster analysis (CA) was performed to identify
170 homogeneous groups of leptospire-negative pools that have similar chemical characteristics to
171 leptospire-positive pool samples. The degree of association between pond sampling sites is represented
172 by the euclidian distance between groups.

173

174 **3. RESULTS**

175

176 **3.1 Centrifugation vs. filtration-based approaches**

177

178 Our experimental comparison of centrifugation- vs. filtration-based approaches showed that spinning
179 allowed us to recover only $24.45\% \pm 10.14$ of the initial inoculated number of bacteria, while $98.30\% \pm$
180 23.78% was recovered by filtration (see Suppl. Fig. 1), thus demonstrating that the latter method
181 provides an excellent opportunity to survey environmental leptospires. Importantly, the recovered
182 bacteria were unevenly distributed among the $0.45\mu\text{m}$ and $0.1\mu\text{m}$ filter membranes, with $87.50\% \pm$

183 22.26% and $10.80\% \pm 5.36\%$, respectively. Importantly, there was a high retention of leptospires on the
184 0.45 μ m filter papers, which was also observed by Kaboosi et al. (2010). We did not detect any major
185 differences in recovery rates between the different amounts of inoculated bacteria, except for very low
186 initial concentrations that resulted in higher variations between experimental replicates as well as lower
187 recovery rates by filtration (65% of recovery of the inoculum of 2,000 bacteria in 500mL) – which still
188 remained much better than with centrifugation (<10%; Suppl. Fig. 1).

189

190 **3.2 *Leptospira*-positive waters**

191

192 In total, six water samples were found pathogenic *Leptospira*-positive, with Ct ranging from
193 37.3 to 43.6 (i.e., 9 to 282 *Leptospira* genome equivalents per litre; gEq/L). Such Ct values may appear
194 high. However, we considered them as unambiguous true positive signals that clearly differ from all other
195 187 negative samples that displayed perfectly flat fluorescence curves despite up to 60 amplification
196 cycles (data not shown). Though close to (and even sometimes beyond) theoretical detection limit, such
197 high Ct values are regularly observed in rodent samples in which 16S rDNA *Leptospira* sequence can
198 yet be obtained (P. Gauthier & G. Dobigny, unpublished data). However, they probably correspond to
199 quite low bacterial load, thus precluding quantification that would make sense.

200 Five positive samples (9, 18, 44, 270 and 282 gEq/L, respectively) originated from pond water
201 in Agla, whilst a sixth positive sample (109 gEq/L) was from groundwater collected inside a well at
202 Ladji (Table 1). In Agla, *Leptospira*-positive waters were from both temporary and permanent ponds.
203 For temporary ponds the total prevalence was 9.5% (2/21). In comparison, for the permanent ponds the
204 prevalence was 27.3 % (3/11) (Table 1). All positive samples were obtained in June (2 temporary ponds
205 in Agla and one groundwater well in Ladji in June 2017; 3 permanent ponds in Agla in June 2018;
206 Fig.1a, 1d and 1e). No positive samples were found in Saint Jean, and no leptospires were detected in
207 the 83 tap waters analyzed.

208

209 3.3 Chemical characteristics of *Leptospira*-positive and -negative waters

210

211 The results of the physicochemical analysis of the early wet season (June) *Leptospira*-positive
212 pond waters in Agla, Ladji and Saint-Jean indicated there was little distinction with early wet season
213 pond waters that were *Leptospira*-negative (data presented in Suppl. Figure 1). *Leptospira*-positive
214 ponds in Agla had water temperatures ranging between 28.2 and 31.9 °C, which is within the range for
215 all pond waters analysed (24.9-35.9°C). The pH of the *Leptospira*-positive pond waters (6.72-8.67) were
216 also within the range of pH for all other ponds sampled (6.69-9.85). This only partly overlaps the pH
217 range for waters usually recognized for pathogenic *Leptospira* survival (i.e. pH 5.5 - 7.6; reviewed in
218 Barragan et al., 2017). The DO and EC values of the Agla *Leptospira*-positive pond waters ranged from
219 0-7.5 mg/L and 62-796 µS/cm, respectively, which are within the same range for all of the pond waters
220 sampled (0-12.4 mg/L and 28-4090 µS/cm respectively). Major ion and trace element concentrations of
221 positive ponds are also within the same ranges as ponds that were *Leptospira*-negative. For example,
222 the concentrations of the cations Na and Ca in *Leptospira*-positive ponds ranged between 6.8-78.9 and
223 0.4-62.7 mg/L, respectively, and between 1.1-8415.5 and 0.2-276.9 mg/L, respectively, for negative
224 ponds. The concentrations of the anions in the positive ponds include HCO₃ and Cl that range between
225 26.0-284.3 and 4.5-74.6 mg/L, respectively, and in the negative ponds these range from 11.7-1115.1 and
226 0.7-16114.9 mg/L. The results of selected trace elements for *Leptospira*-positive ponds includes 37.6-
227 65.9 µg/L for Al, 3.1-194.2 µg/L for Mn, 7.9-3099 µg/L for Fe, 13.8-51.1 µg/L for B and 3.0-24.6 µg/L
228 for Zn.

229 The univariate plots also highlight that the *Leptospira*-containing waters have T°C, pH, EC, DO,
230 major ions and trace elements that overlap with the range of values observed for all urban pond water
231 samples. Similarly, the multivariate analyses shows overlaps of the four first principal components (that
232 together represent 62% of the total variance) between *Leptospira*-positive waters and other urban pond
233 waters (Fig. 2). The results of the hierarchical cluster analysis (AC; illustrated in the dendrogram on Fig.
234 3) also show that the positive pond samples (N38, N21, N18 and N25) are grouped within the same class
235 as several other negative pond samples (e.g. A2, A1, A7, A3, A17, A16, A10, A12, A14).

236 The only groundwater sample to be *Leptospira*-positive also had values of chemical parameters
237 that were similar to ranges observed in groundwater sampled from the shallow aquifer that were
238 *Leptospira*-negative. In Ladji, the *Leptospira*-positive groundwater sample had a temperature of 29.4°C
239 and a pH of 7.69, which is within the range of values for other groundwater samples (26.2-31.1°C and
240 5.85-7.84 respectively). In addition, the DO and EC of the *Leptospira*-positive groundwater (4.4 mg/L
241 and 1357 µS/cm respectively) was also within the range for all other *Leptospira*-negative groundwater
242 values (0.1-8.5 mg/L and 286-2480 µS/cm respectively).

243

244

245 **3.4 Seasonality of *Leptospira*-positive waters**

246

247 *Leptospira*-positive ponds (both temporary and permanent) were found only in the early wet
248 season month of June (2/10 in June 2017, and 3/5 in June 2018), whilst those from the dry season month
249 of October 2017 (N=6) were all *Leptospira*-negative. In addition, the results of the stable isotope data
250 from Agla waters (Fig. 4) show that for four of the *Leptospira*-positive ponds (stable isotope data for
251 the fifth pond is not available) the pond water was recharged by light to moderate rainfall (≤ 250
252 mm/month). Two of the *Leptospira*-positive permanent ponds have stable isotope values close to rainfall
253 values when rainfall levels are between 50 and 250 mm/month. For the other two *Leptospira*-positive
254 ponds (one permanent and one temporary) the stable isotope values are close to rainfall values when the
255 volumes are < 50 mm/month (Fig. 4). The stable isotope values of these 4 ponds (-3.6 to -1.3 ‰ for $\delta^{18}\text{O}$
256 and -0.3 to -16.6 ‰ for $\delta^2\text{H}$) also show that the pond waters are neither evaporated nor affected by
257 mixing with the saline lake water.

258

259 In addition, the hydrochemical results indicate low or insignificant groundwater inflows to these
260 *Leptospira*-positive ponds during the wet season in June. For example, based on the (Na+Ca)/Cl ratios

261 and Cl concentrations (data from Houéménou et al. 2019b), the four *Leptospira*-positive ponds (the fifth
262 pond does not have major ion data) from Agla have waters with relatively higher (Na+Ca)/Cl molar
263 ratios and lower Cl concentrations compared with most groundwater samples (Fig. 5). Therefore, despite
264 the permanent ponds being sustained by groundwater inflows during the dry season, during the wet
265 season the *Leptospira*-positive permanent ponds have received greater inflows from recent rainfall and
266 show a greater resemblance to the *Leptospira*-positive temporary ponds in terms of major ion
267 composition. This is also illustrated by the time series data of one of the *Leptospira*-positive permanent
268 ponds. For example, when this pond is a groundwater discharge site and *Leptospira*-negative during the
269 dry season, the EC is 2,900 $\mu\text{S}/\text{cm}$ (February 2018). In comparison, when the pond is *Leptospira*-
270 positive and has higher rainfall contributions in the wet season, the EC has decreased by over 80% (June
271 2018, EC = 537 $\mu\text{S}/\text{cm}$) and is closer to the EC values of temporary *Leptospira*-positive ponds (62-862
272 $\mu\text{S}/\text{cm}$).

273

274 3.5 Cartography of pond waters

275

276 In Agla, GIS-based evaluation of land use at the beginning of the rainy reason shows that ponds
277 represent quite wide surfaces with 14.7% and 14.1% of the whole surface of the study area in June 2017
278 and June 2018, respectively. Ratios were similar (16% in June 2017) or lower (4% in June 2018), and
279 close to null (1.6% in June 2017 and 0.17% in June 2018) in Saint-Jean. The other values range between
280 0.01% (in Agla in October 2016) to 2.99% (in Ladjji in March 2018), with one exception in Agla in
281 October 2017 (10.4%).

282

283

284

285

286 4. DISCUSSION

287

288 Leptospirosis is considered as one of the most widely distributed zoonosis, but its impact is
289 probably greatest the tropics, especially in Africa where it remains poorly documented (Pappas et al.,
290 2008; Costa et al., 2015). Therefore, studies of leptospirosis in Africa are urgent to improving public
291 health. The filtration-based protocol developed in this study may be particularly useful in remote areas
292 and/or in areas suffering from the lack of well-equipped technical facilities. First, it shows that, when
293 investigating waterborne leptospires in samples that have been filtered, the leptospire screening should
294 include both the water sample and the filter paper (e.g. Wojcik-Fatla et al., 2014). Second, although our
295 method may potentially be less sensitive than those involving specific DNA extraction kits, high-speed
296 centrifugation, bacterial culture and/or high-throughput genetics (e.g., Lall et al. 2016; Riediger et al.
297 2016; Sato et al. 2019), it appears simple, cheap and easily implementable in the field. However, it may
298 further be improved by being coupled with higher efficiency DNA extraction kits (Riediger et al. 2016),
299 something that we did not test here. In addition, we did not investigate possible negative impacts of
300 suspended matter in field water samples on our filtration approach as well as on our qPCR-based
301 detection method (Riedeger et al. 2016). In some instances, we noted that after filtration, filter papers
302 appeared quite coloured, thus suggesting the presence of undetermined suspended particles. The latter
303 may have inhibited to some extent subsequent DNA extractions and qPCRs. However, *Leptospira*-
304 positive samples were among the most turbid water samples (not shown), thus providing confidence on
305 our ability to indeed detect leptospires in environmental samples. To highlight the possible presence of
306 inhibitory compounds in these coloured filter paper samples, a known amount of leptospire DNA
307 obtained from a bacterial culture was added after extraction. These *Leptospira*-enriched samples
308 remained qPCR-negative, suggesting the presence of qPCR inhibitors co-extracted with the DNA (data
309 not shown). Consequently, we cannot definitely exclude the possibility of false negatives where the
310 waters were very turbid, and our results from the field should be cautiously interpreted as potentially
311 underestimated. Additional *in vitro* tests would be useful to address this particular issue and to validate
312 and optimize our centrifugation-based approach for further studies of environmental samples. In a more

313 general manner, our experience confirms previous results that highlight the importance of testing for the
314 possible existence of PCR inhibitors when genetically investigating for the presence or quantity of
315 leptospires in environmental samples (Casanovas-Massana et al., 2018).

316

317 The results retrieved from Cotonou waters demonstrate for the first time that water-borne
318 leptospires circulate in the heart of this large African city (3.1% of total prevalence; 5.5% when tap
319 waters are excluded; see Table 1).

320 It is noteworthy that leptospire-infected waters were found in the two most insalubrious areas,
321 namely Agla and Ladji, whilst none were observed in the popular but non-floodable Saint Jean district.
322 A survey of 780 rodents was conducted in the exact same areas of these three districts, and during the
323 exact same periods (Dossou et al. unpublished data). Interestingly, data in rodents fit well to patterns
324 retrieved in waters since 11.8% and 14% of rodent-borne leptospire prevalence were observed in Agla
325 and Ladji, respectively, while only 0.4% were positive in Saint Jean. This expectedly points towards a
326 higher environmental *Leptospira* contamination in poor, lowland and lake-edge districts where both
327 commensal rodents and surface water are omnipresent (Houéménou et al. 2014; Dobigny et al. 2018).

328 However, different patterns are observed between the high-risk districts of Agla and Ladji. The
329 positive water sample found in Ladji was collected in a groundwater well, thus suggesting that either
330 groundwater in the aquifer was contaminated from infiltration of *Leptospira*-positive water from the
331 surface, or by accident (e.g., use of an infected bucket or rope, or an infected rodent falling down the
332 well). At this stage, we cannot decipher between the two hypotheses, and since the sample size is low
333 we cannot draw robust conclusions on the potential of groundwater contamination by leptospires.
334 Further investigation of leptospires in groundwater in Cotonou deserves special attention, whatever its
335 origin, since one well out of two (50%) was affected in Ladji in June 2017. It is also worth noting that
336 although the groundwater in Agla is shallow, and in contact with permanent ponds water via recharge
337 in the wet season and discharge in the dry season, no leptospires were detected in the Agla groundwater.
338 In comparison, other studies have found groundwater well to be contaminated by leptospires (i.e. in

339 rural Chile; Munoz-Zanzi et al. 2014). In Cotonou, human activities in the immediate proximity of
340 groundwater wells are plentiful, with many children and women searching water daily with their bare
341 feet in close contact with humid soils and the water surrounding the well. Therefore, on-going
342 investigations of leptospires in groundwater is of critical importance.

343

344 In Agla, all *Leptospira*-positive waters originated from ponds, both temporary and permanent,
345 and all were collected in June. Although more intensive sampling is required during different rainfall
346 events, such a pattern strongly suggests that the water-mediated risk of leptospirosis infection may be
347 higher at the beginning of the rainy season through ponds. Our results agree with the conclusions drawn
348 from other studies where leptospiral risk is variable in both time and space, and is strongly dependent
349 on local environmental conditions (reviewed in Barragan et al. 2017). Similar to the results in Cotonou,
350 seasonality patterns were also observed in several regions, such as the Pau da Lima slum (Brazil), where
351 sewage and puddle samples were detected positive more often during the rainy season (Casanovas-
352 Massana et al. 2018). Pathogenic leptospires were also detected in several other urban surface waters
353 (see Tab. 1 in Barragan et al. 2017) where spatial heterogeneity is observed. For example, on the south
354 Andaman Island, the water-borne leptospires were only found in household drainage (18.7%) and
355 sewage (9.8%) waters and, unlike in Cotonou, there were no leptospire found in the pond waters (Lall
356 et al. 2016). The variable results from each study once again highlights the importance of understanding
357 the local socio-environmental context in determining leptospire transmission pathways to humans (e.g.,
358 Maciel et al. 2008; Himsforth et al. 2013; Mwachui et al. 2015; Houéménou et al. 2019a; reviewed in
359 Barragan et al. 2017).

360 The hydrochemical analysis of contaminated waters strongly suggest there are key hydrological
361 processes that may favor the presence of leptospires in the surface waters of Cotonou. Indeed, stable
362 isotopes and major ion data (from Houéménou et al., 2019b) indicate that leptospires are found in
363 temporary and permanent ponds at the beginning of the rainy season that have recently formed as a
364 result of low to moderate precipitation, and have relatively high runoff inputs compared to groundwater
365 or lake water inputs. This agrees with the expectation that there will be larger concentrations of the

366 pathogen in small water bodies (Barragan et al. 2017). It also corresponds well with the proposed model
367 of soil-to-water dispersion of leptospires recently recapitulated by Bierque and colleagues (2020).
368 Finally, and although the causal relationship remains untested, these results are strikingly consistent
369 with previous data on south Benin urban rodents in which *Leptospira* prevalence was found higher
370 during (or soon after) periods of moderate rains (100-200 mm; Houéménou et al. 2019b).

371

372 Though data are quite scarce, some physico-chemical parameters are known to have an influence
373 on *Leptospira* presence, survival and virulence (e.g., Chang et al. 1948; Khairani-Bejo et al. 2014;
374 André-Fontaine et al. 2015; Lall et al. 2018). Our observations strongly suggest that most negative pond
375 samples have similar physico-chemical characteristics to those positive for pathogenic *Leptospira*. In
376 particular, it was recently suggested that iron, manganese and copper could influence the survival of
377 leptospires in Andamanese soils (Lall et al. 2018). However, our data do not confirm such a trend since
378 leptospire-positive and negative water samples did not show any noticeable differences in these ions
379 concentrations (see Suppl. 2). Overlaps observed with both univariate and multivariate approaches mean
380 that no association between physicochemical parameters and presence (Agla, Ladji) or absence (Saint
381 Jean) of water-borne leptospires can be identified (Suppl. Fig. 3). This means also that the possible
382 ecological spectrum of water-borne leptospires extensively covers the physico-chemical spectrum of
383 Cotonou waters as a whole. For instance, pH and temperature of Cotonou waters, which are suspected
384 to be of primary importance for leptospires biology (Chang et al. 1948; Gordon-Smith & Turner, 1961;
385 Khairani-Bejo et al. 2004; André-Fontaine et al. 2015), are perfectly compatible with pH and
386 temperature ranges that allow for the long-term survival of pathogenic leptospires (André-Fontaine et
387 al. 2015). In other words, the potential distribution of waters compatible with *Leptospira* survival, hence
388 human contamination risk, is quite wide in the city, meaning that, though quite polluted (Lapworth et
389 al. 2017; Houéménou et al. 2019b), urban waters in Cotonou are widely compatible with the leptospire
390 life cycle. In the city and, beyond, along the West African Abidjan-Lagos urban corridor, surface waters
391 inundate many zones of the public space, including areas that are used daily by the inhabitants: walking
392 pathways and streets, courtyards and even homes may be flooded for several days if not weeks.

393 Consequently, the population has extensive and on-going contact with potentially contaminated water.
394 Altogether, this indicates that millions of people living along the extensively urbanizing Guinea Gulf
395 are at risk for falling sick of leptospirosis.

396

397 **5. CONCLUSION**

398

399 Future studies should include two major aspects of leptospire ecology, hence leptospiral
400 contamination risk, that were not taken into consideration here: water microbiota and biofilms (e.g.,
401 Trueba et al. 2004; Kumar et al. 2015) on the one hand, and soil-associated leptospire on the other hand
402 (e.g., Thibeaux et al. 2017; Bierque et al. 2020). Nevertheless, our filtration-based results, though
403 potentially underestimated, clearly show that the Cotonou inhabitants live in an urban environment that
404 exposes them to leptospirosis infection risk, potentially at a wide spatial scale. Everyday uses and habits
405 as well as a lack of street and sanitation infrastructures results in many people in close contact with
406 standing waters on a regular basis, especially during the rainy season and associated flooding events.
407 Spatio-temporal investigations of water-borne leptospire point towards a higher risk at the beginning
408 of the rainy season, especially via ponds that are recharged by rainfall. We believe that awareness
409 increase about leptospirosis should be a public health priority since the disease is very poorly
410 documented and obviously overlooked in Benin and, further, along the rapidly extending Abidjan-Lagos
411 urban corridor.

412

413 **6. ACKNOWLEDGEMENTS**

414 We are in debt to all Cotonou inhabitants who allowed us to enter and collect samples within
415 their households. We are grateful to the two anonymous reviewers who greatly contributed to improve
416 the first version of our manuscript. This research was funded by the UMRs CBGP, EMMAH and G-
417 EAU. Field works in Benin were conducted under the research agreement between Republic of Benin

418 and the French Institute of Research for sustainable Development that was signed on 30th September
419 2010.

420

421 7. REFERENCES

422

423 Addinsoft 2019. XLSTAT statistical and data analysis solution. Paris, France. <https://www.xlstat.com>.

424 Africapolis 2020. Visualise urbanisation in Africa. OECD Sahel and West Africa Club,
425 <https://www.africapolis.org/> (last accessed on 21/04/2020).

426 Allan K.J., Biggs H.M., Halliday J.E.B., Kazwala, R.R., Maro V.P., Cleaveland S. & Crump, J.A. 2015.
427 Epidemiology of leptospirosis in Africa: A systematic review of a neglected zoonosis and a
428 paradigm for ‘One Health’ in Africa. *PLOS Neglected Tropical Diseases*, **9**, e0003899.
429 <https://doi.org/10.1371/journal.pntd.0003899>.

430 André-Fontaine G., Aviat F. & Thorin C. 2015. Waterborne leptospirosis: Survival and preservation of
431 the virulence of pathogenic *Leptospira* spp. in fresh water. *Current Microbiology*, **71**,
432 <https://doi.org/10.1007/s00284-015-0836-4>.

433 Barragan V., Olivas S., Keim P. & Pearson T. 2017. Critical knowledge gaps in our understanding of
434 Environmental cycling and transmission of *Leptospira* spp. *Applied Environmental*
435 *Microbiology*, **83**, <https://doi.org/10.1128/AEM.01190-17>.

436 Bierque E., Thibeaux R., Girault D., Soupé-Gilbert M.-E. & Goarant C. 2020. A systematic review of
437 leptospira in water and soil environments. *PLoS ONE*, **15**, e0227055.

438 Cann K.F., Thomas D.R., Salmon R.L., Wyn-Jones A.P. & Kay D. 2013. Extreme water-related weather
439 events and waterborne disease. *Epidemiology and Infections*, **141**, 671–686.

440 Casanovas-Massana A., Costa F., Riedeger I.N., Cunha M., de Oliveira D., Mota D.C., Sousa E.,
441 Querino V.A., Nery Jr N., Reis M.G., Wunder E.A., Diggle P.J. & Ko A.I. 2018. Spatial and
442 temporal dynamics of pathogenic *Leptospira* in surface waters from the urban slum environment.
443 *Water Research*, **130**, 176-194.

444 .

445 Chang S.L., Buckingham M. & Taylor M.P. 1948. Studies on *Leptospira icterohaemorrhagiae*: IV.
446 Survival in water and sewage: destruction in water by halogen compounds, synthetic detergents
447 and heat. *Journal of Infectious Diseases*, **82**, 256-266.

448 Cornwall W. 2016. A plague of rats. *Science*, **352**, 912–915.

449 Costa F., Hagan J.E., Calcagno J., Kane M., Torgerson, P., Martinez-Silveira M.S., Stein C., Abela-
450 Ridder B. & Ko A.I. 2015. Global morbidity and mortality of leptospirosis: A systematic review.
451 *PLoS Neglected Tropical Diseases*, **9**, e0003898.

452 de Vries S.G., Visser B.J., Nagel I.M., Goris M.G.A., Hartskeerl R.A. & Grobusch M.P. 2014.
453 Leptospirosis in Sub-Saharan Africa: a systematic review. *International Journal of Infectious*
454 *Diseases*, **28**, 47–64.

455 Denis E., Moriconi-Ebrard F., Harre D., Thiam O., Séjourné M. & Chatel C. 2009. Africapolis -
456 dynamiques de l'urbanisation, 1950-2020, une approche géostatistique : Afrique de l'Ouest
457 (*Urbanization dynamics, 1950-2020, a geostatistic approach: West Africa*). [https://hal.archives-](https://hal.archives-ouvertes.fr/hal-00357271/file/Africapolis_1_Rapport.pdf)
458 [ouvertes.fr/hal-00357271/file/Africapolis_1_Rapport.pdf](https://hal.archives-ouvertes.fr/hal-00357271/file/Africapolis_1_Rapport.pdf) (last accessed on 21/04/2020).

459 Dobigny G., Garba M., Tatard C., Loiseau A., Galan M., Kadaouré I., Rossi J.-P., Picardeau M. &
460 Bertherat E. 2015. Urban market gardening and rodent-borne pathogenic *Leptospira* in Arid Zones:
461 A Case Study in Niamey, Niger. *PLoS Neglected Tropical Diseases*, **9**, e0004097.

462 Dobigny G., Gauthier P., Houéménou G., Choplin A., Dossou H.-J., Badou S., Etougbétché J., Bourhy
463 P., Koffi S., Durski K., Bertherat E. & Picardeau M. 2018. Leptospirosis and extensive urbanization
464 in West Africa: A neglected and underestimated threat? *Urban Science*, **2**, e029.

465 Dobigny G., Gauthier P., Houéménou G., Dossou H.-J., Badou S.A., Etougbétché J., Tatard C. & Truc
466 P.H.V. 2019. Spatio-temporal survey of small mammal-borne *Trypanosoma lewisi* in Cotonou,
467 Benin, and the potential risk of human infection. *Infection, Genetics and Evolution*, e103967.

468 Ganoza C.A., Matthias M.A., Collins-Richards D., Brouwer K.C., Cunningham C.B., Segura E.R.,
469 Gilman R.H., Gotuzzo E. & Vinetz J.M. 2006. Determining risk for severe leptospirosis by
470 molecular analysis of environmental surface waters for pathogenic *Leptospira*. *PLoS Medicine*, **3**,
471 e030308.

- 472 Haake D.A. & Levett P.N. 2015. Leptospirosis in humans. *Current Topics in Microbiology and*
473 *Immunology*, **387**, 65–97.
- 474 Himsworth C.G., Bidulka J., Parsons K.L., Feng A.Y.T., Tang P., Jardine C.M., Kerr T., Mak S.,
475 Robinson J. & Patrick D.M. 2013. Ecology of *Leptospira interrogans* in Norway rats (*Rattus*
476 *norvegicus*) in an inner-city neighborhood of Vancouver, Canada. *PLoS Neglected Tropical*
477 *Diseases*, **7**, e2270.
- 478 Houéménou H. 2020. Qualité des eaux souterraines et de surface dans la métropole de Cotonou (Bénin) :
479 Implications pour la leptospirose (*Ground and surface waters quality in Cotonou, Benin:*
480 *implications for leptospirosis*). PhD thesis, Avignon University, France.
- 481 Houéménou G., Ahmed A., Libois R. & Hartskeerl R.A. 2014. *Leptospira* spp. prevalence in small
482 mammal populations in Cotonou, Benin. *International Scholarly Research Notices*, **8**, ID 2013,
483 502638, <http://dx.doi.org/10.5402/2013/502638>.
- 484 Houéménou G., Gauthier P., Etougbétché J., Badou S., Dossou H.-J., Agossou D., Picardeau M. &
485 Dobigny G. 2019a. Pathogenic *Leptospira* in commensal small mammals from the extensively
486 urbanized coastal Benin. *Urban Science*, **3**, e3030099.
- 487 Houéménou H., Tweed S., Dobigny G., Mama D., Alassane A., Silmer R., Babic M., Ruy S., Chaigneau
488 A., Gauthier P., Socohou A., Dossou H.-J., Badou S. & Leblanc, M. 2019b. Degradation of
489 groundwater quality in expanding cities in West Africa: a case study of the unregulated shallow
490 aquifer in Cotonou. *Journal of Hydrology*, **582**, e124438.
- 491 Kaboosi H. & Razavi M.R. 2010. Efficiency of filtration technique for isolation of leptospires from
492 surface waters: role of different membranes with different pore size and materials. *African Journal*
493 *of Microbiology Research* **4**(9), 671-676.
- 494 Khairani-Bejo S., Bahaman A.R.B., Zamri-Saad M. & Mutalib A.R. 2004. The Survival of *Leptospira*
495 *interrogans* Serovar Hardjo in the Malaysian environment. *Journal of Animal and Veterinary*
496 *Advances*, **3**(3), 123-129.
- 497 Ko A.I., Reis M.G., Dourado C.M.R., Johnson W.D. & Riley L.W. 1999. Urban epidemic of severe
498 leptospirosis in Brazil. *The Lancet*, **354**, 820–825.

499 Kumar V.K., Lall C., Vimal Raj R., Vedhagiri K. & Vijayachari P. 2015. Co-existence and survival of
500 pathogenic leptospires by formation of biofilm with *Azospirillum*. *FEMS Microbiology and*
501 *Ecology*, **91**, pii:fiv051.

502 Lall C., Vinod Kumar K., Raj R.V., Vedhagiri K., Sunish I.P. & Vijayachari P. 2018. Correlation
503 between physicochemical properties of soil and presence of *Leptospira*. *EcoHealth*, **15**, 670–675.

504 Lapworth D.J., Nkhuwa D.C.W., Okotto-Okotto J., Pedley S., Stuart M.E., Tijani M.N. & Wright, J.
505 2017. Urban groundwater quality in sub-Saharan Africa: current status and implications for water
506 security and public health. *Hydrogeology Journal*, **25**, 1093–1116.

507 Lau C.L., Smythe L.D., Craig S.B. & Weinstein P. 2010. Climate change, flooding, urbanisation and
508 leptospirosis: fuelling the fire? *Transactions of the Royal Society of Tropical Medecine and*
509 *Hygiene*, **104**, 631-638.

510 Maciel E.A.P., Carvalho A.L.F., de Nascimento S.F., Matos R.B., de Gouveia E.L., Reis M.G. & Ko
511 A.I. 2008. Household transmission of *Leptospira* infection in urban slum communities. *PLoS*
512 *Neglected Tropical Diseases*, **2**, e154.

513 Muñoz-Zanzi C., Mason M.R., Encina C., Astroza A. & Romero A. 2014. *Leptospira* contamination in
514 household and environmental water in rural communities in Southern Chile. *International Journal*
515 *of Environmental Research and Public Health*, **11**, 6666–6680.

516 Mwachui M.A., Crump L., Hartskeerl R., Zinsstag J. & Hattendorf J. 2015. Environmental and
517 behavioural determinants of leptospirosis transmission: A systematic review. *PLoS Neglected*
518 *Tropical Diseases*, **9**, e0003843.

519 Pappas G., Papadimitriou P., Siozopoulou V., Christou L. & Akriditis N. 2008. The globalization of
520 leptospirosis: worldwide incidence trends. *International Journal of Infectious Diseases*, **12**, 351-
521 357.

522 Programme de Protection de la Communauté Urbaine de Cotonou Face aux Changements Climatiques.
523 (PCUG3C) 2012. Inondations dans le grand Cotonou : facteurs humains, vulnérabilité des
524 populations et stratégies de lutte et de gestion. (*Program for the protection of the urban community*
525 *of Cotonou against climatic changes: human factors, population vulnerability and management*

526 *strategies*). Final Report, Abomey-Calavi, Benin. Available online: <https://idl-bnc-idrc.dspacedirect.org/handle/10625/50607> (last accessed on 21/04/2020).

527

528 QGIS Development Team 2016. QGIS: Geographic Information System. Open Source Geospatial
529 Foundation Project. <http://qgis.osgeo.org>.

530 Riediger I.N., Hoffmaster A.R., Casanovas-Massana A., Biondo A.W., Ko A.I. & Stoddard, R.A. 2016.
531 An optimized method for quantification of pathogenic *Leptospira* in environmental water samples.
532 *PLoS ONE*, **11**, e160523.

533 RStudio Team 2016. RStudio: Integrated development for R. RStudio Inc., Boston, USA.
534 <http://www.rstudio.com/>.

535 Sato Y., Mizuyama M., Sato M., Minamoto T., Kimura R. & Toma, C. 2019. Environmental DNA
536 metabarcoding to detect pathogenic *Leptospira* and associated organisms in leptospirosis-endemic
537 areas of Japan. *Scientific Reports*, **9**, 1–11.

538 Smith C.E.G. & Turner L.H. 1961. The effect of pH on the survival of leptospires in water. *Bulletin of*
539 *the World Health Organization*, **24**, 35–43.

540 Thaipadungpanit J., Wuthiekanun V., Chantratita N., Yimsamran S., Amornchai P., Boonsilp S.,
541 Maneeboonyang W., Tharnpoophasiam P., Saiprom N., Mahakunkijcharoen Y., Day N.P.J.,
542 Singhasivanon P., Peacock S.J. & Limmathurotsakul D. 2013. *Leptospira* species in floodwater
543 during the 2011 floods in the Bangkok Metropolitan Region, Thailand. *American Journal of*
544 *Tropical Medicine and Hygiene*, **89**, 794–796.

545 Thibeaux R., Geroult S., Benezech C., Chabaud S., Soupé-Gilbert M.S., Girault D., Bierque E. &
546 Goarant, C. 2017. Seeking the environmental source of leptospirosis reveals durable bacterial
547 viability in river soils. *PLoS Neglected Tropical Diseases*, **11**, e5414.

548 Trueba G., Zapata S., Madrid K., Cullen P. & Haake D. 2004. Cell aggregation: a mechanism of
549 pathogenic *Leptospira* to survive in fresh water. *International Microbiology*, **7**, e35–40.

550 UN Habitat 2014. L'urbanisation - un impératif pour la transformation structurelle de l'Afrique : une
551 nouvelle conception de l'urbanisation en appui à l'Agenda 2063. United Nations, Nairobi, Kenya.

552 Wójcik-Fatla A., Zając V., Wasinski B., Sroka J., Cisak E., Sawczyn-Domańska A. & Dutkiewicz J.
553 2014. Occurrence of *Leptospira* DNA in water and soil samples collected in eastern Poland.

554 *Annals of Agricultural and Environmental Medicine of the American Academy of Emergency*
555 *Medicine*, **21**, e730-732.

556 Yabi I. & Afouda F. 2012. Extreme rainfall years in Benin (West Africa). In: West African paleoecology
557 and human responses. *West African Quaternary Research Association*, **262**, 39–43




Figure 1. Sampling sites. Map of Cotonou showing the three neighborhoods Agla, Ladji and Saint Jean (a). Pictures of precarious and stilt habitats in Ladji (b), Saint-Jean (c) and Agla (e and f). A few of the

sampling sites ((c)Google Earth) to illustrate the sampling protocol in Agla (d) with blue, yellow and red dots corresponding to groundwater, *Leptospira*-negative and *Leptospira*-positive ponds, respectively. A19 indicates a *Leptospira*-positive pond presented in the photo (f).




Figure 2. Principal Component Analysis showing the physico-chemical characteristics of the *Leptospira*-positive samples (red dots) among all urban water samples collected in Agla and Ladji.




Figure 3. Hierarchical cluster analysis dendrogram showing the relationship between the *Leptospira*-positive pond samples (*) and *Leptospira*-negative samples in Agla.




Figure 4. Stable isotopes of rainwater, pond water and shallow groundwater samples in Agla during the wet and dry 2017 and 2018 seasons (modified from Houéméno et al. 2019).




Figure 5. (Na+Ca)/Cl molar ratios and Cl concentrations of pond water and shallow groundwater samples in Agla during the wet and dry 2017 and 2018 seasons (modified from Houéménou et al. 2019).

Table 1. *Leptospira* screening results by sampling sites and dates. “ok”: physico-chemical data are available; “NA”: not assessed; “t.ponds”: temporary ponds; “p.ponds”: permanent ponds.

AGLA		Mar-17	Jun-17	Oct-17	Feb-18	Jun-18
N=85	Groundwater	7	7	8		1
	Temporary ponds		10	6		5
	Permanent ponds				3	8
	Tap water	10	10	10		
	Total	17	27	24	3	14
	Positive leptospires		2+(t.ponds.)			3+(p.ponds.)
	Total prevalence	0.00	0.07	0.00	0.00	0.20
	Ponds prevalence	NA	0.20	0.00	0.00	0,21
	Measures		ok	ok	ok	1/2
LADJI		Mar-17	Jun-17	Oct-17	Feb-18	Jun-18
N=53	Groundwater	3	2	3		
	Temporary ponds		5	3		
	Permanent ponds	1			2	3
	Tap water	10	10	8		
	Lake					
	Total	14	18	15	2	4
	Positive leptospires		1+(well)			
	Total prevalence	0.00	0.06	0.00	0.00	0.00
	Groundwater prevalence	0.00	0.50	0.00		
Measures	NA	ok	ok	ok	ok	
St JEAN		Mar-17	Jun-17	Oct-17	Feb-18	Jun-18
N=55	Groundwater	10	10	10		
	Tap water	10	9	6		
	Total	20	19	16		
	Positive leptospires	0	0	0		
	Total prevalence	0.00	0	0.00		
	Measures	NA	ok	ok		

Development of centrifugation method

Dry season


Beginning of the rainy season

Rainy season


Permanent Ponds



Leptospira-negative ponds



Leptospira-positive ponds



Leptospira-negative ponds

Temporary Ponds