

Environmental Plasticity of the RNA Content of Staphylococcus aureus Extracellular Vesicles

Brenda Silva Rosa Da Luz, Aurélie Nicolas, Svetlana Chabelskaya, Vinícius de Rezende Rodovalho, Yves Le Loir, Vasco Ariston de Carvalho Azevedo, Brice Felden, Eric Guédon

▶ To cite this version:

Brenda Silva Rosa Da Luz, Aurélie Nicolas, Svetlana Chabelskaya, Vinícius de Rezende Rodovalho, Yves Le Loir, et al.. Environmental Plasticity of the RNA Content of Staphylococcus aureus Extracellular Vesicles. Frontiers in Microbiology, 2021, 12, 10.3389/fmicb.2021.634226. hal-03169446

HAL Id: hal-03169446 https://hal.inrae.fr/hal-03169446

Submitted on 15 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License





Environmental Plasticity of the RNA Content of Staphylococcus aureus Extracellular Vesicles

Brenda Silva Rosa Da Luz^{1,2}, Aurélie Nicolas¹, Svetlana Chabelskaya³, Vinícius de Rezende Rodovalho^{1,2}, Yves Le Loir¹, Vasco Ariston de Carvalho Azevedo², Brice Felden^{3*†} and Eric Guédon^{1*†}

¹INRAE, Institut Agro, STLO, Rennes, France, ²Laboratory of Cellular and Molecular Genetics, Institute of Biological Sciences, Federal University of Minas Gerais, Belo Horizonte, Brazil, ³BRM [Bacterial Regulatory RNAs and Medicine] UMR_S 1230, University of Rennes, Inserm, Rennes, France

OPEN ACCESS

Edited by:

Carlos Robello, Universidad de la República, Uruguay

Reviewed by:

Maria Rosa Garcia-Silva, Institut Pasteur de Montevideo, Uruguay Lysangela Ronalte Alves, Carlos Chagas Institute (ICC), Brazil

*Correspondence:

Eric Guédon eric.guedon@inrae.fr Brice Felden brice.felden@univ-rennes1.fr; bfelden@univ-rennes1.fr

[†]These authors have contributed equally to this work

Specialty section:

This article was submitted to Infectious Diseases, a section of the journal Frontiers in Microbiology

Received: 27 November 2020 Accepted: 18 February 2021 Published: 11 March 2021

Citation:

Da Luz BSR, Nicolas A, Chabelskaya S, Rodovalho VdR, Le Loir Y, Azevedo VAdC, Felden B and Guédon E (2021) Environmental Plasticity of the RNA Content of Staphylococcus aureus Extracellular Vesicles. Front. Microbiol. 12:634226. doi: 10.3389/fmicb.2021.634226 The roles of bacterial extracellular vesicles (EVs) in cell-to-cell signaling are progressively being unraveled. These membranous spheres released by many living cells carry various macromolecules, some of which influence host-pathogen interactions. Bacterial EVs contain RNA, which may serve in communicating with their infected hosts. *Staphylococcus aureus*, an opportunistic human and animal pathogen, produces EVs whose RNA content is still poorly characterized. Here, we investigated in depth the RNA content of S. aureus EVs. A highthroughput RNA sequencing approach identified RNAs in EVs produced by the clinical S. aureus strain HG003 under different environmental conditions: early- and late-stationary growth phases, and presence or absence of a sublethal vancomycin concentration. On average, sequences corresponding to 78.0% of the annotated transcripts in HG003 genome were identified in HG003 EVs. However, only ~5% of them were highly covered by reads (≥90% coverage) indicating that a large fraction of EV RNAs, notably mRNAs and sRNAs, were fragmented in EVs. According to growth conditions, from 86 to 273 highly covered RNAs were identified into the EVs. They corresponded to 286 unique RNAs, including 220 mRNAs. They coded for numerous virulence-associated factors (hld encoded by the multifunctional sRNA RNAIII, agrBCD, psmβ1, sbi, spa, and isaB), ribosomal proteins, transcriptional regulators, and metabolic enzymes. Twenty-eight sRNAs were also detected, including bona fide RsaC. The presence of 22 RNAs within HG003 EVs was confirmed by reverse transcription quantitative PCR (RT-gPCR) experiments. Several of these 286 RNAs were shown to belong to the same transcriptional units in S. aureus. Both nature and abundance of the EV RNAs were dramatically affected depending on the growth phase and the presence of vancomycin, whereas much less variations were found in the pool of cellular RNAs of the parent cells. Moreover, the RNA abundance pattern differed between EVs and EV-producing cells according to the growth conditions. Altogether, our findings show that the environment shapes the RNA cargo of the S. aureus EVs. Although the composition of EVs is impacted by the physiological state of the producing cells, our findings suggest a selective packaging of RNAs into EVs, as proposed for EV protein cargo. Our study shedds light to the possible roles of potentially functional RNAs in S. aureus EVs, notably in host-pathogen interactions.

Keywords: membrane vesicle, small regulatory RNA, virulence factors, vancomycin, RNA-Seq, extracellular vesicle, RsaC, RNAIII

INTRODUCTION

The release of extracellular vesicles (EVs) by living cells is a well-established phenomenon required for intercellular communications and trans-kingdom interactions (Brown et al., 2015; Toyofuku, 2019). These spherical membranous particles vary from 20 to 300 nm in diameter and contain macromolecules such as nucleic acids, proteins, lipids, and small metabolites. Initially considered to be trash bags to eliminate unwanted material outside of the cells, they are now widely recognized as protective delivery shuttles of bioactive molecules from donor to recipient cells (Brown et al., 2015; Kim et al., 2015; Gill et al., 2019). The functional characterization of bacterial EVs is of interest due to their capacities to affect bacteria-host cell interactions and bacterial pathogenesis (Kaparakis-Liaskos and Ferrero, 2015; Tsatsaronis et al., 2018). Although the formation of outer membrane vesicles (OMVs) in Gram-negative bacteria was early documented in 1966 (Work et al., 1966), the formation of such structures was disregarded in Gram-positive bacteria until recently. The production of EVs by a Gram-positive bacterium, Staphylococcus aureus, was demonstrated in 2009 and, ever since, numereous studies confirmed EV release by other Gram-positive bacteria (Lee et al., 2009, 2013a; Rivera et al., 2010; Prados-Rosales et al., 2011; Brown et al., 2014; Olaya-Abril et al., 2014; Kim et al., 2016a; Liu et al., 2018b).

Staphylococcus aureus commonly colonizes the skin or nasal tract of vertebrates, without causing disease (Wertheim et al., 2005). However, it is also one of the main opportunistic pathogen in humans, and a frequent cause of multi-drug resistant nosocomial infections (Ziebuhr, 2001). S. aureus is responsible for a wide array of diseases, ranging from minor infections in soft tissues to life-threatening diseases, such as sepsis, meningitis, and pneumonia (Salgado-Pabón and Schlievert, 2014; Tong et al., 2015). The type and severity of infections depend on strain-specific virulence factors, mostly expressed from accessory genetic elements (Gill et al., 2011). Secreted and surface-exposed S. aureus virulence factors weaken the host immune response, leading to bacterial immune evasion and pathogenesis (Foster, 2005). EVs could be a vehicle for secretion and surface-display of these molecules and, accordingly, recent studies indicate that S. aureus EVs carry important bacterial survival and virulence factors, such as β-lactamases, toxins, and proteins involved in adhesion to host cells (Lee et al., 2009; Gurung et al., 2011; Jeon et al., 2016; Askarian et al., 2018; Tartaglia et al., 2018, 2020; Wang et al., 2018).

Biologically active β -lactamase in *S. aureus* EVs can confer a transient resistance against ampicillin to surrounding sensible bacteria (Lee et al., 2013b). Furthermore, the presence of α -hemolysin inside EVs accelerates host cell death (Thay et al., 2013; Hong et al., 2014), and EV-associated exfoliative toxin A (ETA) induces a characteristic toxicity onto human epithelial cells (Jeon et al., 2016). Moreover, *S. aureus*-derived EVs facilitate the induction and exacerbation of skin and pulmonary inflammations (Hong et al., 2011, 2014; Kim et al., 2012; Jun et al., 2017). EVs-associated molecules can be more efficient than cytoplasmic proteins to elicit an immune response and host-cell toxicity (Hong et al., 2014). In response to antibiotics exposures, EVs increase *S. aureus* adhesion and cell aggregation, and contribute to biofilm formation (He et al., 2017). Recent data highlight the importance of EVs in staphylococcal pathogenesis since EVs derived from various human and animal strains of *S. aureus* share a conserved EV proteome (Tartaglia et al., 2020).

The vast majority of functional studies on bacterial EVs, however, challenged their proteome. Regarding the presence of DNAs and RNAs in EVs, most studies have been conducted on Gram-negative bacteria (Perez Vidakovics et al., 2010; Blenkiron et al., 2016; Koeppen et al., 2016; Bitto et al., 2017; Choi et al., 2017; Malabirade et al., 2018; Yu et al., 2018; Han et al., 2019). OMV-associated RNAs can include messenger RNAs (mRNA), transfer RNAs (tRNA), ribosomal RNAs (rRNA), or small regulatory RNAs (sRNA; Biller et al., 2014; Ghosal et al., 2015; Ho et al., 2015; Sjöström et al., 2015; Blenkiron et al., 2016; Koeppen et al., 2016; Choi et al., 2017; Dauros-Singorenko et al., 2018; Liu et al., 2018a; Malabirade et al., 2018; Tsatsaronis et al., 2018; Frantz et al., 2019). EV-associated RNA cargo, notably sRNAs, can influence host-pathogen interactions, cell-to-cell communications, and bacterial pathogenesis (Dauros-Singorenko et al., 2018; Tsatsaronis et al., 2018; Lee, 2019; Ahmadi Badi et al., 2020; Lécrivain and Beckmann, 2020). For instance, OMVs from Pseudomonas aeruginosa can transfer an sRNA into the human airway cells, resulting in IL-8 decrease (Koeppen et al., 2016). Likewise, transfection of OMV-associated sRNAs from the periodontal pathogens Aggregatibacter actinomycetemcomitans, Phorphyromonas gingivalis, and Trepanema denticola into human cells reduced host interleukine release (Choi et al., 2017). The presence of RNAs within Gram-positive EVs has been reported for fewer species (Resch et al., 2016; Dauros Singorenko et al., 2017; Frantz et al., 2019; Rodriguez and Kuehn, 2020). Interestingly, Frantz et al. (2019) recently reported that the EV-associated rli32 sRNA of Listeria monocytogenes can trigger the induction of a type I IFN response in host cells. This finding supports that Gram-positive EVs can also participate to host-pathogen interactions by dedicated vesicular RNAs. Data about RNA cargo in EVs released by S. aureus are scarce, with only two recent reports. While the first provided a partial RNA profile of S. aureus MSSA476 EVs without functional analyses (Joshi et al., 2021), the second showed that the uncharacterized RNA content of S. aureus Newman EVs likely stimulate the potent IFN-B response observed in cultured macrophage cells (Rodriguez and Kuehn, 2020).

As far as we know, our work is the first example that provides a detailed RNA profile associated to EVs from a reference clinical *S. aureus* strain, HG003. The staphylococcal EV RNA cargo was unveiled by high-throughput RNA sequencing from purified EVs after release by cells grown under various environmental conditions. They include early- and late-stationary growth phases, with or without a sublethal concentration of vancomycin, an antibiotic used to treat multidrug-resistant infections and that influences *S. aureus* EV biogenesis and functions (Hsu et al., 2011; He et al., 2017). The RNA cargo from the EVs was analyzed and compared to the RNA content of the HG003 parental cells.

MATERIALS AND METHODS

Bacterial Strain and Growth Conditions

The *S. aureus* strain used in this work was the model strain HG003 (Herbert et al., 2010), a NCTC8325 derivative, isolated in 1960 from a sepsis patient. HG003 contains functional *rsbU* and *tcaR* genes, two global regulators that are missing in the NCTC8325 parent strain. The HG003 genome is well documented (Sassi et al., 2014), and this strain is widely used as a reference to investigate staphylococcal regulation and virulence (Liu et al., 2018a). HG003 strain was pre-inoculated in BHI broth and grown overnight at 37°C under 150 rpm/min agitation, and then inoculated 0.1% in 500 ml of fresh BHI (125 rpm/min, at 37°C) on a 1 L Scott flask. Bacterial cultures were retrieved after 6 h and 12 h for early- and late-stationary phases, respectively, in the presence or absence of a sub-inhibitory concentration (0.5 μ g/ml) of vancomycin (**Supplementary Figure S1**).

S. aureus EVs Isolation and Purification

Cultures were submitted to EVs isolation and purification, as previously described (Tartaglia et al., 2018, 2020). In brief, for each condition 1 L of bacterial cell culture was centrifuged at 6,000 × g for 15 min and filtered through 0.22 µm Nalgene top filters (Thermo Scientific). Then, the culture supernatant fraction was concentrated around 100-fold using the Amicon ultrafiltration systems (Millipore) with a 100kDa filter, and ultra-centrifuged for 120 min at 150,000 × g to eliminate the soluble proteins. Next, the suspended pellet was applied to a discontinuous sucrose gradient (8–68%) and ultra-centrifuged at 100,000 × g for 150 min. Fractions containing EVs were recovered and washed in TBS (150 mM NaCl; 50 mM Tris-Cl, pH 7.5) for final ultra-centrifugation at 150,000 × g (120 min). At last, EVs were suspended in cold TBS and kept at -80° C until use.

EVs Visualization by Electron Microscopy

Negative staining electron microscopy was performed as previously described (Rodovalho et al., 2020) to investigate the shape and integrity of purified EVs. EVs samples were diluted, and solutions containing between 10¹⁰ and 10¹¹ particles per ml were analyzed. For this, samples were applied to glow-discharged copper EM grids (FF200-Cu) for 30 s, followed by excess solution removal with filter paper. The same process was repeated with 2% uranyl acetate, and samples were observed with a Jeol 1400 transmission electron microscope (JEOL Ltd.), operating at 120 kV.

Determination of EVs Sizes and Concentrations

Nanoparticle Tracking Analysis (NTA) using an sCMOS camera and a Blue488 laser (Nano Sight NS300) was performed to assess EVs size and concentration. For that, samples were diluted into TBS to achieve optimal concentration and submitted to a constant flux generated by a syringe pump (speed 50), at 25°C. Results were retrieved from 5×60 s videos recorded

RNA Extraction From *S. aureus* **HG003** Whole Cells and Its Derived EVs

RNA extraction was carried out as similar as possible for both cell and EV samples. Bacterial RNA extraction was performed from 10 ml culture pellet. The samples were mixed with glass beads in 300 µl lysis buffer (0.5% SDS w/v, 30 mM sodium acetate; 1 mM EDTA) and 400 µl phenol (acid buffered at pH 5.0) at 65°C. Mechanical lysis was accomplished with 2 cycles of 30 s in Precellys at 6,500 rpm. For EV sample RNA extraction, particles isolated from the equivalent of 800 ml bacterial culture were mixed with 300 µl of lysis buffer and 400 µl phenol at 65°C, the same volumes used for cell RNA extraction. Since EVs lack the thick layer of peptidoglycan (PGN) found in the bacterial cell wall, mechanic lysis was not necessary and was achieved with lysis buffer. EV and EV-producing cell samples were incubated for 10 min at 65°C, being homogenized by vortex every minute. Next, samples were centrifuged during 10 min 13,000 rpm, 4°C, and the upper phase was recovered to a new tube. All samples were mixed with additional 400 µl of phenol at 65°C, and the previous steps were repeated. Then, 400 µl of phenol:chloroform 1:1 was added, followed by two times addition of 400 µl pure chloroform, repeating the step of upper phase recovery, mixture and centrifugation (5 min at 13,000 rpm, 4°C). Subsequently, 1.5 volumes of ice-cold 100% ethanol and 10% volume of NaAc were added and the mix was stored at -20° C overnight. Samples were centrifuged at 13,000 rpm for 30 min at 4°C, and the pellets were washed twice with 1 ml of cold 70% ethanol. Finally, the pellets were dried with a SpeedVac concentrator for 2 min and dissolved in RNase-free water. The quality and quantity of the RNAs were verified by Nano Drop, agarose gel, and Bioanalyzer (Agilent). Samples were kept at -80°C until use. No RNase treatment was applied to bacterial cell or to EV samples before RNA extraction.

RNA Sequencing

The RNA samples were sent to ViroScan3D[®] (Lyon, France) for DNA removal, ribosomal RNA depletion and RNA sequencing. Total RNA samples were submitted to a DNase treatment with RNase-Free DNase Set (Qiagen) according to manufacturer's instructions. Then, the samples quantified using the Quantifluor RNA system (Promega), and qualified using RNA Nano Chip on Bioanalyzer 2100 (Agilent) for the EV-producing cell samples, and on the SS RNA system on Fragment Analyzer (AATI) for the EVs samples. RNA samples were then submitted to the standard protocol Ovation Universal Prokaryotic RNA-Seq, Nugen, Anydeplete rRNA, library preparation. RNA quantity used for library preparation are displayed in Supplementary Table S1. The quality of libraries was assessed with the Quantifluor DNA system (Promega) and qualified with the HS-NGS system on Fragment Analyzer (Aati). The insert mean size of the libraries was 0.34 kp for the EV-producing cell samples, and 0.45 kp for the EV samples (**Supplementary Table S1**). Sequencing was performed with Illumina, NextSeq500, 75 cycles, single-read, High Output. For each experimental condition, three biological replicates were sequenced. EV-producing cell samples ranged from 9 to 27 million pair-end reads per sample, and EV samples ranged from 30 to 67 million pair-end reads per sample. Reads mapping to the reference genome ranged from 8 to 26 million, and from 0.38 to 27 million reads for EV-producing cells and EV samples, respectively. Basic statistics of the RNA-Seq data are displayed in **Supplementary Table S1**.

Transcriptome Analysis

The reads were cleaned and trimmed with Trim-Galore (Martin, 2011) using the default parameters. Reads were mapped with Bowtie2 (Langmead and Salzberg, 2013) in local mode against two staphylococcal genomes used as references: the NCTC 8325 (NC_007795.1) reference genome with sRNA annotation from SRD (Sassi et al., 2015) and the HG003 genome (GCA_000736455.1) for the non-annotated genes in NCTC8325 genome. Genes were counted with FeatureCounts (Liao et al., 2014) with the strand, the multi mapping, and the overlapping options.

A list of differentially expressed RNAs was obtained by EdgeR (Robinson et al., 2009) embedded in SARTools (Varet et al., 2016). The threshold of statistical significance was set to 0.05, with the adjustment method of Benjamini-Hochberg. RNA coverage was calculated with Bedtools coverage (Quinlan and Hall, 2010). RNAs with \geq 90% coverage in at least one EV condition were kept for further indepth analysis. RNA coverage visualization was performed with the Integrative Viewer Software (IGV; Thorvaldsdóttir et al., 2013) on a log scale.

Subcellular location prediction was performed with SurfG+ (Barinov et al., 2009). Clusters of Orthologous Groups (COGs) and KEGG categories were obtained using the eggNOG-mapper v2 web tool (Huerta-cepas et al., 2017, 2019). Functional enrichment analysis was performed with g:Profiler web-server (Raudvere et al., 2019; Reimand et al., 2019). A maximum value of p 0.05 was set as a threshold for significative categories.

A timepoint clustering study was conducted with the R package maSigPro (Conesa et al., 2006; Nueda et al., 2014) on highly covered EV RNAs with normalized counts by EdgeR. In this analysis vancomycin treatment is not taken into consideration. The threshold of statistical significance was set to 0.05, with the adjustment method of Benjamini-Hochberg.

RT-qPCR

Reverse transcription quantitative PCR (RT-qPCR) was used to validate RNA-seq results. EVs were isolated from the cellfree supernatants of three new independent *S. aureus* cultures at late-stationary growth phases (12 h) in the absence of vancomycin. EV RNAs were purified as mentioned above. Around 1.5 μ g of RNAs was treated with DNAse I (Amplification Grade, Invitrogen) according to manufacturer's instructions. cDNA synthesis was performed with the high capacity cDNA Reverse Transcription kit (Applied Biosystems). The primers used for

quantitative PCR (qPCR) are listed in Supplementary Table S2 and were designed using eprimer3 software (EMBOSS). qPCR was carried out in a 16 µl volume containing 15 ng cDNA, specific primers (300 nM), and 8 µl IQ[™] SYBR Green Supermix (Bio-Rad). Reactions were run on a CFX96 real-time system (Bio-Rad, France) using the following cycling parameters: DNA polymerase activation and DNA denaturation 95°C for 5 min, 40 cycles of denaturation at 94°C for 15 s, and extension at 60°C for 30 s. Melting curve analysis was included to check the amplification of single PCR products. Samples setups included biological triplicates and technical duplicates as well as negative controls corresponding to qPCR reactions performed without cDNA (cDNA negative control) and from RT reactions obtained from EV RNAs without reverse transcriptase (RT negative control). Results were analyzed with the GFX Manager software and Ct values were determined. Results with Ct equal or above 40 were considered negative and only experiments with $\Delta Ct \ge 4$ between negative controls and RT samples were considered.

RESULTS

S. aureus HG003 Produces EVs in Different Growth Conditions

Extracellular vesicles secreted by HG003 were isolated from the cell-free supernatants of bacterial cultures at early- and late-stationary growth phases (6 and 12 h, respectively), as well as in the absence (V-) and presence (V+) of a sublethal concentration of vancomycin (0.5 µg/ml). For that purpose, we used centrifugation, filtration, and density gradient ultracentrifugation, the standard method for EV isolation and purification at high purity (Yamada et al., 2012; Dauros Singorenko et al., 2017). EV homogeneity and integrity were evaluated by both negative staining electron microscopy and by NTA. Electron micrographs of purified EVs revealed typical nano-sized vesicular structures, with cup-shaped forms in all tested conditions (Raposo and Stoorvogel, 2013; Figure 1A and Supplementary Figure S2). NTA analyses showed a typical profile of particles for all EV samples (Figure 1B and Supplementary Figure S2). A significant increase of approximately 55% in EV diameter was observed in those purified from late-stationary phase cultures, compared to early-stationary phase cultures (for both 6V- vs. 12V- and 6V+ vs. 12V+), whereas no significant difference was observed in the absence or presence of vancomycin (6V- vs. 6V+ and 12V- vs. 12V+, Figure 1C). EV yield is essentially similar at the two growth phases, irrespective to the presence/absence of vancomycin (Figure 1D). In summary, S. aureus HG003 releases EVs with variable diameters depending on the growth phase. A sublethal concentration of vancomycin, however, does not impact the EV morphology, concentration, or diameter.

The *S. aureus* EVs Harbor all RNA Functional Classes

Total RNA was extracted from HG003 EVs to investigate their compositions. The quality of the RNA preparations was checked and validated, and the samples sequenced. RNA-seq data were



compared between the different growth conditions, and with those obtained from parental HG003 cells that produced the EVs in each condition (i.e., the EV-producing cells). Around 2649 ± 238 RNAs annotated in the HG003 genome were identified in the EVs according to growth conditions, with an average count of over five reads per RNA in each condition, whereas 3120 ± 35 annotated RNAs were identified within the EV-producing cells (Supplementary Tables S3 and S4). All the four main RNA functional classes (tRNAs, rRNAs, mRNAs, and sRNAs) were identified in both the purified EVs and the EV-producing cells (Figure 2A). In both the EVs and the EV-producing cells, ~84% of the mapped RNAs corresponded to protein-coding genes (mRNAs). The sRNAs were the second most abundant mapped RNA class (~12% of the reads). The remaining 4% of the mapped RNAs are tRNAs and residual rRNAs. Note that most of the rRNAs were voluntarily removed during the RNA purification. Most of the annotated mRNAs, tRNAs, and residual rRNAs were identified in the EV-producing parental cells (from 86 to 96%), while this value dropped to 79.8 ± 2.7% for sRNAs (Figure 2B). These percentages were slightly lower for EV samples (from 72 to 89%), although they remained high for the sRNAs (59.1 \pm 10.2%). All these RNAs detected in the purified EVs prompted us to check their coverages, to evaluate their integrity.

The *S. aureus* Purified EVs Contain Both Fragmented and Intact RNAs From Various Functional Classes

For the EV-producing cells, the median values of mRNAs, tRNAs, and residual rRNAs coverages were between 95 and 100%, and were 74 and 92% for the sRNAs (**Figure 3A** and **Supplementary Table S4**), implying that those RNAs were mainly intact, and not degraded. The coverage profile was drastically different for the RNAs recovered from the purified EVs. While the coverage of the residual rRNAs varied from 83 to 92%, the median coverage values for the other RNA functional classes ranged from 61 to 92% for the tRNAs, 15 to 47% for the mRNAs, and 4 to 20% for the sRNAs (**Figure 3A**). These lower coverages suggested that a substantial fraction of mRNAs and sRNAs were fragmented in the EVs compared

to the EV-producing cells. To analyze potentially functional RNAs in EVs, only RNAs with a coverage \geq 90% were considered for further analysis (**Supplementary Table S5**). The distribution of the newly filtered RNAs was depicted in **Figures 3B**,C. Such a harsh quality criterion impacted mainly the RNAs from the EVs, and particularly mRNAs and sRNAs. Only 3.5 ± 2.7 and $3.4 \pm 1.2\%$ of annotated mRNAs and sRNAs, respectively, were identified within EVs with such a threshold, while $67.9 \pm 7.2\%$ and $34.9 \pm 5.0\%$ of annotated mRNAs and sRNA were identified, respectively, for EV-producing cells (**Figure 3C**). Compared to the parental cells, the EVs were slightly depleted into mRNAs ($68.0 \pm 7.0\%$ for the EVs vs. $88.2 \pm 0.3\%$ in the EV-producing cells) but, interestingly, were enriched for the other RNA functional classes including the sRNAs ($14.3 \pm 3.2\%$ for the EVs vs. $8.6 \pm 0.4\%$ in the EV-producing cells, **Figure 3B**).

Functional Characterization of the RNAs From the EVs

According to experimental conditions, from 86 to 273 RNAs with a \geq 90% coverage and an average count of over 5 reads per RNA were identified within EVs from S. aureus HG003 (Figure 4A and Supplementary Table S5). They corresponded to 286 unique RNAs and were either mRNAs (220), tRNAs (28), residual rRNAs (10), and sRNAs (28). The presence of some of these transcrits associated with HG003 EVs and corresponding either to mRNAs or sRNAs was confirmed by RT-qPCR on RNAs extracted from three independent biological replicates (Figure 5). Among the mapped mRNAs, most were implicated in translation, ribosomal structure and biogenesis (17.5%, COG J), energy production and conversion (13.6%, COG C), carbohydrate transport and metabolism (COG G, 7.9%), transcription (5.2%, COG K), and cell wall/membrane/ envelope biogenesis (5.2%, COG M; Figure 4B). Several COG and KEGG categories were notably enriched (p < 0.05) in the EVs compared to the EV-producing bacteria (Figure 4C). mRNAs expressing proteins with a cytoplasmic location prediction were more represented in the EVs (79.5%) than into the producing cells (72.9%; Figure 4D). Interestingly, EVs contained several mRNAs coding for virulence-associated proteins such as the immune evasion protein A and Sbi, the Atl autolysin,



mapped reads (A) and annotated RNAs (B). RNA-Seq data is the average of three independent replicates. Number of reads have been normalized with EdgeR. RNA classes are defined from the S. *aureus* genome annotation NCTC8325/HG003. Early- and late-stationary growth phases (6 and 12, respectively) in absence (V–) or presence (V+) of vancomycin.







the Hld δ-hemolysin encoded by the multifunctional sRNA RNAIII, the PSMB1 Phenol Soluble Modulin, the FntA ironstorage ferritin, and the MntABC iron ABC transporter. Among the 20 tRNAs annotated in the genome, 15 were identified into the EVs (tRNA^{His}, tRNA^{Asn}, tRNA^{Glu}, tRNA^{Arg}, and tRNA^{Asp} were absent). Five copies of the 16S and 23S rRNAs were also detected, implying that our rRNA depletion procedure was incomplete. Finally, 28 annotated potential sRNAs were detected within EVs, and among the 50 or so bona fide sRNAs defined for the HG003 strain (Liu et al., 2018a), only RsaC was identified with a \geq 90% coverage in this study. Note that despite encoding the highly covered Hld transcript, the sRNA RNAIII presented only 71% gene coverage and therefore was excluded from analysis. Around 196 out of the 286 EV-associated RNAs colocalized at the same loci onto the HG003 chromosome, to form 42 clusters of 2 to 29 contiguous genes that were experimentally shown to belong to the same transcriptional units (Mäder et al., 2016; Supplementary Table S6). Among these transcriptional units, 17 displayed a RNA-Seq coverage \geq 90% across the entire operon in both the EVs and the EV-producing bacteria. **Figure 6** illustrates the sequencing coverage of various contiguous genes within the EVs and the EV-producing cells. Long mRNA operons, up to ~14,000 nucleotides, were detected as fully covered by reads into the purified EVs, supporting the presence of highly covered RNAs and operons as full-length transcripts.

EV RNA Composition Varies With Growth Conditions

The RNA composition of the purified EVs was compared between early- and late-stationary phases, and with or without vancomycin. Eighteen percentage (n = 51) of all detected RNAs with $\geq 90\%$ coverage were common to all the EV samples (**Figure 4A** and **Supplementary Table S5**), implying that the RNA content of the EVs highly varied according to the growth conditions. The percentage of RNAs shared by all the EV-producing cell samples, however, was much higher (70%, n = 1761). The shared RNAs among the EVs included mRNAs expressing virulence factors



FIGURE 5 | Reverse transcription quantitative PCR (RT-qPCR) validation of *S. aureus* HG003 EV RNAs. RT-qPCR experiments were performed from RNAs extracted from EV samples isolated from the cell-free supernatants of three independent *S. aureus* cultures at late-stationary growth phases (12 h) in the absence of vancomycin (Rep1, Rep2, and Rep3). Quantitative PCR (qPCR) successfully amplified the coding-sequence of 19 mRNAs, and 3 sRNAs. Samples setups included biological triplicates (Rep1, Rep2, and Rep3) and technical duplicates as well as negative controls corresponding to qPCR reactions performed without reverse transcriptase enzyme. Ct values are expressed as mean ± SD from two independent technical replicates performed in triplicates.

(Atl and Spa), metabolic enzymes (pyruvate dehydrogenase and cytochrome c oxidase complexes, glycolytic enzymes) and transcriptional regulators (SpxA, CggR, and GlnR), as well as RNAs involved in translation (ribosomal proteins, rRNAs, and tRNAs; **Supplementary Table S5**). The 51 common RNAs also included 9 potential sRNAs, notably RsaC involved in *S. aureus* oxidative stress adaptation and nutritional immunity (Lalaouna et al., 2019).

For both the EVs and EV-producing cells, more RNAs were detected at 6 h (275 and 2443 for EVs and EV-producing cells, respectively) than at 12 h (126 and 2161 for EVs and EV-producing cells, respectively, Figure 4A). Likewise, more RNAs were detected in the EVs in the presence (277 and 2,474 for EVs and EV-producing cells, respectively) than in the absence of vancomycin (140 and 2,279 for EVs and EV-producing cells, respectively; Figure 4A), indicating that the antibiotic modifies the RNA cargo of the EVs. These results also highlighted that the growth phase and the antibiotic stress impacted mostly the RNA content of the EVs, but much less that of the parental cells. Indeed, when we considered the RNAs detected in only one condition (i.e., specific RNAs), their fractions were higher in the EVs than in EV-producing cells, and that for all the tested conditions. For example, 58% of RNAs found within EVs at 6 h were specific to this condition, while specific RNAs represented only 14% of all RNAs detected at 6 h in the EV-producing cells.

EV RNA Abundance Varies With Growth Conditions

In addition to the qualitative variations observed, significant differences (Padj $\,<\,0.05)\,$ in EV RNA abundance between the experimental

conditions were also detected (Supplementary Table S7). Among the 286 EV-associated RNAs, 110 were differentially abundant between two conditions. Variations were detected at all times and in the absence or presence of vancomycin, although the growth phase appeared to have a greater impact on RNA abundance (75 and 64 differentially abundant RNAs were detected between early- and late-stationary phase with or without vancomycin, respectively), than the antibiotic stress (8 and 9 differentially abundant RNAs were detected between presence and absence of vancomycin in early- and late-stationary phase, respectively; Supplementary Table S7). A selection of RNAs with a modulation of their abundance according to the growth conditions is displayed in Figure 7. The most modulated RNAs into EVs produced from the two growth conditions were mRNAs coding for virulenceassociated factors, such as agrB, agrC, agrD, psmB1, and hld with a 30- to 1300-fold change, two potential annotated sRNAs, srn_0560, and srn_1000 with a 16- and 190-fold change, respectively, tRNAGly (SAOUHSC_T00025) and tRNAThr (SAOUHSC_T00054), with fold changes greater than 16. Among the differentially abundant RNAs according to the growth phase, 32 were detected both in presence and absence of vancomycin, with similar fold changes highlighting their reproducible variations into EVs across different environmental conditions (Figure 7 and Supplementary Table S7).

Differentially expressed RNAs were also detected for the EV-producing cells when their expression was compared between early- and late-stationary phase both in absence (n = 136) and in presence of vancomycin (n = 147), which was expected since bacterial transcription differs qualitatively



Integrative Viewer Software (IGV) in log scale.

and quantitatively when facing different growth conditions (**Supplementary Table S8**). Note that no significantly differentially expressed RNAs were detected according to the presence of vancomycin. As observed previously for the EV RNA content, the growth conditions, particularly the growth phase, impacted mostly the RNA abundance of the EVs, but much less than of the parental cells. Indeed, 38% of RNAs detected within EVs displayed changes in their abundance between conditions, while the fraction of modulated

RNAs counted for only 2% of the RNAs in the EV-producing cells. The abundance pattern of several RNAs differed between the EVs and the EV-producing cells according to the growth conditions (**Figure 7**). While some RNAs such as *agrBCD*, $psm\beta1$, and *hld* mRNAs displayed the same variations of their abundance pattern in EVs and EV-producing cells regardless the growth conditions. Others, such as *spa* and RsaC, were differentially abundant between the EVs and the EV-producing cells.



(Continued)

FIGURE 7 | EV-producing cells and EVs with a log2 fold change. Comparisons comprised data with at least one of the two samples containing ≥90% coverage. The log2 fold change is displayed as colored squares from -2 (purple) to 10 (red). Early- and late-stationary growth phases (6 and 12, respectively) in absence (V–) or presence (V+) of vancomycin.

Timepoint Clustering Analysis Reveals Different RNA Abundance Profiles Between the EVs and EV Producing Cells

To evaluate the influence of the growth phase on EV and EV producing cell RNA composition, a negative binomial-based approach, with the R package maSigPro (Conesa et al., 2006) was applied. Briefly, maSigPro provides a differentially expressed transcript analysis of serial data between experimental groups (e.g., EV and EV producing cells). maSigPro was applied to the 286 highly covered EV RNAs and identified 91 RNAs with significant temporal profile changes (Padj < 0.05). RNAs were clustered according to their expression profiles (Supplementary Table S9). Figure 8 shows the six RNA clusters obtained. Three clusters grouped transcripts with a similar expression profile between EV and EV-producing cells: cluster 4 with 8 RNAs (including, e.g., agrBD and arcC2) and cluster 3 with 18 RNAs (including, e.g., hld and $psm\beta 1$) contained more abundant transcripts over time in both EVs and EV-producing cells, whereas cluster 6 with 14 RNAs (including, e.g., fusA, tuf, and secY1) contained less abundant transcripts at 12 h than at 6 h in both. Interestingly, the three other clusters grouped RNAs that showed opposite expression profiles over time in EVs and EV-producing cells. Cluster 1 with 10 RNAs (including, e.g., RsaC and pdhA), and cluster 2 with 24 RNAs (including, e.g., ldh1, qoxABC, and rpoBC) grouped, similarly, more abundant transcripts at 12 h in EVs, and less abundant transcripts at 12 h in EV-producing cells. On the contrary, cluster 5 with 17 RNAs (including, e.g., ccgR and sbi) grouped less abundant transcripts in EVs and more abundant transcripts in EV-producing cells at 12 h. Altogether, this analysis highlighted that the transcript expression pattern could temporally differ between EVs and EV-producing cells.

DISCUSSION

Extracellular vesicles are universal carriers of macromolecules including extracellular RNAs all along from bacteria, archaea, and fungi to protists. Recent investigations on bacterial EV biogenesis, release, and trafficking showed their functional importance for bacterial communication and survival (Tsatsaronis et al., 2018). Information regarding *S. aureus* EV RNA cargo, however, is lagging behind. Here, we report the first exploratory work on EVs released by *S. aureus* HG003, the characterization of its EV RNA cargo under different conditions, and an indepth transcriptomic comparison between the EVs and the EV-producing cells.

Environmental conditions, such as growth phase and environmental stresses, reportedly influence the production, content and functions of EVs (Tashiro et al., 2010; Kim et al., 2016b; Orench-Rivera and Kuehn, 2016; Askarian et al., 2018;



Yun et al., 2018; Andreoni et al., 2019). Here, to investigate the impact of environmental changes on S. aureus EV production, the selected conditions were the early- and late-stationary growth phases, with or without a sublethal concentration of vancomycin that does not impact growth. Differences were observed regarding EV sizes. EVs derived from the late-stationary growth phase were larger than those collected during the earlystationary phase. This can be due to cell-wall morphology and peptidoglycan structure that are characteristics of the growth stage in S. aureus (Zhou and Cegelski, 2012). A correlation between the degree of peptidoglycan cross-linking to the cell wall stiffness and EVs release was observed for both Gramnegative and Gram-positive bacteria (Zhou et al., 1998; Deatherage et al., 2009; Schrempf and Merling, 2015; Schwechheimer et al., 2015; Wang et al., 2018). Notably, sub-inhibitory concentrations of penicillin decreases peptidoglycan cross-linking, triggering an increase in S. aureus EV yields and sizes (Wang et al., 2018). Here, we show that sublethal concentrations of vancomycin, an antibiotic that also targets peptidoglycan synthesis in S. aureus, does not impact either EV morphology or EV production yields, but does change their RNA content in terms of composition and abundance. Vancomycin affects S. aureus EV activity starting at 1 µg/ml (He et al., 2017). The sub-inhibitory vancomycin concentration

used here (0.5 $\mu g/ml)$ is probably too low to detect any changes in EV morphology and production.

All RNA classes are detected by RNA-Seq within HG003derived EVs. These include rRNAs, which were still detected, suggesting that the rRNA depletion carried out here was incomplete. In the absence of filtering by coverage of sequencing data, on average, 78.0 \pm 7.0% of the annotated transcripts in HG003 genome were present in the EVs (91.8 \pm 1.0% for EV-producing cells). Of these, a large portion of mapped RNAs corresponded to mRNAs. These results are consistent with RNA-Seq data obtained with similar criteria from OMVs in Salmonella enterica serovar Typhimurium (S. Typhimurium) that harbor around 73% of the annotated transcripts including up to 86% of mRNAs according to growth conditions (Malabirade et al., 2018). A recent study addressing the sRNA content of EVs also pointed out that mRNAs are the more abundant RNA species in EVs derived from S. aureus strain MSSA476 after rRNA depletion (Joshi et al., 2021). On the contrary, studies with Escherichia coli revealed that EVs were enriched mainly with short RNAs, such as tRNAs (Ghosal et al., 2015; Blenkiron et al., 2016). Nevertheless, these variations may be a result of different RNA extraction and library preparation protocols, or simply correspond to singular characteristics of EVs derived from different bacterial species. As expected, data

filtering of RNAs with \geq 90% coverage decreased the number of detected RNAs. However, the RNA content of the EVs, particularly the mRNAs and sRNAs (only 5.1 ± 2.8% of EV RNAs initially detected are still identified after the filtering), was much more affected than that of the EV-producing cells (69.0 ± 7.90% of EV-producing cell RNAs are still detected after the filtering). The low RNA coverage in the HG003 EVs might perhaps reflect the absence of transcription within EVs and, thus, the progressive degradation of a substantial fraction of the EVs-associated RNAs after their formation and/or during their purification. These findings are consistent with a recent report showing that the predominant RNA type in EVs from S. aureus Newman is <300-nucleotide long (Rodriguez and Kuehn, 2020). The presence of numerous processed or degraded RNAs could be a common feature of the bacterial EVs, as Gram-negative S. Typhimurium OMV-associated RNAs are also processed or degraded (Malabirade et al., 2018). Since one of the primary physiological functions attributed to EVs is the removal of unwanted materials from cells, such as misfolded or degraded proteins (McBroom and Kuehn, 2007), the S. aureus EVs may also help removing the degraded RNAs from the bacteria (Groot and Lee, 2020).

Two hundred and eighty-six highly covered RNAs can be identified within HG003 EVs. That number of highly covered EV RNAs varies according to growth conditions, from 86 in late-stationay growth phase with vancomycin to 273 in earlystationay growth phase also with vancomycin. As expected, all the EV transcrits are also detected in the EV-producing cells. Among them, 51 transcripts are shared by EVs collected in all tested conditions. The 286 highly covered EV RNAs encompass short transcripts, such as tRNAs (~75 nucleotides), mRNAs (psm
\$\beta1\$, 135 nucleotides; agrD, 141 nucleotides), and some sRNAs (Teg84, 79 nucleotides; Teg46, 124 nucleotides). Yet, these highly covered EV RNAs also comprise long transcripts, including 15 mRNAs with lengths >2,000 nucleotides (e.g., pbp2, copA, rpoB, rpoC, and atl). 67% of these RNAs are organized into 42 gene clusters that are co-transcribed in S. aureus (Mäder et al., 2016). Among them, 17 are full-length transcripts across entire operons in HG003 EVs, with lengths up to ~14,000 nucleotides. As observed for S. Typhimurium OMV RNAs (Malabirade et al., 2018), these findings support that highly covered RNAs are present as full-length transcripts. These RNAs belong to all annotated classes of RNAs. The mRNAs from the EVs encode proteins involved in transcription, translation, energy production and conversion, carbohydrate metabolism, and cell wall biogenesis. In addition to these housekeeping functions, EVs also harbors mRNAs encoding virulence-associated proteins, such as the *agr* operon responsible of quorum-sensing, autolysin Atl, protein A (Spa), immunoglobulin-binding protein (Sbi), immunodominant staphylococcal antigen B (IsaB), δ hemolysin (Hld) encoded by the multifunctional sRNA RNAIII, and the PSMB1 phenolsoluble modulin, as well as several iron acquisition systems. Besides, most tRNA species are detected within the EVs, as well as residual rRNAs and 28 annotated sRNAs, including bona-fide RsaC involved in S. aureus oxidative stress adaptation and nutritional immunity (Lalaouna et al., 2019). Note that RsaC and RNAIII were also detected within *S. aureus* EVs from strain MSSA476 (Joshi et al., 2021), suggesting their wider occurence in staphylococcal EVs.

The presence of full-length, functional RNAs in EVs raises the question of their biological roles. EVs are produced to transport bioactive molecules to interact and communicate with other cells. So far, most studies on S. aureus EVs investigated the protein cargo. Therefore, the broad spectrum of activities associated with S. aureus-derived EVs was related to their protein content (Jeon et al., 2016). In some S. aureus strains, EVs carry β-lactams that confer transient resistance to ampicillinsusceptible E. coli and S. aureus (Lee et al., 2013b). Likewise, mycobactin-containing Mycobacterium tuberculosis EVs can deliver iron to strains deficient for iron-uptake (Prados-Rosales et al., 2014). The delivery of full-length mRNAs via EVs to the surrounding bacterial cells, notably those involved into energetic and metabolic functions, could improve their responses to environmental stimuli to fasten their adaptation. Likewise, rRNAs and tRNAs could boost translation in EV recipient bacterial cells and improve their fitness. In bacteria, sRNAs fine tune target gene expression, usually at the posttranscriptional level in response to changes in the environment, including antibiotic resistance and tolerance (Mediati et al., 2021). Most bacteria encode dozens of sRNAs that are transcribed as independent transcripts or processed from mRNAs. The presence of sRNAs with regulatory roles within S. aureus EVs could be a relocation strategy in other surrounding bacteria that need more of these sRNAs for adaptation and infection spreading and/or to coordinate bacterial group adaptation, activities and behaviors. The transfer of RsaC could enhance the concentration of that riboregulator to other S. aureus cells when intracellular, especially helping survival within the phagolysosome if the S. aureus EVs are internalized by the host cells together with the bacteria. Transfer of functional RNAs to bacterial cells that do not encode the corresponding genes in their genome could also be part of a transient horizontal phenotype acquisition, which could be of use during infection to disseminate specific virulence-associated factors through the bacterial community. Finally, beside interactions between bacterial cells, S. aureus EV associated RNAs, notably sRNAs, may be involved in the host-pathogen interactions (Eberle et al., 2009; Li and Chen, 2012; Furuse et al., 2014; Sha et al., 2014; Koeppen et al., 2016; Westermann et al., 2016; Choi et al., 2017; Frantz et al., 2019; Han et al., 2019; Lee, 2019; Rodriguez and Kuehn, 2020). The 28 potential annotated sRNA detected within HG003 EVs are potential candidates for further functional characterization, especially during S. aureus-host cell interactions. S. aureus secreted EVs elicit immune responses that mimic those of the EV-producing cells (Gurung et al., 2011; Hong et al., 2011, 2014; Kim et al., 2012, 2019; Thay et al., 2013; Choi et al., 2015; Jeon et al., 2016; Jun et al., 2017; Askarian et al., 2018; Tartaglia et al., 2018; Wang et al., 2018, 2020; Rodriguez and Kuehn, 2020). Strikingly, within their RNA cargo, several mRNAs encode immunomodulatory proteins, as Sbi, Spa and PSMB1, and may participate into the immune response triggered by the protein cargo if they are ultimately translated. Yet, such functions remain to be demonstrated for the RNA cargo of S. aureus EVs. mRNAs expressing PSM β and hemolysin δ toxins from the EVs, if translated into recipient bacteria or host cells, could perhaps facilitate staphylococcal intracellular survival, but this hypothesis should be experimentally challenged.

The RNA cargo of HG003 EVs, in both identity and abundance, depends on the growth conditions. Similarly, the EV RNA cargo of S. Typhimurium is also sensitive to environmental changes indicating that it reflects the bacterial adaptation to its environment (Malabirade et al., 2018). It could be a faster way to transfer information of changes perceived by one cell to surrounding cells even before they sensed the environmental stimuli in order to quickly promote group adaptation. We found, however, that the vancomycin treatment had less impact on RNA abundance compared to the growth phase. Although the composition of HG003 EVs represented the intracellular state of the bacterial transcriptome through global packaging, two main findings, however, reinforce the concept of a potential selective packaging of RNAs into EVs, as proposed for its protein cargo (Haurat et al., 2011; Cahill et al., 2015; Tartaglia et al., 2020). First, we measured an enrichment for several functional and subcellular localization RNA categories in EVs when compared to EV-producing cells. Second, the relative abundance of several RNAs between two environmental conditions was different in the EVs and the EV-producing cells. Other studies also found that some RNA populations were enriched in EVs from Gramnegative and Gram-positive pathogenic bacteria (Ghosal et al., 2015; Koeppen et al., 2016; Resch et al., 2016; Malabirade et al., 2018; Malge et al., 2018; Han et al., 2019; Langlete et al., 2019; Zhang et al., 2020). This notably includes sRNAs, which can play regulatory activity in the host (Koeppen et al., 2016; Malabirade et al., 2018; Langlete et al., 2019; Zhang et al., 2020). The enrichment of RNAs associated with bacterial diseases in EVs derived from many pathogenic bacteria reinforces the physiopathological role of these structures in host-pathogen interaction and host cell invasion, which could be borne by their RNA cargo as well as by their protein cargo. The selective mechanisms of EV RNA content packaging have not yet been elucidated. It has been proposed that RNA packing into EVs could depend on RNA size and location (eg., nearby EVs formation site), as well as on their affinity for other molecules (eg., membrane proteins; Langlete et al., 2019). Nevertheless, such enrichment results should be interpreted with carefulness. Indeed, they could also reflect a difference in RNA half-lives between EVs and EV-producing cells, as well as a difference in RNase activity (Langlete et al., 2019; Lécrivain and Beckmann, 2020), pointing out that RNAs with longer half-lives could be protected from degradation, leading to an artifactual accumulation in the EVs over time.

In summary, our exploratory work provides novel insights in *S. aureus* EVs by the characterization of its RNA cargo and paves the way for further functional studies. Mainly, it sheds light on the possible roles of EV RNA cargo in intra- and inter-species communication, in the virulence and pathogenesis of *S. aureus*, and as trash bags for degraded RNAs. The study of bacterial EV RNA cargo is an emerging area of research. Evidently, as with all emerging fields, each advance raises further questions: Are the full-length RNAs in HG003 EVs functional, and do they

possess similar functions than in the bacterial cytoplasm? What are the rules for RNA sorting into HG003 EVs? What are the roles and functions of the *S. aureus* EV RNA cargo? These exciting questions, among others, should be adressed in further studies.

DATA AVAILABILITY STATEMENT

The datasets presented in this study can be found in online repositories. The names of the repository/repositories and accession number(s) can be found at: https://www.ebi.ac.uk/ena/browser/view/PRJEB40502.

AUTHOR CONTRIBUTIONS

BSRL, BF, SC, YL, VA, and EG conceived and designed the experiments. VA and EG supervised the study. BSRL and SC performed the experiments. AN performed computational analysis. BSRL, BF, SC, VRR, AN, and EG analyzed the data. VA, YL, and EG contributed to funding acquisition. BSRL and EG wrote the original draft. All authors contributed to data interpretation, drafting the manuscript, critically revising the manuscript, and approving its final version.

FUNDING

This work has received financial support from INRAE (Rennes, France) and Institut Agro (Rennes, France). This work was part of the CARAVEL project financed by the MICA division from INRAE. BSRL and VRR were supported by the International Cooperation Program CAPES/COFECUB at the Federal University of Minas Gerais funded by CAPES – the Brazilian Federal Agency for the Support and Evaluation of Graduate Education of the Brazilian Ministry of Education (number 88887.179897/2018-00 and 99999.000058/2017-03, respectively).

ACKNOWLEDGMENTS

This work was conducted in the frame of BactInflam International Associated Laboratory between INRAE (France) and UFMG (Brazil). It benefited from the facilities and expertise of the MRic-TEM platform (https://microscopie.univ-rennes1.fr). We are grateful to Agnes Burel (Univ Rennes, BIOSIT – UMS 3480, US_S 018, Rennes, France) for sessions with the microscope. We are also grateful to the genotoul bioinformatics platform Toulouse Occitanie (Bioinfo Genotoul) for providing computing and storage resources. Sandrine Parayre (INRAE, STLO) and Sandrine Péron (INRAE, STLO) are warmly acknowledged for the technical assistance during RT-qPCR experiments.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: https://www.frontiersin.org/articles/10.3389/fmicb.2021.634226/ full#supplementary-material

REFERENCES

- Ahmadi Badi, S., Bruno, S. P., Moshiri, A., Tarashi, S., Siadat, S. D., and Masotti, A. (2020). Small RNAs in outer membrane vesicles and their function in host-microbe interactions. *Front. Microbiol.* 11:1209. doi: 10.3389/ fmicb.2020.01209
- Andreoni, F., Toyofuku, M., Menzi, C., Kalawong, R., Shambat, S. M., François, P., et al. (2019). Antibiotics stimulate formation of vesicles in *Staphylococcus aureus* in both phage-dependent and -independent fashions and via different routes. *Antimicrob. Agents Chemother.* 63, e01439–e01518. doi: 10.1128/ AAC.01439-18
- Askarian, F., Lapek, J. D., Dongre, M., Tsai, C. M., Kumaraswamy, M., Kousha, A., et al. (2018). *Staphylococcus aureus* membrane-derived vesicles promote bacterial virulence and confer protective immunity in murine infection models. *Front. Microbiol.* 9:262. doi: 10.3389/fmicb.2018.00262
- Barinov, A., Loux, V., Hammani, A., Nicolas, P., Langella, P., Ehrlichh, D., et al. (2009). Prediction of surface exposed proteins in *Streptococcus pyogenes*, with a potential application to other gram-positive bacteria. *Proteomics* 9, 61–73. doi: 10.1002/pmic.200800195.
- Biller, S. J., Schubotz, F., Roggensack, S. E., Thompson, A. W., Summons, R. E., and Chisholm, S. W. (2014). Bacterial vesicles in marine ecosystems. *Science* 343, 183–186. doi: 10.1126/science.1243457
- Bitto, N. J., Chapman, R., Pidot, S., Costin, A., Lo, C., Choi, J., et al. (2017). Bacterial membrane vesicles transport their DNA cargo into host cells. *Sci. Rep.* 7:7072. doi: 10.1038/s41598-017-07288-4
- Blenkiron, C., Simonov, D., Muthukaruppan, A., Tsai, P., Dauros, P., Green, S., et al. (2016). Uropathogenic *Escherichia coli* releases extracellular vesicles that are associated with RNA. *PLoS One* 11:e0160440. doi: 10.1371/journal. pone.0160440
- Brown, L., Kessler, A., Cabezas-Sanchez, P., Luque-Garcia, J. L., and Casadevall, A. (2014). Extracellular vesicles produced by the gram-positive bacterium *Bacillus subtilis* are disrupted by the lipopeptide surfactin. *Mol. Microbiol.* 93, 183–198. doi: 10.1111/mmi.12650
- Brown, L., Wolf, J. M., Prados-Rosales, R., and Casadevall, A. (2015). Through the wall: extracellular vesicles in gram-positive bacteria, mycobacteria and fungi. *Nat. Rev. Microbiol.* 13, 620–630. doi: 10.1038/nrmicro3480
- Cahill, B. K., Seeley, K. W., Gutel, D., and Ellis, T. N. (2015). *Klebsiella pneumoniae* O antigen loss alters the outer membrane protein composition and the selective packaging of proteins into secreted outer membrane vesicles. *Microbiol. Res.* 180, 1–10. doi: 10.1016/j.micres.2015.06.012
- Choi, J. W., Kim, S. C., Hong, S. H., and Lee, H. J. (2017). Secretable small RNAs via outer membrane vesicles in periodontal pathogens. J. Dent. Res. 96, 458–466. doi: 10.1177/0022034516685071
- Choi, S. J., Kim, M. H., Jeon, J., Kim, O. Y., Choi, Y., Seo, J., et al. (2015). Active immunization with extracellular vesicles derived from *Staphylococcus aureus* effectively protects against staphylococcal lung infections, mainly via Th1 cell-mediated immunity. *PLoS One* 10:e0136021. doi: 10.1371/journal. pone.0136021
- Conesa, A., Nueda, M. J., Ferrer, A., and Talón, M. (2006). maSigPro: a method to identify significantly differential expression profiles in time-course microarray experiments. *Bioinformatics* 22, 1096–1102. doi: 10.1093/bioinformatics/btl056
- Dauros Singorenko, P., Chang, V., Whitcombe, A., Simonov, D., Hong, J., Phillips, A., et al. (2017). Isolation of membrane vesicles from prokaryotes: a technical and biological comparison reveals heterogeneity. J. Extracell. Vesicles 6:1324731. doi: 10.1080/20013078.2017.1324731
- Dauros-Singorenko, P., Blenkiron, C., Phillips, A., and Swift, S. (2018). The functional RNA cargo of bacterial membrane vesicles. *FEMS Microbiol. Lett.* 365, 1–9. doi: 10.1093/femsle/fny023
- Deatherage, B. L., Lara, J. C., Bergsbaken, T., Barrett, S. L. R., Lara, S., and Cookson, B. T. (2009). Biogenesis of bacterial membrane vesicles. *Mol. Microbiol.* 72, 1395–1407. doi: 10.1111/j.1365-2958.2009.06731.x
- Eberle, F., Sirin, M., Binder, M., and Dalpke, A. H. (2009). Bacterial RNA is recognized by different sets of immunoreceptors. *Eur. J. Immunol.* 39, 2537–2547. doi: 10.1002/eji.200838978
- Foster, T. J. (2005). Immune evasion by staphylococci. Nat. Rev. Microbiol. 3, 948–958. doi: 10.1038/nrmicro1289
- Frantz, R., Teubner, L., Schultze, T., Pietra, L., Müller, C., Gwozdzinski, K., et al. (2019). The secRNome of *Listeria monocytogenes* harbors small noncoding

RNAs that are potent inducers of Beta interferon. Am. Soc. Microbiol. 10, 1-15. doi: 10.1128/mBio.01223-19

- Furuse, Y., Finethy, R., Saka, H. A., Xet-Mull, A. M., Sisk, D. M., Jurcic Smith, K. L., et al. (2014). Search for MicroRNAs expressed by intracellular bacterial pathogens in infected mammalian cells. *PLoS One* 9:e106434. doi: 10.1371/journal.pone.0106434
- Ghosal, A., Upadhyaya, B. B., Fritz, J. V., Heintz-Buschart, A., Desai, M. S., Yusuf, D., et al. (2015). The extracellular RNA complement of *Escherichia* coli. Microbiologyopen 4, 252–266. doi: 10.1002/mbo3.235
- Gill, S., Catchpole, R., and Forterre, P. (2019). Extracellular membrane vesicles in the three domains of life and beyond. *FEMS Microbiol. Rev.* 43, 273–303. doi: 10.1093/femsre/fuy042
- Gill, S. R., McIntyre, L. M., Nelson, C. L., Remortel, B., Rude, T., Reller, L. B., et al. (2011). Potential associations between severity of infection and the presence of virulence-associated genes in clinical strains of *Staphylococcus aureus*. *PLoS One* 6:e18673. doi: 10.1371/journal.pone.0018673
- Groot, M., and Lee, H. (2020). Sorting mechanisms for MicroRNAs into extracellular vesicles and their associated diseases. *Cell* 9, 1–16. doi: 10.3390/ cells9041044.
- Gurung, M., Moon, D. C., Choi, C. W., Lee, J. H., Bae, Y. C., Kim, J., et al. (2011). *Staphylococcus aureus* produces membrane-derived vesicles that induce host cell death. *PLoS One* 6:e27958. doi: 10.1371/journal.pone.0027958
- Han, E. C., Choi, S. Y., Lee, Y., Park, J. W., Hong, S. H., and Lee, H. J. (2019). Extracellular RNAs in periodontopathogenic outer membrane vesicles promote TNF-α production in human macrophages and cross the blood-brain barrier in mice. *FASEB J.* 33, 13412–13422. doi: 10.1096/fj.201901575R
- Haurat, M. F., Aduse-Opoku, J., Rangarajan, M., Dorobantu, L., Gray, M. R., Curtis, M. A., et al. (2011). Selective sorting of cargo proteins into bacterial membrane vesicles. *J. Biol. Chem.* 286, 1269–1276. doi: 10.1074/jbc. M110.185744
- He, X., Yuan, F., Lu, F., Yin, Y., and Cao, J. (2017). Vancomycin-induced biofilm formation by methicillin-resistant *Staphylococcus aureus* is associated with the secretion of membrane vesicles. *Microb. Pathog.* 110, 225–231. doi: 10.1016/j.micpath.2017.07.004
- Herbert, S., Ziebandt, A. K., Ohlsen, K., Schäfer, T., Hecker, M., Albrecht, D., et al. (2010). Repair of global regulators in *Staphylococcus aureus* 8325 and comparative analysis with other clinical isolates. *Infect. Immun.* 78, 2877–2889. doi: 10.1128/IAI.00088-10
- Ho, M., Chen, C., Goodwin, J. S., Wang, B., and Xie, H. (2015). Functional advantages of *Porphyromonas gingivalis* vesicles. *PLoS One* 10:e0123448. doi: 10.1371/journal.pone.0123448
- Hong, S. W., Choi, E. B., Min, T. K., Kim, J. H., Kim, M. H., Jeon, S. G., et al. (2014). An important role of α -hemolysin in extracellular vesicles on the development of atopic dermatitis induced by *Staphylococcus aureus*. *PLoS One* 9:e100499. doi: 10.1371/journal.pone.0100499
- Hong, S. W., Kim, M. R., Lee, E. Y., Kim, J. H., Kim, Y. S., Jeon, S. G., et al. (2011). Extracellular vesicles derived from *Staphylococcus aureus* induce atopic dermatitis-like skin inflammation. *Allergy* 66, 351–359. doi: 10.1111/j. 1398-9995.2010.02483.x
- Hsu, C. Y., Lin, M. H., Chen, C. C., Chien, S. C., Cheng, Y. H., Su, I. N., et al. (2011). Vancomycin promotes the bacterial autolysis, release of extracellular DNA, and biofilm formation in vancomycin-non-susceptible *Staphylococcus aureus. FEMS Immunol. Med. Microbiol.* 63, 236–247. doi: 10.1111/j.1574-695X.2011.00846.x
- Huerta-cepas, J., Forslund, K., Coelho, L. P., Szklarczyk, D., Jensen, L. J., Mering, C.Von, et al. (2017). Fast genome-wide functional annotation through orthology assignment by eggNOG-mapper. *Mol. Biol. Evol.* 34, 2115–2122. doi: 10.1093/molbev/msx148
- Huerta-Cepas, J., Szklarczyk, D., Heller, D., Hernández-Plaza, A., Forslund, S. K., Cook, H., et al. (2019). EggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Res.* 47, D309–D314. doi: 10.1093/nar/gky1085
- Jeon, H., Oh, M. H., Jun, S. H., Kim, S. I., Choi, C. W., Kwon, H. I., et al. (2016). Variation among *Staphylococcus aureus* membrane vesicle proteomes affects cytotoxicity of host cells. *Microb. Pathog.* 93, 185–193. doi: 10.1016/j. micpath.2016.02.014
- Joshi, B., Singh, B., Nadeem, A., Askarian, F., Wai, S. N., Johannessen, M., et al. (2021). Transcriptome profiling of *Staphylococcus aureus* associated

extracellular vesicles reveals presence of small RNA-cargo. Front. Mol. Biosci. 7:566207. doi: 10.3389/fmolb.2020.566207

- Jun, S. H., Lee, J. H., Kim, S. I., Choi, C. W., Park, T. I., Jung, H. R., et al. (2017). Staphylococcus aureus-derived membrane vesicles exacerbate skin inflammation in atopic dermatitis. *Clin. Exp. Allergy* 47, 85–96. doi: 10.1111/cea.12851
- Kaparakis-Liaskos, M., and Ferrero, R. L. (2015). Immune modulation by bacterial outer membrane vesicles. *Nat. Rev. Immunol.* 15, 375–387. doi: 10.1038/nri3837
- Kim, J., Bin, B. H., Choi, E. J., Lee, H. G., Lee, T. R., and Cho, E. G. (2019). *Staphylococcus aureus*-derived extracellular vesicles induce monocyte recruitment by activating human dermal microvascular endothelial cells in vitro. *Clin. Exp. Allergy* 49, 68–81. doi: 10.1111/cea.13289
- Kim, Y., Edwards, N., and Fenselau, C. (2016b). Extracellular vesicle proteomes reflect developmental phases of *Bacillus subtilis*. *Clin. Proteomics* 13, 1–8. doi: 10.1186/s12014-016-9107-z.
- Kim, M. R., Hong, S. W., Choi, E. B., Lee, W. H., Kim, Y. S., Jeon, S. G., et al. (2012). *Staphylococcus aureus*-derived extracellular vesicles induce neutrophilic pulmonary inflammation via both Th1 and Th17 cell responses. *Allergy Eur. J. Allergy Clin. Immunol.* 67, 1271–1281. doi: 10.1111/all.12001
- Kim, J. H., Jeun, E. J., Hong, C. P., Kim, S. H., Jang, M. S., Lee, E. J., et al. (2016a). Extracellular vesicle-derived protein from *Bifidobacterium longum* alleviates food allergy through mast cell suppression. *J. Allergy Clin. Immunol.* 137, 507–516. doi: 10.1016/j.jaci.2015.08.016
- Kim, J. H., Lee, J., Park, J., and Gho, Y. S. (2015). Gram-negative and grampositive bacterial extracellular vesicles. *Semin. Cell Dev. Biol.* 40, 97–104. doi: 10.1016/j.semcdb.2015.02.006
- Koeppen, K., Hampton, T. H., Jarek, M., Scharfe, M., Gerber, S. A., Mielcarz, D. W., et al. (2016). A novel mechanism of host-pathogen interaction through sRNA in bacterial outer membrane vesicles. *PLoS Pathog.* 12:e1005672. doi: 10.1371/journal.ppat.1005672
- Lalaouna, D., Baude, J., Wu, Z., Tomasini, A., Chicher, J., Marzi, S., et al. (2019). RsaC sRNA modulates the oxidative stress response of *Staphylococcus aureus* during manganese starvation. *Nucleic Acids Res.* 47, 9871–9887. doi: 10.1093/nar/gkz728
- Langlete, P., Krabberød, A. K., and Winther-Larsen, H. C. (2019). Vesicles from Vibrio cholerae contain AT-rich DNA and shorter mRNAs that do not correlate with their protein products. *Front. Microbiol.* 10:2708. doi: 10.3389/fmicb.2019.02708
- Langmead, B., and Salzberg, S. (2013). Fast gapped-read alignment with bowtie 2. Nat. Methods 9, 357-359. doi: 10.1038/nmeth.1923.Fast
- Lécrivain, A. L., and Beckmann, B. M. (2020). Bacterial RNA in extracellular vesicles: a new regulator of host-pathogen interactions? *Biochim. Biophys. Acta-Gene Regul. Mech.* 1863:194519. doi: 10.1016/j.bbagrm.2020.194519
- Lee, H. J. (2019). Microbe-host communication by small RNAs in extracellular vesicles: vehicles for transkingdom RNA transportation. *Int. J. Mol. Sci.* 20:1487. doi: 10.3390/ijms20061487
- Lee, E. Y., Choi, D. Y., Kim, D. K., Kim, J. W., Park, J. O., Kim, S., et al. (2009). Gram-positive bacteria produce membrane vesicles: proteomics-based characterization of *Staphylococcus aureus*-derived membrane vesicles. *Proteomics* 9, 5425–5436. doi: 10.1002/pmic.200900338
- Lee, J. H., Choi, C. W., Lee, T., Kim, S. I., Lee, J. C., and Shin, J. H. (2013a). Transcription factor σB plays an important role in the production of extracellular membrane-derived vesicles in *Listeria monocytogenes*. *PLoS One* 8:e73196. doi: 10.1371/journal.pone.0073196
- Lee, J., Lee, E. Y., Kim, S. H., Kim, D. K., Park, K. S., Kim, K. P., et al. (2013b). *Staphylococcus aureus* extracellular vesicles carry biologically active β-lactamase. *Antimicrob. Agents Chemother.* 57, 2589–2595. doi: 10.1128/ AAC.00522-12.
- Li, X. -D., and Chen, Z. J. (2012). Sequence specific detection of bacterial 23S ribosomal RNA by TLR13. *elife* 1:e00102. doi: 10.7554/eLife.00102.
- Liao, Y., Smyth, G. K., and Shi, W. (2014). FeatureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* 30, 923–930. doi: 10.1093/bioinformatics/btt656
- Liu, Y., Defourny, K. A. Y., Smid, E. J., and Abee, T. (2018b). Gram-positive bacterial extracellular vesicles and their impact on health and disease. *Front. Microbiol.* 9:1502. doi: 10.3389/fmicb.2018.01502
- Liu, W., Rochat, T., Toffano-Nioche, C., Le Lam, T. N., Bouloc, P., and Morvan, C. (2018a). Assessment of bona fide sRNAs in *Staphylococcus aureus*. Front. Microbiol. 9:228. doi: 10.3389/fmicb.2018.00228

- Mäder, U., Nicolas, P., Depke, M., Pané-Farré, J., Debarbouille, M., van der Kooi-Pol, M. M., et al. (2016). *Staphylococcus aureus* Transcriptome architecture: from laboratory to infection-mimicking conditions. *PLoS Genet*. 12:e1005962. doi: 10.1371/journal.pgen.1005962
- Malabirade, A., Habier, J., Heintz-Buschart, A., May, P., Godet, J., Halder, R., et al. (2018). The RNA complement of outer membrane vesicles from *Salmonella enterica* Serovar Typhimurium under distinct culture conditions. *Front. Microbiol.* 9:2015. doi: 10.3389/fmicb.2018.02015
- Malge, A., Ghai, V., Reddy, P. J., Baxter, D., Kim, T., Moritz, R. L., et al. (2018). mRNA transcript distribution bias between *Borrelia burgdorferi* bacteria and their outer membrane vesicles. *FEMS Microbiol. Lett.* 365, 1–9. doi: 10.1093/femsle/fny135
- Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet. J.* 17, 10–22. doi: 10.14806/ej.17.1.200
- McBroom, A. J., and Kuehn, M. J. (2007). Release of outer membrane vesicles by gram-negative bacteria is a novel envelope stress response. *Mol. Microbiol.* 63, 545–558. doi: 10.1111/j.1365-2958.2006.05522.x
- Mediati, D. G., Wu, S., Wu, W., and Tree, J. J. (2021). Networks of resistance: small RNA control of antibiotic resistance. *Trends Genet.* 37, 35–45. doi: 10.1016/j.tig.2020.08.016
- Nueda, M. J., Tarazona, S., and Conesa, A. (2014). Next maSigPro: updating maSigPro bioconductor package for RNA-seq time series. *Bioinformatics* 30, 2598–2602. doi: 10.1093/bioinformatics/btu333
- Olaya-Abril, A., Prados-Rosales, R., McConnell, M. J., Martín-Peña, R., González-Reyes, J. A., Jiménez-Munguía, I., et al. (2014). Characterization of protective extracellular membrane-derived vesicles produced by *Streptococcus pneumoniae*. J. Proteome 106, 46–60. doi: 10.1016/j.jprot.2014.04.023.
- Orench-Rivera, N., and Kuehn, M. (2016). Environmentally-controlled bacterial vesicle-mediated export. *Cell. Microbiol.* 18, 1525–1536. doi: 10.1111/ cmi.12676
- Perez Vidakovics, M. L. A., Jendholm, J., Mörgelin, M., Månsson, A., Larsson, C., Cardell, L. O., et al. (2010). B cell activation by outer membrane vesicles - a novel virulence mechanism. *PLoS Pathog.* 6:e1000724. doi: 10.1371/ journal.ppat.1000724
- Prados-Rosales, R., Baena, A., Martinez, L. R., Luque-Garcia, J., Kalscheuer, R., Veeraraghavan, U., et al. (2011). Mycobacteria release active membrane vesicles that modulate immune responses in a TLR2-dependent manner in mice. J. Clin. Invest. 121, 1471–1483. doi: 10.1172/JCI44261
- Prados-Rosales, R., Weinrick, B. C., Piqué, D. G., Jacobs, W. R., Casadevall, A., and Rodriguez, G. M. (2014). Role for *Mycobacterium tuberculosis* membrane vesicles in iron acquisition. *J. Bacteriol.* 196, 1250–1256. doi: 10.1128/ JB.01090-13
- Quinlan, A. R., and Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* 26, 841–842. doi: 10.1093/ bioinformatics/btq033
- Raposo, G., and Stoorvogel, W. (2013). Extracellular vesicles: Exosomes, microvesicles, and friends. J. Cell Biol. 200, 373–383. doi: 10.1083/jcb.201211138
- Raudvere, U., Kolberg, L., Kuzmin, I., Arak, T., Adler, P., Peterson, H., et al. (2019). G:profiler: a web server for functional enrichment analysis and conversions of gene lists (2019 update). *Nucleic Acids Res.* 47, W191–W198. doi: 10.1093/nar/gkz369
- Reimand, J., Isserlin, R., Voisin, V., Kucera, M., Tannus-Lopes, C., Rostamianfar, A., et al. (2019). Pathway enrichment analysis and visualization of omics data using g:profiler, GSEA, Cytoscape and EnrichmentMap. *Nat. Protoc.* 14, 482–517. doi: 10.1038/s41596-018-0103-9
- Resch, U., Tsatsaronis, J. A., Le Rhun, A., Stübiger, G., Rohde, M., Kasvandik, S., et al. (2016). A two-component regulatory system impacts extracellular membrane-derived vesicle production in group a streptococcus. *MBio* 7, 1–10. doi: 10.1128/mBio.00207-16.
- Rivera, J., Cordero, R. J. B., Nakouzi, A. S., Frases, S., Nicola, A., and Casadevall, A. (2010). *Bacillus anthracis* produces membrane-derived vesicles containing biologically active toxins. *Proc. Natl. Acad. Sci. U. S. A.* 107, 19002–19007. doi: 10.1073/pnas.1008843107
- Robinson, M. D., McCarthy, D. J., and Smyth, G. K. (2009). edgeR: a bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26, 139–140. doi: 10.1093/bioinformatics/btp616
- Rodovalho, V. D. R., Luz, B. S. R. D., Rabah, H., do Carmo, F. L. R., Folador, E. L., Nicolas, A., et al. (2020). Extracellular vesicles produced by the probiotic *Propionibacterium freudenreichii* CIRM-BIA 129 mitigate inflammation by

modulating the NF-κB pathway. Front. Microbiol. 11:1544. doi: 10.3389/ fmicb.2020.01544

- Rodriguez, B. V., and Kuehn, M. J. (2020). Staphylococcus aureus secretes immunomodulatory RNA and DNA via membrane vesicles. Sci. Rep. 10, 1–22. doi: 10.1038/s41598-020-75108-3.
- Salgado-Pabón, W., and Schlievert, P. M. (2014). Models matter: the search for an effective *Staphylococcus aureus* vaccine. *Nat. Rev. Microbiol.* 12, 585–591. doi: 10.1038/nrmicro3308
- Sassi, M., Augagneur, Y., Mauro, T., Ivain, L., Chabelskaya, S., Hallier, M., et al. (2015). SRD: a staphylococcus regulatory RNA database. *RNA* 21, 1005–1017. doi: 10.1261/rna.049346.114
- Sassi, M., Felden, B., and Augagneur, Y. (2014). Draft genome sequence of Staphylococcus aureus subsp. aureus strain HG003, an NCTC8325 derivative. Genome Announc. 2, 6–7. doi: 10.1128/genomeA.00855-14.Copyright
- Schrempf, H., and Merling, P. (2015). Extracellular Streptomyces lividans vesicles: composition, biogenesis and antimicrobial activity. Microb. Biotechnol. 8, 644–658. doi: 10.1111/1751-7915.12274
- Schwechheimer, C., Rodriguez, D. L., and Kuehn, M. J. (2015). NlpI-mediated modulation of outer membrane vesicle production through peptidoglycan dynamics in *Escherichia coli*. *Microbiology* 4, 375–389. doi: 10.1002/mbo3.244
- Sha, W., Mitoma, H., Hanabuchi, S., Bao, M., Weng, L., Sugimoto, N., et al. (2014). Human NLRP3 Inflammasome senses multiple types of bacterial RNAs. *Proc. Natl. Acad. Sci. U. S. A.* 111, 16059–16064. doi: 10.1073/ pnas.1412487111
- Sjöström, A. E., Sandblad, L., Uhlin, B. E., and Wai, S. N. (2015). Membrane vesicle-mediated release of bacterial RNA. *Nat. Publ. Gr.* 5:15329. doi: 10.1038/ srep15329
- Tartaglia, N. R., Breyne, K., Meyer, E., Cauty, C., Jardin, J., Chrétien, D., et al. (2018). *Staphylococcus aureus* extracellular vesicles elicit an immunostimulatory response in vivo on the murine mammary gland. *Front. Cell. Infect. Microbiol.* 8:277. doi: 10.3389/fcimb.2018.00277
- Tartaglia, N. R., Nicolas, A., Rodovalho, V. de R., Luz, B. S. R. D., Briard-Bion, V., Krupova, Z., et al. (2020). Extracellular vesicles produced by human and animal *Staphylococcus aureus* strains share a highly conserved core proteome. *Sci. Rep.* 10:8467. doi: 10.1038/s41598-020-64952-y.
- Tashiro, Y., Ichikawa, S., Shimizu, M., Toyofuku, M., Takaya, N., Nakajima-Kambe, T., et al. (2010). Variation of physiochemical properties and cell association activity of membrane vesicles with growth phase in *Pseudomonas aeruginosa. Appl. Environ. Microbiol.* 76, 3732–3739. doi: 10.1128/ AEM.02794-09
- Thay, B., Wai, S. N., and Oscarsson, J. (2013). Staphylococcus aureus α -toxindependent induction of host cell death by membrane-derived vesicles. *PLoS One* 8:e54661. doi: 10.1371/journal.pone.0054661
- Thorvaldsdóttir, H., Robinson, J. T., and Mesirov, J. P. (2013). Integrative genomics viewer (IGV): high-performance genomics data visualization and exploration. *Brief. Bioinform.* 14, 178–192. doi: 10.1093/bib/bbs017
- Tong, S. Y. C., Davis, J. S., Eichenberger, E., Holland, T. L., and Fowler, V. G. (2015). *Staphylococcus aureus* infections: epidemiology, pathophysiology, clinical manifestations, and management. *Clin. Microbiol. Rev.* 28, 603–661. doi: 10.1128/CMR.00134-14
- Toyofuku, M. (2019). Bacterial communication through membrane vesicles. Biosci. Biotechnol. Biochem. 83, 1599–1605. doi: 10.1080/09168451.2019.1608809
- Tsatsaronis, J. A., Franch-Arroyo, S., Resch, U., and Charpentier, E. (2018). Extracellular vesicle RNA: a universal mediator of microbial communication? *Trends Microbiol.* 26, 401–410. doi: 10.1016/j.tim.2018.02.009

- Varet, H., Brillet-Guéguen, L., Coppée, J. Y., and Dillies, M. A. (2016). SARTools: a DESeq2- and edgeR-based R pipeline for comprehensive differential analysis of RNA-Seq data. *PLoS One* 11:e0157022. doi: 10.1371/journal.pone.0157022
- Wang, X., Eagen, W. J., and Lee, J. C. (2020). Orchestration of human macrophage NLRP3 inflammasome activation by *Staphylococcus aureus* extracellular vesicles. *Proc. Natl. Acad. Sci. U. S. A.* 117, 3174–3184. doi: 10.1073/ pnas.1915829117
- Wang, X., Thompson, C. D., Weidenmaier, C., and Lee, J. C. (2018). Release of *Staphylococcus aureus* extracellular vesicles and their application as a vaccine platform. *Nat. Commun.* 9:1379. doi: 10.1038/s41467-018-03847-z
- Wertheim, H. F. L., Melles, D. C., Vos, M. C., Van Leeuwen, W., Van Belkum, A., Verbrugh, H. A., et al. (2005). The role of nasal carriage in *Staphylococcus aureus* infections. *Lancet Infect. Dis.* 5, 751–762. doi: 10.1016/S1473-3099(05)70295-4
- Westermann, A. J., Förstner, K. U., Amman, F., Barquist, L., Chao, Y., Schulte, L. N., et al. (2016). Dual RNA-seq unveils noncoding RNA functions in hostpathogen interactions. *Nature* 529, 496–501. doi: 10.1038/nature16547
- Work, E., Knox, K. W., and Vesk, M. (1966). The chemistry and electron microscopy of an extracellular lipopolysaccharide from *Escherichia coli. Ann.* N. Y. Acad. Sci. 133, 438–449. doi: 10.1111/j.1749-6632.1966.tb52382.x
- Yamada, T., Inoshima, Y., Matsuda, T., and Ishiguro, N. (2012). Comparison of methods for isolating exosomes from bovine milk. J. Vet. Med. Sci. 74, 1523–1525. doi: 10.1292/jvms.12-0032
- Yu, Y. J., Wang, X. H., and Fan, G. C. (2018). Versatile effects of bacteriumreleased membrane vesicles on mammalian cells and infectious/inflammatory diseases. *Acta Pharmacol. Sin.* 39, 514–533. doi: 10.1038/aps.2017.82
- Yun, S. H., Park, E. C., Lee, S. Y., Lee, H., Choi, C. W., Yi, Y. S., et al. (2018). Antibiotic treatment modulates protein components of cytotoxic outer membrane vesicles of multidrug-resistant clinical strain, *Acinetobacter baumannii* DU202. *Clin. Proteomics* 15, 1–11. doi: 10.1186/s12014-018-9204-2.
- Zhang, H., Zhang, Y., Song, Z., Li, R., Ruan, H., Liu, Q., et al. (2020). sncRNAs packaged by *Helicobacter pylori* outer membrane vesicles attenuate IL-8 secretion in human cells. *Int. J. Med. Microbiol.* 310:151356. doi: 10.1016/j. ijmm.2019.151356
- Zhou, X., and Cegelski, L. (2012). Nutrient-dependent structural changes in Staphylococcus aureus peptidoglycan revealed by solid-state NMR spectroscopy. Biochemistry 51, 8143–8153. doi: 10.1021/bi3012115
- Zhou, L., Srisatjaluk, R., Justus, D. E., and Doyle, R. J. (1998). On the origin of membrane vesicles in gram-negative bacteria. *FEMS Microbiol. Lett.* 163, 223–228. doi: 10.1111/j.1574-6968.1998.tb13049.x
- Ziebuhr, W. (2001). *Staphylococcus aureus* and *Staphylococcus epidermidis*: emerging pathogens in nosocomial infections. *Contrib. Microbiol.* 8, 102–107. doi: 10.1159/000060402.

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Da Luz, Nicolas, Chabelskaya, Rodovalho, Le Loir, Azevedo, Felden and Guédon. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.