

Effet of exogenous matter on soil biodiversity and ecosystem services in tomato plantations

Coraline Moulin, Victor Vaillant, Philippe Julianus, Chantal Flereau, Pascale Bade, Fred Burner, Gladys Loranger-Merciris

▶ To cite this version:

Coraline Moulin, Victor Vaillant, Philippe Julianus, Chantal Flereau, Pascale Bade, et al.. Effet of exogenous matter on soil biodiversity and ecosystem services in tomato plantations. 52. Annual meeting of the Caribbean Food Crops Society (CFCS), Jul 2016, Le Gosier, Guadeloupe, France. CFCS San Juan, Puerto Rico, 00940. hal-03170026

HAL Id: hal-03170026 https://hal.inrae.fr/hal-03170026

Submitted on 15 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Effet of exogenous matter on soil biodiversity and ecosystem services in tomato plantations

Coraline Moulin, Victor Vaillant, Philippe Julianus, Chantal Fléreau, Pascale Bade, Fred Burner, Gladys Loranger-Merciris Université des Antilles / INRA, UR ASTRO Agrosystèmes TROpicaux, F-97170, Petit-Bourg, Guadeloupe.

Email: coraline.moulin@antilles.inra.fr; victor.vaillant@antilles.inra.fr; philippe.julianus@antilles,inra,fr; chantal.flereau@antilles.inra.fr; fred.burner@antilles.inra.fr; gmerciris@antilles.inra.fr







INTRODUCTION

After the Green revolution, the use of chemical fertilizers and pesticides allowed the increase of agriculture productivity in order to respond to global food demand. However, these practices have negative side effects on the environment and human health. Consequently, the use of alternative practices more respectful of the environment is preconized nowadays. Among these alternatives practices, compost and vermicompost are known to preserve soil biodiversity which had a tremendous influence on ecosystem services.

The objective of this study was to investigate the impact of chemical fertilizers, compost and vermicompost on key ecosystem services (primary production, nutrient recycling and decomposition of organic matter) provided by soil biodiversity in tomatoes (Solanum lycopersicum) plantations.

MATERIALS AND METHODS

Experimental design

Soil: Forest Ferralsol (Domaine Duclos, Petit-Bourg).



Tomato plants: Solanum lycopersicum cv. Heatmaster.

Four different substrates were used:

T: soil forest + Ca; **SPE**: soil forest + chemical fertilizers+ Ca; **SPC**: soil forest + compost+ Ca; **SPV**: soil forest + vermicompost+ Ca.





This study was carried out in a greenhouse during 4 months.

Organic matter decomposition






Organic matter decomposition was measured using the litterbags method. Five bags were installed at the soil surface of each pot and were randomly collected 7 days, 15 days, 30 days, 2 months and 4 months after the experiment setting up.

Primary production

Plant aboveground biomass was determined 30 days after the experiment setting up.





Soil chemical properties

Total P, K were determined after soil mineralisation.

Plant chemical properties

The P and K content were determined from dry plant material (leaves and fruits).

RESULTS

Primary production Organic matter decomposition and Nutrient recycling 20 15 Leaf biomass (g of dry ■Total P in leaves (mg) 10 Compost Vermicompost

Figure 1: Leaf biomass and Total P in leaves in chemical fertilizers, compost and vermicompost at 30 days.


Chemica

fertilizers

Control

Figure 2: Organic matter decomposition in tomato plantations under different treatments, at 4 months.

Compost






Figure 3: Total P (% of dry material) in fruits (on the left) and Total K in fruits(on the right) at 4 months for chemical fertilizers and organic fertilizers.

CONCLUSION

In general, the availability of nutrients was not identical for all treatments based on the decomposition of organic matter. We observed that plants, which were treated with vermicompost and compost, had an easily access to nutrients especially for the P. These plants had a tendency to accumulate more nutrients in fruits compared to plants which were treated with chemical fertilizers.

Vermicompost



Tel: 0590 25 59 00 Fax: 0590 25 59 98

