



HAL
open science

Prospective association between dietary pesticide exposure profiles and postmenopausal breast-cancer risk in the NutriNet-Santé cohort

Pauline Rebouillat, Rodolphe Vidal, Jean Pierre Cravedi, Bruno Taupier-Letage, Laurent Debrauwer, Laurence Gamet-Payrastre, Mathilde Touvier, Mélanie Deschasaux-Tanguy, Paule Latino-Martel, Serge Hercberg, et al.

► To cite this version:

Pauline Rebouillat, Rodolphe Vidal, Jean Pierre Cravedi, Bruno Taupier-Letage, Laurent Debrauwer, et al.. Prospective association between dietary pesticide exposure profiles and postmenopausal breast-cancer risk in the NutriNet-Santé cohort. *International Journal of Epidemiology*, 2021, 50 (4), pp.1184-1198. 10.1093/ije/dyab015 . hal-03170770

HAL Id: hal-03170770

<https://hal.inrae.fr/hal-03170770v1>

Submitted on 22 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Prospective association between dietary pesticide exposure profiles and postmenopausal breast cancer risk in the NutriNet-Santé cohort

Pauline Rebouillat¹, Rodolphe Vidal², Jean-Pierre Cravedi³, Bruno Taupier-Letage², Laurent Debrauwer³, Laurence Gamet-Payraastre³, Mathilde Touvier¹, Mélanie Deschasaux-Tanguy¹, Paule Latino-Martel¹, Serge Hercberg^{1,5}, Denis Lairon⁴, Julia Baudry¹, Emmanuelle Kesse-Guyot¹.

¹ Sorbonne Paris Nord University, Inserm, INRAE, Cnam, Nutritional Epidemiology Research Team (EREN), Epidemiology and Statistics Research Center – University of Paris (CRESS), 93017 Bobigny, France

² Institut de l’Agriculture et de l’Alimentation Biologiques (ITAB), 75595 Paris, France

³ Toxalim (Research Centre in Food Toxicology), Université de Toulouse, INRAE, ENVT, INP-Purpan, UPS, 31027 Toulouse, France

⁴ Aix Marseille Université, INSERM, INRAE, C2VN, 13005 Marseille, France

⁵ Département de Santé Publique, Hôpital Avicenne, 93017 Bobigny, France

Names for PubMed indexing: Rebouillat, Vidal, Cravedi, Taupier-Letage, Debrauwer, Gamet-Payraastre, Touvier, Deschasaux, Latino-Martel, Hercberg, Lairon, Baudry, Kesse-Guyot

Disclaimers: The authors declare no conflict of interest.

***Corresponding author:**

Pauline REBOUILLAT, Equipe de Recherche en Epidémiologie Nutritionnelle (EREN)

SMBH Université Sorbonne Paris Nord, 74 rue Marcel Cachin, 93017 Bobigny, France.

p.rebouillat@eren.smbh.univ-paris13.fr

+33 1 48 38 89 79

ORCID Identifiers:

Pauline Rebouillat : 0000-0002-7270-6032

Mathilde Touvier : 0000-0002-8322-8857

Emmanuelle Kesse-Guyot : 0000-0002-9715-3534

Denis Lairon : 0000-0001-9941-3742

Short running head: Dietary pesticide exposure and post-menopausal breast cancer risk

Abbreviations:

ADI: Acceptable Daily Intake; AhR: Aromatic hydrocarbon Receptor; ANSES: Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail ; BMR:

Basal Metabolic Rate ; BC: Breast Cancer; β -HCH : β -Hexachlorocyclohexane ; BMI: Body Mass Index; CépiDC : French Centre for Epidemiology Medical Causes of Death database ;

CI: Confidence Interval ; CNIL: Commission Nationale de l'Informatique et des Libertés ;

CVUA: Chemisches und Veterinäruntersuchungsamt; DDE:

Dichlorodiphenyldichloroethylene ; DDT: Dichlorodiphenyltrichloroethane ; DNA:

Deoxyribonucleic acid ; EDI: Estimated Daily Intake; EFSA: European Food and Safety

Authority; ER-/PR-: Estrogen Receptor Negative/Progesterone Receptor Negative ; FFQ:

Food Frequency Questionnaire; HCB: Hexachlorobenzene ; HR: Hazard Ratio; ICD-10:

International Statistical Classification of Diseases and Related Health Problems 10th

Revision; IPAQ: International Physical Activity Questionnaire; IRB INSERM: Institutional

Review Board of the French Institute for Health and Medical Research; NMF: Non-negative

Matrix Factorization; ; OC: Organochlorine; OP: Organophosphorous;; PCTA:

Pentachlorothioanisole ; PNNS: Programme National Nutrition Santé; SD: Standard

Deviation; SNIIRAM: Système National d'Information Inter-Régimes de l'Assurance

Maladie; sPNNS-GS2: simplified Programme National Nutrition Santé Guideline Score 2;

WHO: World Health Organization.

Clinical Trial Registry: NCT03335644

URL for Trial Registration : <https://clinicaltrials.gov/ct2/show/NCT03335644>

Manuscript Word Count: 4318 words

Data Share Statement:

Data of the study are protected under the protection of health data regulation set by the French National Commission for Information Technology and Liberties (Commission Nationale de l'Informatique et des Libertés, CNIL). The data are available upon reasonable request to the study's operational manager, Nathalie Pecollo (n.pecollo@eren.smbh.univ-paris13.fr), for review by the steering committee of the NutriNet-Santé study.

1 **Abstract:**

2 **Background:**

3 Some pesticides, used in large quantities in current agricultural practices all over Europe, are
4 suspected of adverse effects on human reproductive health (breast and prostate cancers),
5 through mechanisms of endocrine disruption and possible carcinogenic properties, as
6 observed in agricultural settings.

7 However, evidence on dietary pesticide exposure and breast cancer (BC) is lacking for
8 general population. We aimed to assess the associations between dietary exposure to
9 pesticides and BC risk among postmenopausal women of the NutriNet-Santé cohort.

10 **Methods:**

11 In 2014, participants completed a self-administered semi-quantitative Food Frequency
12 Questionnaire, distinguishing conventional and organic foods. Exposures to 25 active
13 substances used in EU plant protection products were estimated using a pesticide residue
14 database accounting for farming practices, from Chemisches und Veterinäruntersuchungsamt
15 Stuttgart, Germany.

16 Non-Negative Matrix Factorization (NMF), adapted for data with excess zeros, was used to
17 establish exposure profiles. The four extracted NMF components' quintiles were introduced
18 into Cox models estimating Hazard Ratio (HR) and 95% confidence interval (95% CI),
19 adjusted for known confounding factors.

20 **Results :**

21 A total of 13,149 postmenopausal women were included in the analysis(169 BC cases,
22 median follow-up=4.83 years). Negative associations between component 3, reflecting low
23 exposure to synthetic pesticides, and post-menopausal BC risk were found (HR_{Q5}=0.57;
24 95%CI(0.34;0.93), p-trend=0.006). Positive association between component 1 score (highly
25 correlated to chlorpyrifos, imazalil, malathion, thiabendazole) and postmenopausal BC risk

26 was found specifically among overweight and obese women ($HR_{Q5}=4.13$; 95%CI(1.50;11.44),
27 p-trend=0.006). No associations were detected for the other components.

28 **Conclusions:**

29 These associations suggest a potential role of dietary pesticide exposure on BC risk. Further
30 research is needed to investigate mechanisms and confirm these results in other populations.

31 **Keywords:** dietary exposure; pesticides; organic farming; epidemiology; breast cancer;
32 environmental health.

33

Key Messages:

- Diet is considered as the main exposure route for pesticide exposure in the general population. Dietary pesticide exposure has been rarely studied in relation with cancers.
- Non-Negative Matrix Factorization (NMF), a method adapted for data with excess zeros, was used to characterise dietary pesticide exposure profiles.
- We observed a reduction in the risk of postmenopausal breast cancer for NMF Component 3 (reflecting low exposure to several synthetic pesticides).
- A positive association between NMF component 1 score (highly correlated to chlorpyrifos, imazalil, malathion, thiabendazole) and postmenopausal breast cancer risk was observed specifically among overweight and obese women.
- For NMF Components 2 and 4, Hazard Ratios (HR) were HR_{Quintile 5 vs Quintile 1} 0.96, 95% Confidence Interval (0.59; 1.56), p for trend : 0.30 and HR_{Quintile 5 vs Quintile 1} 0.65, 95% Confidence Interval (0.38; 1.12), p for trend : 0.13.

35 **Introduction:**

36 Large quantities of plant protection products are used in current European agricultural
37 practices ¹. In particular, France has high usage of pesticides, synthetic or natural, both in
38 global tonnages (80 000 tons in 2018) and by surface area (4.45 kg/ha in 2018) ²⁻⁵.
39 Deleterious impacts of pesticides on human health have been evidenced. Various effects of
40 pesticide active substances have been documented , including genetic material alteration,
41 endocrine disrupting effects, cell apoptosis and cell signaling dysregulation, and oxidative
42 stress induction ^{4,6-8}. These mechanisms have been shown to be involved in carcinogenesis ⁹.
43 Recently, IARC classified many pesticides as “probably carcinogenic to humans” (Group 2A)
44 and “possibly carcinogenic to humans” (Group 2B) ¹⁰. In addition, many pesticides exhibit
45 endocrine disruptors properties ⁷.
46 Indeed, cancer is nowadays the first or second leading cause of premature death in many
47 European countries. It is the first cause of mortality in France ¹¹⁻¹³, breast cancer being the
48 most common and leading cause of cancer death for women in France. Associations between
49 occupational pesticide exposure in agricultural settings (involving respiratory and cutaneous
50 exposure routes) and the occurrence of some locations of cancers (myeloma, non-Hodgkin
51 lymphoma, prostate) were found in several studies¹⁴⁻¹⁹. Associations in agricultural settings
52 with other cancer locations have been reported (stomach, esophagus, liver, colorectal...), and
53 especially reproductive system cancers (prostate, breast), potentially induced by endocrine
54 disruption mechanisms ^{4,7,20}. Notably, associations between breast cancer risk and
55 organophosphorus pesticide exposure were found for farmer’s wives in some studies ^{21,22}.
56 However, in the general population, although food is considered as the first exposure
57 pathway, data is lacking on associations between dietary exposure to pesticides and cancer
58 ^{23,24}. This may be explained by three main challenges. Firstly, measuring pesticide residue
59 concentrations in food is expensive and tedious. In addition, it is difficult to measure
60 pesticides mixtures (opposed to compounds taken separately), but necessary, as it can

61 potentially lead to synergistic effects. Finally, data existing so far generally lacks of precision
62 regarding the production system (conventional vs organic), limiting proper estimation of
63 pesticide exposure.

64 Recently, a study conducted in the NutriNet-Santé cohort showed protective associations
65 between the high proportion of organic food in the diet and different types of cancers,
66 including postmenopausal breast cancer²². An hypothesis advanced to explain this association
67 was the potentially lower concentrations of pesticides residues in plant organic foods ²⁵.

68 In that context, the purpose of this work was to study the associations between dietary
69 pesticide exposure profiles and breast cancer risk among postmenopausal women included in
70 the NutriNet-Santé cohort.

71

72 **Material and Methods:**

73 *Study population*

74 The NutriNet-Santé study is a web-based prospective cohort of adults launched in France in
75 May 2009²⁶. Inclusion criteria was to be aged 18 years old and over and to speak French. A
76 set of self-administered validated questionnaires^{27–29} was completed online by participants at
77 baseline and repeated every year. Complementary questionnaires were regularly proposed
78 concerning dietary behaviors and specific health issue during follow-up.

79 *Dietary intake assessment*

80 A 264-item web-based self-administered semi-quantitative food frequency questionnaire
81 (Org-FFQ) distinguishing organic and conventional foods was sent to the participants
82 between June and December 2014. The Org-FFQ has been extensively described elsewhere³⁰.
83 Briefly, it was elaborated on the basis of an existing validated FFQ³¹ to which a 5-point
84 ordinal scale was added to measure the frequency of organic food consumption. For each
85 item, participants provided their frequency of consumption and the quantity consumed
86 helping with photographs showing different portion sizes³². For food and beverages with an
87 existing organic version (labelled), participants answered the question “How often was the
88 product of organic origin?” by selecting 1 of the 5 following frequency modalities: never,
89 rarely, half-of-time, often, or always. The organic food consumption was then obtained by
90 attributing the respective percentages, 0, 25, 50, 75, and 100, to the modalities. Weighting and
91 sensitivity analyses for the Org-FFQ have been published elsewhere³⁰.

92 All food and beverage items were aggregated into 33 food groups. Nutritional values were
93 obtained from a published food composition database³³. A global proportion (as weight) of
94 organic food in the diet was calculated as well as the proportion of organic food for each food
95 group.

96

97 *Pesticide exposure assessment*

98 Dietary pesticide exposure was estimated by combining dietary intakes of each adult with
 99 pesticide residue concentration values in foods using contamination data from Chemisches
 100 und Veterinäruntersuchungsamt (CVUA) Stuttgart, a European Union reference laboratory for
 101 pesticides³⁴. The database comprised contamination data for conventional and organic food
 102 products. Twenty-five commonly used pesticides were selected among components available
 103 in this database, given both their frequency of detection above the Maximum Residue Levels
 104 (MRL) when sufficient data were available, and their frequency above Acceptable Daily
 105 Intake (ADI) otherwise, as detailed in Baudry et al. 2019 study³⁵. Pesticides commonly used
 106 in organic agricultural systems (e.g. natural pyrethrins, spinosad) were also selected. These
 107 criteria made it possible to take into account a broad spectrum of classes of pesticides. The
 108 264 Org-FFQ items were decomposed into 442 ingredients (comprising at least 5% of at least
 109 one food item). Animal-based ingredients were excluded, as CVUA encompassed plant-based
 110 ingredients only. Indeed, plant-based foods have markedly more frequent and higher
 111 pesticides residues levels than foods of animal origin³⁶. The resulting 180 plant ingredients
 112 were matched to CVUA database and then were attributed a contamination value in organic
 113 and conventional farming modes (as the mean of corresponding data point). A flowchart of
 114 the different steps for the decomposition and matching is shown in **Supplementary Material**
 115 **1**.

116 For each ingredient/pesticide pair in conventional and organic farming, a frequency of
 117 detection and a frequency of quantification were determined using the formula as follows:

$$118 \quad \textit{Frequency of detection} = 100 \times \frac{\textit{Number of analyses} - \textit{Number of undetected}}{\textit{Number of analyses}}$$

$$119 \quad \textit{Frequency of quantification}$$

$$120 \quad = 100 \times \frac{\textit{Number of analyses} - \textit{Number of unquantified}}{\textit{Number of analyses}}$$

121 Treatment of data below detection limit has been extensively described elsewhere³⁵.

122 As food consumption data from NutriNet-Santé referred to edible foods (bone-free, peeled or
123 cooked products), edibility and cooking factors were allocated to each ingredient when
124 necessary. The same conversion factors were used for both conventional and organic
125 products. Cooking or peeling effects on pesticide residue levels were not accounted for as
126 dilution factors are not available for all food/pesticide couples ³⁷. For each pesticide, the
127 estimated daily intake (EDI) (in µg/kg of weight/day) under both lower- and upper-bound
128 scenarios was calculated using methods recommended by EFSA and WHO ^{38,39}. Lower-bound
129 (optimistic) scenario was used for this work, as more in line with available literature
130 comparing both production systems ^{35,40}.

131 *Covariates*

132 Baseline and yearly questionnaires collected sociodemographic and lifestyle characteristics
133 such as sex, date of birth, occupation, educational level, smoking status, number of children.
134 Monthly income by household unit was obtained was obtained using both the household
135 income and composition. Anthropometric measures (height, weight), physical activity (using
136 the validated Physical Activity Questionnaire ⁴¹) and health status (menopausal status, family
137 history of cancer, treatments) were also collected.

138 A specific questionnaire on environmental exposure collected the type of environment in
139 which participants lived: agricultural or urban area.

140 The simplified Programme National Nutrition Santé Guideline Score 2 (sPNNS-GS2), based
141 on the level of adherence to 2017 French dietary guidelines proposed by the High Council of
142 Public Health ^{42,43}, the provegetarian score ⁴⁴ and the percentage of ultra-processed foods in
143 the diet ⁴⁵ were computed to be used as adjustment factors. Briefly, the sPNNS-GS2 includes
144 13 components. One point was allocated for following the guideline (and 0 otherwise), and
145 conversely for moderation components. Component with several subcomponents were
146 standardized and a penalty for overconsumption was applied. The score ranged from $-\infty$ to

147 14.25. Component, cut-off, scoring system and ponderation are presented in **Supplementary**
148 **Material 2**.

149 The provegetarian score was computed as follows ⁴⁴: 7 vegetable food groups and 5 animal
150 food groups were defined and sex-specific quintiles adjusted for total energy intake were
151 calculated. For each plant component, 1 to 5 points were allocated to quintile 1 to 5 and for
152 animal food groups the scoring was reversed. The provegetarian score was obtained by
153 summing each quintile value of vegetable food group and each reverse quintile value of
154 animal food group thus ranging from 12 (low plant food consumption) to 60 (high plant food
155 consumption).

156 Percentage of ultra-processed foods consumed was computed after classification of foods
157 using NOVA categories ⁴⁵, by a committee of dietitians and researchers ⁴⁶. NOVA
158 classification is described in details in **Supplementary Material 3**. Data used to calculate the
159 proportion of ultra-processed foods in the diet were the closest to the Org-FFQ completion
160 date.

161 *Cancer cases*

162 Health events were declared by participants through a yearly health status questionnaire and a
163 dedicated web-interface at any time of the study. All medical records were collected and
164 analyzed by dedicated physicians. Physicians of participants declaring major health events
165 were contacted to collect additional information if necessary. Validation of these major health
166 events was carried out by a medical expert committee.

167 Overall, medical records were obtained for more than 90% of self-reported cancer cases.

168 Moreover, we performed a linkage between our declared health data to medico-administrative
169 registers of the national health insurance system (Système National d'Information Inter-
170 Régimes de l'Assurance Maladie [SNIIRAM] databases). Mortality data were also used from
171 the French Centre for Epidemiology Medical Causes of Death database (CépiDC). Cancer

172 cases were classified using the *International Statistical Classification of Diseases and Related*
173 *Health Problems, 10th Revision, Clinical Modification*⁴⁷. In this study, we considered all first
174 primary breast cancers (ICD-10 C50) diagnosed between baseline (i.e. the date of completion
175 of the Org-FFQ in 2014 or the menopause date, whichever occurred last) and 18 July 2019 to
176 be cases.

177 *Statistical analyses*

178 A flowchart for the study sample selection is presented in Figure 1.

179 For the present study, postmenopausal female participants who completed the Org-FFQ
180 between June and December 2014 (N = 28,445), with no missing covariates for basal
181 metabolic rate computation (N = 28,137), who were not detected as under- or over-reporters
182 (N = 27,158), who were postmenopausal and free of breast cancer when they completed the
183 Org-FFQ, were selected (N=13,149).

184 Regarding under or overreporters, only participants with a plausible energy intake were
185 included in the analyses. The detection method for under and overreporters was based on the
186 comparison between energy intake and energy requirement and is extensively described in a
187 previous article by Baudry et al.³⁰

188 Dietary pesticide exposure profiles were analyzed using Non-Negative Matrix factorization
189 (NMF) (detailed in **Supplementary Material 4**), specially adapted for non-negative data with
190 excess zeros, developed by Lee et al⁴⁸. In total, four components were computed for the NMF
191 procedure using 25 selected pesticide exposure values, reflecting various pesticide exposure
192 patterns.

193 Sociodemographic and lifestyle characteristics were compared between cases and non-cases,
194 and also across NMF-extracted component quintiles using Chi², Mantel-Haenzel, Wilcoxon
195 and Kruskal-Wallis tests, as appropriate.

196 Associations between dietary pesticide exposure, using NMF components divided into
197 quintiles (first quintile used as reference) and breast cancer were assessed using Cox
198 proportional hazards regression models. Participants contributed person time until the date of
199 diagnosis of cancer, the date of last completed questionnaire, the date of death, or 18th July
200 2019, whichever occurred first.

201 NMF component scores were divided into quintiles and introduced into separate models, with
202 age as time scale, and first quintile as reference.

203 Cox models were adjusted for known confounders such as smoking status, alcohol intake,
204 educational level, physical activity (measured with International Physical Activity
205 Questionnaire), Body Mass Index (BMI), height, family history of cancer, menopausal
206 treatment and parity and overall quality of the diet (measured by the PNNS-GS2 score⁴³).

207 Interactions between potential modulating factors and components were tested by introducing
208 the multiplicative interaction term into the models, namely body mass index, sPNNS-GS2
209 (overall nutritional quality of the diet) and the level of plant-based consumption (using the
210 provegetarian score). Interactions with $p < 0.10$ were further investigated.

211 Schoenfeld residuals were used to test the proportional hazard assumption of the Cox model.
212 Potential nonlinear effects of continuous exposure variables were evaluated using martingale
213 residuals.

214 Tests for linear trend were performed using quintiles of the NMF components as ordinal
215 variables.

216 Sensitivity analyses were carried out. A model was performed excluding early cases (1 year
217 after baseline) and two other models were computed with additional adjustments for the level
218 of ultra-processed foods in the diet, and the provegetarian score. Two-sided tests were used..

219 Data management and statistical analyses were performed using SAS (version 9.4; SAS
220 Institute, Inc.). NMF was performed using R's NMF package⁴⁹.

221 **Results:**

222 *Characteristics of the participants*

223 Sociodemographic characteristics of the studied participants are presented in **Table 1**. A total
224 of 13,149 postmenopausal women were included in the analyses; 169 postmenopausal breast
225 cancer were diagnosed during the follow-up (mean \pm SD: 4.35 \pm 1.06 years; median: 4.83
226 years). Mean age at baseline was 60.5 years (SD=7.39). The majority of individuals had a
227 graduate educational attainment, was retired and lived in more than 200,000 inhabitant urban
228 units and were never smokers for 49% of them. One third of the sample was overweight
229 (BMI>25 kg/m²). Most frequent physical activity levels were 'high' and 'moderate'. Overall,
230 no significant differences on sociodemographic characteristics were found between cases and
231 non-cases. The nutritional characteristics of the cases and non-cases are presented
232 in **Supplementary Table S1**. Overall, no differences were observed between cases and non-
233 cases except for organic food proportion in the diet.

234 The absolute estimated dietary pesticide exposure for cases and non-cases is presented
235 in **Table 2**. Among others, the pesticide exhibiting the highest means for exposures in cases
236 and non-cases were boscalid, iprodione, spinosad, thiabendazole, and imazalil.

237 The correlations between the 4 NMF-extracted components and pesticide exposure are shown
238 in **Table 3**. Pesticides such as chlorpyrifos, imazalil, malathion, profenofos, thiabendazole
239 were highly correlated with NMF Component 1. For NMF Component 2, highly correlated
240 pesticides were azoxystrobin, boscalid, cyprodinil, difenoconazole, fenhexamid, iprodione,
241 tebuconazole, lambda cyhalothrin.

242 NMF Component 3 was characterized by low correlations with synthetic pesticides and high
243 correlation with organic pesticide spinosad. For NMF Component 4, high correlations with
244 acetamiprid, carbendazim, chlorpyrifos, cypermethrin, dimethoate/omethoate were
245 observed.

246 Each NMF Component exhibited specific correlates. For information, profiles and dietary

247 patterns are presented in **Supplementary Tables S2 to S9: Supplementary Tables S2-S3** for
248 NMF Component 1, **S4-S5** for NMF Component 2, **S6-S7** for NMF Component 3, **S8-S9** for
249 NMF Component 4. Main findings are the negative and positive linear associations between
250 proportion of organic food and NMF components 1-2 and 3 respectively.
251 The absolute estimated dietary pesticide exposures compared across components quintiles are
252 presented in **Supplementary Table S10** and **Supplementary Table S11**.
253 Correlations between dietary intakes for 33 food groups and NMF components are shown in
254 **Supplementary Table S12**.

255 *Associations between pesticide dietary exposure and breast cancer risk*

256 **Table 4** presents Hazard Ratios (HR) for the associations between NMF components and the
257 risk of postmenopausal breast cancer, with several levels of adjustments. Positive and
258 significant association was found for the fifth quintile of NMF component 1, HR=1.73,
259 95%CI(1.05;2.84). With regard to NMF component 3, participants in the fifth quintile had
260 significantly lower risks (HR=0.57, 95%CI(0.34;0.93), p trend=0.006) of postmenopausal
261 breast cancer than the first quintile (p<0.05). HR for the fifth quintiles of NMF Components 2
262 and 4 were HR 0.96, 95% CI(0.59;1.56), p-trend : 0.30 and HR 0.65, 95% CI(0.38;1.12), p-
263 trend : 0.13.

264 Further adjustments for the quality of the diet (with the sPNNS-GS2 score, Model 2), and
265 residing in an agricultural area (Model 3) did not modify the findings (**Table 4**).

266 Several interactions between NMF components and other variables were tested in the models
267 (provegetarian score, sPNNS-GS2, overweight vs non-overweight). A significant interaction
268 was found between BMI and NMF component 1 (p for interaction with BMI in 2 categories =
269 0.004) on the risk of postmenopausal breast cancer. Therefore, stratified analyses were
270 performed, with a threshold of 25 kg/m², and results are shown in **Table 5**. Associations
271 between NMF Component 1 and post-menopausal breast cancer risk were significant among

272 individuals with a BMI>25 kg/m² only, with higher risk for the fifth quintile and fourth
273 quintile compared to the first quintile, HR_{Quintile 5 vs Quintile 1}: 4.13 (95%CI(1.50;11.44) and
274 HR_{Quintile 4 vs Quintile 1}: 3.02 (95%CI(1.08;8.47)), p trend=0.006 (**Table 5**).

275 *Sensitivity analyses*

276 After exclusion of cancer cases occurring less than 1 year after baseline, HR were similar but
277 the loss of statistical power did not allow to reach significance (**Table 6**).

278 Further adjustments for the percentage of ultra-processed foods, or provegetarian score, did
279 not modify the results substantially.

280 **Discussion:**

281 In this large population of French postmenopausal women, we found significant negative
282 associations between NMF component 3 (reflecting low exposure to several synthetic
283 pesticides) and post-menopausal breast cancer risk. When analyses were stratified on BMI
284 (threshold 25 kg/m²), positive association between NMF Component 1 (reflecting exposure to
285 chlorpyrifos, imazalil, malathion, thiabendazole) and postmenopausal breast cancer risk was
286 found among overweight women. No significant associations were detected for the other
287 components.

288 To our knowledge, the present study is the first to investigate various pesticide exposure
289 patterns, accounting for farming practices in relation with breast cancer risk in the general
290 population. Thus, our findings cannot be directly compared to previous scientific literature.

291 However, some studies have been conducted to investigate associations between
292 occupational, residential or domestic pesticide exposure and breast cancer risks. Studies
293 largely focused on organochlorine (OC) pesticides and related metabolites (for instance
294 Dichlorodiphenyltrichloroethane (DDT), Dichlorodiphenyldichloroethylene (DDE), β -
295 Hexachlorocyclohexane (β -HCH), Hexachlorobenzene (HCB), Pentachloroanisole
296 (PCTA), now banned in European Union, reporting higher breast cancer risk for users
297 (personal or occupational) ⁵⁰⁻⁵².

298 Breast cancer risks were found higher with exposure to OC (use vs never use) in a study
299 published by Engel et al (2005) conducted in farmer's wives population ²¹. In this study,
300 breast cancer risks also appeared elevated regarding organophosphorus pesticide family (OP)
301 as a whole. When analysis was performed on compounds taken separately, association was
302 significant only for malathion. Stratification on menopausal status was performed and showed
303 higher risks among postmenopausal women whose husbands used OC and also OP such as
304 chlorpyrifos, diazinon and malathion. In our study, NMF component 1, positively correlated
305 with malathion and chlorpyrifos (respective correlation coefficients 0.76 and 0.73), was

306 significantly associated with breast cancer risk for participants with a BMI > 25 kg/m².
307 Moreover, in analyses conducted in 2015 by Lerro *et al.* in the Agricultural Health Study ²²,
308 spouses whose husband used OPs had higher breast cancer risk compared to spouses whose
309 husbands never used OPs (RR= 1.20, 95%CI(1.01; 1.43)). However, in that study, when
310 considering pesticides molecules separately (malathion, chlorpyrifos, terbufos), associations
311 with breast cancer risk were no longer significant, except for chlorpyrifos, and especially for
312 Estrogen Receptor Negative/Progesterone Receptor Negative (ER-/PR-) breast cancer risk.
313 These observations could be interpreted in light of some kind of synergistic effects evidenced
314 in toxicological studies when exposed to pesticide residue mixtures ^{6,53,54}. In the same study
315 by Lerro *et al.*, after stratification on menopausal status, significant association between
316 higher breast cancer risks and use of any OPs was observed among the postmenopausal
317 women. Again, no significant associations between OP pesticides taken separately and breast
318 cancer risk were found. Another recent study also found elevated risks in women exposed to
319 chlorpyrifos compared with those not exposed (OR = 3.22; 95%CI(1.38,7.53) ⁵⁵. These results
320 are consistent with our results suggesting an association between NMF component 1
321 (reflecting exposure to chlorpyrifos, imazalil, malathion, thiabendazole) and
322 postmenopausal breast cancer risk.

323 It is important to note that after exclusion of ‘early cases’ (<1 year after baseline), an
324 important drop was observed in the HR_{Quintile 5 vs Quintile 1} for Component 1 from 1.73 to 1.37. It
325 is possible that excluded cases exhibited very specific nutritional and health characteristics
326 linked to their imminent diagnosis and probable health deterioration linked to it. We should
327 also note that Quintile 5 lost more cases than other quintiles (42 cases to 31) and this could
328 somehow influence the analysis.

329 In our study, possible hypotheses to explain the negative associations between NMF
330 Component 3 and postmenopausal breast cancer risk rely on the fact that besides being highly

331 correlated with some pesticides used in organic farming (i.e. natural pyrethrins, spinosad), this
332 component is also negatively correlated with several synthetic pesticides (azoxystrobin,
333 chlorpropham, methamidophos). Participants with high component 3 score, seemed generally
334 less exposed to the synthetic studied pesticides but also less exposed to pesticides with high
335 suspected toxicity such as chlorpyrifos, imazalil, malathion. These results are consistent
336 with those of another study, conducted in the NutriNet-Santé cohort in 2018, that reported a
337 negative association between high organic food score and postmenopausal breast cancer risk
338 ($HR_{\text{Quintile4}} = 0.66$; 95% CI(0.45-0.96))⁵⁶. One formulated hypothesis for this association was
339 that organic farming regulations lead to a lower frequency or an absence of pesticide residues
340 in organic foods compared with conventional foods²⁵. Thus, our present results are consistent
341 with this hypothesis. Moreover, effects had comparable magnitude.

342 Mechanisms underlying these associations could be related to carcinogenic properties of some
343 organophosphate pesticides provoking DNA damage, cell apoptosis dysregulation, epigenetic
344 changes⁵⁷, cell signaling disruption⁵⁸, nuclear receptor binding (Aromatic hydrocarbon
345 Receptor, AhR)⁵⁹ or oxidative stress induction^{4,8,9}. It can be noted that IARC classified some
346 organophosphate pesticides as “probably carcinogenic to humans” (Group 2A) and “possibly
347 carcinogenic to humans” (Group 2B)¹⁰. Endocrine disruption potential of pesticides has also
348 been described in toxicological studies and recently in a review by Yang et al.^{7,60}, and could
349 be particularly involved in hormone-dependent breast carcinogenesis, as some pesticides are
350 known to mimic estrogen functions^{57,61}. Indeed, azole fungicides, including imazalil, for
351 which we found high correlations with NMF Component 1, have been related to inhibition of
352 estrogen biosynthesis in some studies⁶². These pesticides are also known to affect
353 mitochondrial activity and oxidoreduction status⁶³.

354 When considering stratified analyses on BMI (threshold 25 kg/m²), a positive association
355 between component 1 and post-menopausal breast cancer risk was observed in overweight

356 individuals.

357 Several studies have found positive associations between body fat and OC pesticide blood
358 levels, overweight subjects having higher blood levels of these pesticides ^{64,65}. However, it is
359 unlikely this would be the case for OPs, which are not accumulated in adipose tissue. It is
360 possible that there is cumulative effect between obesity and pesticide exposure on cancer
361 risks. The specific association in overweight women could also be explained by differences in
362 paraoxonase 1 (PON1) activity, as this enzyme is involved in lipid metabolism, but also
363 participates in hydrolysis of organophosphate compounds ⁶⁶. Indeed, some studies have
364 shown lower levels of PON1 activity in overweight and obese patients ^{67,68}. In consequence,
365 toxicity of these pesticides could be higher for this subgroup.

366 More data are needed on OP pesticides, in order to fully understand underlying mechanisms
367 of this association and potential modifying effects of BMI on breast cancer.

368 It should be noted that associations between NMF Components 2 and 4 and postmenopausal
369 breast cancer risk were not statistically significant in our study. Given the lack of evidence on
370 specific pesticide mixtures in relation to human health, it is difficult to know whether this
371 could be due to specific non-carcinogenic patterns of the studied pesticides or to the fact that
372 the population may not be exposed enough to experience deleterious health effects.

373 *Limitations and strengths*

374 Some limitations of this study should be mentioned. Firstly, the NutriNet-Santé cohort is
375 composed of volunteers, mostly highly educated, who can be more interested in their health
376 and dietary intakes than the general French population ⁶⁹. This implies cautions when
377 generalizing our results to other populations.

378 It is important to mention that dietary intakes were self-reported through a food frequency
379 questionnaire (FFQ) and this may have caused overestimation of organic food consumption.

380 Other limitations come from the database used to estimate dietary pesticide exposures, since

381 data were not available for animal products and the database did not contain measures for
382 copper or sulfur-based products, widely used in organic farming, but not known as
383 carcinogenic compounds. Measures were performed in Germany, but products from all over
384 European Union were tested.

385 Another limitation that should be mentioned was that the dietary pesticide exposure was
386 estimated and is therefore not as accurate as measuring biomarkers. It is to note that
387 biomarkers can rarely be measured on very large samples given feasibility and cost
388 constraints. Another disadvantage of using biomarkers is that it would not give precision on
389 the active substances to which individuals are exposed since biomarkers are not specific to
390 one molecule.

391 We should acknowledge that pesticide exposure have probably been overestimated as
392 potential concentration or dilution effects during washing, cooking or peeling on pesticide
393 residue levels were not accounted for³⁷. Finally, for this study, follow-up duration was short
394 and the number of cases limited, given a high estimated latency period for this type of disease.
395 This can be a limitation for causal inference and statistical power of the analyses. However,
396 we hypothesize that dietary habits change marginally over time, therefore dietary habits four
397 years before diagnosis were probably very similar to those 10 years before. In the same way,
398 patterns of pesticide use in France were similar during this period, as there were changes in
399 authorizations only for three selected pesticides (anthraquinone, methamidophos and
400 profenofos)⁷⁰. Nevertheless, it will be interesting to reassess cancer risks after several years,
401 in order estimate long-term effects.

402 Some strengths of this study can also be put forward.

403 Cox regression models were adjusted for a wide range of covariates, including major
404 confounders such as diet quality indicators. Despite a limited number of cases, the sample size

405 still allowed us to perform some stratification and sensitivity analyses in order to deepen the
406 understanding of these results and reduce confounding bias.

407

408 **Conclusion**

409 We observed a negative association between low synthetic pesticide exposure profile (through
410 NMF component 3) and post-menopausal breast cancer risk. Positive association between
411 component 1 (highly correlated to chlorpyrifos, imazalil, malathion, thiabendazole) and
412 postmenopausal breast cancer risk was also found specifically among overweight and obese
413 women. If confirmed by other studies, some pesticides profiles may constitute risk factors
414 among subgroups such as those with overweight. Observed associations should be
415 investigated in other prospective studies, in different settings, coupled with experimental
416 studies to complement these observational studies in order to validate estimated dietary
417 pesticide exposure. A better understanding of the impact of dietary pesticides on human
418 health could unlock prevention strategies for the whole population through regulation.

419

420 **Ethics approval :**

421 The NutriNet-Santé study is conducted in accordance with the Declaration of Helsinki, and all
422 procedures were approved by the Institutional Review Board of the French Institute for
423 Health and Medical Research (IRB Inserm 0000388FWA00005831) and the Commission
424 Nationale de l'Informatique et des Libertés (CNIL 908,450 and 909,216). All participants
425 provided their informed consent with an electronic signature, and this study is registered in
426 ClinicalTrials.gov (NCT03335644).

427 **Funding:**

The NutriNet-Santé study is supported by the following public institutions: French Ministry of Health (DGS), Santé Publique France, the National Institute for Health and Medical Research (INSERM), the French National Research Institute for Agriculture, Food and Environment (INRAE), the National Conservatory of Arts and Crafts (CNAM) and the Sorbonne Paris Nord University. The BioNutriNet project was supported by the French

National Research Agency (ANR) in the context of the 2013 Programme de Recherche Systèmes Alimentaires Durables (ANR-13-ALID-0001). Pauline Rebouillat is supported by a doctoral fellowship from Sorbonne Paris Nord University.

428

429 **Data availability**

430 Data of the study are protected under the protection of health data regulation set by the French
431 National Commission for Information Technology and Liberties (Commission Nationale de
432 l'Informatique et des Libertés, CNIL). The data are available upon reasonable request to the
433 study's operational manager, Nathalie Pecollo (n.pecollo@eren.smbh.univ-paris13.fr), for
434 review by the steering committee of the NutriNet-Santé study.

435 **Acknowledgements**

436 We especially thank Cédric Agaesse, Vristi Desan and Cynthia Perlin (dietitians); Thi Hong
437 Van Duong, Younes Esseddik (IT manager), Paul Flanzky, Régis Gatibelza, Jagatjit Mohinder
438 and Aladi Timera (computer scientists); Julien Allegre, Nathalie Arnault, Laurent Bourhis and
439 Fabien Szabo de Edelenyi, PhD (supervisor) (data-manager/statisticians) for their technical
440 contribution to the NutriNet-Santé study and Nathalie Druesne-Pecollo, PhD (operational
441 manager).

442 We also thank the CVUAS for the pesticide residue database and Noémie Soton for her
443 contribution to the data management of the CVUA database. We warmly thank all of the
444 dedicated and conscientious volunteers involved in the NutriNet-Santé cohort.

445 **Author contributions**

446 RV, DL, JB, SH and EK-G conducted the research.

447 PR performed statistical analyses and drafted the manuscript.

448 All authors critically helped in the interpretation of results, revised the manuscript and
449 provided relevant intellectual input. They all read and approved the final manuscript. EK-G
450 supervised the study, had primary responsibility for the final content, she is the guarantor.

451 **Conflict of Interest:**

452 None declared.

453

454

455

References

1. Popp J, Petó K, Nagy J. Pesticide productivity and food security. A review. *Agron Sustain Dev*. 2013 Jan 1;**33**(1):243–255.
2. Eurostat - Tables, Graphs and Maps Interface (TGM) map [Internet]. [cited 2019 Jun 28]. Available from: <https://ec.europa.eu/eurostat/tgm/mapToolClosed.do?tab=map&init=1&plugin=1&language=en&pcod e=tai02&toolbox=types>
3. L'utilisation des pesticides en France : état des lieux et perspectives de réduction [Internet]. [cited 2019 Oct 9]. Available from: <https://agriculture.gouv.fr/lutilisation-des-pesticides-en-france-etat-des-lieux-et-perspectives-de-reduction>
4. Collectif INSERM. Pesticides : Effets sur la santé, une expertise collective de l'Inserm. Salle Presse Inserm. 2013. <http://presse.inserm.fr/pesticides-effets-sur-la-sante-une-expertise-collective-de-linserm/8463/>.
5. Food and Agriculture Organization of the United Nations (FAO). Agri-environmental indicator on the Use of pesticides per area of cropland at national level for the period 1990 to 2016. [Internet]. 2020 [cited 2020 Nov 17]. Available from: <http://www.fao.org/faostat/en/#data/EP>
6. Lukowicz C, Ellero-Simatos S, Régnier M, et al. Metabolic Effects of a Chronic Dietary Exposure to a Low-Dose Pesticide Cocktail in Mice: Sexual Dimorphism and Role of the Constitutive Androstane Receptor. *Environ Health Perspect*. 2018 Jun 15;**126**(6):067007.
7. Mnif W, Hassine AIH, Bouaziz A, Bartegi A, Thomas O, Roig B. Effect of Endocrine Disruptor Pesticides: A Review. *Int J Environ Res Public Health*. 2011 Jun;**8**(6):2265–2303.
8. Mostafalou S, Abdollahi M. Pesticides and human chronic diseases: evidences, mechanisms, and perspectives. *Toxicol Appl Pharmacol*. 2013 Apr 15;**268**(2):157–177.
9. Vakonaki E, Androutsopoulos VP, Liesivuori J, Tsatsakis AM, Spandidos DA. Pesticides and oncogenic modulation. *Toxicology*. 2013 May 10;**307**:42–45.
10. Guyton KZ, Loomis D, Grosse Y, et al. Carcinogenicity of tetrachlorvinphos, parathion, malathion, diazinon, and glyphosate. *Lancet Oncol*. 2015 May;**16**(5):490–491.
11. Ferlay J, Soerjomataram I, Dikshit R, et al. Cancer incidence and mortality worldwide: Sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer*. 2015;**136**(5):E359–E386.
12. Wild CP, Weiderpass E, Stewart BW, editors. World Cancer Report: Cancer Research for Cancer Prevention. : International Agency for Research on Cancer. [Internet]. Lyon, France; 2020. Available from: <http://publications.iarc.fr/586>.
13. Cancers : les chiffres clés - Qu'est-ce qu'un cancer ? [Internet]. [cited 2020 Mar 27]. Available from: <https://www.e-cancer.fr/Comprendre-prevenir-depister/Qu-est-ce-qu-un-cancer/Chiffres-cles>
14. Alavanja MCR, Samanic C, Dosemeci M, et al. Use of Agricultural Pesticides and Prostate Cancer Risk in the Agricultural Health Study Cohort. *Am J Epidemiol*. Oxford Academic; 2003 May 1;**157**(9):800–814.
15. Alavanja MCR, Dosemeci M, Samanic C, et al. Pesticides and Lung Cancer Risk in the Agricultural Health Study Cohort. *Am J Epidemiol*. Oxford Academic; 2004 Nov 1;**160**(9):876–885.

16. De Roos Anneclaire J., Blair Aaron, Rusiecki Jennifer A., et al. Cancer Incidence among Glyphosate-Exposed Pesticide Applicators in the Agricultural Health Study. *Environ Health Perspect.* Environmental Health Perspectives; 2005 Jan 1;**113**(1):49–54.
17. Boulanger M, Tual S, Lemarchand C, et al. Lung cancer risk and occupational exposures in crop farming: results from the AGRiculture and CANcer (AGRICAN) cohort. *Occup Environ Med.* BMJ Publishing Group Ltd; 2018 Nov 1;**75**(11):776–785.
18. Lemarchand C, Tual S, Boulanger M, et al. Prostate cancer risk among French farmers in the AGRICAN cohort. *Scand J Work Environ Health.* [Scandinavian Journal of Work, Environment & Health, Danish National Research Centre for the Working Environment, Norwegian National Institute of Occupational Health, Finnish Institute of Occupational Health]; 2016;**42**(2):144–152.
19. Piel C, Pouchieu C, Tual S, et al. Central nervous system tumors and agricultural exposures in the prospective cohort AGRICAN. *Int J Cancer.* 2017;**141**(9):1771–1782.
20. Mostafalou S, Abdollahi M. Pesticides: an update of human exposure and toxicity. *Arch Toxicol.* 2017 Feb;**91**(2):549–599.
21. Engel LS, Hill DA, Hoppin JA, et al. Pesticide Use and Breast Cancer Risk among Farmers' Wives in the Agricultural Health Study. *Am J Epidemiol.* Oxford Academic; 2005 Jan 15;**161**(2):121–135.
22. Lerro CC, Koutros S, Andreotti G, et al. Organophosphate insecticide use and cancer incidence among spouses of pesticide applicators in the Agricultural Health Study. *Occup Environ Med.* 2015 Oct;**72**(10):736–744.
23. Lu Chensheng, Toepel Kathryn, Irish Rene, Fenske Richard A., Barr Dana B., Bravo Roberto. Organic Diets Significantly Lower Children's Dietary Exposure to Organophosphorus Pesticides. *Environ Health Perspect.* Environmental Health Perspectives; 2006 Feb 1;**114**(2):260–263.
24. Luo Y, Zhang M. Multimedia transport and risk assessment of organophosphate pesticides and a case study in the northern San Joaquin Valley of California. *Chemosphere.* 2009 May 1;**75**(7):969–978.
25. The 2015 European Union report on pesticide residues in food. *EFSA J.* 2017;**15**(4):e04791.
26. Hercberg S, Castetbon K, Czernichow S, et al. The Nutrinet-Santé Study: a web-based prospective study on the relationship between nutrition and health and determinants of dietary patterns and nutritional status. *BMC Public Health.* 2010 May 11;**10**:242.
27. Touvier M, Méjean C, Kesse-Guyot E, et al. Comparison between web-based and paper versions of a self-administered anthropometric questionnaire. *Eur J Epidemiol.* 2010 May 1;**25**(5):287–296.
28. Touvier M, Kesse-Guyot E, Méjean C, et al. Comparison between an interactive web-based self-administered 24 h dietary record and an interview by a dietitian for large-scale epidemiological studies. *Br J Nutr.* 2011 Apr;**105**(7):1055–1064.
29. Vergnaud A-C, Touvier M, Méjean C, et al. Agreement between web-based and paper versions of a socio-demographic questionnaire in the NutriNet-Santé study. *Int J Public Health.* 2011 Aug 1;**56**(4):407–417.
30. Baudry J, Méjean C, Allès B, et al. Contribution of Organic Food to the Diet in a Large Sample of French Adults (the NutriNet-Santé Cohort Study). *Nutrients.* 2015 Oct 21;**7**(10):8615–8632.
31. Kesse-Guyot E, Castetbon K, Touvier M, Hercberg S, Galan P. Relative validity and reproducibility of a food frequency questionnaire designed for French adults. *Ann Nutr Metab.* 2010;**57**(3–4):153–162.

32. Le Moullec N, Deheeger M, Herberg S, et al. Validation du manuel-photos utilisé pour l'enquête alimentaire de l'étude SU.VI.MAX. *Cah nutr diét*. Paris: Masson; 1996;**31**(3):158–164.
33. Etude Nutrinet-Santé (2013) Table de Composition des Aliments de l'étude Nutrinet-Santé (Nutrinet-Santé Study Food Composition Database). Paris: Economica.
34. UA-BW | CVUA Stuttgart (Startseite) [Internet]. [cited 2019 May 29]. Available from: <http://www.cvuas.de/pub/default.asp?subid=1>
35. Baudry J, Pointereau P, Seconda L, et al. Improvement of diet sustainability with increased level of organic food in the diet: findings from the BioNutriNet cohort. *Am J Clin Nutr*. 2019 Apr 1;**109**(4):1173–1188.
36. The 2014 European Union Report on Pesticide Residues in Food. *EFSA J*. 2016;**14**(10):e04611.
37. Yigit N, Velioglu YS. Effects of processing and storage on pesticide residues in foods. *Crit Rev Food Sci Nutr* [Internet]. 2019 Dec 20 [cited 2020 Jan 8]; Available from: <http://www.tandfonline.com/doi/abs/10.1080/10408398.2019.1702501>
38. GEMS/Food Euro. Second workshop on reliable evaluation of low-level contamination of food. Report on a workshop in the frame of GEMS/Food-Euro, EUR/HFA target 22. . 1995. Kulmbach, German. Geneva: WHO; 1995 May.
39. European Food Safety Authority. Management of left-censored data in dietary exposure assessment of chemical substances: Management of left-censored data in dietary exposure assessment of chemical substances. *EFSA J*. 2010;**8**:1557.
40. Barański M, Średnicka-Tober D, Volakakis N, et al. Higher antioxidant and lower cadmium concentrations and lower incidence of pesticide residues in organically grown crops: a systematic literature review and meta-analyses. *Br J Nutr*. 2014 Jun 26;**1**–18.
41. Hagströmer M, Oja P, Sjöström M. The International Physical Activity Questionnaire (IPAQ): a study of concurrent and construct validity. *Public Health Nutr*. 2006 Sep;**9**(6):755–762.
42. High Council for Public Health. French Nutrition and Health Programme's dietary guidelines for adults for the period 2017–2021. 2017 p. 7.
43. Chaltiel D, Adjibade M, Deschamps V, et al. Programme National Nutrition Santé – guidelines score 2 (PNNS-GS2): development and validation of a diet quality score reflecting the 2017 French dietary guidelines. *Br J Nutr*. Cambridge University Press; 2019 Aug;**122**(3):331–342.
44. Martínez-González MA, Sánchez-Tainta A, Corella D, et al. A provegetarian food pattern and reduction in total mortality in the Prevención con Dieta Mediterránea (PREDIMED) study. *Am J Clin Nutr*. 2014 Jul;**100** Suppl 1:320S–8S.
45. Monteiro CA, Cannon G, Levy R, et al. NOVA. The star shines bright. *World Nutr*. 2016 Jan 7;**7**(1–3):28–38.
46. Srour B, Fezeu LK, Kesse-Guyot E, et al. Ultraprocessed Food Consumption and Risk of Type 2 Diabetes Among Participants of the NutriNet-Santé Prospective Cohort. *JAMA Intern Med*. American Medical Association; 2020 Feb 1;**180**(2):283–291.
47. ICD - ICD-10-CM - International Classification of Diseases, Tenth Revision, Clinical Modification [Internet]. 2020 [cited 2020 Mar 9]. Available from: <https://www.cdc.gov/nchs/icd/icd10cm.htm>

48. Seung D, Lee L. Algorithms for non-negative matrix factorization. *Adv Neural Inf Process Syst*. 2001;**13**:556–562.
49. Gaujoux R, Seoighe C. A flexible R package for nonnegative matrix factorization. *BMC Bioinformatics*. 2010 Jul 2;**11**(1):367.
50. Arrebola JP, Belhassen H, Artacho-Cordón F, et al. Risk of female breast cancer and serum concentrations of organochlorine pesticides and polychlorinated biphenyls: A case–control study in Tunisia. *Sci Total Environ*. 2015 Jul 1;**520**:106–113.
51. Parada H, Wolff MS, Engel LS, et al. Organochlorine insecticides DDT and chlordane in relation to survival following breast cancer. *Int J Cancer*. 2016;**138**(3):565–575.
52. Charlier C. Breast cancer and serum organochlorine residues. *Occup Environ Med*. 2003 May 1;**60**(5):348–351.
53. Rizzati V, Briand O, Guillou H, Gamet-Payraastre L. Effects of pesticide mixtures in human and animal models: An update of the recent literature. *Chem Biol Interact*. 2016 Jul 25;**254**:231–246.
54. Hernández AF, Parrón T, Tsatsakis AM, Requena M, Alarcón R, López-Guarnido O. Toxic effects of pesticide mixtures at a molecular level: Their relevance to human health. *Toxicology*. 2013 May 10;**307**:136–145.
55. Tayour C, Ritz B, Langholz B, et al. A case–control study of breast cancer risk and ambient exposure to pesticides. *Environ Epidemiol*. 2019 Oct;**3**(5):e070.
56. Baudry J, Assmann KE, Touvier M, et al. Association of Frequency of Organic Food Consumption With Cancer Risk. *JAMA Intern Med*. 2018 Dec;**178**(12):1597–1606.
57. Ventura C, Zappia CD, Lasagna M, et al. Effects of the pesticide chlorpyrifos on breast cancer disease. Implication of epigenetic mechanisms. *J Steroid Biochem Mol Biol*. 2019 Feb 1;**186**:96–104.
58. Ventura C, Venturino A, Miret N, et al. Chlorpyrifos inhibits cell proliferation through ERK1/2 phosphorylation in breast cancer cell lines. *Chemosphere*. 2015 Feb;**120**:343–350.
59. Moyano P, García J, García JM, et al. Chlorpyrifos-induced cell proliferation in human breast cancer cell lines differentially mediated by estrogen and aryl hydrocarbon receptors and KIAA1363 enzyme after 24 h and 14 days exposure. *Chemosphere*. 2020 Jul;**251**:126426.
60. Yang KJ, Lee J, Park HL. Organophosphate Pesticide Exposure and Breast Cancer Risk: A Rapid Review of Human, Animal, and Cell-Based Studies. *Int J Environ Res Public Health*. 2020 Jul 13;**17**(14):5030.
61. Ventura C, Núñez M, Miret N, et al. Differential mechanisms of action are involved in chlorpyrifos effects in estrogen-dependent or -independent breast cancer cells exposed to low or high concentrations of the pesticide. *Toxicol Lett*. 2012 Sep 3;**213**(2):184–193.
62. Egbuta C, Lo J, Ghosh D. Mechanism of Inhibition of Estrogen Biosynthesis by Azole Fungicides. *Endocrinology*. Oxford Academic; 2014 Dec 1;**155**(12):4622–4628.
63. Nakagawa Y, Moore GA. Cytotoxic effects of postharvest fungicides, ortho-Phenylphenol, thiabendazole and imazalil, on isolated rat hepatocytes. *Life Sci*. 1995 Sep 1;**57**(15):1433–1440.
64. Schildkraut JM, Demark-Wahnefried W, DeVoto E, Hughes C, Laseter JL, Newman B. Environmental Contaminants and Body Fat Distribution. *Cancer Epidemiol Prev Biomark*. American Association for Cancer Research; 1999 Feb 1;**8**(2):179–183.

65. Pelletier C, Imbeault P, Tremblay A. Energy balance and pollution by organochlorines and polychlorinated biphenyls. *Obes Rev.* 2003;**4**(1):17–24.
66. Eckerson HW, Wyte CM, La Du BN. The human serum paraoxonase/arylesterase polymorphism. *Am J Hum Genet.* 1983 Nov;**35**(6):1126–1138.
67. Ferretti G, Bacchetti T, Moroni C, et al. Paraoxonase activity in high-density lipoproteins: a comparison between healthy and obese females. *J Clin Endocrinol Metab.* 2005 Mar;**90**(3):1728–1733.
68. Bajnok L, Seres I, Varga Z, et al. Relationship of serum resistin level to traits of metabolic syndrome and serum paraoxonase 1 activity in a population with a broad range of body mass index. *Exp Clin Endocrinol Diabetes Off J Ger Soc Endocrinol Ger Diabetes Assoc.* 2008 Nov;**116**(10):592–599.
69. Andreeva VA, Salanave B, Castetbon K, et al. Comparison of the sociodemographic characteristics of the large NutriNet-Santé e-cohort with French Census data: the issue of volunteer bias revisited. *J Epidemiol Community Health.* BMJ Publishing Group Ltd; 2015 Sep 1;**69**(9):893–898.
70. EU Pesticides database - European Commission [Internet]. [cited 2019 Sep 27]. Available from: <https://ec.europa.eu/food/plant/pesticides/eu-pesticides-database/public/?event=activesubstance.selection&language=EN>

Tables and Figures:

Table 1: Characteristics of the participants, NutriNet-Santé Study, 2014 (N=13,149).....	33
Table 2: Estimated pesticide exposure for cases and non-cases, lower-bound scenario, NutriNet-Santé Study, 2014 (N=13,149).....	34
Table 3: Spearman Correlations between 25 selected pesticides and NMF Components, NutriNet-Santé Study, 2014 (N=13,149)	35
Table 4: Cox models for associations between dietary pesticide exposure and postmenopausal breast cancer risk, NutriNet-Santé Study, 2014 (N=13,149).....	36
Table 5: Cox models for associations between dietary pesticide exposure and postmenopausal breast cancer risk, stratified analyses on BMI, NutriNet-Santé Study, 2014 (N=13,149) ¹	37
Table 6: Sensitivity analyses for associations between dietary pesticide exposure and postmenopausal cancer risk, NutriNet-Santé Study, 2014 (N=13,149).....	38
Figure 1: Flowchart for the sample selection, NutriNet-Santé Study, 2014 (N=13,149)	

Table 1: Characteristics of the participants, NutriNet-Santé Study, 2014 (N=13,149)

	All participants	Non-cases	Cases	P ¹
N	13,149	12,980	169	
Age , mean (SD)	60.49 (7.39)	60.48 (7.40)	61.15 (6.43)	0.24
Monthly income per household unit , %				0.27
<€1200	9.29	9.30	8.88	
€1200-1800	21.63	21.69	16.57	
€1800-2700	28.09	28.11	26.04	
>€2700	33.61	33.54	38.46	
Unwilling to answer	7.38	7.35	10.06	
Educational level , %				0.17
Less than high-school diploma	26.73	26.81	20.71	
High school diploma	17.58	17.53	20.71	
Post Graduate	55.69	55.65	58.58	
Occupational status , %				0.26
Employee, manual worker	12.29	12.36	7.10	
Intermediate profession	12.91	12.90	13.61	
Managerial staff, intellectual	14.15	14.18	11.83	
Retired	48.92	48.82	56.21	
Self-employed, farmer	1.59	1.59	1.18	
Unemployed or never employed	10.14	10.14	10.06	
Place of residence , %				0.85
Rural community	22.98	23.00	21.30	
Urban unit with a population <20,000 inhabitants	15.77	15.76	16.57	
Urban unit with a population between 20,000 and 200,000	19.45	19.48	17.75	
Urban unit with a population >200,000 inhabitants	41.80	41.76	44.38	
Smoking habits , %				0.20
current smoker	9.00	9.04	5.92	
former smoker	42.29	42.33	39.64	
never smoker	48.71	48.64	54.44	
Body Mass Index (kg/m ²), mean (SD)	24.22 (4.64)	24.22 (4.64)	24.57 (4.45)	0.16
Body Mass Index > 25kg/m ² , %	34.19	34.17	35.50	0.72
Physical activity , %				0.12
High	35.49	35.57	29.59	
Moderate	36.06	36.05	37.28	
Low	17.07	17.08	16.57	
Missing data	11.37	11.30	16.57	
Use of hormonal treatment for menopause , %				0.04
Yes	10.92	10.87	14.20	
No	83.30	83.29	84.02	
Missing data	5.78	5.83	1.78	
Parity , %				0.33
No children	14.47	14.43	17.75	
One child	17.56	17.59	15.38	
2 children	40.12	40.18	35.50	
More than 2 children	27.84	27.80	31.36	
Family history of cancer , %	52.25	52.19	57.40	0.18

¹P-values for comparisons between cases and non-cases using Chi-square tests or Wilcoxon tests, as appropriate.

Table 2: Estimated pesticide exposure for cases and non-cases, lower-bound scenario, NutriNet-Santé Study, 2014 (N=13,149)

Variable	Non-cases N=12,980		Cases N=169	
	Mean	SD	Mean	SD
Acetamiprid	0.0598	0.0778	0.0548	0.0966
Anthraquinone	0.0005	0.0015	0.0005	0.0010
Azadirachtin	0.0004	0.0005	0.0003	0.0004
Azoxystrobin	0.0451	0.0453	0.0457	0.0520
Boscalid	0.1312	0.1114	0.1218	0.1096
Carbendazim	0.0564	0.0581	0.0535	0.0708
Chlorpropham	0.0607	0.0616	0.0674	0.0632
Chlorpyrifos	0.0753	0.0663	0.0758	0.0751
Cypermethrin	0.0881	0.1107	0.0815	0.1370
Cyprodinil	0.0821	0.0892	0.0782	0.0790
Difenoconazole	0.0190	0.0177	0.0175	0.0157
Dimethoate Ometoate	0.0106	0.0134	0.0090	0.0138
Fenhexamid	0.1067	0.1455	0.0897	0.1060
Glyphosate	0.0035	0.0048	0.0040	0.0057
Lambda-Cyhalothrin	0.0116	0.0112	0.0117	0.0122
Imazalil	0.8459	1.0395	0.9367	1.1376
Imidacloprid	0.0791	0.0750	0.0831	0.0750
Iprodione	0.1591	0.1833	0.1552	0.1706
Malathion	0.0003	0.0004	0.0003	0.0003
Methamidophos	0.0002	0.0003	0.0003	0.0005
Profenofos	0.0000	0.0001	0.0000	0.0001
Pyrethrins	0.0020	0.0017	0.0018	0.0015
Spinosad	0.1717	0.1870	0.1447	0.1660
Tebuconazole	0.0385	0.0471	0.0373	0.0399
Thiabendazole	0.2882	0.3247	0.3239	0.3322

Table 3: Spearman Correlations between 25 selected pesticides and NMF Components, NutriNet-Santé Study, 2014 (N=13,149)

Compounds	NMF Component 1	NMF Component 2	NMF Component 3	NMF Component 4
Acetamiprid	0.34	0.41	0.26	0.87
Anthraquinone	0.17	0.19	-0.06	0.18
Azadirachtin	-0.09	*-0.01	0.53	*-0.01
Azoxystrobin	0.59	0.71	-0.18	0.16
Boscalid	0.51	0.90	-0.13	0.19
Carbendazim	0.31	0.38	0.31	0.89
Chlorpropham	0.35	0.53	-0.30	0.12
Chlorpyrifos	0.73	0.44	0.11	0.60
Cypermethrin	0.29	0.27	0.36	0.93
Cyprodinil	0.50	0.91	-0.12	0.16
Difenoconazole	0.52	0.68	*0.02	0.47
Dimethoate Ometoate	0.36	0.42	0.26	0.79
Fenhexamid	0.46	0.79	-0.11	0.12
Glyphosate	0.38	0.45	-0.12	0.17
Imazalil	1.00	0.37	-0.11	0.14
Imidacloprid	0.51	0.24	0.20	0.56
Iprodione	0.52	0.91	-0.10	0.15
Lambda-Cyhalothrin	0.56	0.84	-0.08	0.24
Malathion	0.76	0.49	-0.10	0.17
Methamidophos	0.32	0.35	-0.19	0.17
Profenofos	0.95	0.38	-0.12	0.17
Pyrethrins	0.04	0.02	0.30	0.03
Spinosad	-0.07	-0.09	0.99	0.35
Tebuconazole	0.55	0.84	-0.10	0.19
Thiabendazole	0.98	0.36	-0.11	0.16

*p-value for Spearman correlation >0.05

NMF: Non-negative Matrix Factorization

Bold values denote correlation coefficients >0.60.

Table 4: Cox models for associations between dietary pesticide exposure and postmenopausal breast cancer risk, NutriNet-Santé Study, 2014 (N=13,149)

	Quintile 1	Quintile 2	Quintile 3	Quintile 4	Quintile 5	Total	P value for trend
NMF Component 1							
Number of participants	2629	2630	2630	2630	2630	13,149	
Incident Cases	27	36	34	30	42	169	
Person-years	11,385.52	11,507.50	11,566.09	11,477.81	11,266.78	57,203.70	
Model 1 ¹ , HR (95% CI)	1	1.34 (0.81; 2.21)	1.24 (0.75; 2.06)	1.14 (0.68; 1.93)	1.73 (1.05; 2.84)		0.09
Model 2 ² , HR (95% CI)	1	1.33 (0.80; 2.19)	1.25 (0.75; 2.07)	1.15 (0.68; 1.95)	1.78 (1.08; 2.93)		0.07
Model 3 ³ , HR (95% CI)	1	1.32 (0.80; 2.18)	1.24 (0.75; 2.06)	1.15 (0.68; 1.94)	1.77 (1.07; 2.91)		0.08
NMF Component 2							
Number of participants	2629	2630	2630	2630	2630	13,149	
Incident cases	38	43	28	26	34	169	
Person-years	11,504.90	11,453.11	11,459.27	11,499.86	11,286.57	57,203.70	
Model 1 ¹ , HR (95% CI)	1	1.11 (0.72; 1.72)	0.72 (0.44; 1.17)	0.69 (0.42; 1.15)	0.96 (0.59; 1.56)		0.30
Model 2 ² , HR (95% CI)	1	1.09 (0.70; 1.69)	0.71 (0.44; 1.16)	0.70 (0.42; 1.16)	1.00 (0.61; 1.63)		0.38
Model 3 ³ , HR (95% CI)	1	1.08 (0.70; 1.68)	0.71 (0.43; 1.16)	0.69 (0.42; 1.15)	0.99 (0.61; 1.62)		0.37
NMF Component 3							
Number of participants	2629	2630	2630	2630	2630	13,149	
Incident cases	47	42	23	31	26	169	
Person-years	11,304.65	11,374.78	11,437.14	11,558.22	11,528.91	57,203.70	
Model 1 ¹ , HR (95% CI)	1	0.88 (0.58; 1.34)	0.47 (0.29; 0.78)	0.64 (0.40; 1.01)	0.57 (0.34; 0.93)		0.006
Model 2 ² , HR (95% CI)	1	0.89 (0.59; 1.35)	0.48 (0.29; 0.80)	0.66 (0.41; 1.04)	0.59 (0.36; 0.97)		0.01
Model 3 ³ , HR (95% CI)	1	0.89 (0.59; 1.35)	0.48 (0.29; 0.80)	0.66 (0.41; 1.04)	0.59 (0.36; 0.98)		0.01
NMF Component 4							
Number of participants	2629	2630	2630	2630	2630	13,149	
Incident cases	36	41	33	36	23	169	
Person-years	11,294.95	11,425.20	11,459.16	11,442.46	11,581.94	57,203.70	
Model 1 ¹ , HR (95% CI)	1	1.14 (0.72; 1.78)	0.93 (0.58; 1.50)	1.01 (0.64; 1.62)	0.65 (0.38; 1.12)		0.13
Model 2 ² , HR (95% CI)	1	1.15 (0.73; 1.80)	0.95 (0.59; 1.53)	1.02 (0.64; 1.63)	0.66 (0.39; 1.12)		0.13
Model 3 ³ , HR (95% CI)	1	1.15 (0.73; 1.80)	0.95 (0.59; 1.53)	1.02 (0.64; 1.62)	0.66 (0.39; 1.12)		0.13

Abbreviations: NMF:Non-negative Matrix Factorization; HR: Hazard Ratio; 95% CI : 95% Confidence Interval; sPNNS-GS2: Simplified Programme National Nutrition Santé Guideline Score 2

¹Adjusted for smoking practices, educational level, physical activity, alcohol intake, alcohol-free energy intake, Body Mass Index, height, family history of cancer, menopausal treatment, parity

²Adjusted for Model 1 + sPNNS-GS2 score

³Adjusted for Model 1 + sPNNS-GS2 score + residing currently in an agricultural area

Table 5: Cox models for associations between dietary pesticide exposure and postmenopausal breast cancer risk, stratified analyses on BMI, NutriNet-Santé Study, 2014 (N=13,149)¹

	BMI <25kg/m ² N=8654, 109 cases					BMI >25kg/m ² N= 4495, 60 cases						
	Quintile 1	Quintile 2	Quintile 3	Quintile 4	Quintile 5	Quintile 1	Quintile 2	Quintile 3	Quintile 4	Quintile 5		
N	1731	1732	1732	1732	1732	901	901	901	901	901		
	HR	HR (95% CI)	HR (95% CI)	HR (95% CI)	HR (95% CI)	P-value for trend	HR	HR (95% CI)	HR (95% CI)	HR (95% CI)	HR (95% CI)	P-value for trend
NMF		1.05	0.65	1.02	0.98		2.83	2.58	3.02	4.13		
Component 1	1	(0.60; 1.85)	(0.34; 1.23)	(0.57; 1.81)	(0.53; 1.81)	0.88	1	(0.83; 6.87)	(0.90; 7.38)	(1.08; 8.47)	(1.50; 11.44)	0.006

Abbreviations: NMF: Non-negative Matrix Factorization; HR: Hazard Ratio; 95% CI: 95% Confidence Interval; BMI: Body Mass Index; sPNNS-GS2: Simplified Programme National Nutrition Santé Guideline Score 2

¹All models adjusted for sPNNS-GS2 score, smoking practices, educational level, physical activity, alcohol intake, alcohol-free energy intake, BMI, height, family history of cancer, menopausal treatment, parity.

Table 6: Sensitivity analyses for associations between dietary pesticide exposure and postmenopausal cancer risk, NutriNet-Santé Study, 2014 (N=13,149)

Model excluding early cases (1 year)	Quintile 1	Quintile 2		Quintile 3		Quintile 4		Quintile 5		P-value for trend
N=13,120; 140 cases	HR	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI	
NMF Component 1	1	1.11	(0.65; 1.90)	1.19	(0.70; 2.03)	1.02	(0.58; 1.78)	1.37	(0.80; 2.36)	0.38
NMF Component 2	1	0.96	(0.60; 1.55)	0.65	(0.38; 1.11)	0.66	(0.38; 1.14)	0.86	(0.50; 1.47)	0.23
NMF Component 3	1	0.96	(0.61; 1.50)	0.46	(0.26; 0.81)	0.62	(0.37; 1.04)	0.62	(0.36; 1.07)	0.016
NMF Component 4	1	1.22	(0.75; 2.01)	1.15	(0.69; 1.91)	1.00	(0.59; 1.69)	0.59	(0.32; 1.10)	0.10
Model with additional adjustment for ultra-processed foods ²	Quintile 1 N=2629	Quintile 2 N=2630		Quintile 3 N=2630		Quintile 4 N=2630		Quintile 5 N=2630		P-value for trend
N=13,149; 169 cases	HR	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI	
NMF Component 1	1	1.33	(0.80; 2.19)	1.24	(0.75; 2.07)	1.15	(0.68; 1.94)	1.77	(1.08; 2.92)	0.08
NMF Component 2	1	1.09	(0.70; 1.69)	0.71	(0.43; 1.16)	0.70	(0.42; 1.16)	0.99	(0.61; 1.62)	0.37
NMF Component 3	1	0.88	(0.58; 1.34)	0.47	(0.29; 0.79)	0.64	(0.40; 1.03)	0.58	(0.35; 0.96)	0.009
NMF Component 4	1	1.15	(0.73; 1.80)	0.94	(0.59; 1.52)	1.01	(0.64; 1.62)	0.65	(0.38; 1.11)	0.12
Model with additional adjustment for provegetarian Score ³	Quintile 1 N=2629	Quintile 2 N=2630		Quintile 3 N=2630		Quintile 4 N=2630		Quintile 5 N=2630		P-value for trend
N=13,149; 169 cases	HR	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI	
NMF Component 1	1	1.33	(0.80; 2.19)	1.23	(0.74; 2.04)	1.13	(0.67; 1.91)	1.72	(1.04; 2.82)	0.009
NMF Component 2	1	1.09	(0.70; 1.70)	0.71	(0.43; 1.16)	0.69	(0.41; 1.14)	0.96	(0.59; 1.56)	0.30
NMF Component 3	1	0.88	(0.58; 1.34)	0.47	(0.29; 0.78)	0.64	(0.40; 1.02)	0.57	(0.34; 0.95)	0.008
NMF Component 4	1	1.14	(0.73; 1.79)	0.94	(0.58; 1.51)	1.01	(0.64; 1.62)	0.65	(0.38; 1.12)	0.12

Abbreviations: NMF: Non-negative Matrix Factorization; HR: Hazard Ratio; 95 % CI: 95% Confidence Interval; BMI: Body Mass Index; sPNNS-GS2: Simplified

Programme National Nutrition Santé Guideline Score 2

¹Adjusted for sPNNS-GS2 score, smoking practices, educational level, physical activity, alcohol intake, alcohol-free energy intake, BMI, height, family history of cancer, menopausal treatment, parity

²Adjusted for sPNNS-GS2 score, smoking practices, educational level, physical activity, alcohol intake, alcohol-free energy intake, BMI, height, family history of cancer, menopausal treatment, parity and percentage of ultra-processed foods in the diet

³Adjusted for provegetarian score, smoking practices, educational level, physical activity, alcohol intake, alcohol-free energy intake, BMI, height, family history of cancer, menopausal treatment, parity