



HAL
open science

Molecular characterization of *Cryptosporidium* isolates from diarrheal dairy calves in France

Mohamed Mammeri, Aurélie Chevillot, Ilham Chenafi, Myriam Thomas, Christine Julien, Isabelle Vallée, Bruno Polack, Jérôme Follet, Karim Tarik
Adjou

► To cite this version:

Mohamed Mammeri, Aurélie Chevillot, Ilham Chenafi, Myriam Thomas, Christine Julien, et al.. Molecular characterization of *Cryptosporidium* isolates from diarrheal dairy calves in France. *Veterinary Parasitology: Regional Studies and Reports*, 2019, 18, pp.100323. 10.1016/j.vprsr.2019.100323 . hal-03176056

HAL Id: hal-03176056

<https://hal.inrae.fr/hal-03176056v1>

Submitted on 25 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

20 **Abstract**

21 *Cryptosporidium* is an obligate intracellular protist parasite infecting a wide range of
22 vertebrate hosts and causes significant intestinal disease in both animals and humans, as some
23 species are zoonotic. Cattle and especially calves have been identified as one of the most
24 common reservoirs of this protist. However, little is known about the genetics of
25 *Cryptosporidium* in calves in some regions of France. The aim of this study was to detect and
26 isolate *Cryptosporidium* spp. in faecal samples from naturally infected pre-weaned calves (\leq
27 45 days-old) in France. A total of 35 diarrhoeic pre-weaned calf faecal samples were collected
28 from 26 dairy cattle farms in six departments (French administrative provinces).
29 *Cryptosporidium* presence was established by microscopically screening samples for oocysts
30 with an immunofluorescent (DFA) staining method. DFA-positive samples were then
31 analysed by PCR-RFLP and *18S rRNA* gene sequencing to determine species.
32 *Cryptosporidium parvum*-positive samples were subtyped via nested PCR analysis of a partial
33 fragment of the 60 kDa glycoprotein (*gp60*) gene product. Data were then integrated into
34 phylogenetic tree analysis. DFA revealed the presence of *Cryptosporidium* oocysts in 31 out
35 of 35 (88%) samples. Combined with *18S rRNA* gene analysis results, *C. parvum* was
36 detected in 30 samples. Subtyping analysis in 27/30 samples (90%) of the *C. parvum* isolates
37 revealed two zoonotic subtype families, IIa (24/27) and IId (3/27). Four subtypes were
38 recognised within the subtype family IIa, including the hypertransmissible IIaA15G2R1
39 subtype that is the most frequently reported worldwide (21/27), IIaA17G3R1 (1/27),
40 IIaA17G1R1 (1/27), and IIaA19G1R1 (1/27). Two subtypes were recognised within the IId
41 subtype family including IIdA22G1 (2/27) and IIdA27G1 (1/27). These findings illustrate the
42 high occurrence of *Cryptosporidium* in calves in dairy herds and increase the diversity of
43 molecularly characterised *C. parvum* isolates with the first description of IIaA17G3R1,
44 IIaA19G1R1, and IId subtypes in France. The presence of zoonotic *C. parvum* subtype

45 families (IIa, IIb) in this study suggests that pre-weaned calves are likely to be a significant
46 reservoir of zoonotic *C. parvum*, and highlights the importance of animal to human
47 cryptosporidiosis transmission risk. Further molecular studies in calves and small ruminants
48 from other French regions are required to better understand the epidemiology of
49 cryptosporidiosis in France.

50 **Keywords:** *Cryptosporidium parvum*, PCR-RFLP, 18S rRNA gene, gp60 subtype, France,
51 calves, zoonotic parasite.

52

53

54

55

56

57

58

59

60

61

62

63

64

65

67 1. Introduction

68 *Cryptosporidium* is an obligate intracellular protist parasite infecting a wide range of
69 vertebrate hosts, including humans (Bouzid et al., 2013). The clinical importance of
70 *Cryptosporidium* spp. was highlighted when this parasite was associated with both waterborne
71 and foodborne outbreaks (Efstratiou et al., 2017; Ryan et al., 2018). Currently, more than 30
72 species have been described and validated (Osman et al., 2017). However, due to strong inter-
73 species similarities in microscopic size and shape, species within this genus require additional
74 identification via genetic characterisation. Molecular biology techniques have enabled the
75 description of species that are highly host-specific, as well as others that are capable of
76 infecting many hosts. *Cryptosporidium parvum* is considered to be the most prevalent species
77 worldwide and a major zoonotic transmission risk (Xiao, 2010). Using molecular approaches
78 to genetically characterise *Cryptosporidium* spp. has facilitated an improved understanding of
79 cryptosporidiosis epidemiology (Xiao, 2010). Subtype analysis using the *C. parvum* 60 kDa
80 glycoprotein locus (*gp60*) has revealed both human- and zoonotic-specific subtypes
81 (Sulaiman et al., 2005).

82 Within this context, cattle—particularly pre-weaned dairy calves—are recognised as common
83 hosts of *Cryptosporidium* spp., with both high infection rates and extremely high rates of
84 oocyst excretion in the faeces (Thomson et al., 2017). Cryptosporidiosis in neonatal calves is
85 mainly due to *C. parvum* infection, and is characterised by diarrhoea, dehydration, delayed
86 growth, and weight loss, resulting in considerable economic losses associated with calf
87 morbidity and mortality (Olson et al., 2004; Thomson et al., 2017). In addition, young calves
88 have been considered as a potential source of human cryptosporidiosis in several outbreaks
89 (Xiao, 2010). Besides *C. parvum*, cattle can also be infected by bovine specific species

90 including *C. bovis*, *C. andersoni* and *C. ryanae* with a marked aged-related pattern (Fayer et
91 al., 2006; Santín et al., 2008). Asymptomatic adult cattle can also be sporadically infected by
92 other species including *C. felis* (Cardona et al., 2015). It is well known that not all
93 *Cryptosporidium* species excreted by cattle are zoonotic. Consequently, the molecular
94 characterization of these species is vital to better understanding cattle cryptosporidiosis
95 epidemiology and the zoonotic potential of specific *Cryptosporidium* isolates (Fayer et al.,
96 2000).

97 Since the first report of the prevalence of bovine cryptosporidiosis in suckling (50-95%) and
98 dairy calves (16.8-51,8%) in France in 1999 (Naciri et al., 1999), many cryptosporidiosis
99 epidemiology studies have been solely based on faecal sample microscopy reporting different
100 prevalences: 41.5 % (Delafosse et al., 2015) and 17.9 % (Lefay et al., 2000). However, these
101 traditional staining or flotation screening methods can lead to suboptimal *Cryptosporidium*
102 detection especially with reduced oocyst shedding intensity thus underestimating the real
103 parasitic prevalence in herds. Thus far, only a few studies have used molecular tools to
104 characterise *Cryptosporidium* spp. in cattle in 15 of the 95 metropolitan French departments
105 (administrative regions): Allier, Côtes-d'Armor, Ille-et-Vilaine, Landes, Mayenne, Morbihan,
106 Moselle, Orne, Pas-de-Calais, Puy-de-Dôme, Pyrénées-Atlantiques, Hautes-Pyrénées, Deux-
107 Sèvres, Tarn, and Vendée (Follet et al., 2011; Ngouanesavanh et al., 2006; Rieux et al., 2014,
108 2013b, 2013c, 2013a). Little is known about which *Cryptosporidium* spp. infects calves in
109 other French departments. However, subtype analysis solely on the basis of *C. parvum gp60*
110 amplicons is insufficient (Follet et al., 2011; Rieux et al., 2014, 2013c, 2013a), thus there is a
111 strong need for more molecular epidemiological data on French bovine cryptosporidiosis to
112 fully determine the breadth of *C. parvum* genetic diversity. Thus, the aim of this study was to
113 characterise *Cryptosporidium* isolates in calves from different French departments: Allier,
114 Ardèche, Côte-d'Or, Moselle, Saône-et-Loire, and Yonne. Furthermore, through genetic

115 characterization, this study led the authors to investigate the potential of calves as a zoonotic
116 reservoir for human infection.

117 **Materials and methods**

118 **1.1. Sample collection and study design**

119 In order to generate a global understanding of the different *Cryptosporidium* species and *C.*
120 *parvum* genotypes in calf populations, mostly from uninvestigated French departments, a total
121 of 35 unique faecal samples were collected from pre-weaned calves (\leq 45 days-old) on 26
122 dairy farms (with or without high density breeding, and/or the presence of diarrhoea), located
123 in six departments, between November 2017 and April 2018, with one sample per studied
124 animal. In order to perform anonymous sampling, farms were arbitrarily numbered from E1 to
125 E26 and collected faecal samples were labelled B1 to B35. A map (**Figure 1**) representing the
126 origin of the samples was edited using *Cartes et Données*-[®] Articque
127 (<https://www.articque.com/solutions/cartes-et-donnees/>). Samples were directly collected
128 from the rectum using plastic gloves and transferred into a sterile plastic container. Samples
129 were maintained at 4°C for a maximum of 48 h prior to processing and analysis. For each
130 animal, the sampling date, animal identification number, age, and faecal score were recorded.
131 Faecal consistency was scored on a scale from 0 to 4 (0: normal without mucus; 1: pasty and
132 thick, formed or not, viscous 2: creamy; 3: semi-fluid; 4: liquid) as modified from Koudela
133 and Jirí (1997). The apparent occurrence (percentage of *Cryptosporidium* infection) was
134 calculated by dividing the number of *Cryptosporidium*-positive calves as assessed with
135 different techniques by the total number of tested animals, multiplied by 100.

136 **1.2. Microscopy**

137 One gram of faeces was either suspended in 10 mL of distilled water or concentrated using
138 the diethyl ether-PBS concentration method as previously described (Castro-Hermida et al.,
139 2005). The supernatant was removed and the pellet was resuspended in distilled water up to
140 500 μ L. *Cryptosporidium* oocysts were detected from 20 μ L of both concentrated and non-
141 concentrated oocyst solutions via direct immunofluorescence assay (DFA) using the
142 commercial Merifluor *Cryptosporidium/Giardia* immunofluorescence assay (Meridian
143 Diagnostics, Inc., Milano, Italy) as indicated by the manufacturer, and including previously
144 described modifications (Mammeri et al., 2018). Stained slides were observed with a Leica
145 fluorescent microscope using the Leica Application Suite software (version 4.5.0; Leica
146 Microsystems; Inc., Switzerland) at x40 magnification. The number of *Cryptosporidium*
147 oocysts per gram of faeces (OPG) was obtained by multiplying the total number of oocysts by
148 the dilution factor.

149 **1.3. DNA extraction**

150 Genomic DNA was extracted from 200 μ L of concentrated oocysts using the QIAmp DNA
151 Stool Mini Kit (Qiagen, France) according to manufacturer's instructions. Briefly, samples
152 were suspended in lysis buffer and the oocysts were disrupted by subjecting them to an
153 additional step of 3 freeze-thaw cycles (freezing in liquid nitrogen for 1 min followed by
154 heating in a 90°C water bath for 1 min) before DNA extraction. DNA samples were then
155 stored at -20°C until molecular analysis.

156 **1.4. *Cryptosporidium* genotyping with nested 18S rRNA and gp60 PCR**

157 *Cryptosporidium* species were further analysed in DFA-positive samples with nested PCR
158 which amplified a 847 bp fragment from the 18S rRNA gene (Xiao et al., 1999).
159 Amplification products were subsequently visualised by electrophoresis in 2% agarose gels

160 stained with ethidium bromide (0.5 µg/mL). *Cryptosporidium* species were identified by
161 performing restriction fragment length polymorphism (RFLP) analysis with *SspI* and *MboII*
162 endonucleases on *18S rRNA* PCR products (New England BioLabs, France) as previously
163 described (Feng et al., 2007). Fragments were subsequently visualised by electrophoresis in
164 3% MetaPhor Agarose (Ozyme, France) gels stained with ethidium bromide solution (0.5
165 µg/mL). For example, in the case of *C. parvum*, *SspI* and *MboII* would generate three bands
166 (449, 267, and 108 bp) and two bands (771 and 769 bp), respectively (Feng et al., 2007).

167 *C. parvum* samples were subtyped by nested PCR-sequence analysis of the partial *gp60* gene
168 sequence as previously described (Gatei et al., 2006). Amplification products were
169 subsequently visualised by electrophoresis in 2% agarose gels stained with ethidium bromide.

170 In each PCR reaction, both positive and negative control samples were included. The positive
171 control consisted of DNA extracted from 10⁶ *C. parvum* Iowa strain oocysts (Waterborne Inc.,
172 New Orleans, Louisiana, USA) while the negative control was purified water. In addition,
173 Bovine Serum Albumin (BSA) was added to all PCR mixtures at the final concentration of
174 400 ng/µL as described before (Jiang et al., 2005).

175 **1.5. DNA sequence analysis**

176 PCR products of the two target genes (*18S rRNA* and *gp60*) were sequenced on both strands
177 using internal primer sets by Eurofins Genomics (France) and Genoscreen (France),
178 respectively. Consensus sequences were edited using the BioEdit Sequence Alignment Editor
179 software and compared with published GenBank sequences using the freely-available Basic
180 Local Alignment Search Tool (BLAST) from the National Center for Biotechnology
181 Information (NCBI) (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). Representative nucleotide
182 consensus sequences generated in this study were deposited into the GenBank database under
183 accession numbers as indicated in **Table 1**. *C. parvum* subtypes were recognised based on the

184 number of trinucleotide repeats (TCA or TCG) or rare (R) repeats in the *gp60* sequence as
185 previously described (Sulaiman et al., 2005).

186 **1.6. Statistical analysis**

187 The Pearson correlation test was used to analyse data by comparing infection occurrence with
188 animal age, faecal score, and *C. parvum* subtypes. The results were considered statistically
189 significant when the *P*-value was < 0.05.

190 **2. Results**

191 **2.1. DFA screening**

192 Among the 35 samples included in this study, microscopic examinations revealed the
193 presence of *Cryptosporidium* oocysts in 88.5% of the samples from pre-weaned calves
194 (31/35) (**Table 1**).

195 Immunofluorescent-positive samples indicated that calves excreted from 10^4 to 3.41×10^6 OPG
196 via direct oocyst detection, or from 1.25×10^2 to 2.23×10^6 OPG following oocyst sample
197 concentration (**Figure. 2**). Results indicated oocyst loss in 20 of 30 (66.66%) of the samples
198 (the small amount of faeces from the B26 sample was concentrated and used to perform DFA
199 and PCRs). No correlation existed between age and infection intensity when DFA was used to
200 directly detect oocysts in faeces ($P = 0.68$), whereas a significant correlation was observed
201 between the faecal consistency score and infection intensity using this method ($P = 0.03$).

202 **2.2. Molecular analysis of the *18S rRNA* gene**

203 The *18S rRNA* gene fragment was amplified from concentrated oocysts for all 31 DFA-
204 positive samples (100%). *C. parvum* was identified in 29 of 31 (93.54%) DNA samples by
205 sequence analyses of the *18S rRNA* PCR products, as sequencing results were not usable for
206 the two remaining samples (B11 and B16). *C. parvum* was the only species identified in 29 of

207 31 (93.54%) samples by *18S rRNA* PCR-RFLP, and no mixed infections were observed. PCR
208 amplification failed for two samples (B16 and B30). Combining *18 S rRNA* sequence analysis
209 and PCR-RFLP results indicated that *C. parvum* was detected in 30 of 31 samples (96.7%).
210 Both techniques returned negative results for the B16 sample.

211 **2.3. Molecular analysis of *gp60* gene**

212 Of the 30 *Cryptosporidium 18S rRNA* gene-positive samples, all isolates yielded a *gp60* PCR
213 product. Successful sequencing analysis of the *gp60* gene in 27 (90%) *Cryptosporidium*
214 isolates identified two subtype families: Ila and IId (**Table 1**). Twenty-four out of 27 isolates
215 (88.8%) belonged to subtype family Ila and the remaining three isolates (11.2%) belonged to
216 IId.

217 Four subtypes were recognised within the Ila subtype family including IlaA15G2R1 (21/27),
218 IlaA17G3R1 (1/27), IlaA17G1R1 (1/27), and IlaA19G1R1 (1/27); while two subtypes were
219 identified as part of the IId subtype family, including IIdA22G1 (2/27) and IIdA27G1 (1/27).

220 Three PCR product for sequencing of the *gp60* loci from isolate which was successfully
221 identified as *C. parvum* by PCR-RFLP of the *18S rRNA* (isolate B11), sequence analyses of
222 the *18S rRNA* PCR products (isolate B30), or both techniques (B20), were not usable.

223 Several samples could not be used for all analyses; isolate B11 with successful *C. parvum*
224 identification via *18S rRNA* PCR-RFLP had a failed *gp60* PCR; sequencing of *18S rRNA*
225 PCRs could not be performed on isolate B30; and neither technique could be performed on
226 the B20 isolate.

227 Among the five farms infected with *C. parvum* and which contained more than two calves in
228 this study, three farms carried a single subtype (E1, E2, E17), which in this case was
229 IlaA15G2R1. For the E4 herd (B10 and B11 samples), one of the samples was linked to the
230 subtype IIdA22G1, while the other sample could not be associated with a specific subtype as

231 it likely carried a mixed infection of at least two subtypes. In herd E23, two circulating
232 subtypes were identified from samples B31 and B32, corresponding to subtypes IIaA17G1R1
233 and IIaA15G2R1, respectively.

234 No correlation was reported between *C. parvum* subtype and age ($P = 0.33$), faecal score
235 ($P=0.4$), or infection intensity ($P=0.8$).

236 3. Discussion

237 Several different techniques currently exist to detect *Cryptosporidium* infection in animal
238 and/or human faecal samples (Chalmers and Katzer, 2013). In the present study, DFA
239 screening was used to detect the presence of *Cryptosporidium* oocysts. This technique is
240 recommended as a screening method in epidemiological studies (Thompson and Ash, 2016)
241 due to its high sensitivity and rapidity, however a major disadvantage is that it is unable to
242 differentiate between *Cryptosporidium* species (Amar et al., 2004). In our study, the apparent
243 occurrence was estimated at 88.6% using DFA as a diagnostic screening technique. This high
244 value suggests that *Cryptosporidium* is a common parasite in pre-weaned calves in France,
245 and is also similar to that seen in other French epidemiological studies. In the Brittany region
246 (north-west France), 70.4% (100/142) of studied calves were reported to be infected by
247 *Cryptosporidium* spp. (Follet et al., 2011) using 18S rRNA nested-PCR techniques. In the
248 Deux-Sèvres department, a 92-100% occurrence of *Cryptosporidium* infection in calves was
249 reported using DFA (Rieux et al., 2014). High *Cryptosporidium* occurrence in calves has also
250 been reported in other countries (e.g. the USA and Vietnam) (Nguyen et al., 2007; Santín et
251 al., 2004). However, a relatively low occurrence was recently described in western France
252 where only 402 out of the 968 (41.5%) analysed samples were positive using the Ziehl-
253 Neelsen fuchsin staining method (Delafosse et al., 2015). A worldwide *Cryptosporidium*
254 occurrence ranging from 3.4 to 96.6% has been reported in calves as previously reviewed

255 (Thomson et al., 2017). This extraordinarily broad variation in *Cryptosporidium* infection
256 occurrence with extreme dissimilarities between countries may be due to geographical
257 distribution (location of study farms), climatic conditions, and different farm management
258 practices, but also may relate to study design (number of specimens collected, number of
259 studied farms, specimen collection season, the diagnostic method used, etc.). Some of these
260 factors may also influence *Cryptosporidium* transmission between animals (Bamaiyi and
261 Redhuan, 2016; Causapé et al., 2002; Maurya et al., 2013; Mohammed et al., 1999; Ranjbar
262 and Fattahi, 2017). In addition to substantial differences in farm management, the lower
263 average age of infected calves in the present study might account for the higher occurrence,
264 since *Cryptosporidium* tends to be more frequently present in younger calves aged less than
265 one month (Santín et al., 2004).

266 Our results also indicated that four samples out of 35 (11.4%) were DFA-negative for the
267 presence of *Cryptosporidium* oocysts despite reported neonatal diarrhoea. This could be due
268 to the presence of other diarrhoea-causing enteropathogens (infection by *Escherichia coli*,
269 Rotavirus, Coronavirus, or Coccidia) (Thomson et al., 2017).

270 DFA-positive faecal samples indicated that calves excreted from between 1×10^4 to 3.41×10^6
271 oocysts when DFA was performed directly on faeces. This high level of oocyst excretion
272 reflects that of other studies performed in France which report that oocyst excretion intensity
273 can reach 8×10^6 oocysts per gram of faeces in some calves (Rieux et al., 2013b). When we
274 compared our DFA method *with* oocyst concentration to that *without*, we observed oocyst loss
275 in 65.5% of samples, which has previously been reported for concentration steps. However,
276 these purification techniques are still acceptable for use in young ruminants because they
277 excrete such a large number of oocysts (Fayer et al., 2000). On the other hand, oocyst

278 concentration may facilitate PCR detection of *Cryptosporidium* by eliminating naturally-
279 occurring PCR inhibitors from faeces (Elwin et al., 2012).

280 We reported a strong correlation between the faecal consistency score and infection intensity,
281 as estimated by DFA performed directly on faeces. This result is in accordance with previous
282 studies (El-Khodery and Osman, 2008; Trotz-Williams et al., 2007) which have reported a
283 strong correlation between *Cryptosporidium* oocyst shedding and calf diarrhoea.
284 Nevertheless, it is important to compare infection intensity (oocyst shedding) with the species
285 or subtypes detected by PCR, and faecal consistency, when generating inferences about the
286 clinical impact of cryptosporidiosis. DFA results highlighted the absence of any correlation
287 between calf age, oocyst shedding intensity, and *C. parvum* subtypes. This observation could
288 be explained by the fact that animals were not all infected at the same time, at the same age,
289 and the infectious dose may not have been the same. In addition, when the intestinal
290 epithelium is severely damaged by *Cryptosporidium* infection, parasitic reproduction is
291 impaired, even though marked diarrhoeal clinical signs are observed.

292 Despite positive microscopic identification of oocysts in samples, PCR analysis of the two
293 *18S rRNA* and *gp60* target genes was only positive in 93.5% and 90% samples, respectively.
294 Although the nested PCR method has been described as very sensitive and specific (Bhat et
295 al., 2014), this failure to yield PCR products was due to the unsuccessful PCR amplification.
296 These false negative PCR results could be explained by many factors: low numbers of oocysts
297 in some samples, the presence of PCR inhibitors in faecal samples (haemoglobin, bilirubin,
298 and bile acids), failed extraction procedures, failed oocyst disruption and lysis, insufficient
299 DNA collected, or nucleic acid degradation (Johnson et al., 1995; Lantz et al., 1997;
300 McLauchlin et al., 1999; Yu et al., 2009). Thus far, four common *Cryptosporidium* species
301 have been identified in cattle: *C. parvum*, *C. bovis*, *C. ryanae*, and *C. andersoni*, but only *C.*

302 *parvum* is associated with clinical disease in neonatal calves, as older animals (> 6 weeks)
303 exhibit asymptomatic oocyst shedding (Silverlås et al., 2013). The recent observation of *C.*
304 *hominis* in symptomatic and asymptomatic calves in France (Razakandrainibe et al., 2018)
305 emphasises the importance of identifying which *Cryptosporidium* species is shed by calves.
306 The calves included in this study were all younger than 45 days, thus the finding that 100% of
307 animals were infected with *C. parvum* via *18S rRNA* gene based-PCR-RFLP was similar to
308 previous studies demonstrating high *C. parvum* occurrence in pre-weaned calves (Kvác et al.,
309 2006; Santín et al., 2004). Our data also confirms that neonatal cryptosporidial diarrhoea in
310 calves is primarily caused by *C. parvum* (de Graaf et al., 1999; Santín et al., 2004). In contrast
311 to a previous reports (Björkman et al., 2015), no mixed infections were detected in the present
312 study, which could be attributed to the significant *C. parvum* presence as opposed to other
313 *Cryptosporidium* species which may represent smaller subpopulations. Species-specific
314 multiplex PCR or real-time PCR could be used to detect any low-level infections (Tanriverdi
315 et al., 2003).

316 Current guidelines suggest that genetic characterization of *Cryptosporidium* isolates should be
317 based on two genetic loci and include at least one conserved *18S rRNA* gene, thus in our
318 study, the *18S rRNA* and *gp60* genes were targeted (Cacciò et al., 2005). Concordant results
319 were obtained for the majority of isolates, with only three exceptions where one or both PCRs
320 failed, possibly due to the above-cited reasons.

321 Sequence analysis of the *gp60* gene showed that four *C. parvum* IIa subgenotype groups exist;
322 IIaA15G2R1, IIaA17G3R1, IIaA17G1R1, and IIaA19G1R1. These results concur with other
323 studies demonstrating that *C. parvum* IIa is a common subtype family in humans as well as
324 calves. Thus, this subtype family is considered to be potentially zoonotic and transmissible
325 from livestock (Xiao, 2010).

326 The predominant IIaA15G2R1 subtype has previously been reported as the most prevalent in

327 calves and humans in many countries (Aita et al., 2015; Alves et al., 2006; Danišová et al.,
328 2016; Díaz et al., 2013; Mawly et al., 2015; Soba and Logar, 2008; Wielinga et al., 2008;
329 Xiao, 2010), including France (Follet et al., 2011; Rieux et al., 2014, 2013c, 2013a), thus
330 highlighting the zoonotic potential of calf reservoirs. It seems that the IIA15G2R1 *C. parvum*
331 subtype is hypertransmissible, which may explain its predominance (Feng et al., 2018). Future
332 studies are needed to determine whether this subtype demonstrates greater infectivity and to
333 what degree, or whether subtype predominance is due to a restricted available host.

334 Genotyping enabled the identification of relatively less dominant subtypes (IIA17G1R1 in
335 the Allier department, IIA17G3R1 in the Moselle, and IIA19G1R1 in the Ardèche). The
336 IIA17G1R1 subtype has previously been described in French beef cattle calves (Follet et al.,
337 2011), as well as in many other countries such as Argentina (Tomazic et al., 2013), Estonia
338 (Santoro et al., 2018), and the USA (Xiao et al., 2007), for example. Our study is the first to
339 report the presence of IIA17G3R1 and IIA19G1R1 subtypes in French calves.

340 Only two *C. parvum* subtypes belonged to the IId family group (IIdA22G1 and IIdA27G1). In
341 contrast to results from China (Cai et al., 2017; Feng and Xiao, 2017), the occurrence of this
342 zoonotic IId family group in calves is rare in Europe and seems likely to have spread from
343 Western Asia to other regions including France (Wang et al., 2014). This subtype family was
344 described in European countries, such as in calves from Italy (Díaz et al., 2018), Belgium
345 (Geurden et al., 2007), but never previously reported in France. Thus, the present study is the
346 first to report the presence of the IId subtype in French calves. It appears that this subtype
347 family is not restricted to a few farms, but could be easily transmitted to other animals
348 (predominantly young ruminants) or to humans (Wang et al., 2014; Xiao, 2010).

349 In our study, only two farms were infected with more than one *C. parvum* family subtype.
350 Mixed *C. parvum* subtype infections have previously been reported in studies with high

351 sampling numbers. In fact, it is possible that a range of *C. parvum* subtypes were circulating
352 in the same farm, but remained undetected when only one sampling was performed from each
353 animal (Mawly et al., 2015; Xiao et al., 2007). Similar studies on a larger geographic scale,
354 with greater numbers and horizontal sampling, are necessary to increase our understanding of
355 cryptosporidiosis transmission dynamics in calves.

356 **4. Conclusion**

357 In conclusion, our findings demonstrate that young calves are a potential reservoir for
358 different *C. parvum* subtypes. The high occurrence of zoonotic *C. parvum* subtype family
359 infections (IIa, IIc) in pre-weaned French calves confirms that calves should be considered as
360 a real source of infection and a potential zoonotic reservoir for human infections. Our results
361 also demonstrate that the *Cryptosporidium* population detected in France is more diverse than
362 previous studies would suggest. As a consequence, molecular studies in other regions,
363 including in calves and small ruminants, are needed to improve our understanding of
364 cryptosporidiosis epidemiology and *C. parvum* subtype diversity in France.

365 **Conflicts of interest**

366 The authors declare that they have no conflicting interests.

367 **Acknowledgements**

368 Mohamed Mammeri is the grateful recipient of a Cifre (Industrial Research Training
369 Agreement) grant. He would like to thank Phileo (Lesaffre Animal Care, France) and the
370 ANRT (National Association for Technical Research), Ministry of Research (France).

371 The authors are grateful to Aurélie Heckmann and Amandine Blaizot for their help in creating
372 the map, and to Dr Grégory Karadjian for help with submitting sequences to NCBI.

373 The authors are especially grateful for all the veterinarians who participated in the collection
374 of faecal samples: Dr Valérie Wolgust, Dr Catherine Bardot, Dr Caroline Pagneux, Dr
375 Edwige Bornot, Dr Radu Blaga, and Dr Typhaine Hebert.

376 **References**

- 377 Aita, J., Ichikawa-Seki, M., Kinami, A., Yaita, S., Kumagai, Y., Nishikawa, Y., Itagaki, T.,
378 2015. Molecular characterization of *Cryptosporidium parvum* detected in Japanese
379 black and Holstein calves in Iwate Prefecture and Tanegashima Island, Kagoshima
380 Prefecture, Japan. *J. Vet. Med. Sci.* 77, 997–999. [https://doi.org/10.1292/jvms.15-](https://doi.org/10.1292/jvms.15-0082)
381 0082
- 382 Alves, M., Xiao, L., Antunes, F., Matos, O., 2006. Distribution of *Cryptosporidium* subtypes
383 in humans and domestic and wild ruminants in Portugal. *Parasitol. Res.* 99, 287–292.
384 <https://doi.org/10.1007/s00436-006-0164-5>
- 385 Amar, C.F.L., Dear, P.H., McLauchlin, J., 2004. Detection and identification by real time
386 PCR/RFLP analyses of *Cryptosporidium* species from human faeces. *Lett. Appl.*
387 *Microbiol.* 38, 217–222.
- 388 Bamaiyi, P.H., Redhuan, N.E.M., 2016. Prevalence and risk factors for cryptosporidiosis: a
389 global, emerging, neglected zoonosis. *Asian Biomed.* 10, 309–325.
390 <https://doi.org/10.5372/1905-7415.1004.493>
- 391 Bhat, S.A., Dixit, M., Juyal, P.D., Singh, N.K., 2014. Comparison of nested PCR and
392 microscopy for the detection of cryptosporidiosis in bovine calves. *J. Parasit. Dis. Off.*
393 *Organ Indian Soc. Parasitol.* 38, 101–105. <https://doi.org/10.1007/s12639-012-0201-5>
- 394 Björkman, C., Lindström, L., Oweson, C., Ahola, H., Troell, K., Axén, C., 2015.
395 *Cryptosporidium* infections in suckler herd beef calves. *Parasitology* 142, 1108–1114.
396 <https://doi.org/10.1017/S0031182015000426>
- 397 Bouzid, M., Hunter, P.R., Chalmers, R.M., Tyler, K.M., 2013. *Cryptosporidium* pathogenicity
398 and virulence. *Clin. Microbiol. Rev.* 26, 115–134.
399 <https://doi.org/10.1128/CMR.00076-12>

400 Cacciò, S.M., Thompson, R.C.A., McLauchlin, J., Smith, H.V., 2005. Unravelling
401 *Cryptosporidium* and *Giardia* epidemiology. Trends Parasitol. 21, 430–437.
402 <https://doi.org/10.1016/j.pt.2005.06.013>

403 Cai, M., Guo, Y., Pan, B., Li, N., Wang, X., Tang, C., Feng, Y., Xiao, L., 2017. Longitudinal
404 monitoring of *Cryptosporidium* species in pre-weaned dairy calves on five farms in
405 Shanghai, China. Vet. Parasitol. 241, 14–19.
406 <https://doi.org/10.1016/j.vetpar.2017.05.005>

407 Cardona, G.A., de Lucio, A., Bailo, B., Cano, L., de Fuentes, I., Carmena, D., 2015.
408 Unexpected finding of feline-specific *Giardia duodenalis* assemblage F and
409 *Cryptosporidium felis* in asymptomatic adult cattle in Northern Spain. Vet. Parasitol.
410 209, 258–263. <https://doi.org/10.1016/j.vetpar.2015.02.028>

411 Castro-Hermida, J.A., Pors, I., Poupin, B., Ares-Mazás, E., Chartier, C., 2005. Prevalence of
412 *Giardia duodenalis* and *Cryptosporidium parvum* in goat kids in western France.
413 Small Rumin. Res. 56, 259–264. <https://doi.org/10.1016/j.smallrumres.2004.06.007>

414 Causapé, A.C., Quílez, J., Sánchez-Acedo, C., del Cacho, E., López-Bernad, F., 2002.
415 Prevalence and analysis of potential risk factors for *Cryptosporidium parvum* infection
416 in lambs in Zaragoza (northeastern Spain). Vet. Parasitol. 104, 287–298.

417 Chalmers, R.M., Katzer, F., 2013. Looking for *Cryptosporidium*: the application of advances
418 in detection and diagnosis. Trends Parasitol. 29, 237–251.
419 <https://doi.org/10.1016/j.pt.2013.03.001>

420 Danišová, O., Valenčáková, A., Petrinčová, A., 2016. Detection and identification of six
421 *Cryptosporidium* species in livestock in Slovakia by amplification of SSU and GP60
422 genes with the use of PCR analysis. Ann. Agric. Environ. Med. 23, 254–258.
423 <https://doi.org/10.5604/12321966.1203886>

424 de Graaf, D.C., Vanopdenbosch, E., Ortega-Mora, L.M., Abbassi, H., Peeters, J.E., 1999. A
425 review of the importance of cryptosporidiosis in farm animals. *Int. J. Parasitol.* 29,
426 1269–1287.

427 Delafosse, A., Chartier, C., Dupuy, M.C., Dumoulin, M., Pors, I., Paraud, C., 2015.
428 *Cryptosporidium parvum* infection and associated risk factors in dairy calves in
429 western France. *Prev. Vet. Med.* 118, 406–412.
430 <https://doi.org/10.1016/j.prevetmed.2015.01.005>

431 Dereeper, A., Audic, S., Claverie, J.-M., Blanc, G., 2010. BLAST-EXPLORER helps you
432 building datasets for phylogenetic analysis. *BMC Evol. Biol.* 10, 8.
433 <https://doi.org/10.1186/1471-2148-10-8>

434 Dereeper, A., Guignon, V., Blanc, G., Audic, S., Buffet, S., Chevenet, F., Dufayard, J.-F.,
435 Guindon, S., Lefort, V., Lescot, M., Claverie, J.-M., Gascuel, O., 2008. Phylogeny.fr:
436 robust phylogenetic analysis for the non-specialist. *Nucleic Acids Res.* 36, W465-469.
437 <https://doi.org/10.1093/nar/gkn180>

438 Díaz, P., Rota, S., Marchesi, B., López, C., Panadero, R., Fernández, G., Díez-Baños, P.,
439 Morrondo, P., Poglayen, G., 2013. *Cryptosporidium* in pet snakes from Italy:
440 Molecular characterization and zoonotic implications. *Vet. Parasitol.* 197, 68–73.
441 <https://doi.org/10.1016/j.vetpar.2013.04.028>

442 Díaz, P., Varcasia, A., Pipia, A.P., Tamponi, C., Sanna, G., Prieto, A., Ruiu, A., Spissu, P.,
443 Díez-Baños, P., Morrondo, P., Scala, A., 2018. Molecular characterisation and risk
444 factor analysis of *Cryptosporidium* spp. in calves from Italy. *Parasitol. Res.*
445 <https://doi.org/10.1007/s00436-018-6000-x>

446 Efstratiou, A., Ongerth, J., Karanis, P., 2017. Evolution of monitoring for *Giardia* and
447 *Cryptosporidium* in water. *Water Res.* 123, 96–112.
448 <https://doi.org/10.1016/j.watres.2017.06.042>

449 El-Khodery, S.A., Osman, S.A., 2008. Cryptosporidiosis in buffalo calves (*Bubalus bubalis*):
450 prevalence and potential risk factors. Trop. Anim. Health Prod. 40, 419–426.
451 <https://doi.org/10.1007/s11250-007-9113-2>

452 Elwin, K., Robinson, G., Hadfield, S.J., Fairclough, H.V., Iturriza-Gómara, M., Chalmers,
453 R.M., 2012. A comparison of two approaches to extracting *Cryptosporidium* DNA
454 from human stools as measured by a real-time PCR assay. J. Microbiol. Methods 89,
455 38–40. <https://doi.org/10.1016/j.mimet.2012.02.006>

456 Fayer, R., Morgan, U., Upton, S.J., 2000. Epidemiology of *Cryptosporidium*: transmission,
457 detection and identification. Int. J. Parasitol. 30, 1305–1322.

458 Fayer, R., Santín, M., Trout, J.M., Greiner, E., 2006. Prevalence of species and genotypes of
459 *Cryptosporidium* found in 1–2-year-old dairy cattle in the eastern United States. Vet.
460 Parasitol. 135, 105–112. <https://doi.org/10.1016/j.vetpar.2005.08.003>

461 Feng, Y., Ortega, Y., He, G., Das, P., Xu, M., Zhang, X., Fayer, R., Gatei, W., Cama, V.,
462 Xiao, L., 2007. Wide geographic distribution of *Cryptosporidium bovis* and the deer-
463 like genotype in bovines. Vet. Parasitol. 144, 1–9.
464 <https://doi.org/10.1016/j.vetpar.2006.10.001>

465 Feng, Y., Ryan, U.M., Xiao, L., 2018. Genetic diversity and population structure of
466 *Cryptosporidium*. Trends Parasitol. 0. <https://doi.org/10.1016/j.pt.2018.07.009>

467 Feng, Y., Xiao, L., 2017. Molecular epidemiology of cryptosporidiosis in China. Front.
468 Microbiol. 8. <https://doi.org/10.3389/fmicb.2017.01701>

469 Follet, J., Guyot, K., Leruste, H., Follet-Dumoulin, A., Hammouma-Ghelboun, O., Certad, G.,
470 Dei-Cas, E., Halama, P., 2011. *Cryptosporidium* infection in a veal calf cohort in
471 France: molecular characterization of species in a longitudinal study. Vet. Res. 42,
472 116. <https://doi.org/10.1186/1297-9716-42-116>

473 Gatei, W., Hart, C.A., Gilman, R.H., Das, P., Cama, V., Xiao, L., 2006. Development of a
474 multilocus sequence typing tool for *Cryptosporidium hominis*. J. Eukaryot. Microbiol.
475 53 Suppl 1, S43-48. <https://doi.org/10.1111/j.1550-7408.2006.00169.x>

476 Geurden, T., Berkvens, D., Martens, C., Casaert, S., Vercruyse, J., Claerebout, E., 2007.
477 Molecular epidemiology with subtype analysis of *Cryptosporidium* in calves in
478 Belgium. Parasitology 134, 1981–1987. <https://doi.org/10.1017/S0031182007003460>

479 Iqbal, A., Lim, Y.A.L., Surin, J., Sim, B.L.H., 2012. High diversity of *Cryptosporidium*
480 subgenotypes identified in Malaysian HIV/AIDS individuals targeting gp60 gene.
481 PloS One 7, e31139. <https://doi.org/10.1371/journal.pone.0031139>

482 Jiang, J., Alderisio, K.A., Singh, A., Xiao, L., 2005. Development of procedures for direct
483 extraction of *Cryptosporidium* DNA from water concentrates and for relief of PCR
484 inhibitors. Appl. Environ. Microbiol. 71, 1135–1141.
485 <https://doi.org/10.1128/AEM.71.3.1135-1141.2005>

486 Johnson, D.W., Pieniazek, N.J., Griffin, D.W., Misener, L., Rose, J.B., 1995. Development of
487 a PCR protocol for sensitive detection of *Cryptosporidium* oocysts in water samples.
488 Appl. Environ. Microbiol. 61, 3849–3855.

489 Koudela, B., Jirí, V., 1997. Experimental cryptosporidiosis in kids. Vet. Parasitol. 71, 273–
490 281.

491 Kvác, M., Kouba, M., Vítovec, J., 2006. Age-related and housing-dependence of
492 *Cryptosporidium* infection of calves from dairy and beef herds in South Bohemia,
493 Czech Republic. Vet. Parasitol. 137, 202–209.
494 <https://doi.org/10.1016/j.vetpar.2006.01.027>

495 Lantz, P.-G., Matsson, M., Wadström, T., Rådström, P., 1997. Removal of PCR inhibitors
496 from human faecal samples through the use of an aqueous two-phase system for

497 sample preparation prior to PCR. J. Microbiol. Methods 28, 159–167.
498 [https://doi.org/10.1016/S0167-7012\(97\)00979-2](https://doi.org/10.1016/S0167-7012(97)00979-2)

499 Lefay, D., Naciri, M., Poirier, P., Chermette, R., 2000. Prevalence of *Cryptosporidium*
500 infection in calves in France. Vet. Parasitol. 89, 1–9.

501 Mammeri, M., Chevillot, A., Thomas, M., Polack, B., Julien, C., Marden, J.-P., Auclair, E.,
502 Vallée, I., Adjou, K.T., 2018. Efficacy of chitosan, a natural polysaccharide, against
503 *Cryptosporidium parvum* *in vitro* and *in vivo* in neonatal mice. Exp. Parasitol. 194, 1–
504 8. <https://doi.org/10.1016/j.exppara.2018.09.003>

505 Maurya, P.S., Rakesh, R.L., Pradeep, B., Kumar, S., Kundu, K., Garg, R., Ram, H., Kumar,
506 A., Banerjee, P.S., 2013. Prevalence and risk factors associated with *Cryptosporidium*
507 spp. infection in young domestic livestock in India. Trop. Anim. Health Prod. 45,
508 941–946. <https://doi.org/10.1007/s11250-012-0311-1>

509 Mawly, J.A., Grinberg, A., Velathanthiri, N., French, N., 2015. Cross sectional study of
510 prevalence, genetic diversity and zoonotic potential of *Cryptosporidium parvum*
511 cycling in New Zealand dairy farms. Parasit. Vectors 8, 240.
512 <https://doi.org/10.1186/s13071-015-0855-9>

513 McLauchlin, J., Pedraza-Díaz, S., Amar-Hoetzeneder, C., Nichols, G.L., 1999. Genetic
514 characterization of *Cryptosporidium* strains from 218 patients with diarrhea diagnosed
515 as having sporadic cryptosporidiosis. J. Clin. Microbiol. 37, 3153–3158.

516 Mohammed, H.O., Wade, S.E., Schaaf, S., 1999. Risk factors associated with
517 *Cryptosporidium parvum* infection in dairy cattle in southeastern New York State.
518 Vet. Parasitol. 83, 1–13.

519 Naciri, M., Paul Lefay, M., Mancassola, R., Poirier, P., Chermette, R., 1999. Role of
520 *Cryptosporidium parvum* as a pathogen in neonatal diarrhoea complex in suckling and

521 dairy calves in France. *Vet. Parasitol.* 85, 245–257. <https://doi.org/10.1016/S0304->
522 4017(99)00111-9

523 Ngouanesavanh, T., Guyot, K., Certad, G., Le Fichoux, Y., Chartier, C., Verdier, R.-I.,
524 Cailliez, J.-C., Camus, D., Dei-Cas, E., Bañuls, A.-L., 2006. *Cryptosporidium*
525 population genetics: evidence of clonality in isolates from France and Haiti. *J.*
526 *Eukaryot. Microbiol.* 53 Suppl 1, S33-36. <https://doi.org/10.1111/j.1550->
527 7408.2006.00166.x

528 Nguyen, S.T., Nguyen, D.T., Le, D.Q., Le Hua, L.N., Van Nguyen, T., Honma, H., Nakai, Y.,
529 2007. Prevalence and first genetic identification of *Cryptosporidium* spp. in cattle in
530 central Viet Nam. *Vet. Parasitol.* 150, 357–361.
531 <https://doi.org/10.1016/j.vetpar.2007.09.010>

532 Olson, M.E., O’Handley, R.M., Ralston, B.J., McAllister, T.A., Thompson, R.C.A., 2004.
533 Update on *Cryptosporidium* and *Giardia* infections in cattle. *Trends Parasitol.* 20,
534 185–191. <https://doi.org/10.1016/j.pt.2004.01.015>

535 Osman, M., El Safadi, D., Benamrouz-Vanneste, S., Cian, A., Moriniere, R., Gantois, N.,
536 Delgado-Viscogliosi, P., Guyot, K., Bosc, S., Chabé, M., Petit, T., Viscogliosi, E.,
537 Certad, G., 2017. Prevalence, transmission, and host specificity of *Cryptosporidium*
538 spp. in various animal groups from two French zoos. *Parasitol. Res.* 116, 3419–3422.
539 <https://doi.org/10.1007/s00436-017-5645-1>

540 Ranjbar, R., Fattahi, R., 2017. Prevalence of *Cryptosporidium* spp. in calves under one year
541 old in Ilam county (Iran), from March 2014 to February 2015. *Electron. Physician* 9,
542 4631–4635. <https://doi.org/10.19082/4631>

543 Razakandrainibe, R., Diawara, E.H.I., Costa, D., Goff, L.L., Lemeteil, D., Ballet, J.J.,
544 Gargala, G., Favennec, L., 2018. Common occurrence of *Cryptosporidium hominis* in

545 asymptomatic and symptomatic calves in France. PLoS Negl. Trop. Dis. 12,
546 e0006355. <https://doi.org/10.1371/journal.pntd.0006355>

547 Rieux, A., Chartier, C., Pors, I., Delafosse, A., Paraud, C., 2013a. Molecular characterization
548 of *Cryptosporidium* isolates from high-excreting young dairy calves in dairy cattle
549 herds in Western France. Parasitol. Res. 112, 3423–3431.
550 <https://doi.org/10.1007/s00436-013-3520-2>

551 Rieux, A., Chartier, C., Pors, I., Paraud, C., 2013b. Dynamics of excretion and molecular
552 characterization of *Cryptosporidium* isolates in pre-weaned French beef calves. Vet.
553 Parasitol. 195, 169–172. <https://doi.org/10.1016/j.vetpar.2012.12.043>

554 Rieux, A., Paraud, C., Pors, I., Chartier, C., 2014. Molecular characterization of
555 *Cryptosporidium* isolates from beef calves under one month of age over three
556 successive years in one herd in western France. Vet. Parasitol. 202, 171–179.
557 <https://doi.org/10.1016/j.vetpar.2014.03.004>

558 Rieux, A., Paraud, C., Pors, I., Chartier, C., 2013c. Molecular characterization of
559 *Cryptosporidium* isolates from pre-weaned calves in western France in relation to age.
560 Vet. Parasitol. 197, 7–12. <https://doi.org/10.1016/j.vetpar.2013.05.001>

561 Ryan, U., Hijjawi, N., Xiao, L., 2018. Foodborne cryptosporidiosis. Int. J. Parasitol. 48, 1–12.
562 <https://doi.org/10.1016/j.ijpara.2017.09.004>

563 Santín, M., Trout, J.M., Fayer, R., 2008. A longitudinal study of cryptosporidiosis in dairy
564 cattle from birth to 2 years of age. Vet. Parasitol. 155, 15–23.
565 <https://doi.org/10.1016/j.vetpar.2008.04.018>

566 Santín, M., Trout, J.M., Xiao, L., Zhou, L., Greiner, E., Fayer, R., 2004. Prevalence and age-
567 related variation of *Cryptosporidium* species and genotypes in dairy calves. Vet.
568 Parasitol. 122, 103–117. <https://doi.org/10.1016/j.vetpar.2004.03.020>

569 Santoro, A., Dorbek-Kolin, E., Jeremejeva, J., Tummeleht, L., Orro, T., Jokelainen, P.,
570 Lassen, B., 2018. Molecular epidemiology of *Cryptosporidium* spp. in calves in
571 Estonia: high prevalence of *Cryptosporidium parvum* shedding and 10 subtypes
572 identified. *Parasitology* 1–7. <https://doi.org/10.1017/S0031182018001348>

573 Silverlås, C., Bosaeus-Reineck, H., Näslund, K., Björkman, C., 2013. Is there a need for
574 improved *Cryptosporidium* diagnostics in Swedish calves? *Int. J. Parasitol.* 43, 155–
575 161. <https://doi.org/10.1016/j.ijpara.2012.10.009>

576 Soba, B., Logar, J., 2008. Genetic classification of *Cryptosporidium* isolates from humans and
577 calves in Slovenia. *Parasitology* 135, 1263–1270.
578 <https://doi.org/10.1017/S0031182008004800>

579 Stenger, B.L.S., Clark, M.E., Kváč, M., Khan, E., Giddings, C.W., Dyer, N.W., Schultz, J.L.,
580 McEvoy, J.M., 2015. Highly divergent 18S rRNA gene paralogs in a *Cryptosporidium*
581 genotype from eastern chipmunks (*Tamias striatus*). *Infect. Genet. Evol. J. Mol.*
582 *Epidemiol. Evol. Genet. Infect. Dis.* 32, 113–123.
583 <https://doi.org/10.1016/j.meegid.2015.03.003>

584 Stensvold, C.R., Elwin, K., Winięcka-Krusnell, J., Chalmers, R.M., Xiao, L., Lebbad, M.,
585 2015. Development and application of a gp60-based typing assay for *Cryptosporidium*
586 *viatorum*. *J. Clin. Microbiol.* 53, 1891–1897. <https://doi.org/10.1128/JCM.00313-15>

587 Sulaiman, I.M., Hira, P.R., Zhou, L., Al-Ali, F.M., Al-Shelahi, F.A., Shweiki, H.M., Iqbal, J.,
588 Khalid, N., Xiao, L., 2005. Unique endemicity of cryptosporidiosis in children in
589 Kuwait. *J. Clin. Microbiol.* 43, 2805–2809. [https://doi.org/10.1128/JCM.43.6.2805-](https://doi.org/10.1128/JCM.43.6.2805-2809.2005)
590 [2809.2005](https://doi.org/10.1128/JCM.43.6.2805-2809.2005)

591 Tanriverdi, S., Arslan, M.O., Akiyoshi, D.E., Tzipori, S., Widmer, G., 2003. Identification of
592 genotypically mixed *Cryptosporidium parvum* populations in humans and calves. *Mol.*
593 *Biochem. Parasitol.* 130, 13–22.

594 Thompson, R.C.A., Ash, A., 2016. Molecular epidemiology of *Giardia* and *Cryptosporidium*
595 infections. *Infect. Genet. Evol.* 40, 315–323.
596 <https://doi.org/10.1016/j.meegid.2015.09.028>

597 Thomson, S., Hamilton, C.A., Hope, J.C., Katzer, F., Mabbott, N.A., Morrison, L.J., Innes,
598 E.A., 2017. Bovine cryptosporidiosis: impact, host-parasite interaction and control
599 strategies. *Vet. Res.* 48. <https://doi.org/10.1186/s13567-017-0447-0>

600 Tomazic, M.L., Maidana, J., Dominguez, M., Uriarte, E.L., Galarza, R., Garro, C., Florin-
601 Christensen, M., Schnittger, L., 2013. Molecular characterization of *Cryptosporidium*
602 isolates from calves in Argentina. *Vet. Parasitol.* 198, 382–386.
603 <https://doi.org/10.1016/j.vetpar.2013.09.022>

604 Trotz-Williams, L.A., Wayne Martin, S., Leslie, K.E., Duffield, T., Nydam, D.V., Peregrine,
605 A.S., 2007. Calf-level risk factors for neonatal diarrhea and shedding of
606 *Cryptosporidium parvum* in Ontario dairy calves. *Prev. Vet. Med.* 82, 12–28.
607 <https://doi.org/10.1016/j.prevetmed.2007.05.003>

608 Wang, R., Zhang, L., Axén, C., Bjorkman, C., Jian, F., Amer, S., Liu, A., Feng, Y., Li, G., Lv,
609 C., Zhao, Z., Qi, M., Dong, H., Wang, H., Sun, Y., Ning, C., Xiao, L., 2014.
610 *Cryptosporidium parvum* IId family: clonal population and dispersal from Western
611 Asia to other geographical regions. *Sci. Rep.* 4, 4208.
612 <https://doi.org/10.1038/srep04208>

613 Widmer, G., Sullivan, S., 2012. Genomics and population biology of *Cryptosporidium*
614 species. *Parasite Immunol.* 34, 61–71. <https://doi.org/10.1111/j.1365-3024.2011.01301.x>

615
616 Wielinga, P.R., de Vries, A., van der Goot, T.H., Mank, T., Mars, M.H., Kortbeek, L.M., van
617 der Giessen, J.W.B., 2008. Molecular epidemiology of *Cryptosporidium* in humans

618 and cattle in The Netherlands. *Int. J. Parasitol.* 38, 809–817.
619 <https://doi.org/10.1016/j.ijpara.2007.10.014>

620 Xiao, L., 2010. Molecular epidemiology of cryptosporidiosis: an update. *Exp. Parasitol.* 124,
621 80–89. <https://doi.org/10.1016/j.exppara.2009.03.018>

622 Xiao, L., Morgan, U.M., Limor, J., Escalante, A., Arrowood, M., Shulaw, W., Thompson,
623 R.C.A., Fayer, R., Lal, A.A., 1999. Genetic diversity within *Cryptosporidium parvum*
624 and related *Cryptosporidium* species. *Appl. Environ. Microbiol.* 65, 3386–3391.

625 Xiao, L., Zhou, L., Santin, M., Yang, W., Fayer, R., 2007. Distribution of *Cryptosporidium*
626 *parvum* subtypes in calves in eastern United States. *Parasitol. Res.* 100, 701–706.
627 <https://doi.org/10.1007/s00436-006-0337-2>


628 Yu, J.-R., Lee, S.-U., Park, W.-Y., 2009. Comparative sensitivity of PCR primer sets for
629 detection of *Cryptosporidium parvum*. *Korean J. Parasitol.* 47, 293–297.
630 <https://doi.org/10.3347/kjp.2009.47.3.293>

631

632

633 **Figures**

634 **Figure 1. Geographical map of pre-weaned calf faecal sampling locations in French**
635 **departments.** The number of samples collected from each department (administrative
636 department number-department name) was: 03-Allier: n=18, 07-Ardèche: n= 2, 21-Côte-d'Or:
637 n=6, 57-Moselle: n=1, 71-Saône-et-Loire: n=1, and 89-Yonne: n=7.



638

Figure 2. Pre-weaned calf oocyst excretion. Samples are classified from the youngest calves (4 days-old) to the oldest calf (45 days-old). DFA: Immunofluorescence Assay. OC: oocyst concentration. OPG: oocysts per gram of faeces.




Table 1. Molecular characterization of *Cryptosporidium* from clinically-affected pre-weaned French calves, including age and faecal consistency data. n=total number of samples; 18S=18S ribosomal RNA; gp60=60kDa glycoprotein; NA=No Amplification; NU=Non-Usable; ND=not done. Department number: 03: Allier, 07: Ardèche, 21: Côte-d'Or, 57: Moselle, 71: Saône-et-Loire, 89: Yonne.

Samples (n=35)	Department	Herd	Age (days)	Faecal score	Sequencing (18S rRNA)	Access number (18S rRNA)	PCR (18S rRNA) + RFLP	Sequencing (gp60)	Access number (gp60)
B1	21	E1	7	3	<i>C. parvum</i>	MK014763	<i>C. parvum</i>	IlaA15G2R1	MK109829
B2	21		7	4	<i>C. parvum</i>	MK014764	<i>C. parvum</i>	IlaA15G2R1	MK109830
B3	21	E2	8	4	<i>C. parvum</i>	MK014765	<i>C. parvum</i>	IlaA15G2R1	MK109831
B4	21		8	2	<i>C. parvum</i>	MK014766	<i>C. parvum</i>	IlaA15G2R1	MK109832
B5	21		8	1	<i>C. parvum</i>	MK014767	<i>C. parvum</i>	IlaA15G2R1	MK109833
B6	21		8	2	<i>C. parvum</i>	MK014768	<i>C. parvum</i>	IlaA15G2R1	MK109834
B7	89	E3	30	4	ND	/	ND	ND	/
B8	89		30	3	ND	/	ND	ND	/
B9	89		30	1	ND	/	ND	ND	/
B10	89	E4	11	4	<i>C. parvum</i>	MK014769	<i>C. parvum</i>	IIdA22G1	MK109835
B11	89		20	2	NU	/	<i>C. parvum</i>	NU	/
B12	07	E5	15	4	<i>C. parvum</i>	MK014770	<i>C. parvum</i>	IlaA15G2R1	MK109836
B13	03	E6	7	4	<i>C. parvum</i>	MK014771	<i>C. parvum</i>	IlaA15G2R1	MK109837
B14	07	E7	7	4	<i>C. parvum</i>	MK014772	<i>C. parvum</i>	IlaA19G1R1	MK109838
B15	03	E8	8	4	<i>C. parvum</i>	MK014773	<i>C. parvum</i>	IlaA15G2R1	MK109839
B16	03	E9	13	NR	NU	/	NA	NU	/
B17	03	E10	6	4	<i>C. parvum</i>	MK014774	<i>C. parvum</i>	IlaA15G2R1	MK109840
B18	71	E11	4	4	<i>C. parvum</i>	MK014775	<i>C. parvum</i>	IIdA22G1	MK109841
B19	57	E12	5	3	<i>C. parvum</i>	MK014776	<i>C. parvum</i>	IlaA17G3R1	MK109842
B20	03	E13	10	2	<i>C. parvum</i>	MK014777	<i>C. parvum</i>	NU	/
B21	03	E14	4	4	<i>C. parvum</i>	MK014778	<i>C. parvum</i>	IIdA27G1	MK109843
B22	89	E15	15	4	<i>C. parvum</i>	MK014779	<i>C. parvum</i>	IlaA15G2R1	MK109844
B23	89	E16	15	4	ND	/	ND	ND	/
B24	03	E17	45	4	<i>C. parvum</i>	MK014780	<i>C. parvum</i>	IlaA15G2R1	MK109845
B25	03		45	4	<i>C. parvum</i>	MK014781	<i>C. parvum</i>	IlaA15G2R1	MK109846
B26	03	E18	10	NR	<i>C. parvum</i>	MK014782	<i>C. parvum</i>	IlaA15G2R1	MK109847
B27	03	E19	7	4	<i>C. parvum</i>	MK014783	<i>C. parvum</i>	IlaA15G2R1	MK109848
B28	03	E20	5	4	<i>C. parvum</i>	MK014784	<i>C. parvum</i>	IlaA15G2R1	MK109849
B29	03	E21	6	4	<i>C. parvum</i>	MK014785	<i>C. parvum</i>	IlaA15G2R1	MK109850
B30	03	E22	7	4	<i>C. parvum</i>	MK014786	NA	NU	/
B31	03	E23	7	4	<i>C. parvum</i>	MK014787	<i>C. parvum</i>	IlaA17G1R1	MK109851
B32	03		7	4	<i>C. parvum</i>	MK014788	<i>C. parvum</i>	IlaA15G2R1	MK109852
B33	03	E24	6	4	<i>C. parvum</i>	MK014789	<i>C. parvum</i>	IlaA15G2R1	MK109853

B34	03	E25	8	4	<i>C. parvum</i>	MK014790	<i>C. parvum</i>	IlaA15G2R1	MK109854
B35	03	E26	7	4	<i>C. parvum</i>	MK014791	<i>C. parvum</i>	IlaA15G2R1	MK109855